

ClinicalTrials.gov Protocol Registration and Results System (PRS) Receipt  
Release Date: 07/15/2013

ClinicalTrials.gov ID: NCT00168298

---

## Study Identification

Unique Protocol ID: 206207-009

Brief Title: A Study of the Safety and Efficacy of a New Treatment for Macular Edema Resulting From Retinal Vein Occlusion

Official Title:

Secondary IDs:

## Study Status

Record Verification: July 2013

Overall Status: Completed

Study Start: November 2004

Primary Completion: March 2008 [Actual]

Study Completion: September 2008 [Actual]

## Sponsor/Collaborators

Sponsor: Allergan

Responsible Party: Sponsor

Collaborators:

## Oversight

FDA Regulated?: Yes

Applicable Trial?: Section 801 Clinical Trial? Yes  
Delayed Posting? No

IND/IDE Protocol?: Yes

IND/IDE Information: Grantor: CDER  
IND/IDE Number: 58,663  
Serial Number:  
Has Expanded Access? No

Review Board: Approval Status:  
Board Name:  
Board Affiliation:  
Phone:  
Email:

Data Monitoring?: No

Plan to Share Data?:

Oversight Authorities: United States: Food and Drug Administration

## Study Description

Brief Summary: This study will evaluate the safety and efficacy of an intravitreal implant of dexamethasone for the treatment of macular edema associated with retinal vein occlusion.

Detailed Description:

## Conditions

Conditions: Macular Edema  
Retinal Vein Occlusion

Keywords:

## Study Design

Study Type: Interventional

Primary Purpose: Treatment

Study Phase: Phase 3

Intervention Model: Parallel Assignment

Number of Arms: 3

Masking: Double Blind (Subject, Caregiver, Investigator, Outcomes Assessor)

Allocation: Randomized

Endpoint Classification: Safety/Efficacy Study

Enrollment: 668 [Actual]

## Arms and Interventions

Arms	Assigned Interventions
Experimental: 700 µg Dexamethasone 700 µg dexamethasone intravitreal implant administered on Day 0 and Day 180.	Drug: 700 µg Dexamethasone 700 µg dexamethasone intravitreal implant administered on Day 0 and/or Day 180.  Other Names: <ul style="list-style-type: none"><li>• Posurdex®</li></ul>
Experimental: 350 µg Dexamethasone followed by 700 µg Dexamethasone 350 µg dexamethasone intravitreal implant administered on Day 0 and 700 µg dexamethasone intravitreal implant on Day 180.	Drug: 700 µg Dexamethasone 700 µg dexamethasone intravitreal implant administered on Day 0 and/or Day 180.  Other Names: <ul style="list-style-type: none"><li>• Posurdex®</li></ul> Drug: 350 µg Dexamethasone 350 µg Dexamethasone intravitreal implant administered on Day 0.  Other Names: <ul style="list-style-type: none"><li>• Posurdex®</li></ul>
Sham Comparator: Sham Injection followed by 700 µg Dexamethasone Sham injection on Day 0 and 700 µg dexamethasone intravitreal implant on Day 180.	Drug: 700 µg Dexamethasone 700 µg dexamethasone intravitreal implant administered on Day 0 and/or Day 180.  Other Names: <ul style="list-style-type: none"><li>• Posurdex®</li></ul> Sham Injection Sham injection on Day 0.

## Outcome Measures

[See Results Section.]

## Eligibility

Minimum Age: 18 Years

Maximum Age:

Gender: Both

Accepts Healthy Volunteers?: No

Criteria: Key Inclusion Criteria:

- 18 years of age or older with macular edema resulting from retinal vein occlusion
- Decrease in visual acuity in at least one eye as a result of macular edema (20/50 or worse)
- Visual acuity in other eye no worse than 20/200

Key Exclusion Criteria:

- Known anticipated need for ocular surgery within next 12 months
- History of glaucoma or current high eye pressure requiring more than 1 medication
- Diabetic retinopathy
- Uncontrolled systemic disease
- Known steroid-responder
- Use of systemic steroids
- Use of warfarin/heparin

## Contacts/Locations

Study Officials: Medical Director  
Study Director  
Allergan, Inc.

Locations: United States, Texas  
Houston, Texas, United States

Canada, Ontario  
Mississauga, Ontario, Canada

United Kingdom  
London, United Kingdom

Brazil  
Sao Paulo, Brazil

Italy  
Udine, Italy

India  
Tamil Nadu, India

Korea, Republic of  
Seoul, Korea, Republic of

New Zealand

Auckland, New Zealand

Spain

Alicante, Spain

Hong Kong

Kowloon, Hong Kong

Poland

Poznan, Poland

Singapore

Singapore, Singapore

Colombia

Bogota, Colombia

## References

Citations:

Links:

Study Data/Documents:

## Study Results



### Participant Flow

Pre-Assignment Details	Patients were randomly assigned during the double-blind period of the study to treatment with 700 µg dexamethasone, 350 µg dexamethasone, or sham injection on Day 0. Patients who qualified to continue in the open-label period of the study received 700 µg dexamethasone on Day 180.
------------------------	--

#### Reporting Groups

	Description
700 µg Dexamethasone	700 µg dexamethasone intravitreal implant administered on Day 0 and Day 180.
350 µg Dexamethasone Followed by 700 µg Dexamethasone	350 µg dexamethasone intravitreal implant administered on Day 0 and 700 µg dexamethasone intravitreal implant on Day 180.

	Description
Sham Injection Followed by 700 µg Dexamethasone	Sham injection on Day 0 and 700 µg dexamethasone intravitreal implant on Day 180.

#### Double-Blind Period

	700 µg Dexamethasone	350 µg Dexamethasone Followed by 700 µg Dexamethasone	Sham Injection Followed by 700 µg Dexamethasone
Started	226	218	224
Completed	214	206	209
Not Completed	12	12	15

#### Open-Label Period

	700 µg Dexamethasone	350 µg Dexamethasone Followed by 700 µg Dexamethasone	Sham Injection Followed by 700 µg Dexamethasone
Started	179 <sup>[1]</sup>	173 <sup>[1]</sup>	168 <sup>[1]</sup>
Completed	172	168	160
Not Completed	7	5	8

[1] Not all patients who completed the double-blind period entered the open-label period.

## Baseline Characteristics

#### Reporting Groups

	Description
700 µg Dexamethasone	700 µg dexamethasone intravitreal implant administered on Day 0 and Day 180.
350 µg Dexamethasone Followed by 700 µg Dexamethasone	350 µg dexamethasone intravitreal implant administered on Day 0 and 700 µg dexamethasone intravitreal implant on Day 180.
Sham Injection Followed by 700 µg Dexamethasone	Sham injection on Day 0 and 700 µg dexamethasone intravitreal implant on Day 180.

## Baseline Measures

	700 µg Dexamethasone	350 µg Dexamethasone Followed by 700 µg Dexamethasone	Sham Injection Followed by 700 µg Dexamethasone	Total
Number of Participants	226	218	224	668
Age, Customized [units: participants]				
<45 years	14	15	13	42
45-65 years	109	108	111	328
>65 years	103	95	100	298
Gender, Male/Female [units: participants]				
Female	115	102	101	318
Male	111	116	123	350



## Outcome Measures

### 1. Primary Outcome Measure:

Measure Title	Number of Patients With 15 or More Letter Improvement in Best Corrected Visual Acuity (BCVA) in the Study Eye
Measure Description	BCVA is measured using an eye chart and is reported as the number of letters read correctly (ranging from 0 to 100 letters). The lower the number of letters read correctly on the eye chart, the worse the vision (or visual acuity). An increase in the number of letters read correctly means that vision has improved. The numbers of patients with at least a 15 or more letter improvement in BCVA in the study eye are presented.
Time Frame	Day 180
Safety Issue?	No

### Analysis Population Description

Intent-to-Treat: all randomized patients

### Reporting Groups

	Description
700 µg Dexamethasone	700 µg Dexamethasone intravitreal implant administered on Day 0.
350 µg Dexamethasone	350 µg Dexamethasone intravitreal implant administered on Day 0.
Sham Injection	Sham injection on Day 0.

## Measured Values

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
Number of Participants Analyzed	226	218	224
Number of Patients With 15 or More Letter Improvement in Best Corrected Visual Acuity (BCVA) in the Study Eye [units: Number of Participants]	53	48	38

## 2. Secondary Outcome Measure:

Measure Title	Change From Baseline in Retinal Thickness in the Study Eye
Measure Description	Retinal thickness is assessed by optical coherence tomography (OCT) in the study eye. The retina is the light-sensitive part of the eye. OCT is a laser-based, noninvasive, diagnostic system providing high-resolution, three-dimensional images of the retina. A negative change from baseline indicates an improvement.
Time Frame	Baseline, Day 90, Day 180
Safety Issue?	No

## Analysis Population Description

Intent-to-Treat: all randomized patients

## Reporting Groups

	Description
700 µg Dexamethasone	700 µg Dexamethasone intravitreal implant administered on Day 0.
350 µg Dexamethasone	350 µg Dexamethasone intravitreal implant administered on Day 0.
Sham Injection	Sham injection on Day 0.

## Measured Values

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
Number of Participants Analyzed	226	218	224
Change From Baseline in Retinal Thickness in the Study Eye [units: Microns (µm)] Mean (Standard Deviation)			
Baseline	573.6 (189.08)	566.6 (219.63)	542.5 (185.64)
Change from Baseline at Day 90	-215.6 (207.62)	-205.5 (216.25)	-91.1 (191.85)



	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
Change from Baseline at Day 180	-132.1 (207.20)	-150.5 (220.78)	-127.4 (197.77)

### 3. Secondary Outcome Measure:

Measure Title	Percentage of Patients With a Change From Baseline in BCVA by Category
Measure Description	BCVA is measured using an eye chart and is reported as the number of letters read correctly (ranging from 0 to 100 letters). The lower the number of letters read correctly on the eye chart, the worse the vision (or visual acuity). An increase in the number of letters read correctly means that vision has improved. Data are grouped into the following 5 categories based on change from baseline: ≥15 Letters Improvement, ≥5 and <15 Letters Improvement, No Change (Between -5 to +5 Letters), ≥5 and <15 Letters Worsening, and ≥15 Letters Worsening.
Time Frame	Baseline, Day 90
Safety Issue?	No

### Analysis Population Description

Intent-to-Treat: all randomized patients

### Reporting Groups

	Description
700 µg Dexamethasone	700 µg Dexamethasone intravitreal implant administered on Day 0.
350 µg Dexamethasone	350 µg Dexamethasone intravitreal implant administered on Day 0.
Sham Injection	Sham injection on Day 0.

### Measured Values

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
Number of Participants Analyzed	226	218	224
Percentage of Patients With a Change From Baseline in BCVA by Category [units: Percentage of Patients]			
≥15 Letters Improvement	21.2	25.7	13.8
≥5 and <15 Letters Improvement	45.1	39.0	37.1
No Change (Between -5 to +5 Letters)	25.7	26.6	29.9
≥5 and <15 Letters Worsening	4.4	6.4	11.2

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
≥15 Letters Worsening	3.5	2.3	8.0

#### 4. Secondary Outcome Measure:

Measure Title	Percentage of Patients With a Change From Baseline in BCVA by Category
Measure Description	BCVA is measured using an eye chart and is reported as the number of letters read correctly (ranging from 0 to 100 letters). The lower the number of letters read correctly on the eye chart, the worse the vision (or visual acuity). An increase in the number of letters read correctly means that vision has improved. Data are grouped into the following 5 categories based on change from baseline: ≥15 Letters Improvement, ≥5 and <15 Letters Improvement, No Change (Between -5 to +5 Letters), ≥5 and <15 Letters Worsening, and ≥15 Letters Worsening.
Time Frame	Baseline, Day 180
Safety Issue?	No

#### Analysis Population Description

Intent-to-Treat: all randomized patients

#### Reporting Groups

	Description
700 µg Dexamethasone	700 µg Dexamethasone intravitreal implant administered on Day 0.
350 µg Dexamethasone	350 µg Dexamethasone intravitreal implant administered on Day 0.
Sham Injection	Sham injection on Day 0.

#### Measured Values

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
Number of Participants Analyzed	226	218	224
Percentage of Patients With a Change From Baseline in BCVA by Category [units: Percentage of Patients]			
≥15 Letters Improvement	23.5	21.6	17.0
≥5 and <15 Letters Improvement	35.0	33.9	28.6
No Change (Between -5 to +5 Letters)	27.0	28.9	28.6
≥5 and <15 Letters Worsening	8.0	10.6	13.8

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
≥15 Letters Worsening	6.6	5.0	12.1

## Reported Adverse Events

Time Frame	Adverse Events (AEs) and Serious Adverse Events (SAEs) are reported for the double-blind treatment period through Month 6.
Additional Description	The safety population included all randomized patients who received at least one dose of study medication and was used to assess AEs and SAEs.

### Reporting Groups

	Description
700 µg Dexamethasone	700 µg Dexamethasone intravitreal implant administered on Day 0.
350 µg Dexamethasone	350 µg Dexamethasone intravitreal implant administered on Day 0.
Sham Injection	Sham injection on Day 0.

### Serious Adverse Events

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
	Affected/At Risk (%)	Affected/At Risk (%)	Affected/At Risk (%)
Total	16/225 (7.11%)	7/215 (3.26%)	17/221 (7.69%)
Cardiac disorders			
Angina pectoris <sup>A *</sup>	0/225 (0%)	1/215 (0.47%)	0/221 (0%)
Angina unstable <sup>A *</sup>	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Cardiac arrest <sup>A †</sup>	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Cardiac failure congestive <sup>A †</sup>	1/225 (0.44%)	0/215 (0%)	1/221 (0.45%)
Myocardial infarction <sup>A †</sup>	1/225 (0.44%)	1/215 (0.47%)	0/221 (0%)
Congenital, familial and genetic disorders			
Atrial septal defect <sup>A †</sup>	0/225 (0%)	1/215 (0.47%)	0/221 (0%)

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
	Affected/At Risk (%)	Affected/At Risk (%)	Affected/At Risk (%)
Eye disorders			
Ocular hypertension <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Gastrointestinal disorders			
Upper gastrointestinal haemorrhage <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
General disorders			
Chest pain <sup>A</sup> *	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Non-cardiac chest pain <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Hepatobiliary disorders			
Cholecystitis <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Cholelithiasis <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Infections and infestations			
Appendicitis <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Cellulitis <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Pneumonia <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Urinary tract infection <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Injury, poisoning and procedural complications			
Foot fracture <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Investigations			
Intraocular pressure increased <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Musculoskeletal and connective tissue disorders			
Aneurysmal bone cyst <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Back pain <sup>A</sup> *	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Intervertebral disc degeneration <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
	Affected/At Risk (%)	Affected/At Risk (%)	Affected/At Risk (%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Breast cancer <sup>A</sup> †	1/225 (0.44%)	1/215 (0.47%)	0/221 (0%)
Chronic lymphocytic leukaemia <sup>A</sup> †	0/225 (0%)	1/215 (0.47%)	0/221 (0%)
Prostate cancer <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Nervous system disorders			
Carotid artery occlusion <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Cerebellar infarction <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Cerebrovascular accident <sup>A</sup> †	0/225 (0%)	1/215 (0.47%)	1/221 (0.45%)
Parkinson's disease <sup>A</sup> †	0/225 (0%)	1/215 (0.47%)	0/221 (0%)
Subarachnoid haemorrhage <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Renal and urinary disorders			
Renal failure acute <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	1/221 (0.45%)
Respiratory, thoracic and mediastinal disorders			
Chronic obstructive pulmonary disease <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Haemothorax <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Pulmonary embolism <sup>A</sup> †	0/225 (0%)	0/215 (0%)	1/221 (0.45%)
Respiratory failure <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)
Vascular disorders			
Aortic aneurysm <sup>A</sup> †	1/225 (0.44%)	0/215 (0%)	0/221 (0%)

† Indicates events were collected by systematic assessment.

\* Indicates events were collected by non-systematic methods.

A Term from vocabulary, MedDRA version 11.0

## Other Adverse Events

Frequency Threshold Above Which Other Adverse Events are Reported: 5%

	700 µg Dexamethasone	350 µg Dexamethasone	Sham Injection
	Affected/At Risk (%)	Affected/At Risk (%)	Affected/At Risk (%)
Total	168/225 (74.67%)	145/215 (67.44%)	94/221 (42.53%)
Eye disorders			
Conjunctival haemorrhage <sup>A †</sup>	46/225 (20.44%)	37/215 (17.21%)	37/221 (16.74%)
Conjunctival hyperaemia <sup>A †</sup>	18/225 (8%)	16/215 (7.44%)	13/221 (5.88%)
Eye pain <sup>A *</sup>	18/225 (8%)	10/215 (4.65%)	10/221 (4.52%)
Maculopathy <sup>A †</sup>	8/225 (3.56%)	15/215 (6.98%)	6/221 (2.71%)
Retinal exudates <sup>A †</sup>	6/225 (2.67%)	2/215 (0.93%)	12/221 (5.43%)
Investigations			
Intraocular pressure increased <sup>A †</sup>	60/225 (26.67%)	56/215 (26.05%)	5/221 (2.26%)
Vascular disorders			
Hypertension <sup>A †</sup>	12/225 (5.33%)	9/215 (4.19%)	11/221 (4.98%)

† Indicates events were collected by systematic assessment.

\* Indicates events were collected by non-systematic methods.

A Term from vocabulary, MedDRA version 11.0

## Limitations and Caveats

[Not specified]

## More Information

### Certain Agreements:

Principal Investigators are NOT employed by the organization sponsoring the study.

There IS an agreement between the Principal Investigator and the Sponsor (or its agents) that restricts the PI's rights to discuss or publish trial results after the trial is completed.

A disclosure restriction on the PI is that the sponsor can review results communications prior to public release and can embargo communications regarding trial results for a period that is less than or equal to 90 days from the time submitted to the sponsor for review. The sponsor cannot require changes to the communication and cannot extend the embargo.

Results Point of Contact:

Name/Official Title: Therapeutic Area Head

Organization: Allergan, Inc.

Phone: (714)246-4500

Email: [clinicaltrials@allergan.com](mailto:clinicaltrials@allergan.com)

---

U.S. National Library of Medicine | U.S. National Institutes of Health | U.S. Department of Health & Human Services