

Results for EudraCT: 2007-002031-92

Histomorphological differentiation of non-erosive reflux disease and functional heartburn in patients with PPI-refractory heartburn.

Kandulski A, Jechorek D, Caro C, Weigt J, Wex T, Mönkemüller K, Malfertheiner P. Aliment Pharmacol Ther. 2013 Sep;38(6):643-51. doi: 10.1111/apt.12428. Epub 2013 Jul 29. PMID: 23895770 Free article.

GERD assessment including pH metry predicts a high response rate to PPI standard therapy.

Kandulski A, Peitz U, Mönkemüller K, Neumann H, Weigt J, Malfertheiner P. BMC Gastroenterol. 2013 Jan 16;13:12. doi: 10.1186/1471-230X-13-12. PMID: 23324360 Free PMC article. Clinical Trial.

Role of tight junction proteins in gastroesophageal reflux disease.

Mönkemüller K, Wex T, Kuester D, Fry LC, Kandulski A, Kropf S, Roessner A, Malfertheiner P. BMC Gastroenterol. 2012 Sep 20;12:128. doi: 10.1186/1471-230X-12-128. PMID: 22994974 Free PMC article.

Assessment of desmosomal components (desmoglein 1-3, plakoglobin) in cardia mucosa in relation to gastroesophageal reflux disease and Helicobacter pylori infection.

Wex T, Kuester D, Mönkemüller K, Stahr A, Fry LC, Kandulski A, Kropf S, Roessner A, Malfertheiner P. Hum Pathol. 2012 Oct;43(10):1745-54. doi: 10.1016/j.humpath.2011.12.024. Epub 2012 Apr 20. PMID: 22521077

Gastro-oesophageal reflux disease is associated with up-regulation of desmosomal components in oesophageal mucosa.

Wex T, Mönkemüller K, Stahr A, Kuester D, Fry LC, Völkel S, Kandulski A, Roessner A, Malfertheiner P. Histopathology. 2012 Feb;60(3):405-15. doi: 10.1111/j.1365-2559.2011.04123.x. PMID: 22276604

Identification of clinical and laboratory markers for predicting eosinophilic esophagitis in adults.

von Arnim U, Wex T, Röhl FW, Neumann H, Küster D, Weigt J, Mönkemüller K, Malfertheiner P. Digestion. 2011;84(4):323-7. doi: 10.1159/000331142. Epub 2011 Nov 9. PMID: 22075653

Helicobacter pylori infection and fundic gastric atrophy are not associated with esophageal squamous cell carcinoma: a case-control study.

Venerito M, Kohrs S, Wex T, Adolf D, Kuester D, Schubert D, Peitz U, Mönkemüller K, Malfertheiner P. Eur J Gastroenterol Hepatol. 2011 Oct;23(10):859-64. doi: 10.1097/MEG.0b013e3283496469. PMID: 21811162

Intercellular space volume is mainly increased in the basal layer of esophageal squamous epithelium in patients with GERD.

Neumann H, Mönkemüller K, Fry LC, Dombrowski F, Kuester D, Beyer M, Malfertheiner P. Dig Dis Sci. 2011 May;56(5):1404-11. doi: 10.1007/s10620-010-1458-0. Epub 2010 Oct 30. PMID: 21053078

[IL-17A and IL-17F gene expression is strongly induced in the mucosa of H. pylori-infected subjects from Kenya and Germany.](#)

Kimang'a A, Revathi G, Kariuki S, Sayed S, Devani S, Vivienne M, Kuester D, Mönkemüller K, Malfertheiner P, Wex T. Scand J Immunol. 2010 Dec;72(6):522-8. doi: 10.1111/j.1365-3083.2010.02469.x. PMID: 21044126 Free article.

[Proteinase-activated receptor-2 in the pathogenesis of gastroesophageal reflux disease.](#)

Kandulski A, Wex T, Mönkemüller K, Kuester D, Fry LC, Roessner A, Malfertheiner P. Am J Gastroenterol. 2010 Sep;105(9):1934-43. doi: 10.1038/ajg.2010.265. Epub 2010 Jun 29. PMID: 20588261

[Multiple cyanoacrylate \(Histoacryl\) emboli after injection therapy of cardia varices.](#)

Neumann H, Scheidbach H, Mönkemüller K, Pech M, Malfertheiner P. Gastrointest Endosc. 2009 Nov;70(5):1025-6; discussion 1026. doi: 10.1016/j.gie.2009.06.013. Epub 2009 Aug 21. PMID: 19699475 No abstract available.

[Zonulin is not increased in the cardiac and esophageal mucosa of patients with gastroesophageal reflux disease.](#)

Wex T, Mönkemüller K, Kuester D, Fry L, Kandulski A, Malfertheiner P. Peptides. 2009 Jun;30(6):1082-7. doi: 10.1016/j.peptides.2009.03.017. Epub 2009 Apr 5. PMID: 19463740

[Chemoprevention of adenocarcinoma associated with Barrett's esophagus: potential options.](#)

Neumann H, Mönkemüller K, Vieth M, Malfertheiner P. Dig Dis. 2009;27(1):18-23. doi: 10.1159/000210099. Epub 2009 May 8. PMID: 19439956 Review.

[\[Catheter-free pH-metry using the Bravo capsule versus standard pH-metry in patients with non-erosive reflux disease \(NERD\)\].](#)

Mönkemüller K, Neumann H, Fry LC, Kolffenbach S, Malfertheiner P. Z Gastroenterol. 2009 Apr;47(4):351-6. doi: 10.1055/s-2008-1027945. Epub 2009 Apr 8. PMID: 19358061 Clinical Trial. German.

[Interleukin-1beta and interleukin-8 expression correlate with the histomorphological changes in esophageal mucosa of patients with erosive and non-erosive reflux disease.](#)

Mönkemüller K, Wex T, Kuester D, Fry LC, Peitz U, Beyer M, Roessner A, Malfertheiner P. Digestion. 2009;79(3):186-95. doi: 10.1159/000211714. Epub 2009 Apr 3. PMID: 19342859

[Gastroesophageal reflux disease does not lead to changes in the secretory leukocyte protease inhibitor expression in esophageal mucosa.](#)

Wex T, Mönkemüller K, Kuester D, Weise S, Kropf S, Fry LC, Stahr A, Völkel S, Roessner A, Malfertheiner P. Eur J Gastroenterol Hepatol. 2009 Feb;21(2):150-8. doi: 10.1097/MEG.0b013e32830e4905. PMID: 19212204

[Efficacy, complications and clinical outcomes of endoscopic sclerotherapy with N-butyl-2-cyanoacrylate for bleeding gastric varices.](#)

Fry LC, Neumann H, Olano C, Malfertheiner P, Mönkemüller K. Dig Dis. 2008;26(4):300-3. doi: 10.1159/000177012. Epub 2009 Jan 30. PMID: 19188718

[Lectin UEA-I-binding proteins are specifically increased in the squamous epithelium of patients with Barrett's esophagus.](#)

Neumann H, Wex T, Mönkemüller K, Vieth M, Fry LC, Malfertheiner P. Digestion. 2008;78(4):201-7. doi: 10.1159/000187724. Epub 2009 Jan 8. PMID: 19129702

[Serum gastrin and pepsinogens do not correlate with the different grades of severity of gastro-oesophageal reflux disease: a matched case-control study.](#)

Monkemuller K, Neumann H, Nocon M, Vieth M, Labenz J, Willich SN, Stolte M, Hocker M, Jaspersen D, Lind T, Malfertheiner P. Aliment Pharmacol Ther. 2008 Aug 15;28(4):491-6. doi: 10.1111/j.1365-2036.2008.03769.x. Epub 2008 Jun 16. PMID: 18557987