

Protocol Registration Receipt  
08/15/2013

Grantor: CDER IND/IDE Number: 074696, 077855 Serial Number: 0052, 0051

A Study to Evaluate Annual Rate of Exacerbations and Safety of 3 Dosage Strengths of Fluticasone Furoate (FF)/GW642444 Inhalation Powder in Subjects With Chronic Obstructive Pulmonary Disease (COPD)

This study has been completed.

Sponsor:	GlaxoSmithKline
Collaborators:	
Information provided by (Responsible Party):	GlaxoSmithKline
ClinicalTrials.gov Identifier:	NCT01017952

► Purpose

The Purpose of this study is to assess the efficacy and safety of three strengths of the FF/GW642444 Inhalation Powder in subject with Chronic Obstructive Pulmonary Disease (COPD)

Condition	Intervention	Phase
Pulmonary Disease, Chronic Obstructive	Drug: FF/GW642444 Inhalation Powder	Phase 3

Condition	Intervention	Phase
	Drug: GW642444 Inhalation Powder	

Study Type: Interventional

Study Design: Treatment, Parallel Assignment, Double Blind (Subject, Investigator), Randomized, Safety/Efficacy Study

Official Title: HZC102970: A 52-week Efficacy and Safety Study to Compare the Effect of Three Dosage Strengths of Fluticasone Furoate/GW642444 Inhalation Powder With GW642444 on the Annual Rate of Exacerbations in Subjects With Chronic Obstructive Pulmonary Disease (COPD)

#### Further study details as provided by GlaxoSmithKline:

##### Primary Outcome Measure:

- Annual Rate of Moderate and Severe COPD Exacerbations Expressed as Least Square Mean [Time Frame: From the start of the double blind study medication until Visit 11 (Week 52)/Early Withdrawal] [Designated as safety issue: No]

The annual rate of moderate and severe chronic obstructive pulmonary disease (COPD) exacerbations during the treatment (trt) period (per participant [par.] per year) was assessed. An exacerbation of COPD, is defined as the worsening of two or more major symptoms (dyspnea, sputum volume, sputum purulence [color]) for at least two consecutive days; or the worsening of any one major symptom together with any one of the minor symptoms (sore throat, cold, fever without other cause, increased cough, increased wheeze) for at least two consecutive days. The COPD exacerbation was categorized as mild, moderate and severe by the investigator. Mild: worsening symptoms of COPD that were self-managed by the par. without the use of oral corticosteroids or antibiotics; Moderate: worsening symptoms of COPD that required treatment with oral corticosteroids and/or antibiotics; Severe: worsening symptoms of COPD that required treatment with in-patient hospitalization.

##### Secondary Outcome Measures:

- Time to First Occurrence of Moderate or Severe COPD Exacerbation [Time Frame: From the start of the double blind study medication until Visit 11 (Week 52)/Early Withdrawal] [Designated as safety issue: No]

Time to first occurrence analyzed by using a Cox proportional hazards model with covariates of treatment, smoking status at screening (stratum), baseline disease severity (pre-dose Day 1 % predicted FEV1) and centre grouping. An exacerbation of COPD is defined as the worsening of two or more major symptoms (dyspnea, sputum volume, sputum purulence [color]) for at least two consecutive days; or the worsening of any one major symptom together with any one of the minor symptoms (sore throat, cold, fever without other cause, increased cough, increased wheeze) for at least two consecutive days. A moderate exacerbation is defined as worsening symptoms of COPD that required treatment with oral corticosteroids and/or antibiotics. A severe exacerbation is defined as worsening symptoms of COPD that required treatment with in-patient hospitalization. The number of participants with a moderate or severe COPD exacerbation while on treatment are presented.

- Annual Rate of Exacerbations Requiring Systemic/Oral Corticosteroids Expressed as Least Square Mean [Time Frame: From the start of the double blind study medication until Visit 11 (Week 52)/Early Withdrawal] [Designated as safety issue: No]

The annual rate of COPD exacerbations during the treatment period (per participant per year) that required systemic/oral corticosteroids was assessed. An exacerbation of COPD is defined as the worsening of two or more major symptoms (dyspnea, sputum volume, sputum purulence [color]) for at least

two consecutive days; or the worsening of any one major symptom together with any one of the minor symptom (sore throat, cold, fever without other cause, increased cough, increased wheeze) for at least two consecutive days. The COPD exacerbation was categorized as mild, moderate, and severe by the investigator. Mild: worsening symptoms of COPD that were self-managed by the participant. Mild exacerbations were not associated with the use of oral corticosteroids or antibiotics. Moderate: worsening symptoms of COPD that required treatment with oral corticosteroids and/or antibiotics. Severe: worsening symptoms of COPD that required treatment with in-patient hospitalization.

- Change From Baseline in Trough FEV1 at Week 52 (Visit 11) [Time Frame: Baseline to Visit 11 (Week 52)/Early Withdrawal] [Designated as safety issue: No]

Pulmonary function was measured by forced expiratory volume in one second (FEV1). Trough FEV1 was defined as the 24-hour post-dose FEV1 assessment, which was obtained at each visit. Analysis performed using a repeated measures model with covariates of treatment, smoking status at Screening (stratum), baseline (pre-dose Day 1), centre grouping, Week, Week by Baseline, and Week by treatment interactions.

Enrollment: 1635

Study Start Date: September 2009

Study Completion Date: October 2011

Primary Completion Date: October 2011

Arms	Assigned Interventions
Experimental: FF/GW642444 Inhalation Powder 100/25 mcg QD Inhaled Corticosteroid (ICS)/ Long Acting Beta Agonist(LABA)	Drug: FF/GW642444 Inhalation Powder Inhaled Corticosteroid (ICS)/ Long Acting Beta Agonist(LABA) for COPD
Experimental: FF/GW642444 Inhalation Powder 50mcg/25mcg QD Inhaled Corticosteroid (ICS)/ Long Acting Beta Agonist(LABA)	Drug: FF/GW642444 Inhalation Powder Inhaled Corticosteroid (ICS)/ Long Acting Beta Agonist(LABA) for COPD
Experimental: FF/GW642444 Inhalation Powder 200/25 mcg QD Inhaled Corticosteroid (ICS)/ Long Acting Beta Agonist(LABA)	Drug: FF/GW642444 Inhalation Powder Inhaled Corticosteroid (ICS)/ Long Acting Beta Agonist(LABA) for COPD
Experimental: GW642444 25mcg QD	Drug: GW642444 Inhalation Powder

Arms	Assigned Interventions
Long Acting Beta Agonist(LABA)	Long Acting Beta Agonist(LABA) Inhalation Powder

## ► Eligibility

Ages Eligible for Study: 40 Years and older

Genders Eligible for Study: Both

Inclusion Criteria:

- Type of subject: outpatient
- Informed consent: Subjects must give their signed and dated written informed consent to participate.
- Gender: Male or female subjects A female is eligible to enter and participate in the study if she is of: Non-child bearing potential (i.e., physiologically incapable of becoming pregnant, including any female who is post-menopausal or surgically sterile). Surgically sterile females are defined as those with a documented hysterectomy and/or bilateral oophorectomy or tubal ligation. Post-menopausal females are defined as being amenorrhoeic for greater than 1 year with an appropriate clinical profile, e.g., age appropriate, history of vasomotor symptoms. However in questionable cases, a blood sample with FSH > 40MIU/ml and estradiol <40pg/ml (<140 pmol/L) is confirmatory. OR

Child bearing potential, has a negative pregnancy test at screening, and agrees to one of the following acceptable contraceptive methods used consistently and correctly (i.e., in accordance with the approved product label and the instructions of the physician for the duration of the study – screening to follow-up contact):

- Complete abstinence from intercourse from screening until the Follow-Up Phone Contact; or
- Male partner is sterile (vasectomy with documentation of azoospermia) prior to female subject entry into the study, and this male partner is the sole partner for that subject; or
- Implants of levonorgestral inserted for at least 1 month prior to the study medication administration but not beyond the third successive year following insertion; or
- Injectable progestogen administered for at least 1 month prior to study medication administration and administered until the Follow-Up Phone Contact; or
- Oral contraceptive (combined or progestogen only) administered for at least one monthly cycle prior to study medication administration; or
- Double barrier method: condom or occlusive cap (diaphragm or cervical/vault caps) plus spermicidal agent (foam/gel/film/cream/suppository); or
- An intrauterine device (IUD), inserted by a qualified physician, with published data showing that the highest expected failure rate is less than 1% per year; or
- Estrogenic vaginal ring; or
- Percutaneous contraceptive patches
  - Age: ≥40 years of age at Screening (Visit 1)
  - COPD diagnosis: Subjects with a clinical history of COPD in accordance with the following definition by the American Thoracic Society/European Respiratory Society [Celli, 2004]: COPD is a preventable and treatable disease characterized by airflow limitation that is not fully reversible. The airflow

limitation is usually progressive and is associated with an abnormal inflammatory response of the lungs to noxious particles or gases, primarily caused by cigarette smoking. Although COPD affects the lungs, it also produces significant systemic consequences.

- Tobacco use: Subjects with a current or prior history of  $\geq 10$  pack-years of cigarette smoking at Screening (Visit 1). Former smokers are defined as those who have stopped smoking for at least 6 months prior to Visit 1. Note: Pipe and/or cigar use cannot be used to calculate pack-year history. Number of pack years = (number of cigarettes per day/20) x number of years smoked
- Severity of Disease:
  - Subject with a measured post-albuterol/salbutamol FEV1/FVC ratio of  $\leq 0.70$  at Screening (Visit 1)
  - Subjects with a measured post-albuterol/salbutamol FEV1  $< 70\%$  of predicted normal values calculated (via centralized vendor equipment) using NHANES III reference equations [Hankinson, 1999] at Screening (Visit 1). Post-bronchodilator spirometry will be performed approximately 10-15 minutes after the subject has self-administered 4 inhalations (i.e., total 400mcg) of albuterol/salbutamol via an MDI with a valved-holding chamber. The study provided central spirometry equipment will calculate the FEV1/FVC ratio and FEV1 percent predicted values.
    - History of Exacerbations: A documented history (e.g., medical record verification) of at least one COPD exacerbation in the 12 months prior to Visit 1 that required either oral corticosteroids, antibiotics and/or hospitalization. Prior use of antibiotics alone does not qualify as an exacerbation history unless the use was associated with treatment of worsening symptoms of COPD, such as increased dyspnea, sputum volume, or sputum purulence (color). Subject verbal reports are not acceptable.

#### Exclusion Criteria:

Subjects meeting any of the following criteria must not be enrolled in the study:

- Pregnancy: Women who are pregnant or lactating or are planning on becoming pregnant during the study.
- Asthma: Subjects with a current diagnosis of asthma. (Subjects with a prior history of asthma are eligible if they have a current diagnosis of COPD)
- $\alpha 1$ -antitrypsin deficiency: Subjects with  $\alpha 1$ -antitrypsin deficiency as the underlying cause of COPD
- Other respiratory disorders: Subjects with active tuberculosis, lung cancer, bronchiectasis, sarcoidosis, lung fibrosis, pulmonary hypertension, interstitial lung diseases or other active pulmonary diseases
- Lung resection: Subjects with lung volume reduction surgery within the 12 months prior to Screening (Visit 1)
- Chest X-ray (or CT scan): Subjects with a chest X-ray (or CT scan) that reveals evidence of clinically significant abnormalities not believed to be due to the presence of COPD. A chest X-ray must be taken at Screening (Visit 1) if a chest X-ray or CT scan is not available within 6 months prior to Visit 1. For sites in Germany, if a chest X-ray (or CT scan) is not available in the 6 months preceding Screening (Visit 1), the subject will not be eligible for the study.
- Risk Factors for Pneumonia: immune suppression (HIV, Lupus, etc) or other risk for pneumonia (e.g. neurological disorders affecting control of the upper airway, such as Parkinson's, Myasthenia Gravis, etc).
- A moderate and severe COPD exacerbation that has not resolved at least 14 days prior to Visit 1 and at least 30 days following the last dose of oral corticosteroids (if applicable).
- Pneumonia and/or moderate and severe COPD exacerbation at Visit 1 Note: Subjects who experience a pneumonia and/or exacerbation at Screening (Visit 1) must be not continue in the study, but may be re-screened at a later time provided the pneumonia and/or COPD exacerbation has resolved prior to the

- re-screening visit. At the Re-screening Visit, the chest x-ray should confirm resolution of pneumonia. The Re-screening Visit must be conducted at least  $\geq 14$  days following the resolution date of the exacerbation and/or pneumonia and at least 30 days following the last dose of oral corticosteroids (if applicable).
- Other diseases/abnormalities: Subjects with historical or current evidence of clinically significant cardiovascular (i.e., pacemaker), neurological, psychiatric, renal, hepatic, immunological, endocrine (including uncontrolled diabetes or thyroid disease) or haematological abnormalities that are uncontrolled. Significant is defined as any disease that, in the opinion of the investigator, would put the safety of the subject at risk through participation, or which would affect the efficacy or safety analysis if the disease/condition exacerbated during the study.
  - Peptic Ulcer disease: Subjects with clinically significant peptic ulcer disease that is uncontrolled.
  - Hypertension: Subjects with clinically significant hypertension that is uncontrolled.
  - Cancer: Subjects with carcinoma that has not been in complete remission for at least 5 years. Carcinoma in situ of the cervix, squamous cell carcinoma and basal cell carcinoma of the skin would not be excluded if the subject has been considered cured within 5 years since diagnosis.
  - Drug/food allergy: Subjects with a history of hypersensitivity to any of the study medications (e.g., beta-agonists, corticosteroid) or components of the inhalation powder (e.g., lactose, magnesium stearate). In addition, subjects with a history of severe milk protein allergy that, in the opinion of the study physician, contraindicates the subject's participation will also be excluded.
  - Drug/alcohol abuse: Subjects with a known or suspected history of alcohol or drug abuse within the last 2 years
  - Medication prior to spirometry: Subjects who are medically unable to withhold their albuterol/salbutamol and/or their ipratropium for the 4-hour period required prior to spirometry testing at each study visit.
  - Additional medication: Unable to stop using certain medications such as bronchodilators and corticosteroids for the protocol-specified times prior to Visit 1 (the Investigator will discuss the specific medications)
  - Oxygen therapy: Subjects receiving treatment with long-term oxygen therapy (LTOT) or nocturnal oxygen therapy required for greater than 12 hours a day. Oxygen prn use (i.e.,  $\leq 12$  hours per day) is not exclusionary.
  - Sleep apnea: Subjects with clinically significant sleep apnea who require use of continuous positive airway pressure (CPAP) device or non-invasive positive pressure ventilation (NIPPV) device.
  - Pulmonary rehabilitation: Subjects who have participated in the acute phase of a Pulmonary Rehabilitation Program within 4 weeks prior to Screening (Visit 1) or who will enter the acute phase of a Pulmonary Rehabilitation Program during the study. Subjects who are in the maintenance phase of a Pulmonary Rehabilitation Program are not excluded.
  - Non-compliance: Subjects at risk of non-compliance, or unable to comply with the study procedures. Any infirmity, disability, or geographic location that would limit compliance for scheduled visits.
  - Questionable validity of consent: Subjects with a history of psychiatric disease, intellectual deficiency, poor motivation or other conditions that will limit the validity of informed consent to participate in the study.
  - Prior use of study medication/other investigational drugs: Subjects who have previously been randomized to treatment with GW642444 Inhalation Powder in the B2C111045 study, randomized to treatment in the HZC111348 study or have participated in the HZC112207, HZC102871, HZC102970, or HZC110946 studies. Subjects who have received an investigational drug within 30 days of entry into this study (Screening), or within 5 drug half-lives of the investigational drug, whichever is longer.
  - Affiliation with investigator site: Study investigators, sub-investigators, study coordinators, employees of a participating investigator or immediate family members of the aforementioned are excluded from participating in this study.

## Contacts and Locations

### Locations

#### United States, Alabama

GSK Investigational Site

Birmingham, Alabama, United States, 35215

GSK Investigational Site

Birmingham, Alabama, United States, 35294

GSK Investigational Site

Mobile, Alabama, United States, 36608

GSK Investigational Site

Ozark, Alabama, United States, 36360

GSK Investigational Site

Tallasse, Alabama, United States, 36078

#### United States, Arizona

GSK Investigational Site

Chandler, Arizona, United States, 85224

GSK Investigational Site

Peoria, Arizona, United States, 85381 - 4828

GSK Investigational Site

Phoenix, Arizona, United States, 85012

GSK Investigational Site

Tucson, Arizona, United States, 85723

#### United States, California

GSK Investigational Site

Fresno, California, United States, 93710

GSK Investigational Site

Fresno, California, United States, 93726

GSK Investigational Site

Long Beach, California, United States, 90808

GSK Investigational Site

Monterey Park, California, United States, 91754

GSK Investigational Site  
National City, California, United States, 91950

GSK Investigational Site  
San Diego, California, United States, 92117

GSK Investigational Site  
San Diego, California, United States, 92128

GSK Investigational Site  
Torrance, California, United States, 90505

GSK Investigational Site  
Torrance, California, United States, 90502

GSK Investigational Site  
Van Nuys, California, United States, 91405

#### United States, Colorado

GSK Investigational Site  
Boulder, Colorado, United States, 80304

GSK Investigational Site  
Thornton, Colorado, United States, 80233

#### United States, Connecticut

GSK Investigational Site  
Hartford, Connecticut, United States, 06105

#### United States, Florida

GSK Investigational Site  
Clearwater, Florida, United States, 33765

GSK Investigational Site  
Cocoa, Florida, United States, 32927

GSK Investigational Site  
Deland, Florida, United States, 32720

GSK Investigational Site  
Fort Lauderdale, Florida, United States, 33316

GSK Investigational Site  
Kissimmee, Florida, United States, 34741

GSK Investigational Site  
Miami, Florida, United States, 33183

GSK Investigational Site

Pensacola, Florida, United States, 32503

GSK Investigational Site

St. Petersburg, Florida, United States, 33707

GSK Investigational Site

Tampa, Florida, United States, 33603

GSK Investigational Site

Winter Park, Florida, United States, 32789

## United States, Georgia

GSK Investigational Site

Gainesville, Georgia, United States

GSK Investigational Site

Martinez, Georgia, United States, 30907

GSK Investigational Site

Riverdale, Georgia, United States, 30274

## United States, Idaho

GSK Investigational Site

Coeur D'Alene, Idaho, United States, 83814

## United States, Illinois

GSK Investigational Site

Aurora, Illinois, United States, 60504

GSK Investigational Site

Elk Grove Village, Illinois, United States, 60007

## United States, Indiana

GSK Investigational Site

Evansville, Indiana, United States, 47714

GSK Investigational Site

Evansville, Indiana, United States, 47713

## United States, Iowa

GSK Investigational Site

Council Bluffs, Iowa, United States, 51503

## United States, Kansas

GSK Investigational Site

Olathe, Kansas, United States, 66061

GSK Investigational Site

Topeka, Kansas, United States, 66606

## United States, Kentucky

GSK Investigational Site

Hazard, Kentucky, United States, 41701

GSK Investigational Site

Madisonville, Kentucky, United States, 42431

## United States, Massachusetts

GSK Investigational Site

Brockton, Massachusetts, United States, 02301

GSK Investigational Site

Fall River, Massachusetts, United States, 02720

GSK Investigational Site

Pittsfield, Massachusetts, United States, 01201

## United States, Michigan

GSK Investigational Site

Cadillac, Michigan, United States, 49601

## United States, Minnesota

GSK Investigational Site

Edina, Minnesota, United States, 55438

GSK Investigational Site

Fridley, Minnesota, United States, 55432

## United States, Missouri

GSK Investigational Site

Columbia, Missouri, United States, 65212

GSK Investigational Site

Springfield, Missouri, United States, 65803

GSK Investigational Site

St. Louis, Missouri, United States, 63143

GSK Investigational Site

St. Louis, Missouri, United States, 63141

## United States, Nebraska

GSK Investigational Site

Lincoln, Nebraska, United States, 68506

GSK Investigational Site

Omaha, Nebraska, United States, 68134

GSK Investigational Site

Papillion, Nebraska, United States, 68046

#### United States, Nevada

GSK Investigational Site

Las Vegas, Nevada, United States, 89106

#### United States, New Jersey

GSK Investigational Site

Cherry Hill, New Jersey, United States, 08003

GSK Investigational Site

Summit, New Jersey, United States, 07091

#### United States, New York

GSK Investigational Site

Albany, New York, United States, 12205

GSK Investigational Site

North Syracuse, New York, United States, 13212

#### United States, North Carolina

GSK Investigational Site

Asheville, North Carolina, United States, 28803

GSK Investigational Site

Burlington, North Carolina, United States, 27215

GSK Investigational Site

Charlotte, North Carolina, United States, 28207

GSK Investigational Site

Statesville, North Carolina, United States, 28625

GSK Investigational Site

Wilmington, North Carolina, United States, 28401

#### United States, Ohio

GSK Investigational Site

Centerville, Ohio, United States, 45459

GSK Investigational Site

Toledo, Ohio, United States, 43614

#### United States, Oregon

GSK Investigational Site

Portland, Oregon, United States, 97220

## United States, Pennsylvania

GSK Investigational Site

Philadelphia, Pennsylvania, United States, 19140

## United States, South Carolina

GSK Investigational Site

Charleston, South Carolina, United States, 29406

GSK Investigational Site

Columbia, South Carolina, United States, 29201

GSK Investigational Site

Easley, South Carolina, United States, 29640

GSK Investigational Site

Greenville, South Carolina, United States, 29615

GSK Investigational Site

Murrells Inlet, South Carolina, United States, 29576

GSK Investigational Site

Pelzer, South Carolina, United States, 29669

GSK Investigational Site

Spartanburg, South Carolina, United States, 29303

GSK Investigational Site

Spartanburg, South Carolina, United States, 29303

## United States, Tennessee

GSK Investigational Site

Nashville, Tennessee, United States, 37232

## United States, Texas

GSK Investigational Site

Corsicana, Texas, United States, 75110

GSK Investigational Site

Dallas, Texas, United States, 75231

GSK Investigational Site

Houston, Texas, United States, 77030

GSK Investigational Site

Houston, Texas, United States, 77030

GSK Investigational Site

New Braunfels, Texas, United States, 78130

GSK Investigational Site

San Antonio, Texas, United States, 78229

GSK Investigational Site

Wichita Falls, Texas, United States, 76309

#### United States, Virginia

GSK Investigational Site

Newport News, Virginia, United States, 23606

GSK Investigational Site

Richmond, Virginia, United States, 23249

#### United States, Wisconsin

GSK Investigational Site

Madison, Wisconsin, United States, 53715

#### Argentina

GSK Investigational Site

Ciudad Autónoma de Buenos Aires, Argentina, C1121ABE

GSK Investigational Site

Tucuman, Argentina, 4000

GSK Investigational Site

Tucumán, Argentina, T4000DGF

GSK Investigational Site

Ciudad Autonoma de Buenos Aires, Buenos Aires, Argentina, C1405BCH

GSK Investigational Site

Mendoza, Mendoza, Argentina, M5500CCG

#### Australia, New South Wales

GSK Investigational Site

Kingswood, New South Wales, Australia, 2747

GSK Investigational Site

Westmead, New South Wales, Australia, 2145

#### Australia, Queensland

GSK Investigational Site

Auchenflower, Queensland, Australia, 4066

GSK Investigational Site

Southport, Queensland, Australia, 4215

## Australia, South Australia

GSK Investigational Site

Adelaide, South Australia, Australia, 5000

## Australia, Tasmania

GSK Investigational Site

Hobart, Tasmania, Australia, 7000

## Australia, Victoria

GSK Investigational Site

Box Hill, Victoria, Australia, 3128

## Australia, Western Australia

GSK Investigational Site

Nedlands, Western Australia, Australia, 6009

## Canada, Manitoba

GSK Investigational Site

Winnipeg, Manitoba, Canada, R2K 3S8

## Canada, Nova Scotia

GSK Investigational Site

Truro, Nova Scotia, Canada, B2N 1L2

## Canada, Ontario

GSK Investigational Site

Grimsby, Ontario, Canada, L3M 1P3

GSK Investigational Site

Hamilton, Ontario, Canada, L8M 1K7

GSK Investigational Site

London, Ontario, Canada, N5W 6A2

GSK Investigational Site

Newmarket, Ontario, Canada, L3Y 5G8

GSK Investigational Site

Sarnia, Ontario, Canada, N7T 4X3

GSK Investigational Site

Toronto, Ontario, Canada, M5G 1N8

## Canada, Quebec

GSK Investigational Site

Gatineau, Quebec, Canada, J8Y 6S8  
GSK Investigational Site  
Mirabel, Quebec, Canada, J7J 2K8  
GSK Investigational Site  
Montreal, Quebec, Canada, H2X 2P4

## Chile

GSK Investigational Site  
Talcahuano, Chile, 4270918  
GSK Investigational Site  
Puente Alto - Santiago, Región Metro De Santiago, Chile, 8207257  
GSK Investigational Site  
Santiago, Región Metro De Santiago, Chile, 7500691

## Denmark

GSK Investigational Site  
Aalborg, Denmark, 9100  
GSK Investigational Site  
Aarhus C, Denmark, 8000  
GSK Investigational Site  
Hvidovre, Denmark, 2650  
GSK Investigational Site  
Kobenhavn NV, Denmark, 2400  
GSK Investigational Site  
Odense C, Denmark, 5000

## Germany

GSK Investigational Site  
Hamburg, Hamburg, Germany, 20354  
GSK Investigational Site  
Frankfurt, Hessen, Germany, 60596  
GSK Investigational Site  
Kassel, Hessen, Germany, 34121  
GSK Investigational Site  
Marburg, Hessen, Germany, 35037  
GSK Investigational Site  
Wiesbaden, Hessen, Germany, 65183

GSK Investigational Site  
Hannover, Niedersachsen, Germany, 30167

GSK Investigational Site  
Essen, Nordrhein-Westfalen, Germany, 45359

## Italy

GSK Investigational Site  
Salerno, Campania, Italy, 84100

GSK Investigational Site  
San Felice a Cancello Caserta, Campania, Italy, 81027

GSK Investigational Site  
Modena, Emilia-Romagna, Italy, 41124

GSK Investigational Site  
Parma, Emilia-Romagna, Italy, 43100

GSK Investigational Site  
Roma, Lazio, Italy, 00168

GSK Investigational Site  
Foggia, Puglia, Italy, 71100

GSK Investigational Site  
Palermo, Sicilia, Italy, 90146

GSK Investigational Site  
Pisa, Toscana, Italy, 56124

GSK Investigational Site  
Pisa, Toscana, Italy, 56124

GSK Investigational Site  
Perugia, Umbria, Italy, 06156

## Mexico

GSK Investigational Site  
Guadalajara, Jalisco, Mexico, 44600

GSK Investigational Site  
Zapopan, Jalisco, Mexico, 45040

GSK Investigational Site  
Monterrey, Nuevo León, Mexico, 64460

## Netherlands

GSK Investigational Site

Almelo, Netherlands, 7609 PP  
GSK Investigational Site

Almere, Netherlands, 1315 RA  
GSK Investigational Site

Beek, Netherlands, 6191 JW  
GSK Investigational Site

Breda, Netherlands, 4819 EV  
GSK Investigational Site

Eindhoven, Netherlands, 5623 EJ  
GSK Investigational Site

Groningen, Netherlands, 9728 NP  
GSK Investigational Site

Hoorn, Netherlands, 1624 NP  
GSK Investigational Site

Horn, Netherlands, 6085 NM  
GSK Investigational Site

Nieuwegein, Netherlands, 3435 CM  
GSK Investigational Site

Zutphen, Netherlands, 7207 AE

## Peru

GSK Investigational Site  
Callao, Peru, Callao 2

GSK Investigational Site  
Lima 18, Lima, Peru

GSK Investigational Site  
San Miguel, Lima, Peru, Lima 32

GSK Investigational Site  
Santiago de Surco, Lima, Peru, Lima 33

## South Africa

GSK Investigational Site  
Bellville, South Africa, 7530

GSK Investigational Site  
Bloemfontein, South Africa, 9301

GSK Investigational Site  
Cape Town, South Africa, 7572

GSK Investigational Site  
Gatesville, South Africa, 7764

GSK Investigational Site  
Gatesville, South Africa, 7764

GSK Investigational Site  
Mowbray, South Africa, 7700

GSK Investigational Site  
Roodepoort, South Africa, 1724

GSK Investigational Site  
Worcester, South Africa, 6850

GSK Investigational Site  
Meyerspark, Gauteng, South Africa, 0184

GSK Investigational Site  
Parktown, Gauteng, South Africa, 2193

GSK Investigational Site  
Waterkloof Ridge, Gauteng, South Africa, 0181

## Spain

GSK Investigational Site  
Alicante, Spain, 03114

GSK Investigational Site  
Cartagena (Murcia), Spain, 30202

GSK Investigational Site  
Cáceres, Spain, 10003

GSK Investigational Site  
Elda, Spain, 03600

GSK Investigational Site  
Galdakano, Spain, 48960

GSK Investigational Site  
Madrid, Spain, 28040

GSK Investigational Site  
Orihuela/Alicante, Spain, 03314

GSK Investigational Site  
Pama de Mallorca, Spain, 07010

GSK Investigational Site  
Pozuelo de Alarcón/Madrid, Spain, 28223

## Sweden

GSK Investigational Site

Bankeryd, Sweden, SE-564 31

GSK Investigational Site

Karlskrona, Sweden, SE-371 41

GSK Investigational Site

Lidingö, Sweden, SE-181 58

GSK Investigational Site

Åtvidaberg, Sweden, SE-597 26

## United Kingdom

GSK Investigational Site

Cambridge, United Kingdom, CB2 0QQ

GSK Investigational Site

Chertsey, Surrey, United Kingdom, KT16 0PZ

GSK Investigational Site

Wythenshawe, Manchester, United Kingdom, M23 9LT

GSK Investigational Site

Southampton, Hampshire, United Kingdom, SO16 6YD

GSK Investigational Site

Bradford on Avon, Wiltshire, United Kingdom, BA15 1DQ

GSK Investigational Site

Trowbridge, Wiltshire, United Kingdom, BA14 8QA

## Investigators

Study Director:

GSK Clinical Trials

GlaxoSmithKline

## More Information

Responsible Party: GlaxoSmithKline

Study ID Numbers: 102970

Health Authority: United States: Food and Drug Administration

---

## Study Results

## Participant Flow

### Pre-Assignment Details

At Visit (V) 1, eligible participants (par.) entered a 4-week, open-label Run-in Period (RIP) to establish a stable Baseline. At V 2, eligible par. were randomized to a 52-week, double-blind Treatment Period. 2635 par. were screened, 2092 par. entered the RIP, and 1635 par. were randomized, out of which 1633 par. received  $\geq 1$  study treatment dose.

### Reporting Groups

	Description
FP/SAL 250/50 µg BID	Participants (Par.) were instructed to take open label Fluticasone Propionate and Salmeterol (FP/SAL) 250/50 microgram (µg) twice daily (BID) from the ACCUHALER/DISKUS, one inhalation each morning and evening with approximately 12 hours between doses. In addition, all par. were provided supplemental albuterol/salbutamol (metered dose inhaler [MDI] and/or nebulers) to be used as needed throughout the study.
VI 25 µg QD	Participants received a Vilanterol (VI) 25 µg dry inhalation powder once daily (QD) in the morning from the Dry Powder Inhaler (DPI) for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 50/25 µg QD	Participants received a Fluticasone Furoate/Vilanterol (FF/VI) 50/25 µg inhalation powder QD in the morning from the NDPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 100/25 µg QD	Participants received a FF/VI 100/25 µg inhalation powder QD in the morning from the NDPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.

	Description
FF/VI 200/25 µg QD	Participants received a FF/VI 200/25 µg inhalation powder QD in the morning from the NDPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulas) to be used as needed throughout the study.

#### 4-week, Open-label Run-In Period

	FP/SAL 250/50 µg BID	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Started	2092	0	0	0	0
Completed	1633	0	0	0	0
Not Completed	459	0	0	0	0
Did Not Meet Continuation Criteria	356	0	0	0	0
Adverse Event	23	0	0	0	0
Study Closed/Terminated	1	0	0	0	0
Lost to Follow-up	6	0	0	0	0
Physician Decision	10	0	0	0	0
Withdrawal by Subject	63	0	0	0	0

#### 52-week, Double-blind Treatment Period

	FP/SAL 250/50 µg BID	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Started	0	409	412	403	409

	FP/SAL 250/50 µg BID	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Completed the Treatment Period	0	285 <sup>[1]</sup>	305 <sup>[2]</sup>	293 <sup>[3]</sup>	307 <sup>[4]</sup>
Completed	0	284 <sup>[5]</sup>	303 <sup>[6]</sup>	291 <sup>[7]</sup>	306 <sup>[8]</sup>
Not Completed	0	125	109	112	103
Adverse Event	0	25	32	35	30
Withdrawal by Subject	0	30	22	25	25
Lack of Efficacy	0	35	14	16	14
Protocol Violation	0	7	11	9	8
Met Protocol-Defined Stopping Criteria	0	11	13	12	9
Study Closed/Terminated	0	1	1	0	0
Lost to Follow-up	0	6	8	6	10
Physician Decision	0	10	8	9	7

[1] Par. completed the treatment period if they attended the last treatment visit (Visit 11).

[2] Par. completed the treatment period if they attended the last treatment visit (Visit 11).

[3] Par. completed the treatment period if they attended the last treatment visit (Visit 11).

[4] Par. completed the treatment period if they attended the last treatment visit (Visit 11).

[5] Par. completed the study if they completed the treatment period and safety follow-up phone contact.

[6] Par. completed the study if they completed the treatment period and safety follow-up phone contact.

[7] Par. completed the study if they completed the treatment period and safety follow-up phone

contact.

[8] Par. completed the study if they completed the treatment period and safety follow-up phone contact.

## ▶ Baseline Characteristics

### Reporting Groups

	Description
VI 25 µg QD	Participants received a Vilanterol (VI) 25 µg dry inhalation powder once daily (QD) in the morning from the Dry Powder Inhaler (DPI) for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 50/25 µg QD	Participants received a Fluticasone Furoate/Vilanterol (FF/VI) 50/25 µg inhalation powder QD in the morning from the NDPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 100/25 µg QD	Participants received a FF/VI 100/25 µg inhalation powder QD in the morning from the NDPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 200/25 µg QD	Participants received a FF/VI 200/25 µg inhalation powder QD in the morning from the NDPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.

### Baseline Measures

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD	Total
Number of Participants	409	412	403	409	1633
Age, Continuous [units: Years] Mean (Standard Deviation)	63.6 (9.29)	63.7 (9.56)	64.0 (9.28)	63.5 (8.84)	63.7 (9.24)
Gender, Male/Female [units: Participants]					
Female	174	181	181	191	727
Male	235	231	222	218	906
Race/Ethnicity, Customized [units: participants]					
White	360	359	353	359	1431
African American/ African Heritage	9	14	7	9	39
Asian	4	3	5	3	15
African American/African Heritage & White	0	1	1	0	2
American Indian or Alaska Native & White	20	19	21	20	80
Asian & White	0	0	0	1	1
Native Hawaiian or other Pacific Islander	0	0	0	1	1
American Indian or Alaska Native	16	16	16	16	64

## ► Outcome Measures

### 1. Primary Outcome Measure:

Measure Title	Annual Rate of Moderate and Severe COPD Exacerbations Expressed as Least Square Mean
Measure Description	The annual rate of moderate and severe chronic obstructive pulmonary disease (COPD) exacerbations during the treatment (trt) period (per participant [par.] per year) was assessed. An exacerbation of COPD, is defined as the worsening of two or more major symptoms (dyspnea, sputum volume, sputum purulence [color]) for at least two consecutive days; or the worsening of any one major symptom together with any one of the minor symptoms (sore throat, cold, fever without other cause, increased cough, increased wheeze) for at least two consecutive days. The COPD exacerbation was categorized as mild, moderate and severe by the investigator. Mild: worsening symptoms of COPD that were self-managed by the par. without the use of oral corticosteroids or antibiotics; Moderate: worsening symptoms of COPD that required treatment with oral corticosteroids and/or antibiotics; Severe: worsening symptoms of COPD that required treatment with in-patient hospitalization.
Time Frame	From the start of the double blind study medication until Visit 11 (Week 52)/Early Withdrawal
Safety Issue?	No

### Analysis Population Description

Intent-to-Treat (ITT) Population: all par. randomized who received at least 1 dose of study drug and with available data for analysis. Analysis used a negative binomial regression model with covariates of trt, smoking status at Screening, Baseline pre-dose Day 1 % predicted FEV1 and region and with logarithm of time on trt as an offset variable.

### Reporting Groups

	Description
VI 25 µg QD	Participants received a Vilanterol (VI) 25 µg dry inhalation powder once daily (QD) in the morning from the Dry Powder Inhaler (DPI) for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 50/25 µg QD	Participants received a Fluticasone Furoate/Vilanterol (FF/VI) 50/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 100/25 µg QD	Participants received a FF/VI 100/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 200/25 µg QD	Participants received a FF/VI 200/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.

### Measured Values

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Number of Participants Analyzed	409	412	403	409
Annual Rate of Moderate and Severe COPD Exacerbations Expressed as Least Square Mean [units: Exacerbations per participant per year] Least Squares Mean (95% Confidence Interval)	1.14 (0.99 to 1.32)	0.92 (0.79 to 1.07)	0.90 (0.77 to 1.05)	0.79 (0.67 to 0.92)

Statistical Analysis 1 for Annual Rate of Moderate and Severe COPD Exacerbations Expressed as Least Square Mean

Groups	VI 25 µg QD, FF/VI 50/25 µg QD
Method	Other [Generalized Linear Model]
P-Value	0.040
Risk Ratio (RR)	0.81
95% Confidence Interval	0.66 to 0.99

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Assuming Negative Binomial distribution with logarithm of time on treatment as an offset variable.

Statistical Analysis 2 for Annual Rate of Moderate and Severe COPD Exacerbations Expressed as Least Square Mean

Groups	VI 25 µg QD, FF/VI 100/25 µg QD
Method	Other [Generalized Linear Model]
P-Value	0.024
Risk Ratio (RR)	0.79
95% Confidence Interval	0.64 to 0.97

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Assuming Negative Binomial distribution with logarithm of time on treatment as an offset variable.

### Statistical Analysis 3 for Annual Rate of Moderate and Severe COPD Exacerbations Expressed as Least Square Mean

Groups	VI 25 µg QD, FF/VI 200/25 µg QD
Method	Other [Generalized Linear Model]
P-Value	<0.001
Risk Ratio (RR)	0.69
95% Confidence Interval	0.56 to 0.85

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Assuming Negative Binomial distribution with logarithm of time on treatment as an offset variable.

## 2. Secondary Outcome Measure:

Measure Title	Time to First Occurrence of Moderate or Severe COPD Exacerbation
Measure Description	Time to first occurrence analyzed by using a Cox proportional hazards model with covariates of treatment, smoking status at screening (stratum), baseline disease severity (pre-dose Day 1 % predicted FEV1) and centre grouping. An exacerbation of COPD is defined as the worsening of two or more major symptoms (dyspnea, sputum volume, sputum purulence [color]) for at least two consecutive days; or the worsening of any one major symptom together with any one of the minor symptoms (sore throat, cold, fever without other cause,

	increased cough, increased wheeze) for at least two consecutive days. A moderate exacerbation is defined as worsening symptoms of COPD that required treatment with oral corticosteroids and/or antibiotics. A severe exacerbation is defined as worsening symptoms of COPD that required treatment with in-patient hospitalization. The number of participants with a moderate or severe COPD exacerbation while on treatment are presented.
Time Frame	From the start of the double blind study medication until Visit 11 (Week 52)/Early Withdrawal
Safety Issue?	No

## Analysis Population Description

ITT Population

### Reporting Groups

	Description
VI 25 µg QD	Participants received a Vilanterol (VI) 25 µg dry inhalation powder once daily (QD) in the morning from the Dry Powder Inhaler (DPI) for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 50/25 µg QD	Participants received a Fluticasone Furoate/Vilanterol (FF/VI) 50/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 100/25 µg QD	Participants received a FF/VI 100/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.

	Description
FF/VI 200/25 µg QD	Participants received a FF/VI 200/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulas) to be used as needed throughout the study.

### Measured Values

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Number of Participants Analyzed	409	412	403	409
Time to First Occurrence of Moderate or Severe COPD Exacerbation [units: Participants]	197	198	177	160

### Statistical Analysis 1 for Time to First Occurrence of Moderate or Severe COPD Exacerbation

Groups	VI 25 µg QD, FF/VI 50/25 µg QD
Method	Regression, Cox
P-Value	0.177
Hazard Ratio (HR)	0.87
95% Confidence Interval	0.71 to 1.06

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

[Not specified.]

### Statistical Analysis 2 for Time to First Occurrence of Moderate or Severe COPD Exacerbation

Groups	VI 25 µg QD, FF/VI 100/25 µg QD
Method	Regression, Cox
P-Value	0.036
Hazard Ratio (HR)	0.88
95% Confidence Interval	0.66 to 0.99

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

[Not specified.]

### Statistical Analysis 3 for Time to First Occurrence of Moderate or Severe COPD Exacerbation

Groups	VI 25 µg QD, FF/VI 200/25 µg QD
Method	Regression, Cox
P-Value	<0.001
Hazard Ratio (HR)	0.66
95% Confidence Interval	0.54 to 0.82

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

[Not specified.]

### 3. Secondary Outcome Measure:

Measure Title	Annual Rate of Exacerbations Requiring Systemic/Oral Corticosteroids Expressed as Least Square Mean
Measure Description	The annual rate of COPD exacerbations during the treatment period (per participant per year) that required systemic/oral corticosteroids was assessed. An exacerbation of COPD is defined as the worsening of two or more major symptoms (dyspnea, sputum volume, sputum purulence [color]) for at least two consecutive days; or the worsening of any one major symptom together with any one of the minor symptom (sore throat, cold, fever without other cause, increased cough, increased wheeze) for at least two consecutive days. The COPD exacerbation was categorized as mild, moderate, and severe by the investigator. Mild: worsening symptoms of COPD that were self-managed by the participant. Mild exacerbations were not associated with the use of oral corticosteroids or antibiotics. Moderate: worsening symptoms of COPD that required treatment with oral corticosteroids and/or antibiotics. Severe: worsening symptoms of COPD that required treatment with in-patient hospitalization.
Time Frame	From the start of the double blind study medication until Visit 11 (Week 52)/Early Withdrawal
Safety Issue?	No

#### Analysis Population Description

Intent-to-Treat (ITT) Population: all par. randomized who received at least 1 dose of study drug and with available data for analysis. Analysis used a negative binomial regression model with covariates of trt, smoking status at Screening, Baseline pre-dose Day 1 % predicted FEV1 and region and with logarithm of time on trt as an offset variable.

#### Reporting Groups

	Description
VI 25 µg QD	Participants received a Vilanterol (VI) 25 µg dry inhalation powder once daily (QD) in the morning from the Dry Powder Inhaler (DPI) for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 50/25 µg QD	Participants received a Fluticasone Furoate/Vilanterol (FF/VI) 50/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 100/25 µg QD	Participants received a FF/VI 100/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 200/25 µg QD	Participants received a FF/VI 200/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.

### Measured Values

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Number of Participants Analyzed	409	412	403	409
Annual Rate of Exacerbations Requiring Systemic/Oral Corticosteroids Expressed as Least Square Mean [units: Exacerbations per participant per year] Least Squares Mean (95% Confidence Interval)	0.86 (0.72 to 1.03)	0.72 (0.60 to 0.86)	0.66 (0.55 to 0.80)	0.56 (0.46 to 0.68)

### Statistical Analysis 1 for Annual Rate of Exacerbations Requiring Systemic/Oral Corticosteroids Expressed as Least Square Mean

Groups	VI 25 µg QD, FF/VI 50/25 µg QD
Method	Other [Generalized Linear Model]
P-Value	0.154
Risk Ratio (RR)	0.84
95% Confidence Interval	0.65 to 1.07

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Assuming Negative Binomial distribution with logarithm of time on treatment as an offset variable.

### Statistical Analysis 2 for Annual Rate of Exacerbations Requiring Systemic/Oral Corticosteroids Expressed as Least Square Mean

Groups	VI 25 µg QD, FF/VI 100/25 µg QD
Method	Other [Generalized Linear Model]
P-Value	0.041
Risk Ratio (RR)	0.77
95% Confidence Interval	0.60 to 0.99

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Assuming Negative Binomial distribution with logarithm of time on treatment as an offset variable

### Statistical Analysis 3 for Annual Rate of Exacerbations Requiring Systemic/Oral Corticosteroids Expressed as Least Square Mean

Groups	VI 25 µg QD, FF/VI 200/25 µg QD
Method	Other [Generalized Linear Model]
P-Value	<0.001
Risk Ratio (RR)	0.65
95% Confidence Interval	0.51 to 0.84

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Assuming Negative Binomial distribution with logarithm of time on treatment as an offset variable

#### 4. Secondary Outcome Measure:

Measure Title	Change From Baseline in Trough FEV1 at Week 52 (Visit 11)
Measure Description	Pulmonary function was measured by forced expiratory volume in one second (FEV1). Trough FEV1 was defined as the 24-hour post-dose FEV1 assessment, which was obtained at each visit. Analysis performed using a repeated measures model with covariates of treatment, smoking status at Screening (stratum), baseline (pre-dose Day 1), centre grouping, Week, Week by Baseline, and Week by treatment interactions.

Time Frame	Baseline to Visit 11 (Week 52)/Early Withdrawal
Safety Issue?	No

### Analysis Population Description

ITT Population. Number of participants presented represent those with data available at the time point being presented, however all participants in the ITT population without missing covariate information and with at least one post Baseline measurement are included in the analysis.

### Reporting Groups

	Description
VI 25 µg QD	Participants received a Vilanterol (VI) 25 µg dry inhalation powder once daily (QD) in the morning from the Dry Powder Inhaler (DPI) for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 50/25 µg QD	Participants received a Fluticasone Furoate/Vilanterol (FF/VI) 50/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 100/25 µg QD	Participants received a FF/VI 100/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 200/25 µg QD	Participants received a FF/VI 200/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.

### Measured Values

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Number of Participants Analyzed	276	304	287	300
Change From Baseline in Trough FEV1 at Week 52 (Visit 11) [units: Liters] Least Squares Mean (Standard Error)	-0.019 (0.0116)	0.015 (0.0113)	0.005 (0.0115)	0.006 (0.0113)

#### Statistical Analysis 1 for Change From Baseline in Trough FEV1 at Week 52 (Visit 11)

Groups	VI 25 µg QD, FF/VI 50/25 µg QD
Method	Mixed Models Analysis
P-Value	0.034
Other Estimated Parameter [Least squares mean difference]	0.034
95% Confidence Interval	0.003 to 0.066

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

Nominal p-value

Other relevant information, such as adjustments or degrees of freedom:

Restricted maximum likelihood (REML)-based repeated measures approach (MMRM).

#### Statistical Analysis 2 for Change From Baseline in Trough FEV1 at Week 52 (Visit 11)

Groups	VI 25 µg QD, FF/VI 100/25 µg QD
Method	Mixed Models Analysis
P-Value	0.143

Other Estimated Parameter [Least squares mean difference]	0.024
95% Confidence Interval	-0.008 to 0.056

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Restricted maximum likelihood (REML)-based repeated measures approach (MMRM).

### Statistical Analysis 3 for Change From Baseline in Trough FEV1 at Week 52 (Visit 11)

Groups	VI 25 µg QD, FF/VI 200/25 µg QD
Method	Mixed Models Analysis
P-Value	0.115
Other Estimated Parameter [Least squares mean difference]	0.026
95% Confidence Interval	-0.006 to 0.057

Additional details about the analysis, such as null hypothesis and power calculation:

[Not specified.]

Additional information, such as whether or not the p-value is adjusted for multiple comparisons and the a priori threshold for statistical significance:

[Not specified.]

Other relevant information, such as adjustments or degrees of freedom:

Restricted maximum likelihood (REML)-based repeated measures approach (MMRM).

## ▶ Reported Adverse Events

### Reporting Groups

	Description
VI 25 µg QD	Participants received a Vilanterol (VI) 25 µg dry inhalation powder once daily (QD) in the morning from the Dry Powder Inhaler (DPI) for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 50/25 µg QD	Participants received a Fluticasone Furoate/Vilanterol (FF/VI) 50/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 100/25 µg QD	Participants received a FF/VI 100/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.
FF/VI 200/25 µg QD	Participants received a FF/VI 200/25 µg inhalation powder QD in the morning from the DPI for the duration of the 52 weeks. In addition, all participants were provided supplemental albuterol/salbutamol (MDI and/or nebulers) to be used as needed throughout the study.

### Time Frame

Serious adverse events (SAEs) and non-serious AEs will be collected from Baseline to the end of the treatment period (up to 52 weeks).

### Additional Description

SAEs and non-serious adverse events are reported for members of the Intent-to-Treat Population, comprised of all randomized

participants who received at least one dose of study medication during the treatment period.

### Serious Adverse Events

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Total # participants affected/at risk	66/409 (16.14%)	71/412 (17.23%)	67/403 (16.63%)	61/409 (14.91%)
Blood and lymphatic system disorders				
Anaemia † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	2/403 (0.5%)	0/409 (0%)
# events				
Cardiac disorders				
Acute myocardial infarction † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	2/403 (0.5%)	0/409 (0%)
# events				
Angina pectoris † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Angina unstable † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Atrial fibrillation † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	1/412 (0.24%)	2/403 (0.5%)	0/409 (0%)
# events				
Atrial flutter † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Bradycardia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	2/409 (0.49%)
# events				
Cardiac arrest † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	1/403 (0.25%)	0/409 (0%)
# events				
Cardiac failure † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Cardiac failure congestive † <sup>A</sup>				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Cardiomyopathy † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Coronary artery disease † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Coronary artery stenosis † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Myocardial infarction † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	1/409 (0.24%)
# events				
Myocardial ischaemia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Supraventricular tachycardia † <sup>A</sup>				
# participants affected/at risk	2/409 (0.49%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Tachyarrhythmia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Tachycardia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	2/403 (0.5%)	0/409 (0%)
# events				
Ventricular tachycardia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Ear and labyrinth disorders				
Vertigo † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Endocrine disorders				
Hyperthyroidism † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	1/409 (0.24%)
# events				
Eye disorders				
Eye haemorrhage † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Gastrointestinal disorders				
Abdominal adhesions † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Abdominal hernia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	1/409 (0.24%)
# events				
Abdominal pain † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	2/412 (0.49%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
risk				
# events				
Constipation † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	2/412 (0.49%)	0/403 (0%)	0/409 (0%)
# events				
Diverticulum intestinal † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Duodenal ulcer † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Gastric ulcer † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Gastroesophageal reflux disease † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Ileus † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Inguinal hernia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Intestinal obstruction † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Intestinal perforation † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Megacolon † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Oesophageal achalasia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Oesophageal stenosis † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Pancreatitis acute † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Small intestinal obstruction † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
General disorders				
Chest pain † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	1/409 (0.24%)
# events				
Fatigue † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Non-cardiac chest pain † <sup>A</sup>				
# participants affected/at risk	2/409 (0.49%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Hepatobiliary disorders				
Cholecystitis † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Cholecystitis acute † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Cholelithiasis † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	1/412 (0.24%)	2/403 (0.5%)	0/409 (0%)
# events				
Immune system disorders				
Anaphylactic shock † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
<b>Infections and infestations</b>				
Abscess limb † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Bacteraemia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Bronchitis † <sup>A</sup>				
# participants affected/at risk	2/409 (0.49%)	2/412 (0.49%)	1/403 (0.25%)	0/409 (0%)
# events				
Cellulitis † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	2/403 (0.5%)	1/409 (0.24%)
# events				
Diverticulitis † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Gastroenteritis † <sup>A</sup>				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Infective exacerbation of chronic obstructive airways diseases † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	2/412 (0.49%)	2/403 (0.5%)	3/409 (0.73%)
# events				
Lobar pneumonia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	2/403 (0.5%)	0/409 (0%)
# events				
Lower respiratory tract infection † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	2/409 (0.49%)
# events				
Lung abscess † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Pelvic inflammatory disease † <sup>A</sup>				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Pneumonia † <sup>A</sup>				
# participants affected/at risk	6/409 (1.47%)	10/412 (2.43%)	12/403 (2.98%)	9/409 (2.2%)
# events				
Pneumonia primary atypical † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Postoperative wound infection † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Pyelonephritis † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Tracheobronchitis † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Urinary tract infection † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	1/409 (0.24%)
# events				
Injury, poisoning and procedural complications				
Acetabulum fracture † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Concussion † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Contusion † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Fall † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Femur fracture † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Hip fracture † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	2/412 (0.49%)	0/403 (0%)	0/409 (0%)
# events				
Humerus fracture † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Muscle strain † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Pelvic fracture † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Rib fracture † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Road traffic accident † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Spinal compression fracture † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Traumatic haematoma † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Investigations				
Blood pressure increased † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Electrocardiogram abnormal † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Metabolism and nutrition disorders				
Gout † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Hyponatraemia † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Musculoskeletal and connective tissue disorders				
Arthralgia † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Arthropathy † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Musculoskeletal chest pain				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
† <sup>A</sup>				
# participants affected/at risk	2/409 (0.49%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Osteoarthritis † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Neoplasms benign, malignant and unspecified (incl cysts and polyps)				
B-cell lymphoma † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Basal cell carcinoma † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Bladder cancer † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Bladder neoplasm † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Bone neoplasm † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Breast cancer † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Bronchial neoplasm † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Laryngeal cancer † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Leukaemia † <sup>A</sup>				
# participants affected/at	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
risk				
# events				
Lung neoplasm † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Lung neoplasm malignant † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Lung squamous cell carcinoma stage unspecified † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	1/403 (0.25%)	1/409 (0.24%)
# events				
Malignant melanoma † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Metastases to bone † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Metastases to gastrointestinal tract † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Metastatic squamous cell carcinoma † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Neoplasm malignant † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Neoplasm recurrence † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Neuroendocrine tumour † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Pelvic neoplasm † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Pituitary tumour benign † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Prostate cancer † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	1/409 (0.24%)
# events				
Renal cancer † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Renal cell carcinoma † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Skin cancer † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Small cell lung cancer stage unspecified † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	1/409 (0.24%)
# events				
Squamous cell carcinoma † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Thyroid cancer † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	2/409 (0.49%)
# events				
Uterine leiomyoma † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Nervous system disorders				
Cerebral ischaemia † <sup>A</sup>				
# participants affected/at	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
risk				
# events				
Cerebrovascular accident † A				
# participants affected/at risk	2/409 (0.49%)	1/412 (0.24%)	0/403 (0%)	1/409 (0.24%)
# events				
Convulsion † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Ischaemic stroke † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Presyncope † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Subarachnoid haemorrhage † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Syncope † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	2/403 (0.5%)	0/409 (0%)
# events				
Transient ischaemic attack † A				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Psychiatric disorders				
Major depression † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	1/409 (0.24%)
# events				
Suicidal ideation † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Renal and urinary disorders				
Calculus ureteric † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
risk				
# events				
Haematuria † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Renal failure acute † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Urinary bladder polyp † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Reproductive system and breast disorders				
Benign prostatic hyperplasia † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Cervical dysplasia † <sup>A</sup>				
# participants affected/at	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
risk				
# events				
Respiratory, thoracic and mediastinal disorders				
Acute respiratory failure † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Chronic obstructive pulmonary disease † <sup>A</sup>				
# participants affected/at risk	25/409 (6.11%)	26/412 (6.31%)	29/403 (7.2%)	23/409 (5.62%)
# events				
Chronic respiratory failure † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Dyspnoea † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	1/403 (0.25%)	1/409 (0.24%)
# events				
Haemoptysis † <sup>A</sup>				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Hypoxia † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Pleural effusion † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Pneumothorax † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	2/412 (0.49%)	1/403 (0.25%)	1/409 (0.24%)
# events				
Pulmonary bulla † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	1/412 (0.24%)	0/403 (0%)	0/409 (0%)
# events				
Pulmonary hypertension † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Pulmonary oedema † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Respiratory distress † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Respiratory failure † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	2/412 (0.49%)	0/403 (0%)	2/409 (0.49%)
# events				
Skin and subcutaneous tissue disorders				
Angioedema † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Urticaria † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Vascular disorders				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Aortic aneurysm † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Arteriosclerosis † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Circulatory collapse † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Deep vein thrombosis † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	0/403 (0%)	1/409 (0.24%)
# events				
Haemorrhage † <sup>A</sup>				
# participants affected/at risk	0/409 (0%)	0/412 (0%)	1/403 (0.25%)	0/409 (0%)
# events				
Labile blood pressure † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Peripheral arterial occlusive disease † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				
Thrombophlebitis † <sup>A</sup>				
# participants affected/at risk	1/409 (0.24%)	0/412 (0%)	0/403 (0%)	0/409 (0%)
# events				

† Indicates events were collected by systematic assessment.

A Term from vocabulary, MedDRA

## Other Adverse Events

Frequency Threshold Above Which Other Adverse Events are Reported: 3%

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
Total # participants affected/at risk	211/409 (51.59%)	229/412 (55.58%)	239/403 (59.31%)	249/409 (60.88%)
Gastrointestinal disorders				
Diarrhoea † <sup>A</sup>				
# participants affected/at risk	4/409 (0.98%)	10/412 (2.43%)	14/403 (3.47%)	15/409 (3.67%)
# events				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
General disorders				
Oedema peripheral † <sup>A</sup>				
# participants affected/at risk	16/409 (3.91%)	12/412 (2.91%)	12/403 (2.98%)	8/409 (1.96%)
# events				
Infections and infestations				
Bronchitis † <sup>A</sup>				
# participants affected/at risk	20/409 (4.89%)	25/412 (6.07%)	16/403 (3.97%)	23/409 (5.62%)
# events				
Influenza † <sup>A</sup>				
# participants affected/at risk	6/409 (1.47%)	18/412 (4.37%)	14/403 (3.47%)	18/409 (4.4%)
# events				
Lower respiratory tract infection † <sup>A</sup>				
# participants affected/at risk	13/409 (3.18%)	7/412 (1.7%)	12/403 (2.98%)	5/409 (1.22%)
# events				
Nasopharyngitis † <sup>A</sup>				
# participants affected/at risk	58/409 (14.18%)	54/412 (13.11%)	68/403 (16.87%)	82/409 (20.05%)

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# events				
Oral candidiasis † <sup>A</sup>				
# participants affected/at risk	29/409 (7.09%)	39/412 (9.47%)	39/403 (9.68%)	40/409 (9.78%)
# events				
Oropharyngeal candidiasis † A				
# participants affected/at risk	3/409 (0.73%)	15/412 (3.64%)	11/403 (2.73%)	5/409 (1.22%)
# events				
Pharyngitis † <sup>A</sup>				
# participants affected/at risk	12/409 (2.93%)	10/412 (2.43%)	10/403 (2.48%)	13/409 (3.18%)
# events				
Pneumonia † <sup>A</sup>				
# participants affected/at risk	6/409 (1.47%)	8/412 (1.94%)	14/403 (3.47%)	11/409 (2.69%)
# events				
Rhinitis † <sup>A</sup>				
# participants affected/at risk	12/409 (2.93%)	14/412 (3.4%)	5/403 (1.24%)	10/409 (2.44%)
# events				
Sinusitis † <sup>A</sup>				

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
# participants affected/at risk	19/409 (4.65%)	26/412 (6.31%)	20/403 (4.96%)	27/409 (6.6%)
# events				
Upper respiratory tract infection † <sup>A</sup>				
# participants affected/at risk	31/409 (7.58%)	37/412 (8.98%)	39/403 (9.68%)	36/409 (8.8%)
# events				
Musculoskeletal and connective tissue disorders				
Arthralgia † <sup>A</sup>				
# participants affected/at risk	17/409 (4.16%)	6/412 (1.46%)	20/403 (4.96%)	13/409 (3.18%)
# events				
Back pain † <sup>A</sup>				
# participants affected/at risk	23/409 (5.62%)	19/412 (4.61%)	30/403 (7.44%)	17/409 (4.16%)
# events				
Nervous system disorders				
Headache † <sup>A</sup>				
# participants affected/at	30/409	34/412	32/403	33/409

	VI 25 µg QD	FF/VI 50/25 µg QD	FF/VI 100/25 µg QD	FF/VI 200/25 µg QD
risk	(7.33%)	(8.25%)	(7.94%)	(8.07%)
# events				
Respiratory, thoracic and mediastinal disorders				
Cough † <sup>A</sup>				
# participants affected/at risk	20/409 (4.89%)	14/412 (3.4%)	15/403 (3.72%)	22/409 (5.38%)
# events				
Dyspnoea † <sup>A</sup>				
# participants affected/at risk	16/409 (3.91%)	12/412 (2.91%)	13/403 (3.23%)	10/409 (2.44%)
# events				
Oropharyngeal pain † <sup>A</sup>				
# participants affected/at risk	18/409 (4.4%)	23/412 (5.58%)	17/403 (4.22%)	26/409 (6.36%)
# events				
Vascular disorders				
Hypertension † <sup>A</sup>				
# participants affected/at risk	16/409 (3.91%)	12/412 (2.91%)	10/403 (2.48%)	15/409 (3.67%)
# events				

† Indicates events were collected by systematic assessment.

## More Information

### Certain Agreements:

Principal Investigators are NOT employed by the organization sponsoring the study.

There IS an agreement between the Principal Investigator and the Sponsor (or its agents) that restricts the PI's rights to discuss or publish trial results after the trial is completed.

GSK agreements may vary with individual investigators, but will not prohibit any investigator from publishing. GSK supports the publication of results from all centers of a multi-center trial but requests that reports based on single-site data not precede the primary publication of the entire clinical trial.

### Limitations and Caveats:

### Results Point of Contact:

Name/Official Title: GSK Response Center

Organization: GlaxoSmithKline

Phone: 866-435-7343

Email: