

**Table 18. MOCART scores at Visits 2 and 4–9 (ITT1 population)**

	Visit 2 (3 months)		Visit 4 (12 months)	
	Treatment group		Treatment group	
	ACT3D-CS <i>N</i> = 31	Microfracture <i>N</i> = 31	ACT3D-CS <i>N</i> = 33	Microfracture <i>N</i> = 36
Mean ± SD	65.6 ± 15.5	61.9 ± 10.6	78.9 ± 14.0	76.1 ± 13.2
Minimum	30	45	50	45
Lower quartile	55	55	70	65
Median	70	60	85	75
Upper quartile	75	65	90	85
Maximum	95	95	100	100
	Visit 5 (18 months)		Visit 6 (24 months)	
	ACT3D-CS <i>N</i> = 42	Microfracture <i>N</i> = 41	ACT3D-CS <i>N</i> = 46	Microfracture <i>N</i> = 43
Mean ± SD	76.8 ± 13.8	76.2 ± 13.5	76.3 ± 15.6	78.8 ± 12.7
Minimum	35	55	25	50
Lower quartile	65	65	70	70
Median	80	75	80	80
Upper quartile	85	85	85	85
Maximum	100	100	100	100
	Visit 7 (36 months)		Visit 8 (48 months)	
	ACT3D-CS <i>N</i> = 47	Microfracture <i>N</i> = 43	ACT3D-CS <i>N</i> = 47	Microfracture <i>N</i> = 43
Mean ± SD	75.0 ± 13.4	78.8 ± 12.8	76.4 ± 13.1	77.3 ± 12.6
Minimum	45	50	40	55
Lower quartile	65	70	70	65
Median	75	75	80	75
Upper quartile	85	90	85	90
Maximum	100	100	100	100
	Visit 9 (60 months)			
	ACT3D-CS <i>N</i> = 47	Microfracture <i>N</i> = 43		
Mean ± SD	71.4 ± 14.8	76.7 ± 14.9		
Minimum	40	40		
Lower quartile	60	65		
Median	75	75		
Upper quartile	85	85		
Maximum	95	100		

Differences from total population sizes are due to missing results. Source: Table 14.2.3.7.1.