

**Clinical trial results:**  
**An Open-Label Study of the Efficacy of Atomoxetine Hydrochloride on Quality of Life of Children and Adolescents with Attention-Deficit/Hyperactivity Disorder with or without comorbid conditions****Summary**

EudraCT number	2005-005701-32
Trial protocol	IT
Global end of trial date	06 October 2008

**Results information**

Result version number	v1 (current)
This version publication date	
First version publication date	

**Trial information****Trial identification**

Sponsor protocol code	B4Z-IT-LYDS
-----------------------	-------------

**Additional study identifiers**

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT00320528
WHO universal trial number (UTN)	-
Other trial identifiers	Trial Number: 9867, Trial Alias: B4Z-IT-LYDS

Notes:

**Sponsors**

Sponsor organisation name	Eli Lilly and Company
Sponsor organisation address	Lilly Corporate Center, Indianapolis, IN, United States, 46285
Public contact	Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877-CTLILLY,
Scientific contact	Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877-285-4559,

Notes:

**Paediatric regulatory details**

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

**Results analysis stage**

Analysis stage	Final
----------------	-------

Date of interim/final analysis	06 October 2008
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	06 October 2008
Was the trial ended prematurely?	No

Notes:

### General information about the trial

Main objective of the trial:

This study aims to assess the effectiveness of atomoxetine on psychosocial functioning and emotional well being of children and adolescents with Attention-Deficit/Hyperactivity Disorder (ADHD) and to evaluate whether and in what measure the presence of comorbid conditions (internalizing and externalizing disorders) influences atomoxetine's ability to improve the quality of life of ADHD subjects.

Protection of trial subjects:

This study was conducted in accordance with International Conference on Harmonisation (ICH) Good Clinical Practice, and the principles of the Declaration of Helsinki, in addition to following the laws and regulations of the country or countries in which a study is conducted.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	29 April 2006
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

### Population of trial subjects

#### Subjects enrolled per country

Country: Number of subjects enrolled	Italy: 269
Worldwide total number of subjects	269
EEA total number of subjects	269

Notes:

#### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	187
Adolescents (12-17 years)	82
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

No text entered.

### Pre-assignment

Screening details:

Study Period I was a 3-day screening period. Study Period II was 12 weeks long. Study Period III was an optional additional 12 week open-label extension. Results are presented for Period II (patients who received at least one dose of study drug).

### Period 1

Period 1 title	Study Period II
Is this the baseline period?	Yes
Allocation method	Non-randomised - controlled
Blinding used	Not blinded

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Pure ADHD (Period II)

Arm description:

Attention-Deficit/Hyperactivity Disorder (ADHD) alone. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

Arm type	Experimental
Investigational medicinal product name	atomoxetine
Investigational medicinal product code	
Other name	LY139603, Strattera
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

<b>Arm title</b>	ADHD+Internalizing Disorders (Period II)
------------------	--

Arm description:

Attention-Deficit/Hyperactivity Disorder (ADHD) plus internalizing disorders. Received atomoxetine: ... more 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeksless

Arm type	Experimental
Investigational medicinal product name	atomoxetine
Investigational medicinal product code	
Other name	LY139603, Strattera
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

<b>Arm title</b>	ADHD+Externalizing Disorders (Period II)
------------------	--

Arm description:

Attention-Deficit/Hyperactivity Disorder (ADHD) plus internalizing disorders. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

Arm type	Experimental
----------	--------------

Investigational medicinal product name	atomoxetine
Investigational medicinal product code	
Other name	LY139603, Strattera
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

Number of subjects in period 1	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)
Started	98	41	130
Received at Least One Dose of Study Drug	97	38	128
Completed	87	31	108
Not completed	11	10	22
Entry Criteria Exclusion	-	2	2
Adverse event, non-fatal	3	-	-
Consent withdrawn by subject	-	1	-
Parent/Caregiver Decision	8	-	-

## Period 2

Period 2 title	Study Period III
Is this the baseline period?	No
Allocation method	Non-randomised - controlled
Blinding used	Not blinded

## Arms

Are arms mutually exclusive?	No
<b>Arm title</b>	Pure ADHD (Period III)

Arm description:

Attention-Deficit/Hyperactivity Disorder (ADHD) alone. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

Arm type	Experimental
Investigational medicinal product name	atomoxetine
Investigational medicinal product code	
Other name	LY139603, Strattera
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

<b>Arm title</b>	ADHD+Internalizing Disorders (Period III)
------------------	---

Arm description:

Arm type	Experimental
Investigational medicinal product name	atomoxetine
Investigational medicinal product code	
Other name	LY139603, Strattera
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

<b>Arm title</b>	ADHD+Externalizing Disorders (Period III)
------------------	---

Arm description:

Attention-Deficit/Hyperactivity Disorder (ADHD) plus externalizing disorders. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

Arm type	Experimental
Investigational medicinal product name	atomoxetine
Investigational medicinal product code	
Other name	LY139603, Strattera
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

<b>Number of subjects in period 2</b>	Pure ADHD (Period III)	ADHD+Internalizing Disorders (Period III)	ADHD+Externalizing Disorders (Period III)
Started	87	31	108
Completed	0	0	2
Not completed	87	31	106
Sponsor Decision	-	-	1
Lack of efficacy	2	-	-
Withdrawal by Subject	-	1	-
Withdrawal by Subject	-	-	2
Atomoxetine Commercially Available	68	-	-
Consent withdrawn by subject	1	-	-
Atomoxetine Commercially Available	-	-	90
Parent/Caregiver Decision	14	3	11
Physician decision	-	1	2
Protocol Violation	-	1	-
Atomoxetine Commercially Available	-	25	-
Adverse event, non-fatal	1	-	-
Lost to follow-up	1	-	-



## Baseline characteristics

### Reporting groups

Reporting group title	Pure ADHD (Period II)
-----------------------	-----------------------

Reporting group description:

Attention-Deficit/Hyperactivity Disorder (ADHD) alone. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

Reporting group title	ADHD+Internalizing Disorders (Period II)
-----------------------	--

Reporting group description:

Attention-Deficit/Hyperactivity Disorder (ADHD) plus internalizing disorders. Received atomoxetine: ... more 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeksless

Reporting group title	ADHD+Externalizing Disorders (Period II)
-----------------------	--

Reporting group description:

Attention-Deficit/Hyperactivity Disorder (ADHD) plus internalizing disorders. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

Reporting group values	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)
Number of subjects	98	41	130
Age categorical Units: Subjects			

Age continuous Units: years			
arithmetic mean	10.12	10.42	10
standard deviation	± 2.51	± 2.41	± 2.57
Gender categorical Units: Subjects			
Female	14	3	9
Male	83	35	119
Not Recorded	1	3	2
Region of Enrollment Units: Subjects			
Italy	98	41	130
Race/Ethnicity Units: Subjects			
Caucasian	90	36	124
Hispanic	1	2	2
African	3	0	0
Native American	1	0	2
East Asian	2	0	0
Not Recorded	1	3	2
Child Health and Illness Profile - Child Edition (CHIP-CE) - Achievement Domain (N=96, N=33, N=121)			

Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to

67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean	29.27	27.72	25.83
standard deviation	± 7.89	± 7.73	± 7.6
Child Health and Illness Profile - Child Edition (CHIP-CE) - Satisfaction Domain (N=96, N=24, N=122)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean	39.26	30.11	31.69
standard deviation	± 13.89	± 14.83	± 14.78
Child Health and Illness Profile - Child Edition (CHIP-CE) - Comfort Domain (N=96, N=35, N=122)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean	51.31	46.76	44.74
standard deviation	± 9.4	± 12.28	± 9.38
Child Health and Illness Profile - Child Edition (CHIP-CE) - Resilience Domain (N=96, N=34, N=123)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean	39.34	34.96	31.4
standard deviation	± 10.15	± 10.89	± 11.85
Child Health and Illness Profile - Child Edition (CHIP-CE) - Risk Avoidance (N=96, N=33, N=121)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean	32.03	30.48	22.85
standard deviation	± 9.7	± 12.29	± 12
Children's Depression Rating Scale-Revised (CDRS-R)			
Measures presence and severity of depression. Consists of 17 items scored on a 1-5 or 1-7 scale. A rating of 1 indicates normal, thus the minimum score is 17. The maximum score is 113. In general, scores below 20 indicate an absence of depression; scores of 20 or 30 indicate borderline depression; scores of 40 to 60 indicate moderate depression.			
Units: units on a scale			
arithmetic mean	37.93	43.76	43.06
standard deviation	± 12.26	± 13.48	± 13.69

Clinical Global Impression-Attention Deficit Hyperactivity Disorder-Severity Scale (CGI-ADHD-S)			
Measures severity of the patient's overall severity of ADHD symptoms (1=normal, not at all ill; 7=among the most extremely ill patients).			
Units: units on a scale			
arithmetic mean	4.64	4.97	4.98
standard deviation	± 0.79	± 0.79	± 0.78
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S)			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale			
arithmetic mean	5.97	4.97	8.62
standard deviation	± 4.93	± 4.66	± 4.83
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S) Cognitive Problems			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale			
arithmetic mean	7.86	7.68	7.96
standard deviation	± 3.73	± 3.8	± 3.61
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S)			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale			
arithmetic mean	11.38	12.03	14.67
standard deviation	± 6.09	± 5.21	± 5.15
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S) ADHD			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale			
arithmetic mean	22.14	22.94	25.18
standard deviation	± 6.86	± 6.01	± 6.07
Pediatric Anxiety Rating Scale (PARS)			
The Pediatric Anxiety Rating Scale (PARS) is used to rate the severity of anxiety in children and adolescents, ages 6 to 17 years. The total score for the PARS is derived by summing 5 of the 7 severity/impairment/interference items (2,3,5,6,7). The total score ranges from 0 (none) to 25 (extreme severity). Items 1 (overall number of anxiety symptoms) and 4 (overall severity of physical symptoms) are not included in the total score calculation.			
Units: units on a scale			
arithmetic mean	2.5	7.5	4.67
standard deviation	± 2.67	± 5.68	± 5.12
Swanson, Nolan, and Pelham Rating Scale Revised (SNAP-IV) Inattention Subscale (N=97, N=38, N=127)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscore range=0-27), and Oppositional (items 19-26: subscore range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale			

arithmetic mean	20.28	21.03	21.87
standard deviation	± 4.19	± 3.72	± 3.48
SNAP-IV Hyperactivity/Impulsivity Subscale (N=97, N=38)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscale range=0-27), and Oppositional (items 19-26: subscale range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale			
arithmetic mean	17.21	19.66	21.04
standard deviation	± 6.1	± 5.54	± 3.48
Swanson, Nolan, and Pelham Rating Scale Revised (SNAP-IV) Combined Type Subscale (N=97, N=38, N=127)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscale range=0-27), and Oppositional (items 19-26: subscale range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale			
arithmetic mean	37.48	40.68	42.91
standard deviation	± 7.78	± 7.87	± 5.51
Swanson, Nolan, and Pelham Rating Scale Revised (SNAP-IV) Oppositional Subscale (N=97, N=38, N=127)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscale range=0-27), and Oppositional (items 19-26: subscale range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale			
arithmetic mean	10.15	12.32	17.99
standard deviation	± 4.54	± 4.67	± 3.57

<b>Reporting group values</b>	Total		
Number of subjects	269		
Age categorical			
Units: Subjects			

Age continuous			
Units: years			
arithmetic mean			
standard deviation	-		
Gender categorical			
Units: Subjects			
Female	26		
Male	237		
Not Recorded	6		
Region of Enrollment			
Units: Subjects			
Italy	269		
Race/Ethnicity			
Units: Subjects			

Caucasian	250		
Hispanic	5		
African	3		
Native American	3		
East Asian	2		
Not Recorded	6		
Child Health and Illness Profile - Child Edition (CHIP-CE) - Achievement Domain (N=96, N=33, N=121)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean			
standard deviation	-		
Child Health and Illness Profile - Child Edition (CHIP-CE) - Satisfaction Domain (N=96, N=24, N=122)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean			
standard deviation	-		
Child Health and Illness Profile - Child Edition (CHIP-CE) - Comfort Domain (N=96, N=35, N=122)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean			
standard deviation	-		
Child Health and Illness Profile - Child Edition (CHIP-CE) - Resilience Domain (N=96, N=34, N=123)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score			
arithmetic mean			
standard deviation	-		
Child Health and Illness Profile - Child Edition (CHIP-CE) - Risk Avoidance (N=96, N=33, N=121)			
Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of			

10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.			
Units: T-Score arithmetic mean standard deviation			
Children's Depression Rating Scale-Revised (CDRS-R)			
Measures presence and severity of depression. Consists of 17 items scored on a 1-5 or 1-7 scale. A rating of 1 indicates normal, thus the minimum score is 17. The maximum score is 113. In general, scores below 20 indicate an absence of depression; scores of 20 or 30 indicate borderline depression; scores of 40 to 60 indicate moderate depression.			
Units: units on a scale arithmetic mean standard deviation			
Clinical Global Impression-Attention Deficit Hyperactivity Disorder-Severity Scale (CGI-ADHD-S)			
Measures severity of the patient's overall severity of ADHD symptoms (1=normal, not at all ill; 7=among the most extremely ill patients).			
Units: units on a scale arithmetic mean standard deviation			
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S)			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale arithmetic mean standard deviation			
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S) Cognitive Problems			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale arithmetic mean standard deviation			
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S)			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale arithmetic mean standard deviation			
Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S) ADHD			
A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.			
Units: units on a scale arithmetic mean standard deviation			
Pediatric Anxiety Rating Scale (PARS)			
The Pediatric Anxiety Rating Scale (PARS) is used to rate the severity of anxiety in children and adolescents, ages 6 to 17 years. The total score for the PARS is derived by summing 5 of the 7 severity/impairment/interference items (2,3,5,6,7). The total score ranges from 0 (none) to 25			

(extreme severity). Items 1 (overall number of anxiety symptoms) and 4 (overall severity of physical symptoms) are not included in the total score calculation.			
Units: units on a scale arithmetic mean standard deviation			
Swanson, Nolan, and Pelham Rating Scale Revised (SNAP-IV) Inattention Subscale (N=97, N=38, N=127)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscale range=0-27), and Oppositional (items 19-26: subscale range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale arithmetic mean standard deviation			
SNAP-IV Hyperactivity/Impulsivity Subscale (N=97, N=38)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscale range=0-27), and Oppositional (items 19-26: subscale range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale arithmetic mean standard deviation			
Swanson, Nolan, and Pelham Rating Scale Revised (SNAP-IV) Combined Type Subscale (N=97, N=38, N=127)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscale range=0-27), and Oppositional (items 19-26: subscale range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale arithmetic mean standard deviation			
Swanson, Nolan, and Pelham Rating Scale Revised (SNAP-IV) Oppositional Subscale (N=97, N=38, N=127)			
The SNAP-IV is a 26-item scale that includes 1 item for each of the 18 symptoms contained in the DSM-IV diagnosis of ADHD and 1 item for each of the 8 symptoms contained in the DSM-IV diagnosis of ODD. Each item is score on 0 to 3 scale (0 = "Not at All", 1 = "Just a Little", 2 = "Pretty Much", 3 = "Very Much"). The SNAP-IV yields scores in three domains: Inattention (items 1-9: subscore range=0-27), Hyperactivity/impulsivity (items 10-18: subscale range=0-27), and Oppositional (items 19-26: subscale range=0-24). Combined type (inattention + hyperactivity/impulsivity) scores range from 0-54.			
Units: units on a scale arithmetic mean standard deviation			

## End points

### End points reporting groups

Reporting group title	Pure ADHD (Period II)
Reporting group description:	Attention-Deficit/Hyperactivity Disorder (ADHD) alone. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks
Reporting group title	ADHD+Internalizing Disorders (Period II)
Reporting group description:	Attention-Deficit/Hyperactivity Disorder (ADHD) plus internalizing disorders. Received atomoxetine: ... more 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeksless
Reporting group title	ADHD+Externalizing Disorders (Period II)
Reporting group description:	Attention-Deficit/Hyperactivity Disorder (ADHD) plus internalizing disorders. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks
Reporting group title	Pure ADHD (Period III)
Reporting group description:	Attention-Deficit/Hyperactivity Disorder (ADHD) alone. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks
Reporting group title	ADHD+Internalizing Disorders (Period III)
Reporting group description:	Attention-Deficit/Hyperactivity Disorder (ADHD) plus internalizing disorders. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks
Reporting group title	ADHD+Externalizing Disorders (Period III)
Reporting group description:	Attention-Deficit/Hyperactivity Disorder (ADHD) plus externalizing disorders. Received atomoxetine: 0.5 milligrams per kilogram per day (mg/kg/day), by mouth (PO) for 1 week then 1.2 mg/kg/day, PO for 11 weeks followed by up to 1.4 mg/kg/day, PO for up to 12 additional weeks

### Primary: 1: Change From Baseline to 12 Week Endpoint in Child Health and Illness Profile - Child Edition (CHIP-CE), Achievement Domain

End point title	1: Change From Baseline to 12 Week Endpoint in Child Health and Illness Profile - Child Edition (CHIP-CE), Achievement Domain
End point description:	Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains and subdomains having a mean score of 50 and standard deviation of 10. Standard scores are expressed in standard deviation units. Achievement Domain Range = -3.1 to 67.7. Higher scores mean greater health or level of functioning in achievement.
End point type	Primary
End point timeframe:	Baseline, 12 Weeks

<b>End point values</b>	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	83 <sup>[1]</sup>	28 <sup>[2]</sup>	97 <sup>[3]</sup>	
Units: T-Score				
arithmetic mean (confidence interval 95%)	2.65 (0.93 to 4.37)	0.51 (-1.38 to 2.39)	3.89 (2.22 to 5.57)	

Notes:

[1] - Number of patients with baseline and at least one nonmissing post-baseline measurement.

[2] - Number of patients with baseline and at least one nonmissing post-baseline measurement.

[3] - Number of patients with baseline and at least one nonmissing post-baseline measurement.

### Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis for End Point 1
Comparison groups	Pure ADHD (Period II) v ADHD+Internalizing Disorders (Period II) v ADHD+Externalizing Disorders (Period II)
Number of subjects included in analysis	208
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.05
Method	LOCF analysis

### Secondary: 2: Change From Baseline to 12 Week Endpoint in the Attention-Deficit/Hyperactivity Disorder (ADHD) Subscales of 18-Item Swanson, Nolan and Pelham Rating Scale (SNAP-IV)

End point title	2: Change From Baseline to 12 Week Endpoint in the Attention-Deficit/Hyperactivity Disorder (ADHD) Subscales of 18-Item Swanson, Nolan and Pelham Rating Scale (SNAP-IV)
-----------------	--

End point description:

Items from the Diagnostic and Statistical Manual of Mental Disorders Fourth Edition (DSM-IV) criteria for ADHD are included for the two subsets of symptoms: inattention (items #1-#9: total score=0-27) and hyperactivity/impulsivity (items #11-#19: total score=0-27). The SNAP-IV is based on a 0 (not at all) to 3 (very much) rating scale. Total combined type (inattention plus hyperactivity/impulsivity) subscale scores range from 0 to 54.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 Weeks

<b>End point values</b>	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	90	33	111	
Units: units on a scale				
arithmetic mean (confidence interval 95%)				
Inattention Subscale	-5.88 (-7.25 to -5.13)	-5.52 (-7.34 to -3.69)	-7.58 (-8.69 to -6.46)	

Hyperactivity/Impulsivity Subscale	-6.19 (-13.9 to -10.3)	-5.67 (-7.41 to -3.92)	-8.08 (-9.09 to -7.07)
Combined Type (ADHD) Subscale	-12.1 (-6.97 to -4.78)	-11.2 (-14.4 to -7.92)	-15.7 (-17.6 to -13.7)

### Statistical analyses

No statistical analyses for this end point

### Secondary: 3: Change From Baseline to 12 Week Endpoint in Clinical Global Impressions-ADHD-Severity (CGI-ADHD-S)

End point title	3: Change From Baseline to 12 Week Endpoint in Clinical Global Impressions-ADHD-Severity (CGI-ADHD-S)
-----------------	---

End point description:

Measures severity of the patient's overall severity of ADHD symptoms (1=normal, not at all ill; 7=among the most extremely ill patients).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 Weeks

End point values	Pure ADHD (Period II)	ADHD+Externalizing Disorders (Period II)	ADHD+Internalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	88	112	32	
Units: units on a scale				
arithmetic mean (confidence interval 95%)	-1.23 (-1.46 to -0.99)	-1.42 (-1.63 to -1.21)	-1.63 (-1.99 to -1.26)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: 4: Change From Baseline to 12 Week Endpoint in CHIP-CE Satisfaction, Comfort, Resilience and Risk Avoidance Domains

End point title	4: Change From Baseline to 12 Week Endpoint in CHIP-CE Satisfaction, Comfort, Resilience and Risk Avoidance Domains
-----------------	---

End point description:

Parent-rated assessment of a child's health status and level of functioning. It consists of 76 items. The majority of items assess frequency of activities or feelings using a five-point response format (for example, 'how good is your child at making friends?' 1=never, 5=always). Standard scores (t-value) were established, with all domains having a mean score of 50 and standard deviation of 10. Satisfaction range=-25.7 to 66.3; Comfort=-28.6 to 67.2; Resilience=-36.3 to 71.8; Risk Avoidance=-23.5 to 62.5. Higher scores mean greater health or level of functioning in that domain.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 Weeks

<b>End point values</b>	Pure ADHD (Period II)	ADHD+Externalizing Disorders (Period II)	ADHD+Internalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	97	128	38	
Units: T-Score				
arithmetic mean (confidence interval 95%)				
Satisfaction Domain (N=88, N=31, N=105)	2.14 (-0.06 to 4.34)	3.78 (1.18 to 6.39)	0.81 (-3.33 to 4.95)	
Comfort Domain (N=86, N=28, N=103)	1.01 (-0.88 to 2.9)	4.94 (3.14 to 6.73)	5.72 (1.24 to 10.19)	
Resilience Domain (N=87, N=30, N=106)	1.21 (-0.85 to 3.27)	3.07 (1.1 to 5.03)	2.18 (-1.31 to 5.67)	
Risk Avoidance Domain (N=82, N=28, N=94)	4.72 (2.93 to 6.51)	9.88 (7.88 to 11.88)	5.7 (1.96 to 9.45)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: 5: Change From Baseline to 12 Week Endpoint in Pediatric Anxiety Rating Scale (PARS)

End point title	5: Change From Baseline to 12 Week Endpoint in Pediatric Anxiety Rating Scale (PARS)
End point description:	
The Pediatric Anxiety Rating Scale (PARS) is used to rate the severity of anxiety in children and adolescents, ages 6 to 17 years. The total score for the PARS is derived by summing 5 of the 7 severity/impairment/interference items (2,3,5,6,7). The total score ranges from 0 (none) to 25 (extreme severity). Items 1 (overall number of anxiety symptoms) and 4 (overall severity of physical symptoms) are not included in the total score calculation.	
End point type	Secondary
End point timeframe:	
Baseline, 12 Weeks	

<b>End point values</b>	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	89	33	112	
Units: units on a scale				
arithmetic mean (confidence interval 95%)	-0.18 (-0.7 to 0.34)	-1.88 (-1.9 to 0.46)	-1.18 (-1.9 to 0.46)	

## Statistical analyses

No statistical analyses for this end point

### Secondary: 6: Change From Baseline to 12 Week Endpoint in Children's Depression Rating Scale-Revised (CDRS-R)

End point title	6: Change From Baseline to 12 Week Endpoint in Children's Depression Rating Scale-Revised (CDRS-R)
-----------------	--

End point description:

Measures presence and severity of depression. Consists of 17 items scored on a 1-5 or 1-7 scale. A rating of 1 indicates normal, thus the minimum score is 17. The maximum score is 113. In general, scores below 20 indicate an absence of depression; scores of 20 or 30 indicate borderline depression; scores of 40 to 60 indicate moderate depression.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 Weeks

End point values	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	97	38	128	
Units: units on a scale				
arithmetic mean (confidence interval 95%)	-1.6 (-3.54 to 0.34)	-2.56 (-6.38 to 1.25)	-3.13 (-4.64 to -1.62)	

## Statistical analyses

No statistical analyses for this end point

### Secondary: 7: Change From Baseline to 12 Week Endpoint in SNAP-IV Oppositional Scale

End point title	7: Change From Baseline to 12 Week Endpoint in SNAP-IV Oppositional Scale
-----------------	---

End point description:

Items are included from the Diagnostic and Statistical Manual of Mental Disorders Fourth Edition (DSM-IV) criteria for Oppositional Defiant Disorder. The SNAP-IV is based on a 0 (not at all) to 3 (very much) rating scale. Total subscale scores range from 0 to 24.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 Weeks

End point values	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	90	33	111	
Units: units on a scale				
arithmetic mean (confidence interval 95%)	-2.01 (-2.86 to -1.16)	-2.12 (-3.95 to -0.29)	-5.54 (-6.43 to -4.65)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: 8: Change From Baseline to 12 Week Endpoint in Adolescent Symptom Inventory-4: Parent Checklist (ASI-4)

End point title	8: Change From Baseline to 12 Week Endpoint in Adolescent Symptom Inventory-4: Parent Checklist (ASI-4)
-----------------	---

End point description:

Parent-completed ASI-4 contains 120 items on 18 emotional and behavioral disorders in adolescents (12-18 years old). Item score range:0 (no symptoms) to 3 (maximum impairment). Categories: A=ADHD (0-54); B=Conduct (0-60); C=Oppositional Defiant (0-24); D=Generalized Anxiety (0-18); E=Specific Phobia/Panic Attack/Obsessions/Compulsions/Somatization (0-30); F=Social Phobia (0-6); G=Separation Anxiety (0-24); H=Schizoid Personality (0-9); I=Schizophrenia (0-6); J=Enuresis (0-18); K=Major Depressive (0-42); L=Bipolar (0-27); M=Anorexia (0-12); N=Bulimia (0-12); O=Substance Abuse (0-18).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 Weeks

End point values	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	26	10	30	
Units: units on a scale				
arithmetic mean (standard deviation)				
Category A - Baseline	32.73 (± 11.42)	38.33 (± 10.32)	38.13 (± 8.3)	
Category A - Change from Baseline	-11.4 (± 9.88)	-10.4 (± 11.55)	-11.6 (± 9.5)	
Category B - Baseline	4.04 (± 4.25)	6 (± 5.62)	10.8 (± 5.41)	
Category B - Change from Baseline	-0.73 (± 2.92)	-0.6 (± 5.83)	-5.57 (± 5.58)	
Category C - Baseline	7.92 (± 4.77)	10.5 (± 3.72)	15.6 (± 5.24)	
Category C - Change from Baseline	-0.88 (± 3.85)	-1.4 (± 6.15)	-5.47 (± 5.51)	
Category D - Baseline	4.73 (± 3.14)	9.2 (± 3.58)	8.37 (± 3.54)	
Category D - Change from Baseline	-0.58 (± 3.09)	-1.7 (± 3.65)	-2.97 (± 3.67)	
Category E - Baseline	3.58 (± 4)	7.6 (± 7.85)	3.63 (± 3.16)	
Category E - Change from Baseline	-0.38 (± 4.01)	-4.5 (± 8.72)	-1.3 (± 2.35)	
Category F - Baseline	1.19 (± 1.27)	2.6 (± 1.43)	1.2 (± 1.37)	
Category F - Change from Baseline	-0.58 (± 0.81)	-1.4 (± 1.43)	-0.53 (± 1.25)	
Category G - Baseline	2.27 (± 3.5)	4.5 (± 5.25)	2.1 (± 2.63)	

Category G - Change from Baseline	-0.46 (± 2.49)	-3.2 (± 4.71)	-0.97 (± 2.4)
Category H- Baseline Category H -	0.73 (± 1.28)	1.6 (± 2.27)	1.7 (± 1.74)
Change from Baseline Category I- Baseline	-0.38 (± 1.2)	0.2 (± 2.86)	-0.67 (± 1.69)
Category I - Change from Baseline	1 (± 1.92)	3.2 (± 3.65)	2.13 (± 2.22)
Category J- Baseline Category J -	-0.19 (± 0.8)	-1.3 (± 3.89)	-0.93 (± 1.7)
Change from Baseline Category K- Baseline	0.42 (± 0.99)	0.4 (± 0.97)	0.5 (± 0.78)
Category K - Change from Baseline	0 (± 0.69)	-0.3 (± 0.67)	-0.23 (± 0.43)
Category L- Baseline Category L -	3.73 (± 3.99)	4.9 (± 2.85)	4.23 (± 4.05)
Change from Baseline Category M- Baseline	-1.04 (± 2.9)	-1 (± 3.5)	-1.6 (± 3.5)
Category M - Change from Baseline	3.54 (± 4.68)	5.1 (± 5)	6.03 (± 5.02)
Category N- Baseline Category N -	-1.08 (± 3.46)	-0.2 (± 6.03)	-1.97 (± 4.3)
Change from Baseline Category O- Baseline	0.73 (± 1.22)	1.3 (± 2.06)	0.5 (± 1.41)
Category O - Change from Baseline	0.15 (± 1.08)	-0.1 (± 3.21)	0.3 (± 1.39)
Category N- Baseline Category N -	1.31 (± 1.54)	2.2 (± 3.05)	2.3 (± 1.84)
Change from Baseline Category O- Baseline	-0.38 (± 1.7)	-0.9 (± 2.23)	-0.8 (± 1.49)
Category O - Change from Baseline	0.23 (± 0.71)	0.3 (± 0.95)	0.53 (± 0.97)
Category O - Change from Baseline	0.15 (± 0.37)	-0.2 (± 0.63)	-0.13 (± 0.51)

## Statistical analyses

No statistical analyses for this end point

### Secondary: 9: Change From Baseline to 12 Week Endpoint in Child Symptom Inventory-4: Parent Checklist (CSI-4)

End point title	9: Change From Baseline to 12 Week Endpoint in Child Symptom Inventory-4: Parent Checklist (CSI-4)
End point description:	The CSI-4 contains 97 items that screen for 15 emotional and behavioral disorders in children between 5 and 12 years old. Item score range:0 (no symptoms) to 3 (maximum impairment). Categories: A=ADHD (0-54); B=Conduct (0-60); C=Oppositional Defiant (0-24); D=Generalized Anxiety (0-18); E=Specific Phobia/Panic Attack/Obsessions/Compulsions/Somatization (0-30); F=Social Phobia (0-6); G=Separation Anxiety (0-24); H=Schizoid Personality (0-9); I=Schizophrenia (0-6); J=Enuresis (0-18).
End point type	Secondary
End point timeframe:	Baseline, 12 Weeks

End point values	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	58	22	78	
Units: units on a scale				
arithmetic mean (standard deviation)				
Category A - Baseline	34 (± 10.43)	36 (± 9.32)	38.95 (± 8.68)	
Category A - Change from Baseline	-10.6 (± 9.76)	-9.68 (± 9.54)	-12.7 (± 12.06)	
Category B - Baseline Category B -	9.29 (± 4.72)	11.05 (± 4.13)	15.65 (± 4.4)	
Change from Baseline Category C - Baseline	-2.12 (± 3.88)	-1.14 (± 5.25)	-3.76 (± 5.56)	
	2.1 (± 2.44)	2.23 (± 2.25)	4.63 (± 3.69)	

Category C - Change from Baseline	-0.88 (± 2.11)	-0.05 (± 3.06)	-1.99 (± 3.17)
Category D - Baseline Category D - Change from Baseline	5.83 (± 3.37)	8.23 (± 3.96)	7.41 (± 3.94)
Category E - Change from Baseline	-0.84 (± 3.16)	-2.23 (± 3.01)	-1.63 (± 3.86)
Category E - Baseline	2.07 (± 1.87)	2.09 (± 1.95)	2.79 (± 2.69)
Category E - Change from Baseline	-0.38 (± 1.9)	-0.41 (± 1.75)	-0.85 (± 2.43)
Category F - Baseline Category F - Change from Baseline	0.48 (± 0.86)	0.32 (± 0.57)	0.91 (± 1.35)
Category G - Change from Baseline	-0.17 (± 0.94)	-0.14 (± 0.64)	-0.24 (± 1.28)
Category G - Baseline	3.02 (± 2.7)	4.05 (± 2.79)	3.51 (± 3.3)
Category G - Change from Baseline	-0.62 (± 2.4)	-0.91 (± 2)	-0.46 (± 2.52)
Category H - Baseline Category H - Change from Baseline	3.66 (± 4.14)	4.64 (± 4.02)	4.45 (± 4.39)
Category I - Change from Baseline	-1.1 (± 2.21)	-1.09 (± 2.93)	-1.54 (± 4.31)
Category I - Baseline	2.76 (± 1.85)	2.77 (± 1.38)	2.94 (± 1.93)
Category I - Change from Baseline	-0.03 (± 1.83)	-0.45 (± 1.47)	-0.32 (± 1.75)
Category J - Baseline	3.92 (± 3.98)	5.58 (± 6.05)	5.22 (± 4.47)
Category J - Change from Baseline	-1.34 (± 3.64)	-0.83 (± 2.61)	-1.55 (± 4.34)

### Statistical analyses

No statistical analyses for this end point

### Secondary: 10: Change From Baseline to 12 Week Endpoint in Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S)

End point title	10: Change From Baseline to 12 Week Endpoint in Conners' Teacher Rating Scale-Revised: Short Form (CTRS-R:S)
-----------------	--

End point description:

A 28-item rating scale (0 [not at all/never] to 3 [very much true/very often]) completed by the teacher to assess problem behaviors related to ADHD. Subscale total scores range from 0 to 15 for Oppositional and Cognitive Problems, 0 to 21 for Hyperactivity, and 0 to 36 for ADHD Index.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 Weeks

End point values	Pure ADHD (Period II)	ADHD+Internalizing Disorders (Period II)	ADHD+Externalizing Disorders (Period II)	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	60	15	53	
Units: units on a scale				
arithmetic mean (confidence interval 95%)				
Oppositional	-1.42 (-2.22 to -0.61)	1.07 (-0.33 to 2.46)	-2.35 (-3.55 to -1.15)	
Cognitive Problems	-1.08 (-1.72 to -0.45)	-0.47 (-1.74 to 0.8)	-0.94 (-1.77 to -0.11)	
Hyperactivity	-2.87 (-4.04 to -1.69)	-2.34 (-4.8 to 0.11)	-4.09 (-5.52 to -2.67)	
ADHD Index	-4.96 (-6.36 to -3.52)	-3.6 (-6.65 to -0.55)	-6.62 (-8.59 to -4.65)	

## **Statistical analyses**

---

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Entire Study

Adverse event reporting additional description:

B4Z-IT-LYDS

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	12.0
--------------------	------

### Reporting groups

Reporting group title	Pure ADHD
-----------------------	-----------

Reporting group description: -

Reporting group title	ADHD plus internalizing disorders
-----------------------	-----------------------------------

Reporting group description: -

Reporting group title	ADHD plus externalizing disorders
-----------------------	-----------------------------------

Reporting group description: -

Serious adverse events	Pure ADHD	ADHD plus internalizing disorders	ADHD plus externalizing disorders
Total subjects affected by serious adverse events			
subjects affected / exposed	1 / 97 (1.03%)	0 / 38 (0.00%)	2 / 128 (1.56%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events	0	0	0
Injury, poisoning and procedural complications			
facial bones fracture			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	0 / 97 (0.00%)	0 / 38 (0.00%)	1 / 128 (0.78%)
occurrences casually related to treatment / all	0 / 0	0 / 0	0 / 1
deaths casually related to treatment / all	0 / 0	0 / 0	0 / 0
fall			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	0 / 97 (0.00%)	0 / 38 (0.00%)	1 / 128 (0.78%)
occurrences casually related to treatment / all	0 / 0	0 / 0	0 / 1
deaths casually related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
idiosyncratic drug reaction			

alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences casually related to treatment / all deaths casually related to treatment / all	0 / 97 (0.00%) 0 / 0 0 / 0	0 / 38 (0.00%) 0 / 0 0 / 0	1 / 128 (0.78%) 1 / 1 0 / 0
<b>Gastrointestinal disorders</b> vomiting alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences casually related to treatment / all deaths casually related to treatment / all	0 / 97 (0.00%) 0 / 0 0 / 0	0 / 38 (0.00%) 0 / 0 0 / 0	1 / 128 (0.78%) 1 / 1 0 / 0
<b>Metabolism and nutrition disorders</b> dehydration alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences casually related to treatment / all deaths casually related to treatment / all	0 / 97 (0.00%) 0 / 0 0 / 0	0 / 38 (0.00%) 0 / 0 0 / 0	1 / 128 (0.78%) 1 / 1 0 / 0
<b>Infections and infestations</b> appendicitis alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences casually related to treatment / all deaths casually related to treatment / all	1 / 97 (1.03%) 0 / 1 0 / 0	0 / 38 (0.00%) 0 / 0 0 / 0	0 / 128 (0.00%) 0 / 0 0 / 0

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	Pure ADHD	ADHD plus internalizing disorders	ADHD plus externalizing disorders
<b>Total subjects affected by non-serious            adverse events</b> subjects affected / exposed	70 / 97 (72.16%)	28 / 38 (73.68%)	87 / 128 (67.97%)
<b>Investigations</b> weight decreased alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)	11 / 97 (11.34%) 15	1 / 38 (2.63%) 1	12 / 128 (9.38%) 15
<b>Respiratory, thoracic and mediastinal disorders</b>			

cough alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)	3 / 97 (3.09%) 3	0 / 38 (0.00%) 0	8 / 128 (6.25%) 10
Nervous system disorders headache alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)  somnolence alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)	20 / 97 (20.62%) 34  22 / 97 (22.68%) 23	10 / 38 (26.32%) 17  11 / 38 (28.95%) 15	26 / 128 (20.31%) 35  34 / 128 (26.56%) 44
General disorders and administration site conditions asthenia alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)  fatigue alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)  irritability alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)  pyrexia alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)	5 / 97 (5.15%) 6  5 / 97 (5.15%) 5  4 / 97 (4.12%) 4  3 / 97 (3.09%) 3	6 / 38 (15.79%) 6  0 / 38 (0.00%) 0  6 / 38 (15.79%) 6  0 / 38 (0.00%) 0	7 / 128 (5.47%) 7  2 / 128 (1.56%) 3  12 / 128 (9.38%) 16  8 / 128 (6.25%) 8
Psychiatric disorders aggression alternative dictionary used: MedDRA 12.0 subjects affected / exposed occurrences (all)	4 / 97 (4.12%) 5	1 / 38 (2.63%) 1	9 / 128 (7.03%) 9

Gastrointestinal disorders			
abdominal pain			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	9 / 97 (9.28%)	8 / 38 (21.05%)	18 / 128 (14.06%)
occurrences (all)	13	9	32
abdominal pain upper			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	5 / 97 (5.15%)	6 / 38 (15.79%)	12 / 128 (9.38%)
occurrences (all)	5	8	17
constipation			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	2 / 97 (2.06%)	2 / 38 (5.26%)	1 / 128 (0.78%)
occurrences (all)	2	2	1
diarrhoea			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	1 / 97 (1.03%)	2 / 38 (5.26%)	3 / 128 (2.34%)
occurrences (all)	3	2	3
nausea			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	12 / 97 (12.37%)	10 / 38 (26.32%)	26 / 128 (20.31%)
occurrences (all)	17	19	36
vomiting			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	13 / 97 (13.40%)	6 / 38 (15.79%)	15 / 128 (11.72%)
occurrences (all)	23	13	20
Metabolism and nutrition disorders			
anorexia			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	24 / 97 (24.74%)	11 / 38 (28.95%)	33 / 128 (25.78%)
occurrences (all)	26	15	39
decreased appetite			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	11 / 97 (11.34%)	7 / 38 (18.42%)	14 / 128 (10.94%)
occurrences (all)	13	7	20
Infections and infestations			

gastroenteritis			
alternative dictionary used: MedDRA 12.0			
subjects affected / exposed	2 / 97 (2.06%)	2 / 38 (5.26%)	0 / 128 (0.00%)
occurrences (all)	2	2	0

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? No

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

The original intent was to also calculate change and 95% confidence intervals at end of Period III; however, due to low number of patients completing Period III, variations from baseline (LOCF), are not suitable for further analysis or description.

Notes: