

**Clinical trial results:**

A phase IV, open, multicentric study to evaluate the immune response to a hepatitis B challenge dose in healthy subjects, 72 to 78 months after they received a primary vaccination course of GSK Biologicals' Engerix-B (thiomersal-free 20 µg or preservative-free 10 µg) vaccine, in the primary study HBV-280

Summary

| | |
|--------------------------|----------------|
| EudraCT number | 2007-000261-38 |
| Trial protocol | BE |
| Global end of trial date | 14 May 2008 |

Results information

| | |
|--------------------------------|---|
| Result version number | v2 (current) |
| This version publication date | 13 April 2023 |
| First version publication date | 04 December 2014 |
| Version creation reason | <ul style="list-style-type: none">• Correction of full data set Correction of full data set and alignment between registries. |

Trial information**Trial identification**

| | |
|-----------------------|--------|
| Sponsor protocol code | 108988 |
|-----------------------|--------|

Additional study identifiers

| | |
|------------------------------------|-------------|
| ISRCTN number | - |
| ClinicalTrials.gov id (NCT number) | NCT00524576 |
| WHO universal trial number (UTN) | - |

Notes:

Sponsors

| | |
|------------------------------|--|
| Sponsor organisation name | GlaxoSmithKline Biologicals |
| Sponsor organisation address | Rue de l'Institut 89, Rixensart, Belgium, B-1330 |
| Public contact | Clinical Trials Call Center, GlaxoSmithKline Biologicals, 044 2089904466, GSKClinicalSupportHD@gsk.com |
| Scientific contact | Clinical Trials Call Center, GlaxoSmithKline Biologicals, 044 2089904466, GSKClinicalSupportHD@gsk.com |

Notes:

Paediatric regulatory details

| | |
|--|-----|
| Is trial part of an agreed paediatric investigation plan (PIP) | No |
| Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial? | No |
| Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial? | Yes |

Notes:

Results analysis stage

| | |
|--|-------------|
| Analysis stage | Final |
| Date of interim/final analysis | 14 May 2008 |
| Is this the analysis of the primary completion data? | Yes |
| Primary completion date | 14 May 2008 |
| Global end of trial reached? | Yes |
| Global end of trial date | 14 May 2008 |
| Was the trial ended prematurely? | No |

Notes:

General information about the trial

Main objective of the trial:

To assess the immune response to a challenge dose of hepatitis B vaccine administered in subjects who previously received a complete hepatitis B primary vaccination course, 72 to 78 months ago.

Protection of trial subjects:

As with all injectable vaccines, appropriate medical treatment was always readily available in case of anaphylactic reactions following the administration of the vaccine.

For this reason, the subjects remained under medical supervision for 30 minutes after vaccination.

Background therapy: -

Evidence for comparator: -

| | |
|---|------------------|
| Actual start date of recruitment | 28 November 2007 |
| Long term follow-up planned | No |
| Independent data monitoring committee (IDMC) involvement? | No |

Notes:

Population of trial subjects

Subjects enrolled per country

| | |
|--------------------------------------|---------------|
| Country: Number of subjects enrolled | Australia: 67 |
| Country: Number of subjects enrolled | Belgium: 77 |
| Worldwide total number of subjects | 144 |
| EEA total number of subjects | 77 |

Notes:

Subjects enrolled per age group

| | |
|---|----|
| In utero | 0 |
| Preterm newborn - gestational age < 37 wk | 0 |
| Newborns (0-27 days) | 0 |
| Infants and toddlers (28 days-23 months) | 0 |
| Children (2-11 years) | 0 |
| Adolescents (12-17 years) | 72 |
| Adults (18-64 years) | 72 |
| From 65 to 84 years | 0 |
| 85 years and over | 0 |

Subject disposition

Recruitment

Recruitment details: -

Pre-assignment

Screening details:

During the screening the following steps occurred: check for inclusion/exclusion criteria, contraindications/precautions, medical history of the subjects and signing informed consent forms.

Period 1

| | |
|------------------------------|--------------------------------|
| Period 1 title | Overall Study (overall period) |
| Is this the baseline period? | Yes |
| Allocation method | Non-randomised - controlled |
| Blinding used | Not blinded |

Arms

| | |
|------------------------------|-----|
| Are arms mutually exclusive? | Yes |
|------------------------------|-----|

| | |
|------------------|----------------------------------|
| Arm title | Engerix 2 doses + challenge dose |
|------------------|----------------------------------|

Arm description:

Subjects received 2 doses of Engerix-B (Month 0 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| | |
|--|-----------------------|
| Arm type | Experimental |
| Investigational medicinal product name | Biological: Engerix-B |
| Investigational medicinal product code | |
| Other name | |
| Pharmaceutical forms | Injection |
| Routes of administration | Intramuscular use |

Dosage and administration details:

Subjects received 2 doses of Engerix-B (Month 0 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| | |
|------------------|----------------------------------|
| Arm title | Engerix 3 doses + challenge dose |
|------------------|----------------------------------|

Arm description:

Subjects received 3 doses of Engerix-B (Month 0, 1 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| | |
|--|-------------------|
| Arm type | Experimental |
| Investigational medicinal product name | Engerix-B |
| Investigational medicinal product code | |
| Other name | |
| Pharmaceutical forms | Injection |
| Routes of administration | Intramuscular use |

Dosage and administration details:

Subjects received 3 doses of Engerix-B (Month 0, 1 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| Number of subjects in period 1 | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose |
|---------------------------------------|----------------------------------|----------------------------------|
| Started | 97 | 47 |
| Completed | 97 | 47 |

Baseline characteristics

Reporting groups

| | |
|-----------------------|----------------------------------|
| Reporting group title | Engerix 2 doses + challenge dose |
|-----------------------|----------------------------------|

Reporting group description:

Subjects received 2 doses of Engerix-B (Month 0 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| | |
|-----------------------|----------------------------------|
| Reporting group title | Engerix 3 doses + challenge dose |
|-----------------------|----------------------------------|

Reporting group description:

Subjects received 3 doses of Engerix-B (Month 0, 1 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| Reporting group values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | Total |
|--|----------------------------------|----------------------------------|-------|
| Number of subjects | 97 | 47 | 144 |
| Age categorical Units: Subjects | | | |
| In utero | | | 0 |
| Preterm newborn infants (gestational age < 37 wks) | | | 0 |
| Newborns (0-27 days) | | | 0 |
| Infants and toddlers (28 days-23 months) | | | 0 |
| Children (2-11 years) | | | 0 |
| Adolescents (12-17 years) | | | 0 |
| Adults (18-64 years) | | | 0 |
| From 65-84 years | | | 0 |
| 85 years and over | | | 0 |
| Age continuous Units: years | | | |
| arithmetic mean | 19.5 | 19.3 | |
| standard deviation | ± 1.22 | ± 1.46 | - |
| Gender categorical Units: Subjects | | | |
| Female | 50 | 23 | 73 |
| Male | 47 | 24 | 71 |

End points

End points reporting groups

| | |
|---|----------------------------------|
| Reporting group title | Engerix 2 doses + challenge dose |
| Reporting group description: Subjects received 2 doses of Engerix-B (Month 0 and 6) in the primary study and a single dose of Engerix-B during the booster study. | |
| Reporting group title | Engerix 3 doses + challenge dose |
| Reporting group description: Subjects received 3 doses of Engerix-B (Month 0, 1 and 6) in the primary study and a single dose of Engerix-B during the booster study. | |

Primary: Number of subjects with immunological response to challenge dose in terms of anti-hepatitis B surface antigen (anti-HBs) antibody concentration

| | |
|--|--|
| End point title | Number of subjects with immunological response to challenge dose in terms of anti-hepatitis B surface antigen (anti-HBs) antibody concentration ^[1] |
| End point description: Immune response defined as: *For initially seronegative subjects (anti-HBs antibody concentration <3.3 milli-international unit per milliliter [mIU/mL] before vaccination) antibody concentration ≥ 10mIU/mL at post booster. *For initially seropositive subjects: antibody concentration at post booster ≥ 4-fold the pre-vaccination antibody concentration. | |
| End point type | Primary |
| End point timeframe: 30 days post-challenge dose | |

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of the primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

| End point values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | | |
|-----------------------------------|----------------------------------|----------------------------------|--|--|
| Subject group type | Reporting group | Reporting group | | |
| Number of subjects analysed | 53 | 21 | | |
| Units: Subjects | | | | |
| (anti-HBs) antibody concentration | 53 | 21 | | |

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects with anti-HBs antibody concentrations above the cut-off value

| | |
|---|--|
| End point title | Number of subjects with anti-HBs antibody concentrations above the cut-off value |
| End point description: Anti-HBs antibody cut-off values assessed include 3.3, 10 and 100 mIU/mL. | |
| End point type | Secondary |

End point timeframe:
30 days post-challenge dose

| End point values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | | |
|-----------------------------|----------------------------------|----------------------------------|--|--|
| Subject group type | Reporting group | Reporting group | | |
| Number of subjects analysed | 53 | 21 | | |
| Units: Subjects | | | | |
| ≥ 3.3 mIU/mL | 53 | 21 | | |
| ≥ 10 mIU/mL | 53 | 21 | | |
| ≥ 100 mIU/mL | 50 | 20 | | |

Statistical analyses

No statistical analyses for this end point

Secondary: Concentration of anti-HBs antibodies

| | |
|------------------------|---|
| End point title | Concentration of anti-HBs antibodies |
| End point description: | Concentrations given as geometric mean concentration (GMC) and expressed in mIU/mL. |
| End point type | Secondary |
| End point timeframe: | 30 days post-challenge dose |

| End point values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | | |
|--|----------------------------------|----------------------------------|--|--|
| Subject group type | Reporting group | Reporting group | | |
| Number of subjects analysed | 53 | 21 | | |
| Units: mIU/mL | | | | |
| geometric mean (confidence interval 95%) | | | | |
| Concentration of anti-HBs antibodies | 6214.1 (3213.1 to 12018) | 16564.3 (6394.9 to 42905.6) | | |

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects reporting solicited local symptoms

| | |
|-----------------|---|
| End point title | Number of subjects reporting solicited local symptoms |
|-----------------|---|

End point description:

Solicited local symptoms assessed include pain, redness and swelling.

End point type Secondary

End point timeframe:

During the 4-day follow-up period (Day 0-3) after the challenge dose

| End point values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | | |
|-----------------------------|----------------------------------|----------------------------------|--|--|
| Subject group type | Reporting group | Reporting group | | |
| Number of subjects analysed | 55 | 22 | | |
| Units: Subjects | | | | |
| Pain | 22 | 4 | | |
| Redness | 11 | 1 | | |
| Swelling | 9 | 0 | | |

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects reporting solicited general symptoms

End point title Number of subjects reporting solicited general symptoms

End point description:

Solicited general symptoms assessed include fatigue, fever, gastrointestinal symptoms, and headache.

End point type Secondary

End point timeframe:

During the 4-day follow-up period (Day 0-3) after the challenge dose

| End point values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | | |
|----------------------------------|----------------------------------|----------------------------------|--|--|
| Subject group type | Reporting group | Reporting group | | |
| Number of subjects analysed | 55 | 22 | | |
| Units: Subjects | | | | |
| Fatigue | 19 | 7 | | |
| Fever \geq 37.5 degree Celsius | 1 | 0 | | |
| Gastrointestinal disorder | 7 | 4 | | |
| Headache | 14 | 4 | | |

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects reporting unsolicited adverse events (AE)

End point title | Number of subjects reporting unsolicited adverse events (AE)

End point description:

An AE was any untoward medical occurrence in a clinical investigation subject, temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product.

End point type | Secondary

End point timeframe:

During the 31-day follow-up period (Day 0-30) after the challenge dose

| End point values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | | |
|-----------------------------|----------------------------------|----------------------------------|--|--|
| Subject group type | Reporting group | Reporting group | | |
| Number of subjects analysed | 55 | 22 | | |
| Units: Subjects | | | | |
| Subjects reporting AE | 19 | 5 | | |

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects reporting serious adverse events (SAE)

End point title | Number of subjects reporting serious adverse events (SAE)

End point description:

An SAE was any untoward medical occurrence that: results in death, was life-threatening, required hospitalization or prolongation of existing hospitalization, resulted in disability/incapacity, was a congenital anomaly/birth defect in the offspring of a study subject, or evolved into one of the outcomes listed above.

End point type | Secondary

End point timeframe:

During the 31-day follow-up period (Day 0-30) after the challenge dose

| End point values | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | | |
|-----------------------------|----------------------------------|----------------------------------|--|--|
| Subject group type | Reporting group | Reporting group | | |
| Number of subjects analysed | 55 | 22 | | |
| Units: Subjects | | | | |
| Subjects reporting SAE | 0 | 0 | | |

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

Serious Adverse events (SAE) = Day 0 to Day 30. Solicited local and general symptoms = During the 4-day (Days 0-3) post-challenge dose period. Unsolicited AEs = during the 31-day (Day 0-30) follow-up period after the challenge dose.

Adverse event reporting additional description:

For the other adverse events: Data from the Australian center were not included following data quality issues detected at the investigator site.

| | |
|-----------------|----------------|
| Assessment type | Non-systematic |
|-----------------|----------------|

Dictionary used

| | |
|-----------------|--------|
| Dictionary name | MedDRA |
|-----------------|--------|

| | |
|--------------------|------|
| Dictionary version | 11.1 |
|--------------------|------|

Reporting groups

| | |
|-----------------------|----------------------------------|
| Reporting group title | Engerix 2 doses + challenge dose |
|-----------------------|----------------------------------|

Reporting group description:

Subjects received 2 doses of Engerix-B (Month 0 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| | |
|-----------------------|----------------------------------|
| Reporting group title | Engerix 3 doses + challenge dose |
|-----------------------|----------------------------------|

Reporting group description:

Subjects received 3 doses of Engerix-B (Month 0, 1 and 6) in the primary study and a single dose of Engerix-B during the booster study.

| Serious adverse events | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | |
|---|----------------------------------|----------------------------------|--|
| Total subjects affected by serious adverse events | | | |
| subjects affected / exposed | 0 / 55 (0.00%) | 0 / 22 (0.00%) | |
| number of deaths (all causes) | 0 | 0 | |
| number of deaths resulting from adverse events | | | |

Frequency threshold for reporting non-serious adverse events: 5 %

| Non-serious adverse events | Engerix 2 doses + challenge dose | Engerix 3 doses + challenge dose | |
|---|----------------------------------|----------------------------------|--|
| Total subjects affected by non-serious adverse events | | | |
| subjects affected / exposed | 39 / 55 (70.91%) | 14 / 22 (63.64%) | |
| Nervous system disorders | | | |
| Headache (AE) | | | |
| subjects affected / exposed | 4 / 55 (7.27%) | 2 / 22 (9.09%) | |
| occurrences (all) | 4 | 2 | |
| General disorders and administration site conditions | | | |

| | | |
|--|------------------|-----------------|
| Pain | | |
| alternative assessment type: Systematic | | |
| subjects affected / exposed | 22 / 55 (40.00%) | 4 / 22 (18.18%) |
| occurrences (all) | 22 | 4 |
| Redness | | |
| alternative assessment type: Systematic | | |
| subjects affected / exposed | 11 / 55 (20.00%) | 1 / 22 (4.55%) |
| occurrences (all) | 11 | 1 |
| Swelling | | |
| alternative assessment type: Systematic | | |
| subjects affected / exposed | 9 / 55 (16.36%) | 0 / 22 (0.00%) |
| occurrences (all) | 9 | 0 |
| Fatigue | | |
| alternative assessment type: Systematic | | |
| subjects affected / exposed | 19 / 55 (34.55%) | 7 / 22 (31.82%) |
| occurrences (all) | 19 | 7 |
| Gastrointestinal disorder | | |
| alternative assessment type: Systematic | | |
| subjects affected / exposed | 7 / 55 (12.73%) | 4 / 22 (18.18%) |
| occurrences (all) | 7 | 4 |
| Headache (Solicited Symptom) | | |
| alternative assessment type: Systematic | | |
| subjects affected / exposed | 14 / 55 (25.45%) | 4 / 22 (18.18%) |
| occurrences (all) | 14 | 4 |

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? No

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported