Clinical trial results:

Effect of strategy for blood pressure control on cerebral oxygen balance during aortic coarctation repair: a randomized study

EudraCT number	2007-002640-19
Trial protocol	BE
Global end of trial date	15 March 2012
Result version number	v1 (current)

Result version number	v1 (current)
This version publication date	07 November 2021
First version publication date	07 November 2021

Sponsor protocol code	AGO/2007/003
ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT00535808
WHO universal trial number (UTN)	-

Sponsor organisation name	Ghent University Hospital		
Sponsor organisation address	C. Heymanslaan 10, Ghent, Belgium, 9000		
Public contact	HIRUZ CTU, Ghent University Hospital, 32 93320500, hiruz.ctu@uzgent.be		
Scientific contact	HIRUZ CTU, Ghent University Hospital, 32 93320500, hiruz.ctu@uzgent.be		

Notes:

Notes:

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Notes:	

Analysis stage	Final
Date of interim/final analysis	01 June 2012
Is this the analysis of the primary completion data?	Νο
Global end of trial reached?	Yes
Global end of trial date	15 March 2012
Was the trial ended prematurely?	No

Notes:

Main objective of the trial:

investigation of the effect of different blood pressure controlling agents on the cerebral oxygen balance between both brain hemispheres during aortic coarctation repair by using near-infrared technology

Protection of trial subjects:

Ethics review and approval, informed consent, supportive care and routine monitoring.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	01 October 2007
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No
Notes:	

Worldwide total number of subjects 30	
EEA total number of subjects 30	

Notes:

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	15
Infants and toddlers (28 days-23 months)	15
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

Recruitment details:

Last patient last visit was on 15MAR2011

Screening details:

All neonates and infants, aged 0 – 18 year, with a rtic coarctation requiring surgical correction without the additional use of cardiopulmonary bypass

Period 1 title	overall trial (overall period)		
Is this the baseline period?	Yes		
Allocation method	Randomised - controlled		
Blinding used	Not blinded		

Are arms mutually exclusive?	Yes		
	Nitroprusside		
Arm description: -			
Arm type	Experimental		
Investigational medicinal product name	Nitroprusside		
Investigational medicinal product code			
Other name	Nitropress		
Pharmaceutical forms	Solution for infusion		
Routes of administration	Intravenous use		
Dosage and administration details:			
50 mg/2 ml vial for intravenous use, 0.1 5% dextrose injection before infusion	$-2 \mu g/kg/min$. The solution must be further diluted in sterile		
	Nitroglycerine		
Arm description: -			
Arm type	Experimental		
Investigational medicinal product name	Nitroglycerine		
Investigational medicinal product code			
Other name	Solinitrina		
Pharmaceutical forms	Solution for infusion		
Routes of administration	Intravenous use		
Dosage and administration details:			
50 mg/10 ml ampoule for intravenous u	se, 0.1 – 2 µg/kg/min		
	Sevoflurane		
Arm description: -			
Arm type	Experimental		
Investigational medicinal product name	•		
Investigational medicinal product code			
Other name	Sevorane		
Pharmaceutical forms	Inhalation vapour		
Routes of administration	Inhalation use		
Dosage and administration details:	·		

250 ml bottle, volatile anesthetic, 0.5 – 5 %

	Nitroprusside	Nitroglycerine	Sevoflurane
Started	10	10	10
Completed	10	10	10

Nitroprusside	
Nitroglycerine	
Sevoflurane	
	Nitroglycerine

Reporting group description: -

	Nitroprusside	Nitroglycerine	Sevoflurane
Number of subjects	10	10	10
Age categorical			
Units: Subjects			
In utero			
Preterm newborn infants (gestational age < 37 wks)			
Newborns (0-27 days)			
Infants and toddlers (28 days-23 months)			
Children (2-11 years)			
Adolescents (12-17 years)			
Adults (18-64 years)			
From 65-84 years			
85 years and over			
Age continuous			
Age of patients			
Units: days			
median	20	90	12
full range (min-max)	3 to 77	5 to 568	4 to 193
Gender categorical			
Units: Subjects			
Female	2	4	3
Male	8	6	7
ASA physical status			
Units: Subjects			
II	6	4	5
III	4	5	5
status I	0	1	0
Weight			
Units: kg			
median	3.6	5.1	3.3
full range (min-max)	1.4 to 5.7	3.2 to 11.3	2.0 to 8.6
BSA			
Units: m2			
median	0.22	0.27	0.21
full range (min-max)	0.12 to 0.28	0.2 to 0.49	0.15 to 0.39
	Tatal		
	Total		

Number of subjects	30		
Age categorical			
Units: Subjects			
In utero	0		
Preterm newborn infants (gestational age < 37 wks)	0		
Newborns (0-27 days)	0		
Infants and toddlers (28 days-23 months)	0		
Children (2-11 years)	0		
Adolescents (12-17 years)	0		
Adults (18-64 years)	0		
From 65-84 years	0		
85 years and over	0		
Age continuous			
Age of patients	_	_	
Units: days			
median			
full range (min-max)	-		
Gender categorical			
Units: Subjects			
Female	9		
Male	21		
ASA physical status			
Units: Subjects			
II	15		
III	14		
status I	1		
Weight			
Units: kg			
median			
full range (min-max)	-		
BSA			
Units: m2			
median			
full range (min-max)	-		

	Nitroprusside	Nitroglycerine	Sevoflurane	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	10	10	10	
Units: percentage				
arithmetic mean (standard deviation)	-64 (± 17)	-34 (± 25)	-55 (± 19)	

	Max change in muscle oxygen saturation
Comparison groups	Nitroglycerine v Nitroprusside v Sevoflurane
Number of subjects included in analysis	30
Analysis specification	Pre-specified
Analysis type	equivalence
P-value	= 0.014
Method	Kruskal-wallis

End point title	Decay rate renal oxygen saturation	
End point description:		
End point type	Primary	

overall	trial

	Nitroprusside	Nitroglycerine	Sevoflurane	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	10	10	10	
Units: percentage per minute				
arithmetic mean (standard deviation)	-7.7 (± 2.7)	-3.9 (± 3.0)	-5.6 (± 3.1)	

	Decay rate renal oxygen saturation
Comparison groups	Nitroprusside v Nitroglycerine v Sevoflurane

Number of subjects included in analysis	30
Analysis specification	Pre-specified
Analysis type	equivalence
P-value	= 0.034
Method	Kruskal-wallis

End point title	Decay rate muscle oxygen saturation	
End point description:		
End point type	Primary	
End point timeframe:		
overall trial		

	Nitroprusside	Nitroglycerine	Sevoflurane	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	10	10	10	
Units: percentage per minute				
arithmetic mean (standard deviation)	-9.3 (± 3.7)	-3.9 (± 2.7)	-6.2 (± 2.4)	

	Decay rate SmO2
Comparison groups	Nitroprusside v Nitroglycerine v Sevoflurane
Number of subjects included in analysis	30
Analysis specification	Pre-specified
Analysis type	equivalence
P-value	= 0.003
Method	Kruskal-wallis

Timeframe for reporting adverse events:		
overall trial		
Assessment type	Non-systematic	
Dictionary name	CTCAE	
Dictionary version	5	

Frequency threshold for reporting non-serious adverse events: 5 %

Notes:

[1] - There are no non-serious adverse events recorded for these results. It is expected that there will be at least one non-serious adverse event reported.Justification: No adverse events have been recorded during the study Were there any global interruptions to the trial? No

None reported