

Supplementary Appendix

This appendix has been provided by the authors to give readers additional information about their work.

Supplement to: Kernan WN, Viscoli CM, Furie KL, et al. Pioglitazone after ischemic stroke or transient ischemic attack. *N Engl J Med*. DOI: 10.1056/NEJMoa1506930

PIOGLITAZONE AFTER ISCHEMIC STROKE OR TRANSIENT ISCHEMIC ATTACK

SUPPLEMENTARY APPENDIX

TABLE OF CONTENTS

Section	Page
List of Investigators	1
Figure S1. Randomization and Follow-up of Study Participants	5
Figure S2. Selected Measures over time by Treatment Group	6
Table S1. Features of Baseline Stroke and TIA Events by Treatment Group	7
Table S2. Baseline Metabolic Characteristics by Treatment Group	8
Table S3. Achievement of Selected Secondary Stroke Prevention Goals by Year of Participation and Treatment Group	9
Table S4. Study Drug Adherence by Year of Participation and at Exit by Treatment Group	10
Table S5. Features of Primary and Secondary Stroke Outcomes by Treatment Group	11
Table S6. Changes in Selected Measures from Baseline to Year 1 by Treatment Group	12
Table S7. All Adverse Events Potentially Related to Pioglitazone by Treatment Group	13

Participating Sites (in order by country and number of participants):

AUSTRALIA (108 Participants): C. Bladin, Z. Ross, Monash University-Box Hill Hospital, Box Hill, Victoria (30); S. Davis, Royal Melbourne Hospital, Parkville, Victoria (29); T. Wijeratne, Western Hospital (University of Melbourne), Footscray, Victoria (16); C. Levi, M. Parsons, M. Russell, John Hunter Hospital (University of Newcastle), New Lambton Heights, Newcastle, NSW (15); A. Brodtmann, L. Walker, Austin Health (National Stroke Research Institute), Heidelberg Heights, Victoria (12); S. Ng, J. Archer, A. Rodda, R. Wong, The Northern Hospital, Epping, Victoria (4); C. Delcourt, E. Ray, The George Institute for International Health-Royal Prince Alfred Hospital, Camperdown, Sydney, NSW (2).

CANADA (543 Participants): T. R. Winder, J. Elliott, C. Walker, Center for Neurologic Research, Lethbridge, Alberta (98); L. Berger, J. Boulanger, Hopital Charles LeMoynes, Greenfield Park, Quebec (73); R. K. Chan, J. D. Spence, L. Fleming, K. Ford, Robarts Research Institute, London, Ontario (60); A. Durocher, CHUM-Centre de recherche, Hopital Notre-Dame, Montreal, Quebec (52); A. Mackey, S. Dubé, A. Hache, C. Lessard, CHU de Québec Hôpital de l'Enfant-Jésus, Quebec, Quebec (51); J. Minuk, C. Schanz, McGill University - Jewish General, Montreal, Quebec (50); A. M. Penn, L. Atkins, Vancouver Island Health Research Centre, Victoria, British Columbia (42); A. Shuaib, Y. Gable, B. Schwindt, University of Alberta, Edmonton, Alberta (37); R. Cote, L. Wadup, McGill University - Montreal General, Montreal, Quebec (35); D. Selchen, N. Bayer, P. Kostyrko, St. Michaels Hospital, University of Toronto, Toronto, Ontario (29); M. Sweet, S. Malik, Intermountain Research Consultants, Thunder Bay, Ontario (12); G. Stotts, S. Robert, Ottawa Hospital-General Campus, Ottawa, Ontario (4).

GERMANY (151 Participants): B. Griewing, H. Soda, R. Weinhardt, B. Behringer, M. Kirchner, A. Märkert, Neurologische Klinik Bad Neustadt, Bad Neustadt (27); J. Berrouschot, A. Stoll, Klinikum Altenburger Land, Altenburg (19); O. W. Witte, Jena University Hospital, Jena (17); U. Bodechtel, I. Dzialowski, K. Haase, J. Kepplinger, T. Siepman, University Hospital Dresden, Dresden (13); U. Schminke, Ernst-Moritz-Arndt-University Greifswald, Greifswald (10); C. Hobohm, Leipzig University, Leipzig (9); A. Hetzel, J. Lambeck, U. Bergmann, Freiburg University, Freiburg (8); K. E. Wartenberg, D. Herale, Martin-Luther-Universität Halle-Wittenberg, Halle (8); H. Huttner, University of Erlangen, Erlangen (8); R. Dittrich, University Hospital Münster, Münster (6); D. G. Nabavi, D. Seraphin, O. Crome, Neukolln Hospital, Berlin (5); K. Gröschel, University Hospital Mainz, Mainz (5); G. Thomalla, M. Rosenkranz, C. Beck, University Medical Center Hamburg-Eppendorf, Hamburg (5); S. Jander, B. Cullman, Heinrich-Heine-University, Medical Faculty, Department of Neurology, Düsseldorf (4); A. Meisel, S. Hoffmann, Charité-Universitätsmedizin Berlin, Berlin (3); A. Ludolph, K. Althaus, R. Huber, C. Adldinger, University of Ulm, Ulm (3); M. Lorenz, I.M. Dos Santos, University Hospital Frankfurt, Frankfurt am Main (1).

ISRAEL (178 Participants): D. Tanne, O. Merzlyak, C. Sasson, Sheba Medical Center, Tel Hashomer (47); N. M. Bornstein, L. Shopin, T. Ulanovsky-Selinger, Tel Aviv Medical Center, Tel Aviv (45); G. Telman, Rambam Health Corporation, Haifa (29); Y. Lampl, M. Israelson, Wolfson Medical Center, Holon (26); J. Streifler, N. Roizen, Rabin Medical Center-Golda Campus, Petach Tikva (18); B. Weller, E. Aubhanov, L. Mazareb, A. Mahagney, O. Nicola, S. Sabetay, Bnai-Zion Medical Center, Haifa (8); G. Ifergane, Y. Wirgin, Soroka University Medical Center, Beer-Sheva (5).

ITALY (48 Participants): A. Carolei, D. Degan, M. Mantenuto, S. Sacco, University of Laquila, L'Aquila (10); D. Toni, M. DiMascio, University of Rome La Sapienza, Rome (9); P. Stanzione, University of Rome Tor Vergata, Roma (7); G. Micieli, I. Canavero, A. Persico, IRCCS Fondazione Istituto Neurologico C. Mondino, Pavia (5); G. Agnelli, V. Caso, C. D'Amore, University of Perugia, Perugia (5); C. Gandolfo, D. Sassos, G. Sivori, Genoa University Hospital, Genoa (4); G. Comi, Hospital San Raffaele S.r.l., Milan (3); D. Consoli, F. Galati, P. Postorino, Jazzolino Hospital, Vibo Valentia (2); M. Rasura, F. Di Lisi, A. Spalloni, S. La Starza, University of Rome (S. Andrea Hospital), Roma (2); V. Di Lazzaro, F. Pilato, Sacred Heart Catholic University, Rome (1).

UNITED KINGDOM (256 Participants): A. Dixit, A. Barkat, G. Ford, K. Storey, Newcastle upon Tyne, Newcastle upon Tyne, Tyne and Wear, England (29); B. Jupp, D. Jenkinson, Royal Bournemouth & Christchurch Hospitals, Bournemouth, England (28); L. Shaw, B. Madigan, Royal United Hospital Bath

NHS Foundation Trust, Avon, England (28); I. Salih, D. Kelly, Torbay Hospital (South Devon Healthcare NHS Foundation Trust), Devon, England (21); B. Esisi, Queen Elizabeth Hospital Gateshead, Sheriff Hill, Gateshead, England (18); M. Power, D. Wilson, B. Wroath, Ulster Hospital, Dundonald, Belfast, Northern Ireland (12); W. D. Strain, S. Elyas, S. Keenan, H. Kingwell, Royal Devon and Exeter, Exeter, England (11); D. Manawadu, L. Kalra, K. Anderson, B. Mistry, Kings College London, London, England (10); E. O'Brien, E. A. Warburton, E. Amis, N. Butler, J. Mitchell, Addenbrookes Foundation NHS Trust (Cambridge), Cambridge, England (9); K. Chatterjee, S. Leason, Countess of Chester Foundation Trust, Chester, England (9); D. R. Hargroves, L. Cowie, William Harvey Hospital, Ashford, Kent, England (9); A. Blight, C. Peixoto, K. Pasco, Royal Surrey County Hospital, Guildford, England (8); B. Moynihan, H. Markus, S. Trippier, Saint Georges University of London, Tooting, England (8); M. Macleod, J. Furnace, H. Gow, A. Joyson, Aberdeen Royal Infirmary, Foresterhill, Aberdeen, Scotland (6); D. L. Broughton, L. Dixon, K. Chapman, James Cook University Hospital, Middlesbrough, Cleveland, England (6); H. Rodgers, A. Smith, North Tyneside General Hospital, Newcastle upon Tyne, Tyne and Wear, England (5); T. Hlaing, University Hospital Aintree, Liverpool, England (5); S. Muir, B. Manak, L. McDonald, Western Infirmary, Glasgow, Scotland (5); M. Sajid, R. Gascoyne, Chesterfield Hospital, Chesterfield, Derbyshire (4); P. M. Bath, J. Clarke, K. Krishnan, University of Nottingham, Nottingham, England (4); C. Price, Wansbeck General Hospital, Ashington, Northumberland, England (4); L. Sekaran, F. Justin, Luton and Dunstable Hospital, Luton, England (3); D. Vahidassr, A. Thompson, Northern Trust, Co Antrim, N Ireland (3); K. W. Muir, A. Welch, Southern General Hospital, Glasgow, Scotland (3); J. McIlmoyle, Blackpool Victoria Hospital, Blackpool, England (2); P. K. Datta, R. Davey, G. Bateman, A. Needle, Mid Yorkshire NHS Trust (Dewsbury), West Yorkshire, England (2); P. Langhorne, D. Stott, R. Graham, Glasgow Royal Infirmary, Glasgow, Scotland (1); P. K. Datta, R. Davey, Mid Yorkshire NHS Trust (Pinderfields), West Yorkshire, England (1); T. J. England, K. Muhiddin, Royal Derby Hospital, Mickleover, Derby, England (1); J. E. O'Connell, N. Majmudar, S. Crawford, E. Osborne, Sunderland Royal Hospital, Sunderland, Tyne and Wear, England (1).

UNITED STATES (2592 Participants): J. Schindler, N. Finch, W. Kernan B. Mendes, K. Shrauger, Yale University, New Haven, CT (339); W. M. Clark, J. Foley, Oregon Health & Science University, Portland, OR (201); P. Sethi, S. Hammonds, D. Parker, Guilford Neurologic Associates, Greensboro, NC (123); G. Rordorf, M. Amatangelo, Massachusetts General Hospital/General Hospital Corp, Boston, MA (106); D. O. Kleindorfer, A. Hodge, J. Osborne, E. Goodall, University of Cincinnati, Cincinnati, OH (100); S. L. Silliman, Y. Shabbir, M. Varghese, University of Florida, Jacksonville, FL (96); M. Gorman, C. Gregory, S. Kenney, University of Vermont, Burlington, VT (89); M. A. Kelly, L. Singleton, M. Ohprecio, John H. Stroger, Jr. Hospital of Cook County, Chicago, IL (73); B. C. Meyer, C. Jackson, R. Chavez, N. Kelly, K. Rapp, University of California, San Diego, San Diego, CA (70); J. Walker, A. Ehtisham, H. Goodpasture, D. Collie, S. Miller, Via Christi Regional Medical Center, Wichita, KS (65); D. Wang, J. Beck, A. Bogner, OSF Saint Francis Medical Center, Peoria, IL (63); P. Fayad, A. Colburn, M. Phillips, University of Nebraska, Omaha, NE (49); S. Cordina, D. Naritoku, R. Backlin, University of South Alabama, Mobile, AL (42); D. Chiu, Methodist Hospital Research Institute, Houston, TX (41); T. Lukovits, R. Goddeau, R. Clark-Arbogast, E. Clough, M. Notestine, D. Sherman, Dartmouth, Lebanon, NH (40); R. Leigh, R. J. Wityk, S. Rice, Johns Hopkins University, Baltimore, MD (40); L. Pettigrew, D. Taylor, University of Kentucky, Lexington, KY (40); A. H. Tayal, M. Tian, Allegheny Singer Research Institute, Pittsburgh, PA (36); G. H. Friday, J. Brown, A. Whittington-Gordon, Lankenau Institute for Medical Research, Wynnewood, PA (36); S. Sen, J. Burdine, H. Kowalczyk, R. Sumner, University of South Carolina, Columbia, SC (36); A. S. Kim, J. S. Elkins, S. C. Johnston, T. Yi, P. Buekea, K. Fukuda, A. Teng, T. Yi, University of California, San Francisco, San Francisco, CA (34); A. M. Barrett, J. Feriod, J. R. Masmela, K. Sandefur, Kessler Foundation, West Orange, NJ (33); E. C. Leira, H. Olalde, J. Sieren, University of Iowa, Iowa City, IA (33); A. Kelly, S. Burgin, D. Rempe, H. Finley, J. Zentner, University of Rochester, Rochester, NY (31); M. R. Jacoby, B. Hughes, Ruan Neuroscience Center/Mercy Medical Center, Des Moines, IA (30); J. Majersik, E. J. Skalabrin, J. Austin, S. Niu, University of Utah, Salt Lake City, UT (28); J. Lee, J. Naylor, J. Newgent, R. Sargent, Washington University, St. Louis, MO (28); S. Sundararajan, D. Korosec, Case Western Reserve University, Cleveland, OH (26); A. Slivka, A. Green, P. Notestine, C. Heckman, Ohio State University,

Columbus, OH (26); A. Minagar, LSU Health Sciences Center, Shreveport, LA (25); R. Alicic, M. Geraghty, R. Franks, K. Worrell, Providence Medical Research Center, Spokane, WA (24); C. S. Kase, H. Lau, V. Babikian, Boston Medical Center Corp, Boston, MA (21); M. Lansberg, G. Albers, M. Garcia, Stanford University, Stanford, CA (21); D. W. Dietrich, L. Armstrong, Advanced Neurology Specialists, Great Falls, MT (20); J. P. Hanna, D. Cook, Metrohealth Medical Center, Cleveland, OH (20); N. T. Gentile, H. Reimer, Temple University, Philadelphia, PA (20); F. Santiago, P. Fuste, University of Puerto Rico, San Juan, Puerto Rico (20); I. Katzan, A. Richmond, L. Strozniak, Cleveland Clinic Foundation, Cleveland, OH (19); M. Ching, A. Crumlish, D. Holler, Research Foundation SUNY, University of Buffalo, Buffalo, NY (18); T. Warwick, R. Garcia, UCSF (Fresno), Fresno, CA (18); E. Yilmaz, D. Lawrence, Ingalls Memorial Hospital, Harvey, IL (17); L. Pedelty, M. Hillman, University of Illinois, Chicago, Chicago, IL (17); M. J. Schneck, T. Lynch, J. Ray, Loyola University Chicago, Maywood, IL (15); B. M. Coull, N. Malekniazi, University of Arizona, Tucson, AZ (15); N. J. Solenski, K. Johnston, C. Harman, University of Virginia, Charlottesville, VA (15); V. Lee, S. Prabhakaran, T. Cole, S. Dahl, Rush University, Chicago, IL (14); M. D. Johnson, J. Cameron-Watts, J. Wright, University of Texas, Southwestern, Dallas, TX (14); I. E. Silverman, M. Ahlquist, Hartford Hospital, Hartford, CT (13); M. W. Salgado, R. Birkhahn, New York Methodist Hospital, Brooklyn, NY (13); R. Strawsburg, Associates in Neurology, P.C., Valparaiso, IN (12); I. Altafullah, Minneapolis Clinic of Neurology, Golden Valley, MN (12); D. A. Cohen, R. Zweifler, S. Bright, P. Hollsten, Sentara Neurology Specialists, Norfolk, VA (12); P. Lee Kwen, N. Kamuda, Southtowns Neurology of WNY, P.C., West Seneca, NY (12); M. D. Hammer, J. Oakley, University of Pittsburgh, Pittsburgh, PA (12); G. E. Tietjen, A. Korsnack, University of Toledo, Toledo, OH (11); E. Albakri, I. Antonek, Florida Neurovascular Institute, Tampa, FL (10); B. K. Dandapani, B. Anthony, G. Tirschner, Health First Physicians, Inc., Melbourne, FL (10); G. Jickling, P. Verro, UC-Davis Medical Center, Sacramento, CA (10); M. J. Roller, R. Hendrickson, R. Sperle, Altru Health System, Grand Forks, ND (9); R. L. Hughes, J. Simpson, Denver Health and Hospital Authority, Denver, CO (9); T. R. Vidic, M. Quakenbush, K. Stanley, Indiana Medical Research, Elkhart, IN (9); S. Lash, B. Sigsbee, J. Sigsbee, Penobscot Bay Neurology, Rockport, ME (9); D. Rosenbaum, S. Cherian, L. Thomas, SUNY Downstate, Brooklyn, NY (9); P. Fonzetti, D. Zondorak, Burke Medical Research Institute, White Plains, NY (8); J. D. Fleck, S. Guingrich, Indiana University, Indianapolis, IN (8); A. J. Goldszmidt, Sinai Hospital of Baltimore, Baltimore, MD (8); A. V. Alexandrov, J. H. Halsey, M. Taylor, University of Alabama, Birmingham, AL (8); R. Hart, L. Moreno, University of Texas, San Antonio, San Antonio, TX (8); J. A. Sattin, A. Harris, University of Wisconsin, Madison, WI (8); S. Kumar, Beth Israel Deaconess, Boston, MA (7); D. Book, M. Torbey, B. Blaney, Medical College of Wisconsin, Milwaukee, WI (7); J. J. Poock, P. Trenkamp, Northeast Iowa Medical Education Foundation, Waterloo, IA (7); M. K. King, G. D. Graham, J. Harnar, University of New Mexico, Albuquerque, NM (7); G. Y. Sung, M. Fernandez, University of Southern California, Los Angeles, CA (6); T. Mirsen, C. Hackmyer, Cooper University Hospital, Camden, NJ (5); A. W. Dromerick, M. Giannetti, L. Priestley, National Rehabilitation Hospital, Washington, DC (5); A. D. Runheim, Salem Neurological Center, Winston-Salem, NC (5); C. M. Jackson, S. Abel, G. Moore, C. Sanchez, Scripps Clinic, La Jolla, CA (5); E. Feen, C. Morefield, St. Louis University, St. Louis, MO (5); R. K. Reichwein, K. Hitz, Penn State-Hershey Medical Center, Hershey, PA (4); M. F. Waters, E. Brooks, University of Florida, Gainesville, Gainesville, FL (4); C. Amory, G. L. Bernardini, S. Rauch, Albany Medical Center, Albany, NY (3); R. D. Bell, Thomas Jefferson University, Philadelphia, PA (2); B. Diamond, Abington Memorial Hospital, Abington, PA (1); D. M. Rosenbaum, Albert Einstein, Bronx, NY (1); D. Palestrant, Cedars-Sinai Medical Center, Los Angeles, CA (1); A. Z. Segal, Cornell University, New York, NY (1); K. Burger, George Washington University, Washington, DC (1); R. L. Schwartz, Hattiesburg Clinic, Hattiesburg, MS (1); P. Mitsias, B. Stalling, Henry Ford Health Sciences Center, Detroit, MI (1); J. Kramer, Jeffrey Kramer, MDSC (formerly Kramer/Mercy), Chicago, IL (1); D. Robbins, Pines Neurological Associates, Pembroke Pines, FL (1); B. Silver, J.D. Easton, E. Feldmann, Rhode Island Hospital, Providence, RI (1); M. M. Rymer, J. McIntire, St. Lukes Brain and Stroke Institute, Kansas City, MO (1); L. Ali, B. Ovbiagele, G. Avila, S. Pineda, University of California, Los Angeles, Los Angeles, CA (1); H. S. Kirshner, D. Brown, Vanderbilt University, Nashville, TN (1).

Operations Committee: W. Kernan (chair), L. Brass (deceased), K. Furie (principal neurologist), M. Gorman (research neurologist), A. Lovejoy (principal coordinator), R. Horwitz (2004-2011), S. Inzucchi (principal endocrinologist), C. Viscoli (principal epidemiologist), L. Young (principal cardiologist); **Steering Committee:** Members of the Operations Committee plus H. Adams, A. Carolei, W. Clark, B. Coull, G. Ford, P. Guarino, M. Parsons, P. Peduzzi, P. Ringleb, G. Schwartz, J. Spence, D. Tanne; **Scientific Advisory Committee:** P. Gorelick (chair), E. Barrett, M. Dyken, R. Nesto, W. Longstreth, Jr., D. Richardson; **Statistical Center:** P. Guarino, J. O’Leary, P. Peduzzi, B. Zhou; **Central Pharmacist:** O. Abdelghany, Sandra Alfano (2004-2005); **Medical Safety Monitor:** L. Golden, D. Lorber (2004-2006); **Cancer Reviewer:** E. Hofstatter; **Cardiology Review Committee:** F. Ling (chair), J. Abbott, J. Curtis, J. Foody, D. Jacoby, D. Kolansky, S. Pfau, J. Lasala (2004-2007); **Endocrinology Review Committee:** M. Korytkowski (chair), S. Engel (2004-2005), F. Ismail-Beigi, D. McCulloch, R. Pratley, P. Berhanu (2006-2007), E. Caballero (2004-2008), S. Dagogo-Jack (2006-2007), M. Schutta (2007); **Fracture Review Committee:** A. Haims, K. Insogna, D. Lindskog; **Macular Edema Reviewer:** R. Adelman; **Neurology Review Committee:** S. Kasner (chair), J. Broderick, R. Cote, C. Kase, M. Alberts (2004-2006); **Data Safety and Monitoring Board:** M. Walker (chair), J. Buse, L. Chambless, D. Faxon, J. Pary; **Funding Agency:** R. Conwit (Project Scientist); C.S. Moy (Program Official); L.S. Janis (DSMB Liaison); L. Gutmann (Program Official), B. Radziszewska (Program Official) National Institutes of Health/National Institute of Neurological Disorders and Stroke (NIH/NINDS), Bethesda, MD, USA.








Figure S1. Randomization and Follow-up of Study Participants. Participants were assessed for eligibility before consenting to be tested for insulin resistance, but the number of patients assessed is unknown because screening logs were not maintained. *Includes 30 participants (19 in pioglitazone group and 11 in placebo group) who were randomized in error (i.e., did not meet entry criteria); these participants were retained in follow-up and are included in analysis.





Pioglitazone (N)= 1939 1659 1484 1290 1029 723
 Placebo (N)= 1937 1709 1519 1314 1027 719

1935 1683 1553 1370 1099
 1934 1723 1588 1384 1102



Pioglitazone (N)= 1934 1682 1552 1370 1098
 Placebo (N)= 1934 1723 1586 1384 1102

1918 1675 1541 1363 1092
 1920 1705 1576 1373 1087



Pioglitazone (N)= 1993 1711 1570 1378 1111 803
 Placebo (N)= 1931 1741 1595 1403 1126 791

1933 1711 1570 1378 1111 803
 1931 1741 1595 1403 1126 790

Figure S2. Selected Measures over time by Treatment Group (Pioglitazone ◆; Placebo ■).

Least square means with 95% confidence intervals are from a repeated measures covariance pattern model assuming missing at random at follow-up time points and for overall estimate. Overall p values are for the treatment effect for years 1 to 4 adjusted for baseline (for lipids) and for years 1 to 5 adjusted for baseline (for glucose and systolic blood pressure). For diastolic blood pressure, p-values are shown separately for each follow-up year because of a significant treatment by year effect.

Table S1. Features of Baseline Stroke and TIA Events by Treatment Group

Subtype	Pioglitazone		Placebo	
	N	%	N	%
Lacunar	584	30%	559	29%
Embolic	147	8%	137	7%
Large vessel	500	26%	504	26%
Uncertain	697	36%	730	38%
NA (not eligible stroke/TIA)	11	1%	7	0%
Distribution				
Left Anterior/Carotid	665	34%	659	34%
Right Anterior/Carotid	601	31%	604	31%
Vertebral/Basilar	470	24%	469	24%
Multiple	55	3%	45	2%
Uncertain	137	7%	153	8%
NA (not eligible stroke/TIA)	11	1%	7	0%
Modified Rankin Scale				
0	761	39%	774	40%
1	653	34%	679	35%
2	345	18%	322	17%
3	125	6%	118	6%
4	44	2%	36	2%
Unknown	0	0%	1	0%
NA (not eligible stroke/TIA)	11	1%	7	0%
Stroke Severity (NIH Stroke Scale)				
Mild impairment (<5)	1835	95%	1843	95%
Moderately severe impairment (5-14)	92	5%	84	4%
Severe impairment (15-24)	0	0%	2	0%
Very severe impairment (>25)	0	0%	0	0%
Unknown	1	0%	1	0%
NA (not eligible stroke/TIA)	11	1%	7	0%

Table S2. Baseline Metabolic Characteristics by Treatment Group

Characteristic	Pioglitazone (N=1939)	Placebo (N=1937)
Fasting glucose, no (%)		
<100 mg/dL	1126 (58.1)	1137 (58.7)
100-109 mg/dL	546 (28.2)	543 (28.0)
110-125 mg/dL	267 (13.8)	257 (13.3)
Insulin (μU/mL), median (Q25,Q75)		
	19 (16,26)	19 (16,25)
HOMA-IR, median (Q25,Q75)		
	4.7 (3.8,6.2)	4.6 (3.7,6.2)
HbA_{1c}, no (%)		
<5.7%	672 (34.7)	690 (35.6)
5.7-6.4%	1150 (59.3)	1118 (57.7)
6.5-6.9%	116 (6.0)	129 (6.7)
Diabetes Status[¶], no (%)		
Euglycemia	457 (23.6)	473 (24.4)
Pre-diabetes	1366 (70.4)	1335 (68.9)
Diabetes	116 (6.0)	129 (6.7)
[¶] According to ADA 2010 criteria: Euglycemia: Fasting Glucose <100 mg/dL and HbA _{1c} <5.7%; Pre-diabetes: Fasting Glucose 100-125 mg/dL or HbA _{1c} 5.7-6.4%; Diabetes: HbA _{1c} 6.5-6.9%		

Table S3. Achievement of Selected Secondary Stroke Prevention Goals by Year of Participation and Treatment Group

Prevention Goal		Pioglitazone			Placebo		
		No. Pts. in Trial ^x	No. Pts. missing data ^y	% Pts. Achieving Goal ^o	No. Pts. in Trial ^x	No. Pts. missing data ^y	% Pts. Achieving Goal ^o
Blood Pressure <140/90	Baseline	1939	6	66	1937	6	67
	Year 1	1894	183	71	1881	140	69
	Year 2	1834	265	73	1838	243	70
	Year 3	1684	307	74	1679	276	71
	Year 4	1422	312	73	1396	270	72
	Year 5	1021	219	75	1008	218	73
On Anti-Thrombotics [†]	Baseline	1939	2	99	1937	1	99
	Year 1	1894	128	97	1881	99	97
	Year 2	1834	159	97	1838	130	97
	Year 3	1684	177	97	1679	154	96
	Year 4	1422	166	95	1396	143	96
	Year 5	1021	106	95	1008	112	95
On Statin	Baseline	1939	7	83	1937	5	82
	Year 1	1894	128	79	1881	103	80
	Year 2	1834	161	80	1838	135	80
	Year 3	1684	176	79	1679	151	78
	Year 4	1422	172	79	1396	144	79
	Year 5	1021	108	77	1008	112	77
Non-Smoker	Baseline	1939	2	83	1937	1	85
	Year 1	1894	120	83	1881	93	84
	Year 2	1834	156	83	1838	130	84
	Year 3	1684	166	84	1679	148	85
	Year 4	1422	161	85	1396	143	85
	Year 5	1021	104	85	1008	108	86

^xNumber of participants in trial (excludes participants who had died, withdrawn consent or completed the final follow-up interview before beginning of year).

^yNumber of participants with missing data on prevention goal.

^oPercent of participants in trial who met prevention goal (denominator is number of participants in trial minus number participants missing data).

[†]Taking an anti-platelet or anti-coagulant medication.

Table S4. Study Drug Adherence by Year of Participation and at Exit by Treatment Group

	Study Drug Status at End of Year (%)						Adherence by Pill Counts by Year		
	No. Pts. ^o	Removed by ISC [†]	Off Drug for Other Reason	15 mgs/day ^v	30 mgs/day	45 mgs/day	Mgs/day, median	%Maximum Dose ^a , mean	%Pts with Good Adherence [‡]
Pioglitazone									
Year 1	1894	2%	22%	5%	4%	67%	37	73	61
Year 2	1834	4%	26%	5%	4%	61%	40	65	55
Year 3	1684	5%	29%	6%	4%	56%	38	61	52
Year 4	1422	7%	32%	6%	3%	51%	35	56	49
Year 5	1021	9%	31%	6%	3%	50%	38	61	52
Exit	1692	8%	32%	7%	3%	49%	29	61	47
Placebo									
Year 1	1881	1%	14%	2%	2%	81%	39	81	75
Year 2	1838	3%	19%	3%	2%	73%	43	75	69
Year 3	1679	4%	24%	3%	1%	68%	42	71	64
Year 4	1396	6%	27%	4%	1%	61%	42	66	61
Year 5	1008	7%	29%	4%	1%	59%	42	68	62
Exit	1689	6%	27%	4%	2%	61%	37	71	60

^oNumber of participants in trial (excludes participants who had died, withdrawn consent or completed the final follow-up interview before beginning of year). Participants who were unable to be located were classified as off study drug.

[†]Participants were only classified as removed if they had been instructed to stop taking the study drug by the IRIS Internal Safety Committee.

^vIncludes participants on less than 15mg/day dosing regimen.

^a%Maximum dose= (mg consumed per day as estimated by pill countbacks)/(maximum mg as stipulated by protocol if dose is never adjusted and treatment is not interrupted for any reason).

[‡]Good Adherence=taking ≥80% of maximum dose as stipulated by protocol.

Table S5. Features of Primary and Secondary Stroke Outcome Events by Treatment Group

Feature	Stroke Events in Primary Outcome (Stroke or MI)				Stroke Events in Secondary Outcome (Stroke alone)				
	Pioglitazone		Placebo		Pioglitazone		Placebo		
	N	%	N	%	N	%	N	%	
Number	123		150		127		154		
Stroke Type									
Ischemic	109	89%	136	91%	112	88%	140	91%	
Non-fatal	104	85%	129	86%	107	84%	132	86%	
Fatal	5	4%	7	5%	5	4%	8	5%	
Hemorrhagic	13	11%	14	9%	14	11%	14	9%	
Non-fatal	10	8%	8	5%	10	8%	8	5%	
Fatal	3	2%	6	4%	4	3%	6	4%	
Uncertain (fatal)	1	1%	0	0%	1	1%	0	0%	
Ischemic Stroke Subtype									
Lacunar	9	8%	17	13%	9	8%	17	12%	
Embolic	15	14%	19	14%	16	14%	20	14%	
Large vessel	17	16%	22	16%	17	15%	22	16%	
Uncertain	68	62%	78	57%	70	63%	81	58%	
Ischemic Stroke Distribution									
Carotid	72	66%	92	68%	74	66%	93	66%	
Vertebral/Basilar	17	16%	31	23%	18	16%	32	23%	
Multiple	7	6%	2	1%	7	6%	3	2%	
Uncertain	13	12%	11	8%	13	12%	12	9%	
NIH Stroke Scale									
0-4	82	67%	88	59%	84	66%	89	58%	
5-14	26	21%	36	24%	27	21%	37	24%	
15-41	3	2%	6	4%	3	2%	7	5%	
42 (fatal)	9	7%	13	9%	10	8%	14	9%	
Missing	3	2%	7	5%	3	2%	7	5%	
Mean (SD)	6.8 (10.9)		7.9 (11.7)		6.9 (11.1)		8.2 (12.0)		
Mean change from Baseline									
All strokes	5.5 (10.8)		6.4 (11.7)		5.7 +11.1		6.8 +12.0		
Non-fatal strokes	2.6 (4.2)		3.0 (4.7)		2.6 +4.1		3.2 +5.1		
Days to NIHSS (median)	15		17		16		17		

Table S6. Changes in Selected Measures from Baseline to Year 1 by Treatment Group

Measure	Pioglitazone			Placebo			P-value for mean change ^Y
	Baseline	Year 1	Change	Baseline	Year 1	Change	
HOMA							
(N)	(1440)			(1458)			<.0001
Mean	5.4	4.1	-1.3	5.3	5.7	0.4	
SD	2.7	2.8	3.1	2.6	6.6	6.5	
Median	4.6	3.4	-1.3	4.5	4.4	-0.3	
IQR	2.4	2.1	2.2	2.5	3.0	2.5	
Insulin, µU/mL							
(N)	(1449)			(1460)			<.0001
Mean	22.3	17.2	-5.1	22.0	22.5	0.5	
SD	10.2	9.9	11.2	9.7	17.8	17.0	
Median	19	15	-5	19	18	-1	
IQR	9	9	8	9	11	10	
Fasting Glucose, mg/dL							
(N)	(1659)			(1709)			<.0001
Mean	98.2	95.1	-3.0	98.3	99.7	1.4	
SD	10.0	11.0	10.5	9.9	16.6	15.0	
Median	97	94	-3	98	97	0	
IQR	14	13	12	14	14	12	
C-Reactive Protein, mg/L							
(N)	(1628)			(1665)			0.02
Mean	4.5	3.3	-1.2	4.7	4.4	-0.3	
SD	8.9	7.5	10.5	8.8	9.8	11.8	
Median	2.1	1.3	-0.4	2.3	2.0	-0.2	
IQR	3.7	2.6	2.1	3.7	3.4	2.0	

^YP-value from T-test for comparison of mean change from baseline to year 1 between treatment groups.

Table S7. All Adverse Events Potentially Related to Pioglitazone by Treatment Group

Event	Pioglitazone (N=1939)	Placebo (N=1937)	P-value
Weight Gain			
> 10 lbs (4.5 kgs) ^y	1013 (52.2)	653 (33.7)	<.001
> 30 lbs (13.5 kgs) ^y	221 (11.4)	88 (4.5)	<.001
At Year 1, mean	3.9 lbs (1.8 kgs)	-0.7 lbs (-0.3 kgs)	<.001
At Year 4, mean	5.8 lbs (2.6 kgs)	-1.2 lbs (-0.5 kgs)	<.001
Edema^a			
Any	691 (35.6)	483 (24.9)	<.001
Severe	348 (17.9)	235 (12.1)	<.001
Myalgia^z			
Any	676 (34.9)	650 (33.6)	0.39
Severe	76 (3.9)	74 (3.8)	0.87
Shortness of Breath^δ			
	342 (17.6)	292 (15.1)	0.03
Bone Fracture^μ			
Any	218 (11.2)	145 (7.5)	<.0001
Serious	99 (5.1)	62 (3.2)	0.003
Incident Cancer			
Any	133 (6.9)	150 (7.7)	0.29
Non-melanoma skin	21 (1.1)	32 (1.6)	0.13
Prostate	28 (1.4)	25 (1.3)	0.68
Breast	10 (0.5)	16 (0.8)	0.24
Lung	13 (0.7)	11 (0.6)	0.68
Bladder	12 (0.6)	8 (0.4)	0.37
Colon and Rectum	7 (0.4)	8 (0.4)	0.79
Melanoma	8 (0.4)	5 (0.3)	0.41
Kidney	3 (0.2)	8 (0.4)	0.13
Lymphoma	5 (0.3)	4 (0.2)	0.74
Uncertain	3 (0.2)	4 (0.2)	0.70
Other (<i>types with n<6</i>)	29 (1.5)	33 (1.7)	0.61
Heart Failure^χ			
Any	74 (3.8)	71 (3.7)	0.80
Serious	51 (2.6)	42 (2.2)	0.35
Alanine aminotransferase			
>upper limit of normal	26 (1.3)	59 (3.0)	<.001
>2.5 upper limit of normal	7 (0.4)	14 (0.7)	0.13
Macular Edema^β			
	3 (0.2)	2 (0.1)	0.66
Other Symptoms and Complaints^v			
Tooth problems	555 (28.6)	601 (31.0)	0.10
Frequent headaches	395 (20.4)	410 (21.2)	0.54
Sinus infection	229 (11.8)	241 (12.4)	0.55
Throat infection	120 (6.2)	133 (6.9)	0.39

^yWeight change from baseline at any time in trial.

^aSelf-report at quarterly interview of new or worse swelling of feet or lower legs; severe=persisting after leg elevation or associated with discomfort, redness, skin breakdown or difficulty getting shoes on feet.

^zSelf-report at quarterly interview of new or worse muscle aches; severe=diffuse muscle aches associated with limitation in mobility or substantial discomfort.

^δSelf-report at quarterly interview of new or worse shortness of breath or difficulty breathing.

^μOther types of cancer with frequency < 6.

^χAdjudicated bone fracture; serious=fracture resulting in hospitalization, surgery or procedure.

^βAdjudicated heart failure; serious=heart failure causing or prolonging hospitalization or causing death.

^βAdjudicated episode of macular edema.

^vSymptoms/complaints by self-report at annual interview: Tooth problems, infections requiring a visit to a dentist/doctor.