

DEATHS WHILE ON TREATMENT AND WITHIN 30 DAYS FROM LAST CETUXIMAB DOSE

Patient ID	Arm	Gender	Primary tumour	Interval on treatment (weeks)	Reason for discontinuation	Cause of death	Interval last cetuximab dose to death (days)
100-01-002	B (standard cetuximab)	Male	Colon	13	Disease progression	Disease progression	20
100-08-002	B (standard cetuximab)	Male	Colon	49	Death	Bronchial infection post serious cardiac event	8
200-06-001	B (standard cetuximab)	Male	Colon	16	Disease progression	Disease progression and intercurrent pneumonia	20
400-02-002	B (standard cetuximab)	Male	Rectum	2	Adverse event	Perforation of the sigmoid with fecaloid peritonitis followed by respiratory insufficiency	19
400-02-010	B (standard cetuximab)	Male	Rectum	12	Death	Unexpected death during sleep due to circulatory failure of unknown origin	11
400-06-007	B (standard cetuximab)	Male	Colon	2	Death	Ileus due to progressive disease	23
600-01-008	A (escalated cetuximab)	Male	Colon	5	Death	Septic shock and multiorgan failure secondary to colonic perforation	14
600-01-018	B (standard cetuximab)	Male	Colon	5	Death	Cardiopulmonary arrest	1
600-01-030	B (standard cetuximab)	Male	Colon	72	Disease progression	Progressive disease, intestinal occlusion and cardiac failure	29