



## Clinical trial results:

### A Long-Term, Open-Label, Safety Study of Oral Olanzapine in Adolescents with Bipolar I Disorder (Manic or Mixed Episodes) or Schizophrenia

#### Summary

EudraCT number	2009-010276-16
Trial protocol	DE
Global end of trial date	07 May 2013

#### Results information

Result version number	v1 (current)
This version publication date	04 July 2016
First version publication date	02 August 2015

#### Trial information

##### Trial identification

Sponsor protocol code	F1D-MC-HGMX
-----------------------	-------------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT00982020
WHO universal trial number (UTN)	-
Other trial identifiers	Trial Number: 12117, Trial Alias: F1D-MC-HGMX

Notes:

##### Sponsors

Sponsor organisation name	Eli Lilly and Company
Sponsor organisation address	Lilly Corporate Center, Indianapolis, IN, United States, 46285
Public contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877-285-4559,
Scientific contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877-CTLilly,

Notes:

##### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	07 May 2013
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	07 May 2013
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

The primary objectives of this study were to assess in adolescent patients with bipolar I disorder (manic or mixed episode) or schizophrenia: 1) the overall safety of olanzapine for up to approximately 52 weeks of treatment, and 2) whether an Intense behavioral weight intervention program was superior to a Standard behavioral weight intervention program in mitigation of weight gain as assessed by overall mean change from baseline body mass index (BMI).

Protection of trial subjects:

This study was conducted in accordance with International Conference on Harmonisation (ICH) Good Clinical Practice, and the principles of the Declaration of Helsinki, in addition to following the laws and regulations of the country or countries in which a study is conducted.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	01 September 2009
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Germany: 3
Country: Number of subjects enrolled	Poland: 18
Country: Number of subjects enrolled	Russian Federation: 55
Country: Number of subjects enrolled	United States: 107
Country: Number of subjects enrolled	Puerto Rico: 20
Worldwide total number of subjects	203
EEA total number of subjects	21

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0

Adolescents (12-17 years)	203
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

No text entered

### Pre-assignment

Screening details:

No text entered

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Not blinded

### Arms

Are arms mutually exclusive?	Yes
------------------------------	-----

<b>Arm title</b>	Olanzapine/standard behavioral weight intervention
------------------	--

Arm description:

Olanzapine: 2.5 milligrams (mg) to 20 mg given orally, daily for 52 weeks.

Standard behavioral weight intervention: One time counseling and basic counseling information on healthy eating and exercise habits at randomization visit only

Arm type	Experimental
Investigational medicinal product name	Olanzapine
Investigational medicinal product code	
Other name	Zyprexa, LY170053
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Olanzapine: 2.5 milligrams (mg) to 20 mg given orally, daily for 52 weeks. Standard behavioral weight intervention: One time counseling and basic counseling information on healthy eating and exercise habits at randomization visit only

<b>Arm title</b>	Olanzapine/intense behavioral weight intervention
------------------	---

Arm description:

Olanzapine: 2.5 mg to 20 mg given orally, daily for 52 weeks.

Intense behavioral weight intervention: Counseling provided at randomization and at all subsequent study visits by appropriately trained individual regarding healthy lifestyle habits. Participants also provided with simple tools to help enable healthy eating and exercise habits

Arm type	Experimental
Investigational medicinal product name	Olanzapine
Investigational medicinal product code	
Other name	Zyprexa, LY170053
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Olanzapine: 2.5 mg to 20 mg given orally, daily for 52 weeks. Intense behavioral weight intervention: Counseling provided at randomization and at all subsequent study visits by appropriately trained individual regarding healthy lifestyle habits. Participants also provided with simple tools to help enable healthy eating and exercise habits

<b>Number of subjects in period 1</b>	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention
Started	102	101
Received at least 1 dose of study drug	102	101
Completed	45	37
Not completed	57	64
Parent/Caregiver Decision	5	5
Physician decision	2	5
Consent withdrawn by subject	11	14
Clinical Relapse	-	1
Adverse event, non-fatal	19	15
Protocol Violation	-	2
Treatment Non-compliance	3	6
Lost to follow-up	5	11
Entry Criteria Not Met	2	1
Lack of efficacy	10	4

## Baseline characteristics

### Reporting groups

Reporting group title	Olanzapine/standard behavioral weight intervention
-----------------------	--

Reporting group description:

Olanzapine: 2.5 milligrams (mg) to 20 mg given orally, daily for 52 weeks.

Standard behavioral weight intervention: One time counseling and basic counseling information on healthy eating and exercise habits at randomization visit only

Reporting group title	Olanzapine/intense behavioral weight intervention
-----------------------	---

Reporting group description:

Olanzapine: 2.5 mg to 20 mg given orally, daily for 52 weeks.

Intense behavioral weight intervention: Counseling provided at randomization and at all subsequent study visits by appropriately trained individual regarding healthy lifestyle habits. Participants also provided with simple tools to help enable healthy eating and exercise habits

Reporting group values	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention	Total
Number of subjects	102	101	203
Age categorical			
Units: Subjects			
Adolescents (12-17 years)	102	101	203
Age Continuous			
Units: years			
arithmetic mean	15.86	15.66	
standard deviation	± 1.49	± 1.52	-
Gender, Male/Female			
Units: participants			
Female	50	47	97
Male	52	54	106
Ethnicity (NIH/OMB)			
Units: Subjects			
Hispanic or Latino	15	9	24
Not Hispanic or Latino	87	92	179
Unknown or Not Reported	0	0	0
Race (NIH/OMB)			
Units: Subjects			
American Indian or Alaska Native	0	1	1
Asian	2	1	3
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	12	14	26
White	84	80	164
More than one race	4	5	9
Unknown or Not Reported	0	0	0
Region of Enrollment			
Units: Subjects			
Germany	2	1	3
Poland	6	12	18
Russian Federation	28	27	55
United States	53	54	107

Puerto Rico	13	7	20
-------------	----	---	----

---

## End points

### End points reporting groups

Reporting group title	Olanzapine/standard behavioral weight intervention
Reporting group description: Olanzapine: 2.5 milligrams (mg) to 20 mg given orally, daily for 52 weeks. Standard behavioral weight intervention: One time counseling and basic counseling information on healthy eating and exercise habits at randomization visit only	
Reporting group title	Olanzapine/intense behavioral weight intervention
Reporting group description: Olanzapine: 2.5 mg to 20 mg given orally, daily for 52 weeks. Intense behavioral weight intervention: Counseling provided at randomization and at all subsequent study visits by appropriately trained individual regarding healthy lifestyle habits. Participants also provided with simple tools to help enable healthy eating and exercise habits	

### Primary: Mean Change From Baseline to 52 Weeks in Body Mass Index (BMI) for All Participants

End point title	Mean Change From Baseline to 52 Weeks in Body Mass Index (BMI) for All Participants
End point description:	
End point type	Primary
End point timeframe: Baseline, 52 weeks	

End point values	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	102 <sup>[1]</sup>	101 <sup>[2]</sup>		
Units: kilograms per meter squared (kg/m <sup>2</sup> )				
least squares mean (standard error)	3.64 (± 0.39)	2.83 (± 0.4)		

Notes:

[1] - Randomized subjects who received ≥ 1 dose of study drug. Mixed model repeated measures (MMRM)

[2] - Randomized subjects who received ≥ 1 dose of study drug. Mixed model repeated measures (MMRM)

### Statistical analyses

Statistical analysis title	Statistical Analysis 1 for End Point 1
Comparison groups	Olanzapine/intense behavioral weight intervention v Olanzapine/standard behavioral weight intervention

Number of subjects included in analysis	203
Analysis specification	Pre-specified
Analysis type	superiority <sup>[3]</sup>
P-value	= 0.15
Method	Mixed models analysis

Notes:

[3] - Mixed model repeated measures analysis terms included baseline BMI, baseline age, gender, intervention group, visit, region, intervention group\*visit.

### Secondary: Mean change from baseline to Endpoint in Body Mass Index (BMI) for participants with duration of treatment of at least 6 months

End point title	Mean change from baseline to Endpoint in Body Mass Index (BMI) for participants with duration of treatment of at least 6 months
-----------------	---

End point description:

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 52 weeks

End point values	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	54 <sup>[4]</sup>	55 <sup>[5]</sup>		
Units: kg/m <sup>2</sup>				
least squares mean (standard error)	3.36 (± 0.41)	2.99 (± 0.4)		

Notes:

[4] - Received ≥ 1 dose of study drug with ≥ 6 months data. Last observation carried forward (LOCF)

[5] - Received ≥ 1 dose of study drug with ≥ 6 months data. Last observation carried forward (LOCF)

### Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1 for End Point 2
Comparison groups	Olanzapine/intense behavioral weight intervention v Olanzapine/standard behavioral weight intervention
Number of subjects included in analysis	109
Analysis specification	Pre-specified
Analysis type	superiority <sup>[6]</sup>
P-value	= 0.52
Method	ANCOVA

Notes:

[6] - ANCOVA model terms included: baseline, baseline age, gender, intervention group, and region.

### Secondary: Time to event for 7%, 15%, and 25% weight gain for all participants

End point title	Time to event for 7%, 15%, and 25% weight gain for all participants
-----------------	---

End point description:

Kaplan-Meier methodology used to estimate time to event. Participants who never reached the target weight gain contributed to the set of patients at risk up to the point at which they discontinued from the

study and were then censored (i.e., removed from the risk set).

65 participants (32%; 34% in Standard Group [SG], 30% in Intense Group [IG]) did not meet 7% weight gain criterion [WGC] by the time they discontinued. 122 participants (60%; 58% in SG, 62% in IG) did not meet 15% WGC by the time they discontinued. 161 participants (79%; 76% in SG, 82% in IG) did not meet 25% WGC by the time they discontinued.

End point type	Secondary
End point timeframe:	
Baseline up to 52 weeks	

<b>End point values</b>	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	102 <sup>[7]</sup>	101 <sup>[8]</sup>		
Units: days				
median (confidence interval 95%)				
Time to First 7% Weight Gain	57 (44 to 84)	57 (44 to 84)		
Time to First 15% Weight Gain	197 (140 to 359)	198 (141 to 999999)		
Time to First 25% Weight Gain	999999 (999999 to 999999)	99999999 (99999999 to 99999999)		

Notes:

[7] - Insufficient # of events of 25% weight gain in the Standard group to compute.

[8] - Insufficient # of events of 15% and 25% weight gain in the Intense group to compute.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Mean change from baseline to 52 weeks in Adolescent Structured YoungMania Rating Scale (YMRS) for participants with Bipolar I disorder

End point title	Mean change from baseline to 52 weeks in Adolescent Structured YoungMania Rating Scale (YMRS) for participants with Bipolar I disorder
-----------------	--

End point description:

The YMRS is an 11-item scale that measures the severity of manic episodes. Four items are rated on a scale from 0 (symptom not present) to 8 (symptom extremely severe). The remaining items are rated on a scale from 0 (symptom not present) to 4 (symptom extremely severe). The YMRS total score ranges from 0 (symptom not present) to 60 (symptom extremely severe).

End point type	Secondary
End point timeframe:	
Baseline, 52 weeks	

<b>End point values</b>	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	58 <sup>[9]</sup>	58 <sup>[10]</sup>		
Units: units on a scale				
least squares mean (standard error)	-17.66 (± 1.75)	-12.05 (± 1.92)		

Notes:

[9] - Subjects with bipolar I disorder, who received ≥ 1 dose of study drug. MMRM used

[10] - Subjects with bipolar I disorder, who received ≥ 1 dose of study drug. MMRM used

### Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1 for End Point 4
Comparison groups	Olanzapine/standard behavioral weight intervention v Olanzapine/intense behavioral weight intervention
Number of subjects included in analysis	116
Analysis specification	Pre-specified
Analysis type	superiority <sup>[11]</sup>
P-value	= 0.008
Method	Mixed models analysis

Notes:

[11] - MMRM analysis terms included baseline, intervention group, visit, region, and intervention group visit.

### Secondary: Mean Clinical Global Impression of Improvement (CGI-I) at 52 weeks for All Participants

End point title	Mean Clinical Global Impression of Improvement (CGI-I) at 52 weeks for All Participants
-----------------	---

End point description:

The Clinical Global Impression of Improvement (CGI-I) is used by the clinician to record the improvement of illness at the time of assessment. The score ranges from 1 (very much improved) to 7 (very much worse).

End point type	Secondary
----------------	-----------

End point timeframe:

52 weeks

<b>End point values</b>	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	102 <sup>[12]</sup>	101 <sup>[13]</sup>		
Units: units on a scale				
least squares mean (standard error)	2.04 (± 0.15)	2.29 (± 0.16)		

Notes:

[12] - Subjects who received ≥ 1 dose of study drug. MMRM used

[13] - Subjects who received ≥ 1 dose of study drug. MMRM used

## Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1 for End Point 5
Comparison groups	Olanzapine/intense behavioral weight intervention v Olanzapine/standard behavioral weight intervention
Number of subjects included in analysis	203
Analysis specification	Pre-specified
Analysis type	superiority <sup>[14]</sup>
P-value	= 0.266
Method	Mixed models analysis

Notes:

[14] - MMRM analysis terms included intervention group, visit, region, and intervention group visit.

---

### Secondary: Mean change from baseline to 52 weeks in waist circumference for all participants

End point title	Mean change from baseline to 52 weeks in waist circumference for all participants
-----------------	---

End point description:

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 52 weeks

<b>End point values</b>	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	102 <sup>[15]</sup>	101 <sup>[16]</sup>		
Units: centimeters (cm)				
least squares mean (standard error)	7.22 (± 1.06)	7.31 (± 1.11)		

Notes:

[15] - Subjects who received ≥ 1 dose of study drug.

[16] - Subjects who received ≥ 1 dose of study drug.

## Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1 for End Point 6
Comparison groups	Olanzapine/intense behavioral weight intervention v Olanzapine/standard behavioral weight intervention
Number of subjects included in analysis	203
Analysis specification	Pre-specified
Analysis type	superiority <sup>[17]</sup>
P-value	= 0.954
Method	Mixed models analysis

Notes:

[17] - MMRM analysis terms included baseline, baseline age, gender, intervention group, visit, region, and intervention group visit.

---

### Secondary: Mean change from baseline to 52 weeks in Clinical Global Impression - Severity (CGI-S) for All Participants

End point title	Mean change from baseline to 52 weeks in Clinical Global Impression - Severity (CGI-S) for All Participants
End point description:	The CGI-S is used by the clinician to record the severity of illness at the time of assessment. The score ranges from 1 = normal, not at all ill to 7 = among the most extremely ill.
End point type	Secondary
End point timeframe:	Baseline, 52 weeks

End point values	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	102 <sup>[18]</sup>	101 <sup>[19]</sup>		
Units: units on a scale				
least squares mean (standard error)	-2.06 (± 0.14)	-1.74 (± 0.14)		

Notes:

[18] - Subjects who received ≥ 1 dose of study drug. MMRM used

[19] - Subjects who received ≥ 1 dose of study drug. MMRM used

### Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1 for End Point 7
Comparison groups	Olanzapine/intense behavioral weight intervention v Olanzapine/standard behavioral weight intervention
Number of subjects included in analysis	203
Analysis specification	Pre-specified
Analysis type	superiority <sup>[20]</sup>
P-value	= 0.103
Method	Mixed models analysis

Notes:

[20] - MMRM analysis terms included baseline, intervention group, visit, region, and intervention group\*visit.

### Secondary: Mean change from baseline to 52 weeks in Anchored Version of the Brief Psychiatric Rating Scale for Children (BPRS-C) for participants with schizophrenia

End point title	Mean change from baseline to 52 weeks in Anchored Version of the Brief Psychiatric Rating Scale for Children (BPRS-C) for participants with schizophrenia
End point description:	The BPRS-C characterizes psychopathology. A total of 21 items are rated on a scale from 0 (not present) to 6 (extremely severe) with a total score ranging from 0 to 126. A decrease in score indicates a reduction in psychopathology.
End point type	Secondary
End point timeframe:	Baseline, 52 weeks

<b>End point values</b>	Olanzapine/standard behavioral weight intervention	Olanzapine/intense behavioral weight intervention		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	44 <sup>[21]</sup>	43 <sup>[22]</sup>		
Units: units on a scale				
least squares mean (standard error)	-28 (± 2.83)	-30.96 (± 2.67)		

Notes:

[21] - Subjects with schizophrenia who received ≥ 1 dose of study drug. MMRM used

[22] - Subjects with schizophrenia who received ≥ 1 dose of study drug. MMRM used

### Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1 for End Point 8
Comparison groups	Olanzapine/standard behavioral weight intervention v Olanzapine/intense behavioral weight intervention
Number of subjects included in analysis	87
Analysis specification	Pre-specified
Analysis type	superiority <sup>[23]</sup>
P-value	= 0.436
Method	Mixed models analysis

Notes:

[23] - MMRM analysis terms included baseline, intervention group, visit, region, and intervention group visit.

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Entire Study

Adverse event reporting additional description:

F1D-MC-HGMX

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	16.0
--------------------	------

### Reporting groups

Reporting group title	Olanzapine/intense behavioral weight intervention
-----------------------	---

Reporting group description: -

Reporting group title	Olanzapine/standard behavioral weight intervention
-----------------------	--

Reporting group description: -

<b>Serious adverse events</b>	Olanzapine/intense behavioral weight intervention	Olanzapine/standard behavioral weight intervention	
Total subjects affected by serious adverse events			
subjects affected / exposed	20 / 101 (19.80%)	13 / 102 (12.75%)	
number of deaths (all causes)	0	0	
number of deaths resulting from adverse events	0	0	
Injury, poisoning and procedural complications			
acetabulum fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
clavicle fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
craniocerebral injury			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
fall			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
facial bones fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
hip fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
humerus fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
jaw fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
multiple injuries			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

overdose alternative dictionary used: MedDRA 16.0 subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
rib fracture alternative dictionary used: MedDRA 16.0 subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
traumatic liver injury alternative dictionary used: MedDRA 16.0 subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
traumatic lung injury alternative dictionary used: MedDRA 16.0 subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders neuroleptic malignant syndrome alternative dictionary used: MedDRA 16.0 subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
General disorders and administration site conditions irritability alternative dictionary used: MedDRA 16.0 subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			

gastroduodenitis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 101 (0.99%) 1 / 1 0 / 0	0 / 102 (0.00%) 0 / 0 0 / 0	
Respiratory, thoracic and mediastinal disorders pneumothorax alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 101 (0.00%) 0 / 0 0 / 0	1 / 102 (0.98%) 0 / 1 0 / 0	
Psychiatric disorders abnormal behaviour alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 101 (0.99%) 0 / 2 0 / 0	0 / 102 (0.00%) 0 / 0 0 / 0	
aggression alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 101 (0.99%) 0 / 1 0 / 0	0 / 102 (0.00%) 0 / 0 0 / 0	
bipolar i disorder alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	4 / 101 (3.96%) 0 / 4 0 / 0	0 / 102 (0.00%) 0 / 0 0 / 0	
bipolar disorder alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	2 / 101 (1.98%) 0 / 2 0 / 0	0 / 102 (0.00%) 0 / 0 0 / 0	
drug abuse			

alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
hallucination, auditory			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
mental disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
psychotic disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	4 / 101 (3.96%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	2 / 5	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
schizophrenia, paranoid type			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
schizophrenia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
suicide attempt			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	0 / 101 (0.00%)	2 / 102 (1.96%)	
occurrences causally related to treatment / all	0 / 0	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
suicidal ideation			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 101 (1.98%)	3 / 102 (2.94%)	
occurrences causally related to treatment / all	1 / 2	3 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
gastroenteritis viral			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 0 %

<b>Non-serious adverse events</b>	Olanzapine/intense behavioral weight intervention	Olanzapine/standard behavioral weight intervention	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	81 / 101 (80.20%)	81 / 102 (79.41%)	
Vascular disorders			
haematoma			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
hypotension			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)	
occurrences (all)	1	1	
Surgical and medical procedures			
cyst removal			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	

wisdom teeth removal alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
Pregnancy, puerperium and perinatal conditions pregnancy alternative dictionary used: MedDRA 16.0 subjects affected / exposed <sup>[1]</sup> occurrences (all)	1 / 47 (2.13%) 1	1 / 50 (2.00%) 1	
General disorders and administration site conditions asthenia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  chills alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  chest pain alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  fatigue alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  gait disturbance alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  influenza like illness alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	2 / 101 (1.98%) 2  2 / 101 (1.98%) 2  0 / 101 (0.00%) 0  6 / 101 (5.94%) 6  0 / 101 (0.00%) 0  1 / 101 (0.99%) 1	3 / 102 (2.94%) 3  2 / 102 (1.96%) 2  2 / 102 (1.96%) 2  9 / 102 (8.82%) 10  1 / 102 (0.98%) 1  0 / 102 (0.00%) 0	

irritability alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	3 / 101 (2.97%) 5	5 / 102 (4.90%) 5	
oedema peripheral alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
pain alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	1 / 102 (0.98%) 2	
pyrexia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	3 / 101 (2.97%) 3	4 / 102 (3.92%) 4	
thirst alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
Immune system disorders seasonal allergy alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	2 / 101 (1.98%) 3	2 / 102 (1.96%) 2	
Reproductive system and breast disorders amenorrhoea alternative dictionary used: MedDRA 16.0 subjects affected / exposed <sup>[2]</sup> occurrences (all)	1 / 47 (2.13%) 1	2 / 50 (4.00%) 2	
breast pain alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
breast enlargement			

alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences (all)	0	1	
dysmenorrhoea			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed <sup>[3]</sup>	1 / 47 (2.13%)	3 / 50 (6.00%)	
occurrences (all)	4	18	
menstruation irregular			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed <sup>[4]</sup>	0 / 47 (0.00%)	1 / 50 (2.00%)	
occurrences (all)	0	1	
metrorrhagia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed <sup>[5]</sup>	0 / 47 (0.00%)	1 / 50 (2.00%)	
occurrences (all)	0	1	
premenstrual cramps			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed <sup>[6]</sup>	0 / 47 (0.00%)	1 / 50 (2.00%)	
occurrences (all)	0	1	
Respiratory, thoracic and mediastinal disorders			
asthma			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
cough			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	3 / 101 (2.97%)	7 / 102 (6.86%)	
occurrences (all)	5	8	
dyspnoea			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
epistaxis			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	5 / 101 (4.95%)	2 / 102 (1.96%)
occurrences (all)	6	2
nasal congestion		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	4 / 101 (3.96%)	4 / 102 (3.92%)
occurrences (all)	4	4
oropharyngeal pain		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	4 / 101 (3.96%)	4 / 102 (3.92%)
occurrences (all)	5	4
pharyngeal inflammation		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
respiratory disorder		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
respiratory tract congestion		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
rhinitis allergic		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1
rhinorrhoea		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1
sinus congestion		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	2 / 102 (1.96%)
occurrences (all)	1	2

throat tightness alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
Psychiatric disorders abnormal behaviour alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	2 / 102 (1.96%) 2	
affect lability alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
affective disorder alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
aggression alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	2 / 101 (1.98%) 3	1 / 102 (0.98%) 1	
agitation alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	2 / 102 (1.96%) 2	
anhedonia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
anxiety alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	3 / 101 (2.97%) 4	6 / 102 (5.88%) 6	
delusion alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	2
depressed mood		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1
depression		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	2 / 102 (1.96%)
occurrences (all)	2	2
dermatillomania		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
distractibility		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
dysphoria		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
emotional distress		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
emotional poverty		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
hallucination, visual		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1

hallucination		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	2 / 102 (1.96%)
occurrences (all)	0	2
impulsive behaviour		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
initial insomnia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	2 / 102 (1.96%)
occurrences (all)	1	2
insomnia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	5 / 101 (4.95%)	5 / 102 (4.90%)
occurrences (all)	5	6
intentional self-injury		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	4 / 102 (3.92%)
occurrences (all)	1	5
middle insomnia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
mood altered		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
mood swings		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
nightmare		
alternative dictionary used: MedDRA 16.0		

subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1
obsessive thoughts		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
panic attack		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	2	0
psychotic disorder		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
restlessness		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	3 / 101 (2.97%)	4 / 102 (3.92%)
occurrences (all)	4	4
schizophrenia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	2 / 102 (1.96%)
occurrences (all)	2	2
self injurious behaviour		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
stereotypy		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
thought blocking		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1

tachyphrenia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
Investigations			
alanine aminotransferase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	3 / 101 (2.97%) 3	3 / 102 (2.94%) 3	
aspartate aminotransferase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	3 / 102 (2.94%) 3	
blood alkaline phosphatase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
blood cholesterol increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	2 / 102 (1.96%) 2	
blood creatine phosphokinase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	5 / 101 (4.95%) 6	7 / 102 (6.86%) 9	
blood glucose decreased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
blood glucose increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
blood iron decreased			

alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
blood insulin increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	7 / 101 (6.93%)	10 / 102 (9.80%)
occurrences (all)	7	11
blood insulin decreased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	2 / 102 (1.96%)
occurrences (all)	2	2
blood pressure increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
blood prolactin increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	4 / 102 (3.92%)
occurrences (all)	2	4
blood triglycerides increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	5 / 101 (4.95%)	4 / 102 (3.92%)
occurrences (all)	5	5
body mass index increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
body temperature increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	3 / 102 (2.94%)
occurrences (all)	1	4
cardiac murmur		
alternative dictionary used: MedDRA 16.0		

subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
eosinophil count increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	3 / 102 (2.94%)
occurrences (all)	1	3
erythroblast morphology abnormal		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
gamma-glutamyltransferase increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
glycosylated haemoglobin increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
hepatic enzyme increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
heart rate increased		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
hepatitis c antibody positive		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
liver function test abnormal		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	2 / 102 (1.96%)
occurrences (all)	0	2

low density lipoprotein increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 3	0 / 102 (0.00%) 0	
neutrophil count decreased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
nitrite urine present alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 2	0 / 102 (0.00%) 0	
transaminases increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
waist circumference increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
weight increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	21 / 101 (20.79%) 21	31 / 102 (30.39%) 31	
white blood cell count decreased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
Injury, poisoning and procedural complications arthropod bite alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	1 / 102 (0.98%) 1	
concussion alternative dictionary used:			

MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
contusion		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	1 / 102 (0.98%)
occurrences (all)	2	1
face injury		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
foot fracture		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	2	0
hand fracture		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
intentional overdose		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	2 / 102 (1.96%)
occurrences (all)	3	2
joint injury		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
laceration		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
ligament sprain		
alternative dictionary used: MedDRA 16.0		

subjects affected / exposed	0 / 101 (0.00%)	3 / 102 (2.94%)	
occurrences (all)	0	3	
limb injury			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	3 / 101 (2.97%)	0 / 102 (0.00%)	
occurrences (all)	4	0	
meniscus injury			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
nerve injury			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
scar			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
sunburn			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences (all)	0	1	
wound			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
wrist fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
Cardiac disorders			
sinus tachycardia			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
tachycardia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	5 / 102 (4.90%)	
occurrences (all)	1	5	
<b>Nervous system disorders</b>			
akathisia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	4 / 101 (3.96%)	5 / 102 (4.90%)	
occurrences (all)	5	6	
bradykinesia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
convulsion			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 101 (1.98%)	0 / 102 (0.00%)	
occurrences (all)	3	0	
dizziness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 101 (1.98%)	7 / 102 (6.86%)	
occurrences (all)	2	7	
drooling			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	2 / 102 (1.96%)	
occurrences (all)	0	2	
dyskinesia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	2 / 102 (1.96%)	
occurrences (all)	2	2	
disturbance in attention			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	1 / 101 (0.99%)	2 / 102 (1.96%)
occurrences (all)	1	2
extrapyramidal disorder		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	3 / 102 (2.94%)
occurrences (all)	2	7
headache		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	20 / 101 (19.80%)	19 / 102 (18.63%)
occurrences (all)	30	49
head titubation		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
hypersomnia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	4 / 102 (3.92%)
occurrences (all)	2	5
lethargy		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
migraine		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1
petit mal epilepsy		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
restless legs syndrome		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1

sedation alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	3 / 101 (2.97%) 5	8 / 102 (7.84%) 8	
somnolence alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	24 / 101 (23.76%) 28	19 / 102 (18.63%) 22	
tremor alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	5 / 101 (4.95%) 5	3 / 102 (2.94%) 3	
<b>Blood and lymphatic system disorders</b>			
agranulocytosis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
leukopenia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	4 / 101 (3.96%) 4	2 / 102 (1.96%) 2	
lymphadenitis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
lymphocytosis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
lymphadenopathy alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
neutropenia alternative dictionary used: MedDRA 16.0			

subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
Ear and labyrinth disorders			
ear disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
ear pain			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	2 / 102 (1.96%) 2	
external ear disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
vertigo			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
Eye disorders			
blepharospasm			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	1 / 102 (0.98%) 1	
conjunctivitis			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	2 / 102 (1.96%) 2	
eye discharge			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
eye haemorrhage			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
lacrimation increased alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
<b>Gastrointestinal disorders</b>			
abdominal pain upper alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	4 / 101 (3.96%) 4	5 / 102 (4.90%) 5	
abdominal pain alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	2 / 101 (1.98%) 2	0 / 102 (0.00%) 0	
abdominal discomfort alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	4 / 102 (3.92%) 6	
abnormal faeces alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	0 / 102 (0.00%) 0	
anal fissure alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
constipation alternative dictionary used: MedDRA 16.0			
subjects affected / exposed occurrences (all)	4 / 101 (3.96%) 4	1 / 102 (0.98%) 1	
diarrhoea alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	3 / 101 (2.97%)	5 / 102 (4.90%)
occurrences (all)	5	6
dry mouth		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	4 / 101 (3.96%)	6 / 102 (5.88%)
occurrences (all)	4	6
dysphagia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
dyspepsia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	3 / 102 (2.94%)
occurrences (all)	2	3
gastrooesophageal reflux disease		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1
glossitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
lip disorder		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
nausea		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	4 / 101 (3.96%)	3 / 102 (2.94%)
occurrences (all)	4	3
salivary hypersecretion		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	0 / 102 (0.00%)
occurrences (all)	2	0

toothache alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	3 / 102 (2.94%) 3	
vomiting alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	4 / 101 (3.96%) 7	7 / 102 (6.86%) 8	
Skin and subcutaneous tissue disorders			
acne alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	4 / 102 (3.92%) 4	
alopecia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
hyperhidrosis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
pruritus alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	2 / 101 (1.98%) 2	0 / 102 (0.00%) 0	
pruritus generalised alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 101 (0.00%) 0	1 / 102 (0.98%) 1	
rash alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 101 (0.99%) 1	3 / 102 (2.94%) 5	
seborrhoeic dermatitis alternative dictionary used: MedDRA 16.0			

<p>subjects affected / exposed occurrences (all)</p> <p>skin irritation alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)</p> <p>urticaria alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)</p>	<p>0 / 101 (0.00%) 0</p> <p>0 / 101 (0.00%) 0</p> <p>1 / 101 (0.99%) 2</p>	<p>1 / 102 (0.98%) 1</p> <p>1 / 102 (0.98%) 1</p> <p>0 / 102 (0.00%) 0</p>	
<p>Renal and urinary disorders</p> <p>dysuria alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)</p> <p>enuresis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)</p> <p>pollakiuria alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)</p>	<p>1 / 101 (0.99%) 1</p> <p>0 / 101 (0.00%) 0</p> <p>1 / 101 (0.99%) 1</p>	<p>0 / 102 (0.00%) 0</p> <p>1 / 102 (0.98%) 1</p> <p>0 / 102 (0.00%) 0</p>	
<p>Musculoskeletal and connective tissue disorders</p> <p>arthralgia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)</p> <p>back pain alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)</p> <p>joint stiffness alternative dictionary used: MedDRA 16.0</p>	<p>2 / 101 (1.98%) 2</p> <p>1 / 101 (0.99%) 1</p>	<p>0 / 102 (0.00%) 0</p> <p>2 / 102 (1.96%) 2</p>	

subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)	
occurrences (all)	1	1	
medial tibial stress syndrome			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	1	0	
muscle rigidity			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences (all)	0	1	
muscle spasms			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 101 (1.98%)	1 / 102 (0.98%)	
occurrences (all)	3	1	
musculoskeletal pain			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences (all)	0	1	
musculoskeletal stiffness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	3 / 101 (2.97%)	2 / 102 (1.96%)	
occurrences (all)	3	2	
myalgia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)	
occurrences (all)	1	1	
pain in extremity			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	2 / 102 (1.96%)	
occurrences (all)	0	2	
Infections and infestations			
acute tonsillitis			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
bacterial infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
bronchitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	4 / 101 (3.96%)	3 / 102 (2.94%)
occurrences (all)	5	3
chronic tonsillitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
cystitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	0 / 102 (0.00%)
occurrences (all)	2	0
ear infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	1 / 102 (0.98%)
occurrences (all)	2	1
gastroenteritis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	2
gastroenteritis rotavirus		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
gastroenteritis viral		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	3 / 102 (2.94%)
occurrences (all)	1	3

herpes zoster		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
influenza		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	3 / 102 (2.94%)
occurrences (all)	2	3
localised infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
lower respiratory tract infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
nasopharyngitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	11 / 101 (10.89%)	14 / 102 (13.73%)
occurrences (all)	11	16
otitis externa		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
pharyngitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	1 / 102 (0.98%)
occurrences (all)	1	1
pharyngitis streptococcal		
alternative dictionary used: MedDRA 16.0		

subjects affected / exposed	1 / 101 (0.99%)	4 / 102 (3.92%)
occurrences (all)	1	4
pneumonia		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
respiratory tract infection viral		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	1 / 102 (0.98%)
occurrences (all)	2	1
sinusitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	3 / 102 (2.94%)
occurrences (all)	1	4
tinea pedis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)
occurrences (all)	0	1
tonsillitis		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)
occurrences (all)	1	0
upper respiratory tract infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	2 / 101 (1.98%)	4 / 102 (3.92%)
occurrences (all)	2	4
urinary tract infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed	3 / 101 (2.97%)	2 / 102 (1.96%)
occurrences (all)	3	3
vulvovaginal mycotic infection		
alternative dictionary used: MedDRA 16.0		
subjects affected / exposed <sup>[7]</sup>	0 / 47 (0.00%)	1 / 50 (2.00%)
occurrences (all)	0	1

Metabolism and nutrition disorders			
decreased appetite			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 101 (1.98%)	2 / 102 (1.96%)	
occurrences (all)	2	2	
hypercholesterolaemia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 101 (0.99%)	0 / 102 (0.00%)	
occurrences (all)	2	0	
hyperinsulinaemia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences (all)	0	1	
hyperlipidaemia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	2 / 102 (1.96%)	
occurrences (all)	0	2	
hypertriglyceridaemia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 101 (1.98%)	2 / 102 (1.96%)	
occurrences (all)	2	2	
increased appetite			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	13 / 101 (12.87%)	16 / 102 (15.69%)	
occurrences (all)	14	16	
insulin resistance			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 101 (0.00%)	1 / 102 (0.98%)	
occurrences (all)	0	1	
obesity			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 101 (1.98%)	2 / 102 (1.96%)	
occurrences (all)	2	2	

Notes:

[1] - The number of subjects exposed to this adverse event is less than the total number of subjects

exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

[2] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

[3] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

[4] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

[5] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

[6] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

[7] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

## **More information**

### **Substantial protocol amendments (globally)**

Were there any global substantial amendments to the protocol? No

---

### **Interruptions (globally)**

Were there any global interruptions to the trial? No

### **Limitations and caveats**

None reported