

**Table 50. Modified Lysholm score (ITT population)**

Dose group: Visit	<b>Low</b> N = 24	<b>Medium</b> N = 25	<b>High</b> N = 24	<b>All</b> N = 73
<b>Values at each visit</b>				
Pre-arthroscopy (baseline)	17.6	17.5	15.8	16.9
Pre-implantation (Day 0')	17.6	17.3	15.0	16.6
Visit 1 (6 weeks)	14.8	14.2	14.0	14.3
Visit 2 (3 months)	18.2	17.8	16.7	17.6
Visit 3 (6 months)	20.9	20.1	17.6	19.5
Visit 4 (12 months)	20.5	20.5	18.5	19.8
Visit 5 (18 months)	21.1	21.0	18.8	20.3
Visit 6 (24 months)	20.7	21.1	18.5	20.1
Visit 7 (36 months)	21.3	21.9	18.7	20.8
Visit 8 (48 months)	21.5	21.8	18.9	20.8
Visit 9 (60 months)	20.7	21.9	19.2	20.6
<b>Changes from baseline</b>				
Pre-implantation (Day 0')	0.0	-0.2	-0.8	-0.3
Visit 1 (6 weeks)	-2.8	-3.2	-1.8	-2.6
Visit 2 (3 months)	0.6	0.4	0.9	0.6
Visit 3 (6 months)	3.3	2.6	1.8	2.6
Visit 4 (12 months)	2.9	3.0	2.8	2.9
Visit 5 (18 months)	3.5	3.5	3.0	3.3
Visit 6 (24 months)	3.1	3.6	2.7	3.2
Visit 7 (36 months)	3.7	4.4	2.9	3.7
Visit 8 (48 months)	3.9	4.4	3.2	3.8
Visit 9 (60 months)	3.1	4.4	3.4	3.7
<b>Changes from Day 0'</b>				
Visit 1 (6 weeks)	-2.8	-3.0	-1.0	-2.3
Visit 2 (3 months)	0.6	0.6	1.7	0.9
Visit 3 (6 months)	3.3	2.8	2.6	2.9
Visit 4 (12 months)	2.9	3.2	3.5	3.2
Visit 5 (18 months)	3.5	3.7	3.8	3.7
Visit 6 (24 months)	3.0	3.8	3.5	3.5
Visit 7 (36 months)	3.6	4.6	3.7	4.0
Visit 8 (48 months)	3.8	4.6	4.0	4.1
Visit 9 (60 months)	3.1	4.6	4.2	4.0

Mean values are shown throughout. Source: Tables 14.2.2.16.1 and 14.2.2.16.3.