

**Table 20. KOOS ‘pain’ at each visit (ITT population)**

Visit	Dose group:	<b>Low</b> N = 24	<b>Medium</b> N = 25	<b>High</b> N = 24	<b>All</b> N = 73
<b>Values at each visit</b>					
Pre-arthroscopy (baseline)	mean ± SD	68.2 ± 14.6	66.0 ± 18.9	55.3 ± 19.7	63.2 ± 18.5
	median	71	67	56	67
Visit 1 (6 weeks)	mean ± SD	70.4 ± 19.6	68.0 ± 18.1	63.4 ± 18.8	67.3 ± 18.8
	median	74	64	63	67
Visit 2 (3 months)	mean ± SD	77.4 ± 13.6	75.8 ± 11.6	64.7 ± 18.3	72.7 ± 15.6
	median	79	75	65	72
Visit 3 (6 months)	mean ± SD	83.0 ± 11.7	82.1 ± 12.5	68.7 ± 21.6	78.0 ± 16.9
	median	86	86	71	81
Visit 4 (12 months)	mean ± SD	84.1 ± 12.6	85.0 ± 12.9	69.3 ± 23.2	79.6 ± 18.2
	median	86	92	67	86
Visit 5 (18 months)	mean ± SD	84.8 ± 11.7	88.8 ± 9.5	69.3 ± 22.7	81.1 ± 17.6
	median	89	92	67	86
Visit 6 (24 months)	mean ± SD	82.9 ± 16.1	86.8 ± 14.3	67.2 ± 20.9	79.1 ± 19.0
	median	90	92	67	83
Visit 7 (36 months)	mean ± SD	84.8 ± 13.8	89.8 ± 11.8	69.1 ± 22.8	81.4 ± 18.8
	median	90	94	69	89
Visit 8 (48 months)	mean ± SD	84.5 ± 17.0	90.1 ± 13.5	70.9 ± 22.8	82.0 ± 19.6
	median	90	94	74	89
Visit 9 (60 months)	mean ± SD	83.9 ± 13.6	90.6 ± 10.7	69.2 ± 24.8	81.4 ± 19.4
	median	89	94	75	89

Baseline = Day 0. Source: Table 14.2.2.2.1.

**Table 21. KOOS ‘other symptoms’ at each visit (ITT population)**

Visit	Dose group:	<b>Low</b> N = 24	<b>Medium</b> N = 25	<b>High</b> N = 24	<b>All</b> N = 73
<b>Values at each visit</b>					
Pre-arthroscopy (baseline)	mean ± SD	72.5 ± 17.4	72.9 ± 15.9	66.5 ± 14.7	70.6 ± 16.1
	median	75	75	68	75
Visit 1 (6 weeks)	mean ± SD	70.8 ± 18.6	68.1 ± 12.9	66.5 ± 14.5	68.5 ± 15.4
	median	75	64	68	68
Visit 2 (3 months)	mean ± SD	77.1 ± 12.8	77.3 ± 10.8	69.8 ± 15.8	74.8 ± 13.5
	median	77	79	71	75
Visit 3 (6 months)	mean ± SD	85.3 ± 10.1	83.7 ± 11.0	74.1 ± 16.5	81.1 ± 13.6
	median	84	86	75	82
Visit 4 (12 months)	mean ± SD	86.5 ± 10.2	87.0 ± 10.4	76.9 ± 16.8	83.5 ± 13.4
	median	86	89	77	86
Visit 5 (18 months)	mean ± SD	86.0 ± 9.8	88.1 ± 10.6	76.6 ± 18.1	83.7 ± 14.0
	median	86	93	79	86
Visit 6 (24 months)	mean ± SD	87.6 ± 9.2	89.0 ± 10.9	74.9 ± 16.8	83.9 ± 14.1
	median	89	93	73	89
Visit 7 (36 months)	mean ± SD	89.3 ± 8.3	89.9 ± 10.7	76.9 ± 17.5	85.4 ± 13.9
	median	89	93	80	89
Visit 8 (48 months)	mean ± SD	89.9 ± 9.0	87.3 ± 12.3	76.8 ± 17.6	84.7 ± 14.4
	median	91	93	80	89
Visit 9 (60 months)	mean ± SD	86.9 ± 12.2	89.4 ± 12.8	74.6 ± 19.3	83.7 ± 16.3
	median	88	93	77	89

Baseline = Day 0. Source: Table 14.2.2.3.1.

**Table 22. KOOS ‘function in daily living’ at each visit (ITT population)**

Visit	Dose group:	<b>Low</b> N = 24	<b>Medium</b> N = 25	<b>High</b> N = 24	<b>All</b> N = 73
<b>Values at each visit</b>					
Pre-arthroscopy (baseline)	mean ± SD	75.3 ± 16.5	77.0 ± 17.3	68.3 ± 19.4	73.6 ± 17.9
	median	81	87	68	76
Visit 1 (6 weeks)	mean ± SD	61.5 ± 22.1	60.4 ± 16.4	57.1 ± 19.5	59.7 ± 19.2
	median	57	65	53	63
Visit 2 (3 months)	mean ± SD	79.4 ± 14.1	82.9 ± 11.5	72.0 ± 21.4	78.2 ± 16.6
	median	81	88	78	81
Visit 3 (6 months)	mean ± SD	86.8 ± 12.2	88.4 ± 8.2	76.8 ± 20.5	84.0 ± 15.2
	median	90	91	79	87
Visit 4 (12 months)	mean ± SD	89.1 ± 10.8	89.8 ± 9.9	79.0 ± 22.3	86.0 ± 15.9
	median	91	94	84	93
Visit 5 (18 months)	mean ± SD	89.5 ± 11.6	93.2 ± 8.4	78.1 ± 20.3	87.0 ± 15.5
	median	93	96	78	94
Visit 6 (24 months)	mean ± SD	88.2 ± 13.8	91.2 ± 15.1	77.9 ± 18.5	85.8 ± 16.7
	median	93	96	77	91
Visit 7 (36 months)	mean ± SD	88.8 ± 13.2	93.2 ± 11.7	79.4 ± 20.3	87.2 ± 16.3
	median	92	97	80	94
Visit 8 (48 months)	mean ± SD	89.9 ± 13.2	93.8 ± 12.5	79.5 ± 19.4	87.8 ± 16.3
	median	95	99	80	96
Visit 9 (60 months)	mean ± SD	87.3 ± 14.6	93.7 ± 10.7	79.7 ± 21.3	87.0 ± 16.9
	median	93	99	85	96

Baseline = Day 0. Source: Table 14.2.2.4.1.

**Table 23. KOOS ‘function in sport and recreation’ at each visit (ITT population)**

Visit	Dose group:	<b>Low</b> <i>N</i> = 24*	<b>Medium</b> <i>N</i> = 25	<b>High</b> <i>N</i> = 24	<b>All</b> <i>N</i> = 73*
<b>Values at each visit</b>					
Pre-arthroscopy (baseline)	mean ± SD	51.1 ± 21.6	47.8 ± 21.9	38.3 ± 24.5	45.7 ± 23.0
	median	50	50	40	50
Visit 1 (6 weeks)	mean ± SD	28.5 ± 30.8	21.4 ± 23.5	23.1 ± 19.7	24.3 ± 24.9
	median	15	15	23	15
Visit 2 (3 months)	mean ± SD	47.3 ± 25.3	43.0 ± 19.0	44.8 ± 25.8	45.0 ± 23.3
	median	43	40	38	40
Visit 3 (6 months)	mean ± SD	63.3 ± 21.6	64.4 ± 16.1	45.0 ± 25.3	57.7 ± 22.8
	median	65	70	40	60
Visit 4 (12 months)	mean ± SD	69.4 ± 22.4	66.2 ± 20.5	54.2 ± 31.2	63.3 ± 25.6
	median	70	65	48	65
Visit 5 (18 months)	mean ± SD	70.4 ± 19.3	73.8 ± 17.6	52.7 ± 31.2	65.8 ± 24.9
	median	70	80	43	70
Visit 6 (24 months)	mean ± SD	70.4 ± 23.2	71.2 ± 22.4	48.8 ± 29.9	63.6 ± 27.1
	median	73	75	40	70
Visit 7 (36 months)	mean ± SD	74.4 ± 19.6	80.8 ± 15.7	55.0 ± 30.8	70.2 ± 25.1
	median	78	85	58	75
Visit 8 (48 months)	mean ± SD	72.3 ± 21.4	78.4 ± 20.5	55.6 ± 29.4	68.9 ± 25.6
	median	73	85	60	75
Visit 9 (60 months)	mean ± SD	70.2 ± 22.4	82.2 ± 17.5	54.2 ± 32.1	69.0 ± 27.0
	median	73	85	55	75

Baseline = Day 0. [\* The baseline value for overall KOOS of one patient in the low-dose group was missing, because of incomplete response to the ePRO questions. Therefore, *N* = 23 for this group (and *N* = 72 total) at baseline.] Source: Table 14.2.2.5.1

**Table 24. KOOS ‘knee-related quality of life’ at each visit (ITT population)**

Visit	Dose group:	<b>Low</b> N = 24	<b>Medium</b> N = 25	<b>High</b> N = 24	<b>All</b> N = 73
<b>Values at each visit</b>					
Pre-arthroscopy (baseline)	mean ± SD	34.6 ± 16.7	34.3 ± 16.8	26.8 ± 18.6	31.9 ± 17.5
	median	38	31	25	31
Visit 1 (6 weeks)	mean ± SD	31.5 ± 19.5	28.5 ± 15.3	24.5 ± 17.4	28.2 ± 17.5
	median	28	25	25	25
Visit 2 (3 months)	mean ± SD	47.1 ± 24.0	38.3 ± 17.0	38.8 ± 20.4	41.4 ± 20.7
	median	38	44	44	38
Visit 3 (6 months)	mean ± SD	57.8 ± 22.7	48.3 ± 17.7	44.3 ± 24.9	50.1 ± 22.3
	median	56	50	44	50
Visit 4 (12 months)	mean ± SD	60.2 ± 23.9	53.5 ± 19.2	50.0 ± 27.9	54.5 ± 23.9
	median	59	56	44	56
Visit 5 (18 months)	mean ± SD	62.0 ± 24.3	58.0 ± 19.6	45.6 ± 29.5	55.2 ± 25.4
	median	69	63	38	56
Visit 6 (24 months)	mean ± SD	63.5 ± 25.2	59.3 ± 21.9	47.7 ± 26.1	56.8 ± 25.0
	median	66	63	44	56
Visit 7 (36 months)	mean ± SD	63.8 ± 25.1	66.8 ± 19.7	51.6 ± 28.3	60.8 ± 25.1
	median	69	69	50	56
Visit 8 (48 months)	mean ± SD	63.5 ± 25.6	70.5 ± 23.3	51.6 ± 26.2	62.0 ± 25.9
	median	63	75	44	56
Visit 9 (60 months)	mean ± SD	65.4 ± 24.2	71.3 ± 24.5	53.9 ± 28.3	63.6 ± 26.4
	median	69	81	50	63

Baseline = Day 0. Source: Table 14.2.2.6.1.