

Table 31. MOCART scores at Visits 2 and 4–9 (ITT population)

	Visit 2 (3 months)				Visit 4 (12 months)			
	Dose group				Dose group			
	Low N = 20	Medium N = 19	High N = 19	All N = 58	Low N = 22	Medium N = 19	High N = 21	All N = 62
Mean ± SD	59.8 ± 10.9	64.5 ± 10.3	64.7 ± 9.4	62.9 ± 10.3	74.1 ± 13.0	74.5 ± 14.5	68.8 ± 11.4	72.4 ± 13.0
Minimum	45	50	45	45	55	50	45	45
Lower quartile	53	60	60	60	60	60	65	60
Median	60	60	65	60	75	75	70	70
Upper quartile	68	75	70	70	85	90	75	80
Maximum	85	85	90	90	95	100	100	100
	Visit 5 (18 months)				Visit 6 (24 months)			
	N = 23	N = 22	N = 21	N = 66	N = 23	N = 25	N = 21	N = 69
Mean ± SD	72.2 ± 11.7	72.0 ± 17.0	69.0 ± 13.4	71.1 ± 14.0	72.2 ± 11.5	76.0 ± 14.2	71.2 ± 9.7	73.3 ± 12.1
Minimum	50	20	50	20	35	55	55	35
Lower quartile	65	65	60	65	70	65	65	65
Median	70	73	65	70	75	70	70	70
Upper quartile	80	85	75	80	80	90	75	80
Maximum	95	95	100	100	85	100	90	100
	Visit 7 (36 months)				Visit 8 (48 months)			
	N = 23	N = 25	N = 21	N = 69	N = 23	N = 25	N = 21	N = 69
Mean ± SD	72.8 ± 12.0	79.6 ± 13.1	72.6 ± 14.2	75.2 ± 13.4	73.9 ± 13.1	78.0 ± 12.4	74.3 ± 14.0	75.5 ± 13.1
Minimum	50	50	30	30	30	55	30	30
Lower quartile	65	70	65	70	70	70	65	70
Median	70	80	70	75	75	75	75	75
Upper quartile	85	90	85	85	80	90	85	85
Maximum	90	100	95	100	95	100	95	100
	Visit 9 (60 months)							
	Dose group							
	N = 20	N = 19	N = 19	N = 58				
Mean ± SD	75.0 ± 14.9	76.4 ± 14.2	73.6 ± 15.0	75.1 ± 14.5				
Minimum	30	45	30	30				
Lower quartile	70	65	65	65				
Median	75	75	80	75				
Upper quartile	85	90	85	85				
Maximum	100	100	90	100				

Differences in N from total population sizes are due to missing results. Source: Table 14.2.2.7.1.