

9HPT (Dominant hand): Change from Part 1 baseline to each scheduled efficacy visit in Part 2  
 Subgroup Analysis: Confirmed Progression of disability  
 ITT population

Confirmed Progressors

	Placebo to Natalizumab 300 mg	Natalizumab 300 mg	p-value (a)
Change from Baseline to Week 156			
n	158	138	
Mean	10.12	5.01	0.1129
SD	33.675	20.625	

(a) p-value for comparison between the active and placebo groups at Week 156 is based on ANCOVA model, adjusted for baseline EDSS ( $\leq 5.5$  or  $\geq 6$ ) and baseline 9HPT (dominant hand).

9HPT (Dominant hand): Change from Part 1 baseline to each scheduled efficacy visit in Part 2  
 Subgroup Analysis: Confirmed Progression of disability  
 ITT population

Non Progressors

	Placebo to Natalizumab 300 mg	Natalizumab 300 mg	p-value (a)
Change from Baseline to Week 156			
n	99	133	
Mean	-2.60	-0.89	0.2351
SD	9.543	10.602	

(a) p-value for comparison between the active and placebo groups at Week 156 is based on ANCOVA model, adjusted for baseline EDSS (<=5.5 or >=6) and baseline 9HPT (dominant hand).