

<b>Statistical analysis title</b>	Cohort 1, ACP: Fesoterodine 4 mg
Statistical analysis description	P-value was calculated for change from Baseline at Week 12, based on an ANCOVA model with terms for treatment group, baseline number of catheterisations per 24 hours and baseline weight.
Comparison groups	Cohort 1, Active Comparator Phase: Fesoterodine 4 mg
Number of subjects included in analysis	37
Analysis specification	Pre-specified
Analysis type	Superiority
P-value	= 0.0787
Method	ANCOVA
Confidence interval	
<b>Statistical analysis title</b>	Cohort 1, ACP: Fesoterodine 8 mg
Statistical analysis description	P-value was calculated for change from Baseline at Week 12, based on an ANCOVA model with terms for treatment group, baseline number of catheterisations per 24 hours and baseline weight.
Comparison groups	Cohort 1, Active Comparator Phase: Fesoterodine 8 mg

Number of subjects included in analysis	33
Analysis specification	Pre-specified
Analysis type	Superiority
P-value	= 0.0727
Method	ANCOVA
Confidence interval	
<b>Statistical analysis title</b>	Cohort 1, ACP: Oxybutynin
Statistical analysis description	P-value was calculated for change from Baseline at Week 12, based on an ANCOVA model with terms for treatment group, baseline number of catheterisations per 24 hours and baseline weight.
Comparison groups	Cohort 1, Active Comparator Phase: Oxybutynin
Number of subjects included in analysis	31
Analysis specification	Pre-specified
Analysis type	Superiority
P-value	= 0.0666

Method	ANCOVA
Confidence interval	