

Statistical analysis title	Cohort 1, ACP: Fesoterodine 4 mg
Statistical analysis description	P-value was calculated for change from Baseline at Week 12, based on an ANCOVA model with terms for treatment group, baseline detrusor pressure at maximum bladder capacity and baseline weight.
Comparison groups	Cohort 1, Active Comparator Phase: Fesoterodine 4 mg
Number of subjects included in analysis	40
Analysis specification	Pre-specified
Analysis type	Superiority
P-value	= 0.2334
Method	ANCOVA
Confidence interval	
Statistical analysis title	Cohort 1, ACP: Fesoterodine 8 mg
Statistical analysis description	P-value was calculated for change from Baseline at Week 12, based on an ANCOVA model with terms for treatment group, baseline detrusor pressure at maximum bladder capacity and baseline weight.
Comparison groups	Cohort 1, Active Comparator Phase: Fesoterodine 8 mg

Number of subjects included in analysis	41
Analysis specification	Pre-specified
Analysis type	Superiority
P-value	= 0.5087
Method	ANCOVA
Confidence interval	
Statistical analysis title	Cohort 1, ACP: Oxybutynin
Statistical analysis description	P-value was calculated for change from Baseline at Week 12, based on an ANCOVA model with terms for treatment group, baseline detrusor pressure at maximum bladder capacity and baseline weight.
Comparison groups	Cohort 1, Active Comparator Phase: Oxybutynin
Number of subjects included in analysis	38
Analysis specification	Pre-specified
Analysis type	Superiority
P-value	= 0.3333

Method	ANCOVA
Confidence interval	