

Table 38 : Efficacy - Epworth Sleepiness Scale - Change between baseline and M12 – Without replacement of missing values – ITT population

Epworth Scale at Control Visit at 12 months(V7)		Naive patients N=52	Patients CUP/ ATU N=16	Total N=68
Baseline	N	52	16	68
	Missing	0	0	0
	Mean (SD)	17.46 (3.10)	14.50 (3.22)	16.76 (3.35)
	Median	18.00	13.00	17.00
	Q1; Q3	15.50; 19.00	12.50; 16.00	14.00; 19.00
	Min; Max	12.00; 24.00	9.00; 22.00	9.00; 24.00
Control Visit at 12 months (V7)	N	52	16	68
	Missing	0	0	0
	Mean (SD)	12.21 (5.38)	11.88 (5.12)	12.13 (5.28)
	Median	12.50	10.50	12.00
	Q1; Q3	8.00; 17.00	9.00; 14.50	8.50; 17.00
	Min; Max	1; 21	2; 23	1; 23
Change from baseline	N	52	16	68
	Missing	0	0	0
	Mean (SD)	-5.25 (4.84)	-2.63 (4.76)	-4.63 (4.91)
	Median	-5.50	-1.50	-4.00
	Q1; Q3	-8.00; -2.00	-4.50; 1.00	-7.00; -1.00
	Min; Max	-22.0; 4.00	-16.0; 4.00	-22.0; 4.00

Source data: [Section 14, SAR Part I, table 344](#)