

**A RANDOMISED EVALUATION OF LOW-DOSE ARA-C PLUS TOSEDOSTAT  
VERSUS LOW DOSE ARA-C IN OLDER PATIENTS WITH ACUTE MYELOID  
LEUKAEMIA: RESULTS OF THE LI-1 TRIAL**

Mike Dennis<sup>1</sup>, Alan Burnett<sup>2\*</sup>, Robert Hills<sup>3</sup>, Ian Thomas<sup>4</sup>, Cono Ariti<sup>4</sup>, Marianne Tang  
Severinsen<sup>5,6</sup>, Claire Hemmaway<sup>7</sup>, Paul Greaves<sup>8</sup>, Richard E Clark<sup>9</sup>, Mhairi  
Copland<sup>2</sup>, Nigel Russell<sup>10</sup> on behalf of the NCRI AML Working Group

1. Department of Haematology, Christie Hospital NHS Trust, Manchester, UK

2. Paul O’Gorman Leukaemia Centre, Institute of Cancer Sciences, University of  
Glasgow, Glasgow, UK

3. Nuffield Department of Population Health, University of Oxford, Oxford, U.K

4. Centre for Trials Research, Cardiff University, Cardiff, UK

5 Department of Haematology, Clinical Cancer Research Centre, Aalborg  
University Hospital, Aalborg, Denmark,

6 Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

7 Department of Haematology, Auckland City Hospital, Auckland, New Zealand,

8 Department of Haematology, Queen's Hospital, Romford, UK

9 Department of Molecular & Clinical Cancer Medicine, University of Liverpool.  
UK.

10 Department of Haematology, Nottingham University Hospitals, Nottingham,  
UK

**Abstract:** 206 words

**Text:** 3453 words

**Corresponding author:**

Dr Mike Dennis

The Christie Hospital

Wilmslow Road

MANCHESTER

M20 4BX

Tel: 0161 446 3869

Mike.dennis1@nhs.net

\*Formerly Department of Haematology, Cardiff University, Cardiff, U.K.

### **Abstract**

Older patients with acute myeloid leukaemia account for nearly half of those with the disease. Because they are perceived to be unfit for, unwilling to receive, or unlikely to benefit from conventional chemotherapy they represent an important unmet need. Tosedostat is a selective oral aminopeptidase inhibitor, which in phase I/II trials showed acceptable toxicity and encouraging efficacy. We report the only randomised study of low dose ara-C (LDAC) combined with tosedostat (LDAC-T) versus LDAC in untreated older patients not suitable for intensive treatment. 243 patients were randomised 1:1 as part of the "Pick a Winner" LI-1 trial. There was a non-significant increase in CR rate with the addition of tosedostat (LDAC-T vs LDAC; 19% vs 12%, OR 0.61 (0.30-1.23)  $p=0.17$ ). For overall response (CR+CRi), there was little evidence of a benefit to the addition of tosedostat (25% vs 18%; OR 0.68 (0.37-1.27)  $p=0.22$ ). However overall survival (OS) showed no difference (2-year OS 16% vs 12%, HR 0.97 (0.73-1.28)  $p=0.8$ ). Exploratory analyses failed to identify any subgroup benefitting from tosedostat. Despite promising pre-clinical, early unrandomised clinical data with acceptable toxicity and an improvement in response, we did not find evidence that the

addition of tosedostat to LDAC produced a survival benefit in this group of AML patients.

Trial Reference ISRCTN40571019

## **Introduction**

A major current challenge in the treatment of acute myeloid leukaemia (AML) is to find effective, convenient and safe treatment for older patients<sup>1,2</sup>. Almost half of patients with AML are over 70 years of age. To date, intensive therapy, even for those considered fit enough to receive it, delivers poor survival particularly for patients with co-morbidities, poor performance score or adverse disease biology. Ever since, in the overdue clinical trials in this population, it has been assumed that unless remission was achieved, little benefit was anticipated. Standards of care include low dose ara-C (LDAC)<sup>3</sup> and the hypomethylating agents azacitidine<sup>4</sup> or decitabine<sup>5</sup>, each of which have low remission rates, although the hypomethylating agents may prolong survival without achieving remission. Several new treatments tested in this context have substantially improved remission rates, but not overall survival, although the recently published results of combining venetoclax with azacitidine have for the first time prolonged survival in this patient group with a non-intensive approach<sup>6</sup>

Tosedostat is an example of a new class of orally administered metalloenzyme inhibitors with anti-proliferative and anti-angiogenic activity in vivo and in vitro against a wide range of haematological and solid human cancer cells<sup>7</sup>. Exposure of cells to tosedostat results in the intracellular accumulation of an acid metabolite, CHR-79888, which exerts a powerful inhibitory effect on intracellular metalloenzymes resulting in

anti-proliferative, pro-apoptotic and anti-angiogenic activity<sup>8</sup>. The intracellular metalloenzyme targets for tosedostat are likely to be members of the M1 family of aminopeptidases, so tosedostat is an aminopeptidase inhibitor. Aminopeptidases play a critical role in the final steps of protein recycling downstream of proteasomal degradation and inhibition of aminopeptidases by tosedostat may, like proteasome inhibition, disrupt the turnover of cellular proteins in such a way that it impacts cancer cell growth<sup>9</sup>. Natural product inhibitors of aminopeptidases, particularly bestatin, exhibit similar, albeit weaker, pharmacological actions to tosedostat, including its pro-apoptotic, anti-proliferative and anti-angiogenic effects and its ability to induce amino acid deprivation response (AADR) related gene expression changes<sup>10</sup>. Tosedostat synergises in vitro with a very wide range of chemotherapeutic and targeted agents in inducing anti-proliferative effects in many haematological and non-haematological cancer cell lines. We previously showed evidence of synergy with ara-C in pre-clinical studies with human AML cells<sup>11</sup>.

A number of early stage clinical trials established a daily dose level of 120mg, with little toxicity and some encouraging clinical activity. The initial phase 1 study defined 180mg as the maximum tolerated dose (MTD) with the limitation being protracted thrombocytopenia, and demonstrated good tolerance at a daily dose of 130mg. In a total of 51 patients with relapse/refractory disease in the study, the overall marrow response was 24%<sup>12</sup>. A second study, (OPAL)<sup>13</sup>, also in relapsed/refractory older patients, assessed more prolonged administration at two dose levels (240mg for 2 months then 120mg for 4 months or 120mg for 6 months). Initially 35 patients were allocated to each schedule which resulted in an overall response rate (ORR) of 22%. From this study the dose for prolonged treatment emerged as 120mg once a day. Based on the pre-clinical evidence of synergy Mawad and colleagues<sup>14</sup>, in a phase 2

study which included 26 untreated older patients combined tosedostat (120mg) daily with conventional dose ara-C ( $1\text{g}/\text{m}^2$  days 1-5) or decitibine ( $20\text{mg}/\text{m}^2/\text{days}$  1-5). A subsequent 8 patients received a higher tosedostat dose. Complete remission (CR)/complete remission with incomplete recovery of counts (CRi) CR/CRi was achieved in 53% and it was concluded that the 120mg dose was preferable. Finally Visani and colleagues<sup>15</sup> conducted an unrandomised phase II study on 33 older untreated patients with the LDAC and tosedostat combination and showed a CR/CRi rate of 54%, the majority of which were CRs. Of additional interest was that they suggested that those patients who achieved CR could be predicted with a 212 gene panel. A microarray analysis performed in 29 of 33 patients identified 188 genes associated with clinical response (CR vs no CR). Three of them (CD93, GORASP1, CXCL16) were validated by quantitative polymerase chain reaction<sup>16</sup>. This potential improvement in efficacy and tolerability suggested that it may be especially relevant in the management of older patients who frequently have resistant disease and tolerate traditional therapies poorly. We therefore investigated whether tosedostat combined with LDAC was superior to LDAC alone as first line therapy for older patients with AML who were not considered fit for intensive therapy.

## **Methods.**

This evaluation of tosedostat was a component of our “Pick a Winner” trial strategy in the LI-1 trial (ISRCTN40571019) where patients are randomised between a control arm (LDAC) and one of a number of experimental options<sup>17</sup>. The comparison is only between each experimental option and LDAC, and not between the experimental options. Patients allocated to LDAC only act as controls to patients who have been contemporaneously randomised to an experimental arm.

Patients were eligible if they had de novo or secondary AML or high risk myelodysplastic syndrome (MDS), defined as >10% marrow blasts, and were older than 60 years and considered unfit for intensive chemotherapy. "Unfitness" was determined by the investigator/attending clinician- not specifically protocol defined and documented by collection of co-morbidity using components of the Sorrow index<sup>18</sup>. Patients with a prior diagnosis of MDS (>10% blasts, RAEB 2) who had received azacitidine were not eligible, but patients with a prior diagnosis of MDS with <10% blasts who have failed a demethylation agent and then developed AML were. Patients were categorised for response and survival using the validated multi-parameter Wheatley risk score<sup>19</sup> which predicted survival based on age, performance status, cytogenetics, and de novo or secondary disease. This score has been prospectively validated in older patients treated both non-intensively with LDAC and with intensive chemotherapy. Diagnosis and response definitions described below were designated by the local investigator. Cytogenetics (a minimum of 20 metaphases) and immunophenotypic characterization were carried out in regional reference laboratories which participate in national quality assurance schemes.

In this study patients were randomised 1:1 to LDAC or LDAC combined with tosedostat (LDAC-T). LDAC treatment comprised Ara-C 20mg twice a day for 10 days by subcutaneous injection for 4 courses given at 4 to 6 weeks intervals (there was no placebo). Tosedostat was given orally at 120mg once a day continuously for up to 6 months. Patients who were considered to be benefiting, by demonstrating stable disease or continuing response, were permitted to continue on their allocated treatment.

Patients were required to provide written consent and the trial was sponsored by Cardiff University and approved by the Wales Research Ethics Committee in compliance with the Declaration of Helsinki.

**Endpoints and assessments:** The primary endpoint was overall survival (OS), following international guidelines OS is defined as the time from randomisation to death. The protocol defined complete remission (CR) as a normocellular bone marrow aspirate containing <5% leukaemic blasts and showing evidence of normal maturation of other marrow elements. Persistence of myelodysplastic features did not preclude the diagnosis of CR. To achieve CR, patients required neutrophil recovery to  $\geq 1.0 \times 10^9/l$  and also platelets to  $\geq 100 \times 10^9/l$ , without evidence of extramedullary disease. Patients who achieved CR according to the protocol, but without evidence of adequate count recovery are denoted here as CRi, patients were required to be platelet-transfusion independent indicating sufficient time for marrow regeneration. Overall response was defined as CR/CRi as we do not have complete data on partial response and morphologic leukaemia free state. For remitters, relapse free survival (RFS) was the time from remission (CR or CRi) until relapse or death. Survival from CR is defined as the time from CR/CRi (first report) until death.

**Toxicity:** Adverse events and toxicity were recorded as defined by the National Cancer Institute common terminology criteria for adverse events (NCI CTCAE) version 3.

**Statistical methods:** All analyses are by intention-to-treat. Categorical endpoints (e.g. CR rates) were compared using Mantel-Haenszel tests, giving Peto odds ratios and confidence intervals. Continuous/scale variables were analysed by non-parametric (Wilcoxon rank sum) tests. Time-to-event outcomes were analysed using the log-rank

test, with Kaplan-Meier survival curves. Odds/hazard ratios (OR/HR) less than 1 indicate benefit for the investigational therapy. In the Pick-a-Winner design analyses are performed for each investigational arm separately versus the control arm of LDAC. In addition to overall analyses, exploratory analyses were performed stratified by the randomisation stratification parameters and other important variables, with suitable tests for interaction. Because of the well-known dangers of subgroup analysis, these were interpreted cautiously.

The power calculation for the trial as a whole specified that final analysis was to be performed after 340 events (deaths) had been reported. Under the rules of the “Pick a Winner” design, the data monitoring committee (DMC) initially examined outcomes after response data were available for the first 100 patients in each randomisation (50 patients in each arm). At this point, in order to show sufficient promise to be carried forward, there had to be at least a 2.5% improvement in remission rates (CR+CRi) for the experimental arm over the control arm. At this time, the DMC also assessed survival and toxicity as additional criteria to be satisfied, although there was no formal stopping rule for either of these endpoints. If the DMC believed there was sufficient promise in the arm, the trial would continue to accrue until approximately 100 patients were in each arm. Once 170 deaths had been recorded a further interim analysis was performed and the hazard ratio for survival was required to be less than 0.85 in order for the trial to consider continuing to 400 patients and 340 events. At this point, the decision to stop or continue is made on the basis of the hazard ratio for OS. The aspiration of the study is a doubling of survival from 11% to 22% at two years which is equivalent to an average hazard ratio of 0.69.

At the time of this final analysis the median follow-up for OS is 48 months (range 0.2-40.5). Surviving patients are censored at the date last known to be alive.



194

195 **Results:**

196 **Patient Characteristics:** Between June 2014 and February 2017, 243 patients with  
 197 a median age of 76 years (range 60-88) entered the randomisation, of whom 60%  
 198 were male and 40% female. Sixty-six percent had de novo AML, 28% secondary AML,  
 199 and 6% high risk MDS. Cytogenetic analysis identified 1% had favourable, 65%  
 200 intermediate and 22% adverse cytogenetics (table1). By the validated Wheatley  
 201 index<sup>19</sup>, 5% were good risk, 36% standard risk and 58% poor risk. This validated score  
 202 would predict an expected 12 month survival of 36%, 42% and 14% for LDAC  
 203 monotherapy in the three risk groups based on historical data, and would be equivalent  
 204 to a predicted overall 12-month survival of approximately 25%.

205 The disposition of the patients is shown in figure 1 (CONSORT diagram). A median of  
 206 2 courses (range 1-8) was delivered in either arm. For LDAC-T the mean was 2.9; and  
 207 number of courses was 0=6%; 1=38%; 2=24%; 3=5%; 4=6%; 5=5%; 6=4%; 7=3%;  
 208 8=13%. For LDAC alone, the mean was 2.3, and the number of courses was 0=5%;  
 209 1=34%; 2=18%; 3=4%; 4=10%; 5=3%; 6=9%; 7=2%; 8=15%; p= 0.3).

210 The reasons provided by investigators for not receiving intensive therapy were age in  
 211 90% of cases, fitness in 45% of cases (both together in 38% of cases), and other  
 212 reasons in 5% of cases of which over half were patient choice. The HCT-CI was (0 =  
 213 42%, 1-2 = 30%, 3+ =28%). Of the co-morbidities listed on entry, the most frequent  
 214 were those described as prior tumour (14%), diabetes (13%); cardiac (9%); infection  
 215 (9%), mild-to-moderate pulmonary (8%); rheumatological (8%); obesity (8%) and  
 216 arrhythmia (5%) (table 1). No other co-morbidity was present in more than 5% of  
 217 patients.

218 **Response:**

219 Initial assessment by the DMC after the first 100 patients in September 2015 agreed  
220 that the randomisation should continue. In February 2017, the DMC performed an  
221 outcomes assessment on the LDAC-T versus LDAC arms of the LI-1 trial (n=243), at  
222 which point additional randomisations were suspended pending the review. At the  
223 second interim analysis in November 2017 after 183 events, while there was a benefit  
224 in remission rates, however LDAC-T failed to show a sufficiently promising hazard  
225 ratio for survival, and therefore on the recommendation of the DMC the arm was  
226 closed. Patients who were benefitting from tosedostat were permitted to stay on  
227 treatment. The data presented here represents an analysis undertaken after the DMC  
228 recommendation with cleaner data and more mature follow up.

229 Overall, CR was achieved in 16% of patients with a further 6% achieving a CRi (total  
230 ORR 21%). There was a non-significant increase in CR rate with tosedostat (LDAC-  
231 T vs LDAC) (19% vs 12%, OR 0.61 (0.30-1.23) p=0.17). For the overall response  
232 (CR+CRi), there was little evidence that a benefit of the addition of tosedostat could  
233 be seen (25% vs 18%; OR 0.68 (0.37-1.27) p=0.22). A non-significant reduction in  
234 resistant disease was observed by the addition of tosedostat (60% vs 68% OR 0.68  
235 (0.40-1.16) p=0.16. The thirty-day mortality was not significantly increased (16% vs  
236 14%, HR 1.26 (0.65-2.46) p=0.5. (table 2).;

237 **Treatment Compliance:** Following remission, treatment was given to 19/22 LDAC  
238 patients (5 patients received 1 course, 4 patients 2 courses, 1 patient 3 courses, 2  
239 patients 4 courses, and 7 patients 6 or more courses) and 26/30 Tosedostat patients  
240 treated (3 patients received 1 course, 4 patients 2 courses, 6 patient 3 courses, 4  
241 patients 4 courses, 1 patient 5 courses, and 12 patients 6 or more courses). No patient

allocated to LDAC alone received tosedostat; however 2 patients randomised to receive LDAC-T received 1 and 3 courses of LDAC alone.

**Overall Survival** The OS did not differ by treatment arm (LDAC-T vs LDAC) (2-year OS 16% vs 12%, HR 0.97 (0.73-1.28)  $p=0.8$ ; figure 2a).

**Survival of Responders:** For the total 52 patients who achieved a CR/CRi, the median OS from remission was 21.8 months. Although there was an apparent modest benefit in 2 years survival from response (447% vs 36%), this failed to reach statistical significance (HR 0.88 (0.43-1.80)  $p=0.7$ ) (figure 2b). For patients who relapsed, there was no significant difference in the survival following relapse between treatment arms (1 year survival post relapse 30% vs 17%; HR 0.93 (0.45-1.92)  $p=0.8$ ; (figure 2c). In the patients who did not achieve CR/CRi, the survival was not different between the arms.

**Relapse Free Survival:** Although remission rates were higher in the tosedostat arm, there was no significant difference in duration of remission RFS (HR 0.82 (0.46-1.47)  $p=0.5$ ; figure 2d).

**Toxicity:** Although rates of grade 3+ toxicity were low overall, tosedostat was associated with significantly increased diarrhoea, and cardiac toxicity (2 grade 4 events that led to tosedostat discontinuation- AF and raised troponin) in course 1, and with greater cardiac and liver alanine transaminase (ALT) toxicity in course 2 . Resource usage (blood product support, antibiotics and hospital utilisation) tended to be consistently higher in the tosedostat arm, though the only significant difference between arms was an increased use of platelets in course 1 (mean 5.0 vs 3.5 pools  $p=0.006$ ); (figure 3a and 3b).

## **Exploratory Subgroup Analysis:**

Exploratory analyses were carried out on survival, to find out if there was an identifiable subgroup with a differential effect of treatment. Baseline covariates including age, sex, diagnosis, cytogenetics, white blood count, performance status, and Wheatley risk group were explored (Supplemental Figure 2). Additional analysis by NPM1 and FLT3-ITD/TKD status was additionally explored. More detailed molecular analyses were not available. Although the power of such analyses is limited by small numbers in some subgroups, there were no significant interactions between baseline variables and treatment for survival. In particular, no subgroup could be identified where there was a benefit for LDAC-T.

## **Discussion:**

Compared to younger patients with AML, the decision in treatment strategy is not always obvious. At one end of the spectrum there are patients who have several co-morbidities where even if the prognostic assessment of their disease biology is not adverse, are at high risk of not surviving a version of standard chemotherapy. At the other are patients who are chronologically old but have few co-morbidities combined with good performance status. In these cases intensive chemotherapy may be of benefit, but the decision to offer conventional chemotherapy may be negatively influenced by adverse disease biology, where chemotherapy may have a low chance of success. Some patients who are fit may decline treatment in preference for more time out of hospital, particularly if facilitated by outpatient or oral medication. At the centre of this is the physician – indeed in our previous AML14 trial where an intensive and non-intensive treatment approach were available, the physician emerged as an

independent factor in treatment choice. Many prognostic scoring systems have been developed for younger patients to guide treatment decisions, and such scores can be developed for older patients, but few have been prospectively validated in recipients of non-intensive therapy. We developed the Wheatley Score<sup>19</sup>, which is useful in predicting expected outcomes for non-intensive treatment approaches. In this study based on the Wheatley score 4% of patients were favourable, 31% intermediate and 65% were adverse risk with respective expected 12-month OS 36%, 42% and 14% respectively. The predicted 12 month OS was 25%, which is what was achieved.

We developed LDAC as a standard of care at a time when no other randomised trials in this patient population had suggested an alternative. We found that clinical toxicities were no greater than best supportive care<sup>3</sup>. However durable benefit was only seen in the 18% of patients who entered CR, where median OS was 575 days compared to only 66 days for those that did not respond. This experience led to the development of a “Pick a Winner” design which depended on an initial improvement in remission rate as a surrogate for future survival benefit. A number of novel treatments that produced encouraging results in unrandomised trials have been included, but failed the scrutiny of randomisation<sup>20-23</sup>. Others were able to double the remission rates but did not improve overall survival<sup>24,25</sup>. Another observation has been that in different cohorts of LDAC patients the remission rate varied from 14% to 21% and the 12 month survival from 25% to 32%, without obvious differences in patients’ characteristics<sup>26</sup>. To date 2480 randomisations have been undertaken in 1753 patients to evaluate 13 agents or combinations<sup>21-26</sup>. The evaluation is complete on 11 options, and 2 are ongoing. The use of remission as a surrogate endpoint helps identify and exclude unpromising treatments, but should not replace survival as an endpoint in trials in this population.

314 Mechanistically tosedostat has several properties which could be particularly helpful  
315 in older patients<sup>8</sup>. The developmental phase I/II experience in relapse and in  
316 combination was both feasible from the toxicity point of view, and appeared to offer an  
317 improved clinical response. The oral formulation is also helpful in the elderly  
318 population. We therefore initiated the randomised comparison reported here.  
319 Disappointingly, the combination failed to meet the IDMC criteria to continue the trial.  
320 In reaching their recommendation the IDMC looked not only at the strict continuation  
321 criteria set down, based upon remission, but also relied upon safety data, and in  
322 particular early mortality when deciding whether or not to continue. The IDMC closed  
323 the tosedostat arm based on a failure to improve survival as assessed by the  
324 confidence intervals at the time of their analysis which depended on observing a  
325 hazard ratio of 0.69, representing the requirement to improve 2-year survival from 11%  
326 to 22%. It was therefore concluded that even with more patients included the drug was  
327 unlikely to demonstrate the sort of benefit required by the design of the trial. As is  
328 observed in many such studies the primary reason for discontinuation was refractory  
329 disease. For responding patients the median OS was an impressive 21.8 months,  
330 although we were unable to identify any clinical or laboratory findings which could  
331 reliably identify such patients a recent publication by Visani<sup>15</sup> has proposed a gene  
332 expression profile that could predict such a response and could warrant further  
333 evaluation.

334  
335 The introduction of hypomethylating agents has improved survival without  
336 substantially improving the rate of remission<sup>4</sup> and globally considered the standard of  
337 care for the frail unfit AML patient. New combinations (including venetoclax,  
338 enasidenib, ivosidenib and glasdegib) show considerable promise, and indeed have

received regulatory approval for this patient group, mostly based on unrandomised data<sup>27-31</sup>. As described above there are several examples of early promise which fail in the rigour of randomization. Although recently published data from the VIALE-A study, in perhaps a more selected frail elderly AML population, combining venetoclax with azacitidine has demonstrated a significant improvement in overall survival, this combination may ultimately become considered the new standard of care in this setting<sup>6</sup>.

In conclusion, tosedostat demonstrated promising early data and acceptable tolerability, its addition to LDAC did achieve a modest improvement in response rates, but we did not find evidence that it produced a survival benefit in this group of patients. Strategies other than aminopeptidase inhibition appear to demonstrate more rational approaches for future non intensive combination therapy in AML.

## Acknowledgements

We wish to thank CTI Biopharma for supporting the trial, Blood Cancer UK for research support, and the Haematology Clinical Trials Unit and Centre for Trials Research, Cardiff University for managing the trial.

363

364 The following investigators recruited patients to the trial:

365 **Aalborg University Hospital:** Maria Kallenbach, M Severinsen; **Aberdeen Royal Infirmary:** Dominic  
366 Culligan, Caroline Duncan; **Addenbrooke's Hospital:** Pramila Krishnamurthy; **Airedale General**  
367 **Hospital:** Ann Cuthbert, Chetan Patalappa, Michail Spanoudakis; **Arrowe Park Hospital:** David Galvani;  
368 **Auckland City Hospital:** I Berkahn, N Patton, Lucy Pemberton; **Barnet Hospital:** Andres Virchis;  
369 **Basildon University Hospital:** Parag Jasani; **Beatson West of Scotland Cancer Centre:** Anne Parker;  
370 **Belfast City Hospital:** Claire Arnold, Robert Cuthbert, Damien Finnega; **Birmingham Heartlands**  
371 **Hospital:** Manos Nikolousis, Guy Pratt; **Blackpool Victoria Hospital:** Paul Cahalin; **Bradford Royal**  
372 **Infirmary:** Sam Ackroyd; **Bristol Haematology & Oncology Centre:** Priyanka Mehta, Rachel Protheroe;  
373 **Capital & Coast DHB (Wellington):** Robert Weinkove; **Castle Hill Hospital (Hull):** Sara Ali; **Chesterfield**  
374 **Royal Hospital:** Emma Welch; **Christchurch Hospital:** Mark Smith, Ruth Spearing; **Churchill Hospital:**  
375 **Paresh Vyas;** **Colchester General Hospital:** Gavin Campbell, Mike Hamblin, Marion Wood; **Conquest**  
376 **Hospital:** Sunil Gupta, Simon Westonsmith; **Countess of Chester Hospital:** Arvind Pillai; **Derriford**  
377 **Hospital (Plymouth):** Hannah Hunter, Simon Rule; **Forth Valley Royal Hospital:** Roderick Neilson;  
378 **Gloucestershire Royal Hospital:** Richard Lush; **Huddersfield Royal Infirmary:** Kate Rothwell, Wunna  
379 Swe; **Ipswich Hospital:** Isobel Chalmers, Mahesh Prahladan; **Kettering General Hospital:** Mark Kwan,  
380 Matthew Lyttleton; **Leicester Royal Infirmary:** Ann Hunter, Alexander M Martin; **Lincoln County**  
381 **Hospital:** Kandeepan Saravanamuttu; **Manchester Royal Infirmary:** Muhammad Saif, Eleni Tholouli;  
382 **Medway Maritime Hospital:** Vivienne Andrews; **Middlemore Hospital:** Sharon Jackson, Rajeev  
383 Rajagopal; **New Victoria Hospital:** Anne Morrison; **NHS Ayrshire & Arran (Ayr Hospital):** William  
384 Gordon; **NHS Lanarkshire:** Lindsay Mitchell; **Northampton General Hospital:** Angela Bowen, Alistair  
385 McGrann, Sajjan Mittal; **Northwick Park Hospital:** Nicki Panoskaltsis; **Nottingham University Hospital:**  
386 Jenny Byrne, Emma Das-Gupta, Rohini Radia, Nigel Russell; **Palmerston North Hospital:** Bart Baker,  
387 Allanah Kilfoyle, Elayne Knottenbelt; **Pilgrim Hospital:** Ciro Rinaldi; **Pinderfields Hospital:** Paul



388 Moreton; **Poole General Hospital:** Darshyani Furby, Fergus Jack; **Queen Alexandra Hospital**  
389 **(Portsmouth):** Charles Alderman, Edward Belsham, Robert Corser, Tanya Cranfield, Helen Dignum,  
390 Christopher James; **Queen Elizabeth Hospital, Kings Lynn:** Emma Gudgin; **Queen's Hospital, Romford:**  
391 Claire Hemmaway, Mahalakshmi Mohan; **Rigshospitalet:** Peter Kampman, Lars Kjeldsen, Jindrich  
392 Mourek, Ove Juul Nielsen, Carsten Utoft Niemann; **Royal Berkshire Hospital:** Henri Grech, Asif Khan,  
393 Rebecca Sampson; **Royal Cornwall Hospital:** Desmond Creagh; **Royal Darwin Hospital:** Ferenc Szabo;  
394 **Royal Devon & Exeter Hospital:** J Coppel, D Veale; **Royal Free Hospital:** Panagiotis Kottaridis; **Russells**  
395 **Hall Hospital:** Savio Fernandes; **Salisbury District Hospital:** Jonathan Cullis, Effie Grand; **Sandwell**  
396 **Hospital:** Farooq Wandroo; **Singleton Hospital:** Unmesh Mohite; **Southport and Formby District**  
397 **General Hospital:** David O'Brien; **St Helens & Knowsley (Whiston):** Toby Nicholson; **St Richard's**  
398 **Hospital:** Sarah Janes; **Stoke Mandeville Hospital:** Robin Aitchison; **Sunderland Royal Hospital:** Mike  
399 Galloway, Susanna Mathew; **The Christie:** Mike Dennis; **The Great Western Hospital:** Norbert Blesing,  
400 Sarah Green; **The James Cook University Hospital:** Rajesh Mamadigi, Diane Plews; **The Royal**  
401 **Bournemouth Hospital:** Joseph Chacko; **The Royal Liverpool Hospital:** Rahuman Salim; **The Royal**  
402 **Oldham Hospital:** Allameddine Allameddine, Hayley Greenfield, David Osborne; **Townsville Hospital:**  
403 Georgina Hodges, Hock Choong Lai; **Trafford General Hospital:** Patrick Carrington; **University Hospital**  
404 **Aintree:** Walid Sadik, Lynny Yung; **University Hospital Coventry & Warwickshire:** Anand Lokare;  
405 **University Hospital of Wales:** Caroline Alvares, Steve Knapper; **Waikato Hospital:** Hugh Goodman,  
406 Shahid Islam; **Wycombe Hospital:** Robin Aitchison, Jonathan Pattinson, Beena Pushkaran; **Ysbyty Glan**  
407 **Clywd:** Earnest Heartin; **Ysbyty Gwynedd:** David Edwards, Jim Seale

408

## 409 References

- 410 1. Kantarjian H, Ravandi F, O'Brien S, Cortes J, Faderl S, Garcia-Manero G et  
411 al. Intensive chemotherapy does not benefit most older patients (age 70 years  
412 or older) with acute myeloid leukemia. *Blood* 2011; 116: 4422–4429.

2. Appelbaum, F.R; Gundacker, H.; Head, D.R et al. Age and acute myeloid leukemia. *Blood* 2008; 107(9), 3481-3285
3. Burnett AK, Milligan D, Prentice, AG, et al. A comparison of low-dose cytarabine and hydroxyurea with or without all-trans retinoic acid for acute myeloid leukemia and high-risk myelodysplastic syndrome in patients not considered fit for intensive treatment. *Cancer*. 2007;109 (6):1114-1124.
4. Fenaux P, Mufti GJ, Hellstrom-Liindberg E, et al. Azacytidine prolongs overall survival compared with conventional care regimens in elderly patients with low bone marrow blast count acute myeloid leukemia. *J. Clin.Oncol.* 2010, 28: 562 – 569
5. Kantarjian HM, Thomas XG, Dmoszynska A, et al.. Multicenter, randomized, open-label, phase III trial of decitabine versus patient choice, with physician advice, of either supportive care or low-dose cytarabine for the treatment of older patients with newly diagnosed acute myeloid leukemia. *J.Clin.Oncol.* 2012;30 (21):2670-2677.
6. DiNardo CD, Jonas BA, Pullarkat V, Thirman MJ, Garcia JS, Wei AH, Konopleva M, Döhner H, Letai A, Fenaux P, Koller E, Havelange V, Leber B, Esteve J, Wang J, Pejsa V, Hájek R, Porkka K, Illés Á, Lavie D, Lemoli RM, Yamamoto K, Yoon SS, Jang JH, Yeh SP, Turgut M, Hong WJ, Zhou Y, Potluri J, Pratz KW. Azacitidine and Venetoclax in Previously Untreated Acute

Myeloid Leukemia. N Engl J Med. 2020 Aug 13;383(7):617-629. doi:  
10.1056/NEJMoa2012971. PMID: 32786187.

7. Reid AH, Protheroe A, Attard G, et al. A first-in-man phase i and  
pharmacokinetic study on CHR-2797 (Tosedostat), an inhibitor of M1  
minopeptidases, in patients with advanced solid tumors. *Clin Cancer Res.*  
2009; 15(15):4978-4985

8. Krige D, Needham LA, Bawden LJ, et al. CHR-2797: an antiproliferative  
aminopeptidase inhibitor that leads to amino acid deprivation in human  
leukemic cells. *Cancer Res.* 2008; 68(16):6669-6679.

9. Saric T, Graef CI and Goldberg AL. Pathway for degradation of peptides  
generated by proteasomes: a key role for thimet oligopeptidase and other  
metallopeptidases. *J. Biol. Chem.*2004, 279: 46723-46732

10. Drummond AH, Farmer H, Flores N, Miles LEC, Callaghan J, Bawden LJ and  
Krige D. Antiproliferative aminopeptidase inhibitors inhibit mTOR and alter  
expression of Noxa and Mlc-1 in leukemic cells via amino acid deprivation.  
*AACR Abstract* 2007: 4069

11. Jenkins C, Hewamana S, Krige D, et al. Amino peptidase inhibition by the novel  
agent CHR-2797 (tosedostat) for the therapy of acute myeloid leukemia. *Leuk*  
*Res* 2011;35(5):677-81

12. Löwenberg B<sup>1</sup>, Morgan G, Ossenkoppele GJ, et al. Phase I/II clinical study of Tosedostat, an inhibitor of aminopeptidases, in patients with acute myeloid leukemia and myelodysplasia. *J Clin Oncol*. 2010; 28(28):4333-4338
13. Cortes J, Feldman E, Yee K, et al. Two dosing regimens with relapsed or refractory acute myeloid leukaemia (OPAL): a randomised open-label phase 2 study. *Lancet Oncol*. 2013; 14(4):354-362
14. Mawad R, Becker PS, Hendrie P, et al. Phase II study of tosedostat with cytarabine or decitabine in newly diagnosed older patients with acute myeloid leukaemia or high-risk MDS. *Br J Haematol*. 2016; 172(2):238-245.
15. Visani G, Loscocco F, Fuligini F et al. Tosedostat with Low-Dose Cytarabine combo induces a high rate of responses that can be predicted by genetic profiling in AML: final results of a phase II multicentre study. *Haematologica* 2015; 100703, p 564
16. Visani G, Loscocco F, Dennis M, Zuffa E, Candoni A, Sensi A, Giannini B, Musuraca G, Mianulli AM, Clavio M, Rocchi M, Gibellini D, Navari M, Gilkes A, Piccaluga PP, Isidori A. Gene expression profile predicts response to the combination of tosedostat and low-dose cytarabine in elderly AML. *Blood Adv*. 2020 Oct 27;4(20):5040-5049.
17. Hills RK, Burnett AK. Applicability of a "Pick a Winner" trial design to acute myeloid leukemia. *Blood*. 2011; 118(9): 2389-94

- 488
- 489 18. Sorrow ML, Giralt S, Sandmaier BM, de Lima M, Shahjahan M, Maloney DG et
- 490 al. Hematopoietic cell transplantation-specific comorbidity index as an outcome
- 491 predictor for patients with acute myeloid leukemia in first remission: combined
- 492 FHCRC and MDACC experiences. *Blood* 2007; 110: 4606–4613.
- 493
- 494 19. Wheatley K, Brookes CL, Howman AJ, Goldstone AH, Milligan DW, Prentice
- 495 AG et al. Prognostic factor analysis of the survival of elderly patients with AML
- 496 in the MRC AML11 and LRF AML14 trials. *Br J Haematol* 2009; 145:598–605.
- 497
- 498 20. Burnett AK, Russell NH, Culligan D, Cavanagh J, Kell J, Wheatley K et al. The
- 499 addition of the farnesyl transferase inhibitor, tipifarnib, to low dose cytarabine
- 500 does not improve outcome for older patients with AML. *Br J Haematol* 2012;
- 501 158: 519–522.
- 502
- 503 21. Burnett AK, Hills RK, Hunter A, Milligan D, Kell J, Wheatley K et al. The
- 504 addition of arsenic trioxide to low-dose Ara-C in older patients with AML does
- 505 not improve outcome. *Leukemia* 2011; 25: 1122–1127
- 506
- 507 22. Burnett AK, Russell N, Hills RK, et al. A randomised comparison of the novel
- 508 nucleoside analogue sapacitabine with low-dose cytarabine in older patients
- 509 with acute myeloid leukaemia. *Leukemia*. 2015;29(6):1312-1319.
- 510 doi:10.1038/leu.2015.38
- 511

23. Dennis M, Russell N, Hills RK, et al. Vosaroxin and vosaroxin plus low-dose Ara-C (LDAC) vs low-dose Ara-C alone in older patients with acute myeloid leukemia. *Blood*. 2015;125(19):2923-2932. doi:10.1182/blood-2014-10-608117
24. Burnett, A. K., Hills, R. K., Hunter, A. E. et al, UK National Cancer Research Institute AML Working Group. The addition of gemtuzumab ozogamicin to low-dose Ara-C improves remission rate but does not significantly prolong survival in older patients with acute myeloid leukaemia: results from the LRF AML14 and NCRI AML16 pick-a-winner comparison. *Leukemia*, 2013; 27(1), 75-81
25. Burnett AK, Russell NH, Hunter AE, Milligan D, Knapper S, Wheatley K et al. Clofarabine doubles the response rate in older patients with acute myeloid leukemia but does not improve survival. *Blood* 2013; 122: 1384–1394
26. Dennis M, Hills R, Russell N, Copland M, Thomas I, McMullin MF, Ali S, Dignum H and Burnett AK An Evaluation of 17 years of Low Dose Cytarabine as Therapy for AML Patients Not Fit for Intensive Treatment, Including Patients with Adverse Cytogenetics, Shows Improving Survival and Potential Underutilisation. *Blood* (2017) 130 (Supplement 1): 3874
27. Stein EM, DiNardo CD, Pollyea DA, et al. Enasidenib in mutant-IDH2 relapsed or refractory acute myeloid leukemia. *Blood* 2017;doi.10.1182/blood-2017-04-779405.

27 28DiNardo CD, Stein EM, de Botton S, et al. Durable Remissions with  
Ivosidenib in IDH1-Mutated Relapsed or Refractory AML. N Engl J Med  
2018;378:2386-98.

28 29Wei AH, Jr SAS, Hou J-Z, et al. Venetoclax Combined With Low-Dose  
Cytarabine for Previously Untreated Patients With Acute Myeloid Leukemia:  
Results From a Phase Ib/II Study. J Clin Oncol 2019;37:1277-84.

30 .DiNardo CD, Pratz KW, Letai A, et al. Safety and preliminary efficacy of  
venetoclax with decitabine or azacitidine in elderly patients with previously  
untreated acute myeloid leukaemia: a non-randomised, open-label, phase 1b  
study. The Lancet Oncology 2018;19:216-28

31. Cortes JE, Heidel FH, Hellmann A, et al. Randomized comparison of low  
dosecytarabine with or without glasdegib in patients with newly diagnosed  
acute myeloid leukemia or high risk myelodysplastic syndrome. Leukemia  
2019;33:379-389.

#### **Disclosure of Conflicts of Interest**

AKB was an employee of CTI Biopharma 2015-2017. REC has received research funding and honoraria from Novartis and Bristol Myers Squibb, and honoraria from Pfizer, Jazz Pharmaceuticals and Abbvie. MC has received research funding from Novartis, Bristol-Myers Squibb, Cyclacel and Takeda/Incyte, is/has been an advisory board member for Bristol-Myers Squibb, Novartis, Incyte, Daiichi Sankyo, Jazz and Pfizer and has received honoraria from Astellas, Bristol-Myers Squibb, Novartis, Incyte, Pfizer and Gilead. The other authors have nothing to disclose.

**Author Contributions:** MD: chief investigator; reviewed the data and wrote the manuscript AKB: designed the trial; wrote protocol; chief investigator until Q3 2014; RKH: designed the trial, wrote the protocol, analysed the data. CA analysed the data with extended follow up, IT supervised the data collection, reviewed the data. MTS, CH, and PG were major recruiters. NHR: designed trial; reviewed the data. MC Co-CI, and REC reviewed the data. All authors reviewed the manuscript.