



Clinical trial results:

A randomised, open-label study to assess the immunogenicity and safety of GSK Biologicals' IPV vaccine administered as a three-dose primary vaccination course at 2-3-4 months of age in healthy infants in China.

Summary

EudraCT number	2011-003167-30
Trial protocol	Outside EU/EEA
Global end of trial date	05 July 2010

Results information

Result version number	v3 (current)
This version publication date	19 April 2023
First version publication date	01 July 2015
Version creation reason	<ul style="list-style-type: none">• Correction of full data set Correction of full data set and alignment between registries.

Trial information

Trial identification

Sponsor protocol code	112679
-----------------------	--------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT01021293
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	GlaxoSmithKline Biologicals
Sponsor organisation address	Rue de l'Institut 89, Rixensart, Belgium, B-1330
Public contact	Clinical Trials Call Center, GlaxoSmithKline Biologicals, 44 2089904466, GSKClinicalSupportHD@gsk.com
Scientific contact	Clinical Trials Call Center, GlaxoSmithKline Biologicals, 44 2089904466, GSKClinicalSupportHD@gsk.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	Yes

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	15 February 2011
Is this the analysis of the primary completion data?	Yes
Primary completion date	05 July 2010
Global end of trial reached?	Yes
Global end of trial date	05 July 2010
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

To demonstrate the non-inferiority of GSK Biologicals' IPV vaccine as compared to the Chinese OPV vaccine in terms of the immune response to poliovirus type 1, 2 and 3, one month after the third vaccine dose.

Non-inferiority in terms of immunogenicity to the three poliovirus antigens will be demonstrated if the upper limit of the 95% confidence interval (CI) on the group difference [Control Group minus Poliorix Group] in the percentage of seroprotected subjects is less than or equal to 10%.

Protection of trial subjects:

The subjects were observed closely for at least 30 minutes, with appropriate medical treatment readily available in case of anaphylaxis following the administration of vaccines.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	28 November 2009
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	China: 1101
Worldwide total number of subjects	1101
EEA total number of subjects	0

Notes:

Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	1101
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	0
From 65 to 84 years	0

85 years and over	0
-------------------	---

Subject disposition

Recruitment

Recruitment details:

Of the 1101 enrolled subjects, one subject was not randomised and administered any vaccine as the parents of the subject refused to vaccinate their child after blood collection at Visit 1 and withdrew their consent.

Pre-assignment

Screening details:

During the screening the following steps occurred: check for inclusion/exclusion criteria, contraindications/precautions, medical history of the subjects and signing informed consent forms.

Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
Arm title	Poliorix Group

Arm description:

Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Poliorix (IPV) vaccine at 2, 3 and 4 months of age, administered intramuscularly into the anterolateral side of the right thigh.

Arm type	Experimental
Investigational medicinal product name	Poliorix
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Intramuscular use

Dosage and administration details:

3 doses of IPV vaccine administered intramuscularly into the anterolateral side of the right thigh.

Arm title	Control Group
------------------	---------------

Arm description:

Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Oral Poliomyelitis Vaccine (OPV) at 2, 3 and 4 months of age, according to the vaccination policy recommended in China.

Arm type	Active comparator
Investigational medicinal product name	Oral Poliomyelitis Vaccine (OPV)
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Oral suspension
Routes of administration	Oral use

Dosage and administration details:

3 doses of OPV vaccine administered orally.

Number of subjects in period 1^[1]	Poliorix Group	Control Group
Started	550	550
Completed	538	526
Not completed	12	24
Adverse event, serious fatal	2	1
Consent withdrawn by subject	1	3
Adverse event, non-fatal	2	1
Migrated/moved from study area	6	18
Lost to follow-up	1	1

Notes:

[1] - The number of subjects reported to be in the baseline period are not the same as the worldwide number enrolled in the trial. It is expected that these numbers will be the same.

Justification: Of the 1101 enrolled subjects, one subject was not randomised and administered any vaccine as the parents of the subject refused to vaccinate their child after blood collection at Visit 1 and withdrew their consent.

Baseline characteristics

Reporting groups

Reporting group title	Poliorix Group
-----------------------	----------------

Reporting group description:

Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Poliorix (IPV) vaccine at 2, 3 and 4 months of age, administered intramuscularly into the anterolateral side of the right thigh.

Reporting group title	Control Group
-----------------------	---------------

Reporting group description:

Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Oral Poliomyelitis Vaccine (OPV) at 2, 3 and 4 months of age, according to the vaccination policy recommended in China.

Reporting group values	Poliorix Group	Control Group	Total
Number of subjects	550	550	1100
Age categorical Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	0	0	0
Infants and toddlers (28 days-23 months)	550	550	1100
Children (2-11 years)	0	0	0
Adolescents (12-17 years)	0	0	0
Adults (18-64 years)	0	0	0
From 65-84 years	0	0	0
85 years and over	0	0	0
Age continuous Units: weeks			
arithmetic mean	10	10.1	
standard deviation	± 1.16	± 1.18	-
Gender categorical Units: Subjects			
Female	268	259	527
Male	282	291	573
Race/Ethnicity Units: Subjects			
Asian-Chinese heritage	550	550	1100

End points

End points reporting groups

Reporting group title	Poliorix Group
Reporting group description: Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Poliorix (IPV) vaccine at 2, 3 and 4 months of age, administered intramuscularly into the anterolateral side of the right thigh.	
Reporting group title	Control Group
Reporting group description: Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Oral Poliomyelitis Vaccine (OPV) at 2, 3 and 4 months of age, according to the vaccination policy recommended in China.	

Primary: Number of seroprotected subjects against poliovirus types 1, 2 and 3

End point title	Number of seroprotected subjects against poliovirus types 1, 2 and 3
End point description: A seroprotected subject was defined as a vaccinated subject with anti-polio types 1, 2 and 3 titers greater than or equal to (\geq) 8 effective dose 50 (ED50).	
End point type	Primary
End point timeframe: At Month 3, one month after the third vaccine dose	

End point values	Poliorix Group	Control Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	306	296		
Units: Subjects				
Anti-poliovirus 1	306	296		
Anti-poliovirus 2	306	296		
Anti-poliovirus 3	306	291		

Statistical analyses

Statistical analysis title	Non-inferiority of IPV as compared to OPV
Statistical analysis description: Non-inferiority of IPV vaccine as compared to OPV vaccine in terms of the immune response to poliovirus type 1 one month after the third vaccine dose. Non-inferiority in terms of immunogenicity to poliovirus antigens was demonstrated if the upper limit of the 95% confidence interval (CI) on the group difference [Control Group minus IPV Group] in the percentage of seroprotected subjects was lesser than or equal to (\leq) 10%.	
Comparison groups	Control Group v Poliorix Group

Number of subjects included in analysis	602
Analysis specification	Pre-specified
Analysis type	non-inferiority
Parameter estimate	Difference in seroprotection rate
Point estimate	0
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.28
upper limit	1.24

Statistical analysis title	Non-inferiority of IPV as compared to OPV
-----------------------------------	---

Statistical analysis description:

Non-inferiority of IPV vaccine as compared to OPV vaccine in terms of the immune response to poliovirus type 2 one month after the third vaccine dose. Non-inferiority in terms of immunogenicity to poliovirus antigens was demonstrated if the upper limit of the 95% confidence interval (CI) on the group difference [Control Group minus IPV Group] in the percentage of seroprotected subjects was $\leq 10\%$.

Comparison groups	Poliorix Group v Control Group
Number of subjects included in analysis	602
Analysis specification	Pre-specified
Analysis type	non-inferiority
Parameter estimate	Difference in seroprotection rate
Point estimate	0
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.28
upper limit	1.24

Statistical analysis title	Non-inferiority of IPV as compared to OPV
-----------------------------------	---

Statistical analysis description:

Non-inferiority of IPV vaccine as compared to OPV vaccine in terms of the immune response to poliovirus type 3 one month after the third vaccine dose. Non-inferiority in terms of immunogenicity to poliovirus antigens was demonstrated if the upper limit of the 95% confidence interval (CI) on the group difference [Control Group minus IPV Group] in the percentage of seroprotected subjects was $\leq 10\%$.

Comparison groups	Poliorix Group v Control Group
Number of subjects included in analysis	602
Analysis specification	Pre-specified
Analysis type	non-inferiority
Parameter estimate	Difference in seroprotection rate
Point estimate	-1.69
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.9
upper limit	-0.44

Secondary: Number of seroprotected subjects against poliovirus types 1, 2 and 3

End point title	Number of seroprotected subjects against poliovirus types 1, 2 and 3
-----------------	--

End point description:

A seroprotected subject was defined as a vaccinated subject with anti-poliovirus types 1, 2 and 3 titers greater than or equal to (\geq) 8 ED50.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 0, prior to the first vaccine dose

End point values	Poliorix Group	Control Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	306	296		
Units: Subjects				
Anti-poliovirus 1	131	113		
Anti-poliovirus 2	93	99		
Anti-poliovirus 3	48	52		

Statistical analyses

No statistical analyses for this end point

Secondary: Anti-poliovirus types 1, 2 and 3 antibody titers

End point title	Anti-poliovirus types 1, 2 and 3 antibody titers
-----------------	--

End point description:

Antibody titers were presented as geometric mean titers (GMTs).

End point type	Secondary
----------------	-----------

End point timeframe:

Prior to the first vaccine dose (Day 0) and one month after the third vaccine dose (Month 3)

End point values	Poliorix Group	Control Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	306	296		
Units: Titre				
geometric mean (confidence interval 95%)				
Anti-poliovirus 1; Day 0	8.7 (7.6 to 9.8)	7.8 (6.9 to 8.9)		
Anti-poliovirus 2; Day 0	6.5 (5.9 to 7.1)	7.2 (6.5 to 8.1)		
Anti-poliovirus 3; Day 0	5.2 (4.8 to 5.7)	5.2 (4.8 to 5.7)		

Anti-poliovirus 1; Month 3	485.1 (436.7 to 538.9)	2817 (2479.5 to 3200.4)		
Anti-poliovirus 2; Month 3	234.3 (209 to 262.6)	468.5 (416.6 to 526.9)		
Anti-poliovirus 3; Month 3	824.3 (725.3 to 936.9)	423.4 (363.3 to 493.3)		

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects with any and grade 3 solicited local symptoms

End point title	Number of subjects with any and grade 3 solicited local symptoms ^[1]
-----------------	---

End point description:

Assessed solicited local symptoms were pain, redness and swelling. Any = occurrence of the symptom regardless of intensity grade. Grade 3 pain = Cry when limb is moved/spontaneously painful. Grade 3 redness/swelling = redness/swelling spreading beyond 30 millimeters (mm) of injection site. This outcome measure concerns subjects from the IPV Group only.

End point type	Secondary
----------------	-----------

End point timeframe:

During the 4-day (Days 0–3) post-vaccination period following each vaccine dose and across doses

Notes:

[1] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is only reporting values for the Poliorix Group.

End point values	Poliorix Group			
Subject group type	Reporting group			
Number of subjects analysed	550			
Units: Subjects				
Any Pain, Dose 1 [N=550]	80			
Grade 3 Pain, Dose 1 [N=550]	1			
Any Redness, Dose 1 [N=550]	20			
Grade 3 Redness, Dose 1 [N=550]	0			
Any Swelling, Dose 1 [N=550]	9			
Grade 3 Swelling, Dose 1 [N=550]	0			
Any Pain, Dose 2 [N=543]	58			
Grade 3 Pain, Dose 2 [N=543]	2			
Any Redness, Dose 2 [N=543]	23			
Grade 3 Redness, Dose 2 [N=543]	1			
Any Swelling, Dose 2 [N=543]	8			
Grade 3 Swelling, Dose 2 [N=543]	1			
Any Pain, Dose 3 [N=540]	41			
Grade 3 Pain, Dose 3 [N=540]	0			
Any Redness, Dose 3 [N=540]	15			
Grade 3 Redness, Dose 3 [N=540]	0			
Any Swelling, Dose 3 [N=540]	6			
Grade 3 Swelling, Dose 3 [N=540]	0			
Any Pain, Across doses [N=550]	113			
Grade 3 Pain, Across doses [N=550]	3			

Any Redness, Across doses [N=550]	46			
Grade 3 Redness, Across doses [N=550]	1			
Any Swelling, Across doses [N=550]	18			
Grade 3 Swelling, Across doses [N=550]	1			

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects with any, grade 3 and related solicited general symptoms

End point title	Number of subjects with any, grade 3 and related solicited general symptoms
-----------------	---

End point description:

Assessed solicited general symptoms were drowsiness, gastrointestinal symptoms, irritability/fussiness, loss of appetite and fever [defined as axillary temperature higher than (>) 37.0°C degrees Celsius]. Gastrointestinal symptoms included nausea, vomiting, diarrhoea and/or abdominal pain. Any = occurrence of any general symptom regardless of intensity grade or relationship to vaccination. Grade 3 drowsiness = drowsiness that prevented normal activity. Grade 3 irritability = crying that could not be comforted/ prevented normal activity. Grade 3 loss of appetite = subject did not eat at all. Grade 3 gastrointestinal symptoms = gastrointestinal symptoms that prevented normal activity. Grade 3 fever= temperature > 39°C. Related = symptom assessed by the investigator as causally related to the vaccination.

End point type	Secondary
----------------	-----------

End point timeframe:

During the 4-day (Days 0 -3) post-vaccination period following each vaccine dose and across doses

End point values	Poliorix Group	Control Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	550	550		
Units: Subjects				
Any Drowsiness, Dose 1 [N=550;550]	99	75		
Grade 3 Drowsiness, Dose 1 [N=550;550]	0	1		
Related Drowsiness, Dose 1 [N=550;550]	71	51		
Any Gastrointestinal symptoms, Dose 1 [N=550;550]	102	89		
Grade 3 Gastrointestinal symptoms, Dose 1 [N=550;550]	0	2		
Related Gastrointestinal symptoms, Dose 1 [N=550;550]	44	43		
Any Irritability/Fussiness, Dose 1 [N=550;550]	160	151		
Grade 3 Irritability/Fussiness, Dose 1 [N=550;550]	1	5		
Related Irritability/Fussiness, Dose 1 [N=550;550]	130	105		
Any Loss of appetite, Dose 1 [N=550;550]	82	83		
Grade 3 Loss of appetite, Dose 1 [N=550;550]	0	1		

Related Loss of appetite, Dose 1 [N=550;550]	54	47		
Any Temperature, Dose 1 [N=550;550]	39	20		
Grade 3 Temperature, Dose 1 [N=550;550]	0	0		
Related Temperature, Dose 1 [N=550;550]	28	10		
Any Drowsiness, Dose 2 [N=543;544]	68	50		
Grade 3 Drowsiness, Dose 2 [N=543;544]	2	0		
Related Drowsiness, Dose 2 [N=543;544]	51	25		
Any Gastrointestinal symptoms, Dose 2 [N=543;544]	66	67		
Grade 3 Gastrointestinal symptoms, Dose 2 [N=543;5]	2	1		
Related Gastrointestinal symptoms, Dose 2 [N=543;5]	32	35		
Any Irritability/Fussiness, Dose 2 [N=543;544]	121	86		
Grade 3 Irritability/Fussiness, Dose 2 [N=543;544]	7	2		
Related Irritability/Fussiness, Dose 2 [N=543;544]	105	50		
Any Loss of appetite, Dose 2 [N=543;544]	70	78		
Grade 3 Loss of appetite, Dose 2 [N=543;544]	3	0		
Related Loss of appetite, Dose 2 [N=543;544]	48	47		
Any Temperature, Dose 2 [N=543;544]	39	26		
Grade 3 Temperature, Dose 2 [N=543;544]	2	1		
Related Temperature, Dose 2 [N=543;544]	28	11		
Any Drowsiness, Dose 3 [N=540;534]	47	45		
Grade 3 Drowsiness, Dose 3 [N=540;534]	2	1		
Related Drowsiness, Dose 3 [N=540;534]	36	25		
Any Gastrointestinal sympoms, Dose 3 [N=540;534]	46	67		
Grade 3 Gastrointestinal sympoms, Dose 3 [N=540;5]	3	0		
Related Gastrointestinal sympoms, Dose 3 [N=540;5]	16	25		
Any Irritability/Fussiness, Dose 3 [N=540;534]	87	72		
Grade 3Irritability/Fussiness, Dose 3 [N=540;534]	2	2		
Related Irritability/Fussiness, Dose 3 [N=540;534]	77	44		
Any Loss of appetite, Dose 3 [N=540;534]	52	64		
Grade 3 Loss of appetite, Dose 3 [N=540;534]	0	0		
Related Loss of appetite, Dose 3 [N=540;534]	34	36		
Any Temperature, Dose 3 [N=540;534]	33	32		
Grade 3 Temperature, Dose 3 [N=540;534]	0	2		

Related Temperature, Dose 3 [N=540;534]	18	17		
Any Drowsiness, Across doses [N=550;550]	140	125		
Grade 3 Drowsiness, Across doses [N=550;550]	4	2		
Related Drowsiness, Across doses [N=550;550]	115	80		
Any Gastrointestinal, Across doses [N=550;550]	154	156		
Grade 3 Gastrointestinal, Across doses [N=550;550]	5	3		
Related Gastrointestinal, Across doses [N=550;550]	76	87		
Any Irritability, Across doses [N=550;550]	244	216		
Grade 3 Irritability, Across doses [N=550;550]	10	9		
Related Irritability, Across doses [N=550;550]	214	159		
Any Loss of appetite, Across doses [N=550;550]	147	168		
Grade 3 Loss of appetite, Across doses [N=550;550]	3	1		
Related Loss of appetite, Across doses [N=550;550]	100	109		
Any Fever, Across doses [N=550;550]	99	71		
Grade 3 Fever, Across doses [N=550;550]	2	3		
Related Fever, Across doses [N=550;550]	69	36		

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects with any unsolicited adverse events

End point title	Number of subjects with any unsolicited adverse events
-----------------	--

End point description:

An unsolicited AE covers any untoward medical occurrence in a clinical investigation subject temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product and reported in addition to those solicited during the clinical study and any solicited symptom with onset outside the specified period of follow-up for solicited symptoms. Any was defined as the occurrence of any unsolicited AE regardless of intensity grade or relation to vaccination.

End point type	Secondary
----------------	-----------

End point timeframe:

Within the 31-day (Days 0-30) post-vaccination period

End point values	Poliorix Group	Control Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	550	550		
Units: Subjects				
Any AE(s)	155	162		

Statistical analyses

No statistical analyses for this end point

Secondary: Number of subjects with serious adverse events (SAEs)

End point title	Number of subjects with serious adverse events (SAEs)
End point description:	
Serious adverse events (SAEs) assessed include medical occurrences that results in death, are life threatening, requires hospitalization or prolongation of hospitalization or results in disability/incapacity.	
End point type	Secondary
End point timeframe:	
During the entire study period (Day 0 to Month 3)	

End point values	Poliorix Group	Control Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	550	550		
Units: Subjects				
Any SAE(s)	3	6		

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

Solicited local and general symptoms: during the 4-day (Days 0-3) post-vaccination period. Unsolicited AEs: within the 31-day (Days 0-30) post-vaccination period. SAEs: during the entire study period (from Day 0 up to Month 3).

Adverse event reporting additional description:

The number of occurrences reported for adverse events were not available for posting. The number of subjects affected by each specific event was indicated as the number of occurrences.

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	13.1
--------------------	------

Reporting groups

Reporting group title	Control Group
-----------------------	---------------

Reporting group description:

Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Oral Poliomyelitis Vaccine (OPV) at 2, 3 and 4 months of age, according to the vaccination policy recommended in China.

Reporting group title	Poliorix Group
-----------------------	----------------

Reporting group description:

Healthy male and female Chinese infants between, and including 60 and 90 days of age, who received 3 doses of Poliorix (IPV) vaccine at 2, 3 and 4 months of age, administered intramuscularly into the anterolateral side of the right thigh.

Serious adverse events	Control Group	Poliorix Group	
Total subjects affected by serious adverse events			
subjects affected / exposed	6 / 550 (1.09%)	3 / 550 (0.55%)	
number of deaths (all causes)	0	0	
number of deaths resulting from adverse events	0	0	
Cardiac disorders			
Cardiac failure			
alternative assessment type: Non-systematic			
subjects affected / exposed	1 / 550 (0.18%)	0 / 550 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
Epilepsy			
alternative assessment type: Non-systematic			
subjects affected / exposed	1 / 550 (0.18%)	0 / 550 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Hydrocephalus alternative assessment type: Non-systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 550 (0.00%) 0 / 0 0 / 0	1 / 550 (0.18%) 0 / 1 0 / 0	
Gastrointestinal disorders Abdominal distension alternative assessment type: Non-systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 550 (0.18%) 0 / 1 0 / 0	0 / 550 (0.00%) 0 / 0 0 / 0	
Diarrhoea alternative assessment type: Non-systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 550 (0.00%) 0 / 0 0 / 0	1 / 550 (0.18%) 0 / 1 0 / 0	
Enteritis alternative assessment type: Non-systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 550 (0.18%) 0 / 1 0 / 0	0 / 550 (0.00%) 0 / 0 0 / 0	
Respiratory, thoracic and mediastinal disorders Respiratory failure alternative assessment type: Non-systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 550 (0.18%) 0 / 1 0 / 0	0 / 550 (0.00%) 0 / 0 0 / 0	
Infections and infestations Bronchopneumonia alternative assessment type: Non-systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all Bronchitis	2 / 550 (0.36%) 0 / 2 0 / 0	0 / 550 (0.00%) 0 / 0 0 / 0	

alternative assessment type: Non-systematic			
subjects affected / exposed	1 / 550 (0.18%)	0 / 550 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Herpes zoster			
alternative assessment type: Non-systematic			
subjects affected / exposed	0 / 550 (0.00%)	1 / 550 (0.18%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Upper respiratory tract infection			
alternative assessment type: Non-systematic			
subjects affected / exposed	1 / 550 (0.18%)	0 / 550 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 5 %

Non-serious adverse events	Control Group	Poliorix Group	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	366 / 550 (66.55%)	391 / 550 (71.09%)	
General disorders and administration site conditions			
Pain			
subjects affected / exposed	0 / 550 (0.00%)	113 / 550 (20.55%)	
occurrences (all)	0	113	
Drowsiness			
subjects affected / exposed	125 / 550 (22.73%)	140 / 550 (25.45%)	
occurrences (all)	125	140	
Gastrointestinal Symptoms			
subjects affected / exposed	156 / 550 (28.36%)	154 / 550 (28.00%)	
occurrences (all)	156	154	
Irritability/Fussiness			
subjects affected / exposed	216 / 550 (39.27%)	244 / 550 (44.36%)	
occurrences (all)	216	244	
Loss of appetite			

subjects affected / exposed occurrences (all)	168 / 550 (30.55%) 168	147 / 550 (26.73%) 147	
Fever (Axillary) subjects affected / exposed occurrences (all)	71 / 550 (12.91%) 71	99 / 550 (18.00%) 99	
Redness subjects affected / exposed occurrences (all)	46 / 550 (8.36%) 46	0 / 550 (0.00%) 0	
Infections and infestations			
Upper respiratory tract infection subjects affected / exposed occurrences (all)	97 / 550 (17.64%) 97	99 / 550 (18.00%) 99	
Nasopharyngitis subjects affected / exposed occurrences (all)	38 / 550 (6.91%) 38	32 / 550 (5.82%) 32	

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? No

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported