



Clinical trial results:

A Phase Ib/II Study of GDC-0068 or GDC-0980 with Abiraterone Acetate versus Abiraterone Acetate in Patients with Castration-Resistant Prostate Cancer Previously Treated with Docetaxel-Based Chemotherapy

Summary

EudraCT number	2011-004126-10
Trial protocol	GR CZ ES GB NL FR IT
Global end of trial date	

Results information

Result version number	v1
This version publication date	07 December 2016
First version publication date	07 December 2016

Trial information

Trial identification

Sponsor protocol code	GO27983
-----------------------	---------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT01485861
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	F. Hoffmann-La Roche AG
Sponsor organisation address	Grenzacherstrasse 124, CH, Basel, Switzerland, 4070
Public contact	Roche Trial Information Hotline, F. Hoffmann-La Roche AG, +41 61 6878333, global.trial_information@roche.com
Scientific contact	Roche Trial Information Hotline, F. Hoffmann-La Roche AG, +41 61 6878333, global.trial_information@roche.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

Results analysis stage

Analysis stage	Interim
Date of interim/final analysis	01 September 2015
Is this the analysis of the primary completion data?	Yes
Primary completion date	01 September 2015
Global end of trial reached?	No

Notes:

General information about the trial

Main objective of the trial:

The primary objectives of the Phase Ib portion of the study were as follows:

To evaluate the safety and tolerability of ipatasertib (Protein kinase B [Akt] inhibitor) administered in combination with abiraterone and of GDC-0980 (phosphatidylinositol 3 kinase [PI3K], and/or mammalian target of rapamycin [mTOR] inhibitor) administered in combination with abiraterone.

To identify dose-limiting toxicities, estimate the maximum tolerated dose, and identify a recommended Phase II dose of ipatasertib administered in combination with abiraterone and of GDC-0980 administered in combination with abiraterone.

The primary objective of the Phase II portion of the study was to estimate the efficacy as measured by radiographic progression-free survival of ipatasertib (dosed at either 400 milligrams [mg] or 200 mg daily) + abiraterone and prednisone/prednisolone versus placebo + abiraterone and prednisone/prednisolone.

Protection of trial subjects:

The study was conducted in accordance with the principles of the "Declaration of Helsinki" and Good Clinical Practice (GCP) according to the regulations and procedures.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	11 January 2012
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Netherlands: 11
Country: Number of subjects enrolled	Spain: 46
Country: Number of subjects enrolled	United Kingdom: 24
Country: Number of subjects enrolled	Czech Republic: 22
Country: Number of subjects enrolled	France: 34
Country: Number of subjects enrolled	Greece: 16
Country: Number of subjects enrolled	Italy: 40
Country: Number of subjects enrolled	Romania: 42
Country: Number of subjects enrolled	United States: 38
Worldwide total number of subjects	273
EEA total number of subjects	235

Notes:

Subjects enrolled per age group	
In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	87
From 65 to 84 years	182
85 years and over	4

Subject disposition

Recruitment

Recruitment details: -

Pre-assignment

Screening details:

The study consisted 2 stages: Phase Ib determined recommended Phase II dose (RP2D) for ipatasertib and GDC-0980 in combination with abiraterone and prednisone/prednisolone. Phase II stage compared ipatasertib (400 mg or 200 mg daily) with abiraterone and prednisone/prednisolone versus placebo with abiraterone and prednisone/prednisolone.

Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

Blinding implementation details:

The Phase Ib portion of the study was open-label. In Phase II, participants and investigators were blinded with regard to treatment status (i.e., ipatasertib vs. placebo).

Arms

Are arms mutually exclusive?	No
Arm title	Phase Ib: Ipatasertib 400 mg

Arm description:

Participants received ipatasertib 400 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg twice daily (bid) orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Arm type	Experimental
Investigational medicinal product name	Ipatasertib
Investigational medicinal product code	
Other name	GDC-0068
Pharmaceutical forms	Capsule, hard
Routes of administration	Oral use

Dosage and administration details:

Ipatasertib was administered orally once daily beginning on Day 1 of Cycle 1 until disease progression or intolerable toxicity. Ipatasertib could be dosed with or without food. If a dose was missed (not taken within 8 hours after the scheduled dosing time), the participant had to resume dosing with the next scheduled dose.

Investigational medicinal product name	Abiraterone acetate
Investigational medicinal product code	
Other name	Zytiga®
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Abiraterone had to be taken on an empty stomach. No food was to be consumed for greater than or equal to (\geq) 2 hours before the dose of abiraterone was taken and for \geq 1 hour after the dose of abiraterone had been taken.

Investigational medicinal product name	Prednisone/Prednisolone
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Prednisone/prednisolone was to be taken per local clinical practice/investigator recommendation.

Arm title	Phase Ib: GDC-0980 30 mg
------------------	--------------------------

Arm description:

Participants received GDC-0980 30 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Arm type	Experimental
Investigational medicinal product name	GDC-0980
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

GDC-0980 was administered daily beginning on Day 1 of Cycle 1 until disease progression or intolerable toxicity. GDC-0980 could be dosed with or without food. If a dose was missed (not taken within 8 hours after the scheduled dosing time), the participant had to resume dosing with the next scheduled dose.

Investigational medicinal product name	Abiraterone acetate
Investigational medicinal product code	
Other name	Zytiga®
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Abiraterone had to be taken on an empty stomach. No food was to be consumed for ≥ 2 hours before the dose of abiraterone was taken and for ≥ 1 hour after the dose of abiraterone had been taken.

Investigational medicinal product name	Prednisone/Prednisolone
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Prednisone/prednisolone was to be taken per local clinical practice/investigator recommendation.

Arm title	Phase II: Ipatasertib 400 mg + Abiraterone
------------------	--

Arm description:

Participants received ipatasertib 400 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Arm type	Experimental
Investigational medicinal product name	Ipatasertib
Investigational medicinal product code	
Other name	GDC-0068
Pharmaceutical forms	Capsule, hard
Routes of administration	Oral use

Dosage and administration details:

Ipatasertib was administered orally once daily beginning on Day 1 of Cycle 1 until disease progression or intolerable toxicity. Ipatasertib could be dosed with or without food. If a dose was missed (not taken within 8 hours after the scheduled dosing time), the participant had to resume dosing with the next scheduled dose.

Investigational medicinal product name	Abiraterone acetate
Investigational medicinal product code	
Other name	Zytiga®
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Abiraterone had to be taken on an empty stomach. No food was to be consumed for ≥ 2 hours before the dose of abiraterone was taken and for ≥ 1 hour after the dose of abiraterone had been taken.

Investigational medicinal product name	Prednisone/Prednisolone
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Prednisone/prednisolone was to be taken per local clinical practice/investigator recommendation.

Arm title	Phase II: Ipatasertib 200 mg + Abiraterone
------------------	--

Arm description:

Participants received ipatasertib 200 mg orally once daily, abiraterone 1000 mg once orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Arm type	Experimental
Investigational medicinal product name	Ipatasertib
Investigational medicinal product code	
Other name	GDC-0068
Pharmaceutical forms	Capsule, hard
Routes of administration	Oral use

Dosage and administration details:

Ipatasertib was administered orally once daily beginning on Day 1 of Cycle 1 until disease progression or intolerable toxicity. Ipatasertib could be dosed with or without food. If a dose was missed (not taken within 8 hours after the scheduled dosing time), the participant had to resume dosing with the next scheduled dose.

Investigational medicinal product name	Abiraterone acetate
Investigational medicinal product code	
Other name	Zytiga®
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Abiraterone had to be taken on an empty stomach. No food was to be consumed for ≥ 2 hours before the dose of abiraterone was taken and for ≥ 1 hour after the dose of abiraterone had been taken.

Investigational medicinal product name	Prednisone/Prednisolone
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Prednisone/prednisolone was to be taken per local clinical practice/investigator recommendation.

Arm title	Phase II: Placebo + Abiraterone
------------------	---------------------------------

Arm description:

Participants received placebo (matched to ipatasertib 400 mg or 200) orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Arm type	Placebo
----------	---------

Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule, hard
Routes of administration	Oral use

Dosage and administration details:

Placebo (for Ipatasertib 400 mg or 200 mg) was administered daily beginning on Day 1 of Cycle 1 until disease progression or intolerable toxicity. Placebo could be dosed with or without food. If a dose was missed (not taken within 8 hours after the scheduled dosing time), the participant had to resume dosing with the next scheduled dose.

Investigational medicinal product name	Abiraterone acetate
Investigational medicinal product code	
Other name	Zytiga®
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Abiraterone had to be taken on an empty stomach. No food was to be consumed for ≥ 2 hours before the dose of abiraterone was taken and for ≥ 1 hour after the dose of abiraterone had been taken.

Investigational medicinal product name	Prednisone/Prednisolone
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Prednisone/prednisolone was to be taken per local clinical practice/investigator recommendation.

Number of subjects in period 1	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg	Phase II: Ipatasertib 400 mg + Abiraterone
Started	14	6	84
Treated	14	6	84
Completed	0	0	0
Not completed	14	6	84
Physician decision	1	1	-
Consent withdrawn by subject	4	1	5
Death	1	-	29
In survival follow-up	-	-	30
Transferred to other arm/group	-	-	-
Adverse event	-	4	-
Unspecified	3	-	-
Progression of disease	5	-	2
Treatment on-going	-	-	18
Lost to follow-up	-	-	-
Joined	0	0	0
Transferred in from other group/arm	-	-	-

Number of subjects in period 1	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone
Started	86	83
Treated	87	82
Completed	0	0
Not completed	87	83
Physician decision	-	-
Consent withdrawn by subject	1	3
Death	36	38
In survival follow-up	29	25
Transferred to other arm/group	-	1
Adverse event	-	-
Unspecified	1	-
Progression of disease	1	-
Treatment on-going	18	13
Lost to follow-up	1	3
Joined	1	0
Transferred in from other group/arm	1	-

Baseline characteristics

Reporting groups^[1]

Reporting group title	Phase Ib: Ipatasertib 400 mg
-----------------------	------------------------------

Reporting group description:

Participants received ipatasertib 400 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg twice daily (bid) orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase Ib: GDC-0980 30 mg
-----------------------	--------------------------

Reporting group description:

Participants received GDC-0980 30 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase II: Ipatasertib 400 mg + Abiraterone
-----------------------	--

Reporting group description:

Participants received ipatasertib 400 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase II: Ipatasertib 200 mg + Abiraterone
-----------------------	--

Reporting group description:

Participants received ipatasertib 200 mg orally once daily, abiraterone 1000 mg once orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase II: Placebo + Abiraterone
-----------------------	---------------------------------

Reporting group description:

Participants received placebo (matched to ipatasertib 400 mg or 200) orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Notes:

[1] - The number of subjects reported to be in the baseline period is not equal to the worldwide number of subjects enrolled in the trial. It is expected that these numbers will be the same.

Justification: 1 participant in Phase II was randomized to the Placebo + Abiraterone group, but took 5 consecutive days of ipatasertib and therefore was assigned to ipatasertib 200 mg + Abiraterone. A total of 273 participants were included.

Reporting group values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg	Phase II: Ipatasertib 400 mg + Abiraterone
Number of subjects	14	6	84
Age categorical Units: Subjects			

Age continuous			
Number of participants analysed for this parameter were 14, 6, 84, 86, 83, respectively.			
Units: years			
arithmetic mean	68.1	69.3	66.9
standard deviation	± 9.1	± 9.3	± 8.5
Gender categorical Units: Subjects			
Female	0	0	0
Male	14	6	84

Reporting group values	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	Total
Number of subjects	87	83	273

Age categorical			
Units: Subjects			
Age continuous			
Number of participants analysed for this parameter were 14, 6, 84, 86, 83, respectively.			
Units: years			
arithmetic mean	68.8	67.6	
standard deviation	± 7.2	± 7.8	-
Gender categorical			
Units: Subjects			
Female	0	0	0
Male	87	83	273

End points

End points reporting groups

Reporting group title	Phase Ib: Ipatasertib 400 mg
Reporting group description: Participants received ipatasertib 400 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg twice daily (bid) orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.	
Reporting group title	Phase Ib: GDC-0980 30 mg
Reporting group description: Participants received GDC-0980 30 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.	
Reporting group title	Phase II: Ipatasertib 400 mg + Abiraterone
Reporting group description: Participants received ipatasertib 400 mg orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.	
Reporting group title	Phase II: Ipatasertib 200 mg + Abiraterone
Reporting group description: Participants received ipatasertib 200 mg orally once daily, abiraterone 1000 mg once orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.	
Reporting group title	Phase II: Placebo + Abiraterone
Reporting group description: Participants received placebo (matched to ipatasertib 400 mg or 200) orally once daily, abiraterone 1000 mg orally once daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.	

Primary: Percentage of Participants With Dose-Limiting Toxicity (DLTs)

End point title	Percentage of Participants With Dose-Limiting Toxicity
End point description: DLT: 1 of the following toxicities, at least possibly related to ipatasertib or GDC-0980. 1) Grade ≥ 3 non-hematologic, non-hepatic major organ AE; 2) Grade ≥ 3 febrile neutropenia; 3) Grade ≥ 4 neutropenia (absolute neutrophils less than [$<$] 500 per microliter) lasting greater than ($>$) 7 days; 4) Grade ≥ 3 thrombocytopenia associated with acute hemorrhage; 5) Grade ≥ 4 thrombocytopenia; 6) Grade ≥ 4 anemia; 7) 1 episode of fasting Grade ≥ 4 hyperglycemia or 3 episodes of fasting Grade 3 hyperglycemia on separate days within 7 days, as determined by laboratory blood glucose evaluation; 8) Grade ≥ 3 elevation lasting for > 48 hours for hepatic transaminase or liver-specific alkaline phosphatase or total bilirubin. Adverse events were graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events v 4.0. Safety population: all participants who had received ipatasertib, GDC-0980, placebo, or abiraterone treatment on Day 1 of Cycle 1.	
End point type	Primary
End point timeframe: Days 1 to 28 of Cycle 1 (Cycle length = 28 days)	
Notes: [1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point. Justification: No statistical analysis was planned for this endpoint. [2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.	

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	14	6		
Units: percentage of participants				
number (not applicable)	0	0		

Statistical analyses

No statistical analyses for this end point

Primary: Phase Ib: Percentage of Participants by Nature of DLTs

End point title	Phase Ib: Percentage of Participants by Nature of DLTs ^[3] ^[4]
-----------------	--

End point description:

DLT: 1 of the following toxicities, at least possibly related to ipatasertib or GDC-0980. 1) Grade \geq 3 non-hematologic, non-hepatic major organ AE; 2) Grade \geq 3 febrile neutropenia; 3) Grade \geq 4 neutropenia (absolute neutrophils $<$ 500 per microliter) lasting $>$ 7 days; 4) Grade \geq 3 thrombocytopenia associated with acute hemorrhage; 5) Grade \geq 4 thrombocytopenia; 6) Grade \geq 4 anemia; 7) 1 episode of fasting Grade \geq 4 hyperglycemia or 3 episodes of fasting Grade 3 hyperglycemia on separate days within 7 days, as determined by laboratory blood glucose evaluation; 8) Grade \geq 3 elevation lasting for $>$ 48 hours for hepatic transaminase or liver-specific alkaline phosphatase or total bilirubin. Adverse events were graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events v 4.0. Analysis was to be performed on safety population participants who had at least 1 DLT.

End point type	Primary
----------------	---------

End point timeframe:

Days 1 to 28 of Cycle 1 (Cycle length = 28 days)

Notes:

[3] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: No statistical analysis was planned for this endpoint.

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	0 ^[5]	0 ^[6]		
Units: percentage of participants				
number (not applicable)				

Notes:

[5] - No participant experienced any DLT.

[6] - No participant experienced any DLT.

Statistical analyses

No statistical analyses for this end point

Primary: Phase Ib: Recommended Phase II Dose (RP2D) of Ipatasertib

End point title	Phase Ib: Recommended Phase II Dose (RP2D) of
-----------------	---

End point description:

RP2D is a dose of a drug which would be used in Phase II stage of the study. RP2D was to be determined based on maximum tolerated dose (MTD) in Phase Ib stage of the study. The highest dose level (in 3+3 escalation scheme) with an acceptable safety profile and with a minimum of 6 participants at which fewer than one-third of participants experienced a DLT was declared the MTD and RP2D. Analysis was performed on safety population.

End point type	Primary
----------------	---------

End point timeframe:

Days 1 to 28 of Cycle 1 (Cycle length = 28 days)

Notes:

[7] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The endpoint is qualitative in nature. Hence, no statistical analysis was performed.

[8] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	14	0 ^[9]		
Units: milligrams				
number (not applicable)	400			

Notes:

[9] - Based on safety analysis, further testing of GDC-0980 arm was discontinued.

Statistical analyses

No statistical analyses for this end point

Primary: Phase II: Percentage of Participants With Radiographic Progression (as Assessed by Response Evaluation Criteria in Solid Tumors [RECIST] 1.1) or Death on Study - Intent to Treat (ITT) Population

End point title	Phase II: Percentage of Participants With Radiographic Progression (as Assessed by Response Evaluation Criteria in Solid Tumors [RECIST] 1.1) or Death on Study - Intent to Treat (ITT) Population ^{[10][11]}
-----------------	--

End point description:

Tumor assessment was done by investigator via computed tomography (CT) scan and bone scans.

Progression was defined as follows:

Soft tissue mass (RECIST 1.1): at least 20 percent (%) increase in sum of diameters of target lesions compared to smallest sum of diameters on-study and absolute increase of at least 5 millimeters (mm), progression of existing non-target lesions, or presence of new lesions.

Bone: ≥ 2 new bone lesions plus 2 additional at confirmation on a second bone scan ≥ 4 weeks later (< 12 weeks after randomization). ≥ 2 new bone lesions consistent with progression, without need for confirmatory bone scan (≥ 12 weeks after randomization). ITT Population: all randomized participants with the participants allocated to the treatment arm according to the randomization.

End point type	Primary
----------------	---------

End point timeframe:

Screening up to radiographic progression or death, whichever occurred first (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[10] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The endpoint is qualitative in nature. Hence, no statistical analysis was performed.

[11] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: percentage of participants				
number (not applicable)	59.5	68.6	77.1	

Statistical analyses

No statistical analyses for this end point

Primary: Phase II: Radiographic Progression Free Survival as Assessed by RECIST 1.1- ITT Population

End point title	Phase II: Radiographic Progression Free Survival as Assessed by RECIST 1.1- ITT Population ^[12]
-----------------	--

End point description:

Radiographic PFS: Time from randomization to the first occurrence of disease progression, as determined by investigator review of tumor assessments via CT scan and bone scans, or death on study from any cause, whichever occurred first. Progression was defined as follows:

Soft tissue mass (RECIST 1.1): at least 20% increase in sum of diameters of target lesions compared to smallest sum of diameters on-study and absolute increase of at least 5 mm, progression of existing non-target lesions, or presence of new lesions.

Bone: ≥ 2 new bone lesions plus 2 additional at confirmation on a second bone scan ≥ 4 weeks later (< 12 weeks after randomization). ≥ 2 new bone lesions consistent with progression, without need for confirmatory bone scan (≥ 12 weeks after randomization). Analysis was performed on ITT population.

End point type	Primary
----------------	---------

End point timeframe:

Screening up to radiographic progression or death, whichever occurred first (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[12] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: months				
median (confidence interval 90%)	8.18 (6.67 to 10.87)	8.31 (6.44 to 10.48)	6.37 (4.6 to 8.34)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.1606
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.77
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.56
upper limit	1.04

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.53
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.89
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.66
upper limit	1.2

Statistical analysis title	Stratified analysis:Ipatasertib 400 mg v Placebo
Statistical analysis description:	
Strata were: prior enzalutamide (Yes vs. No), progression factor (prostate-specific antigen [PSA] only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.1689
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.75
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.54
upper limit	1.05

Statistical analysis title	Stratified analysis: Ipatasertib 200 mg v Placebo
-----------------------------------	---

Statistical analysis description:

Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).

Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7484
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.94
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.69
upper limit	1.28

Primary: Phase II: Percentage of Participants With Radiographic Progression (as Assessed by RECIST 1.1) or Death in Participants With Institute of Cancer Research (ICR) Phosphatase and Tensin Homolog (PTEN) Loss

End point title	Phase II: Percentage of Participants With Radiographic Progression (as Assessed by RECIST 1.1) or Death in Participants With Institute of Cancer Research (ICR) Phosphatase and Tensin Homolog (PTEN) Loss ^{[13][14]}
-----------------	--

End point description:

Tumor assessment was done by investigator via CT scan and bone scans. Progression was defined as follows:

Soft tissue mass (RECIST 1.1): at least 20% increase in sum of diameters of target lesions compared to smallest sum of diameters on-study and absolute increase of at least 5 mm, progression of existing non-target lesions, or presence of new lesions.

Bone: ≥2 new bone lesions plus 2 additional at confirmation on a second bone scan ≥4 weeks later (< 12 weeks after randomization). ≥2 new bone lesions consistent with progression, without need for confirmatory bone scan (≥12 weeks after randomization). PTEN status was assessed by a research-use-only (RUO) immunohistochemical (IHC) assay that was performed at The Institute of Cancer Research (ICR), United Kingdom (UK). Samples with 100% of the tumor with no PTEN staining were classified as

"ICR PTEN loss". Number of participants analyzed=participants with PTEN loss.

End point type	Primary
----------------	---------

End point timeframe:

Screening up to radiographic progression or death, whichever occurred first (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[13] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The endpoint is qualitative in nature. Hence, no statistical analysis was performed.

[14] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: percentage of participants				
number (not applicable)	60	64	85.7	

Statistical analyses

No statistical analyses for this end point

Primary: Phase II: Radiographic Progression Free Survival (as Assessed by RECIST 1.1) in Participants With ICR PTEN Loss

End point title	Phase II: Radiographic Progression Free Survival (as Assessed by RECIST 1.1) in Participants With ICR PTEN Loss ^[15]
-----------------	---

End point description:

Radiographic PFS: Time from randomization to the first occurrence of disease progression, as determined by investigator review of tumor assessments via CT scan and bone scans, or death on study from any cause, whichever occurred first. Progression was defined as follows:

Soft tissue mass (RECIST 1.1): at least 20% increase in sum of diameters of target lesions compared to smallest sum of diameters on-study and absolute increase of at least 5 mm, progression of existing non-target lesions, or presence of new lesions.

Bone: ≥ 2 new bone lesions plus 2 additional at confirmation on 2nd bone scan ≥ 4 weeks later (< 12 weeks after randomization). ≥ 2 new bone lesions consistent with progression, without need for confirmatory bone scan (≥ 12 weeks after randomization). PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss.

End point type	Primary
----------------	---------

End point timeframe:

Screening up to radiographic progression or death, whichever occurred first (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[15] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: months				
median (confidence interval 90%)	11.53 (6.67 to 13.73)	11.1 (6.34 to 16.36)	4.6 (4.4 to 6.37)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0064
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.39
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.22
upper limit	0.7

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Placebo + Abiraterone v Phase II: Ipatasertib 200 mg + Abiraterone
Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0285
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.46
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.25
upper limit	0.83

Secondary: Phase II: Percentage of Participants Who Died - ITT Population

End point title	Phase II: Percentage of Participants Who Died - ITT Population
-----------------	--

End point description:

Percentage of participants who died due to any cause. Analysis was performed on ITT population.

End point type Secondary

End point timeframe:

Screening up to death (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[16] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: percentage of participants				
number (not applicable)	35.7	41.9	45.8	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II: Overall Survival - ITT Population

End point title Phase II: Overall Survival - ITT Population^[17]

End point description:

Overall survival was defined as the interval between the date of screening and death due to any cause. Overall survival was estimated using Kaplan Meier method. Analysis was performed on ITT population. Here, "99999" indicates that the data was not evaluable due to insufficient number of events.

End point type Secondary

End point timeframe:

Screening up to death (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[17] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: months				
median (confidence interval 90%)	18.92 (17.12 to 99999)	21.52 (13.27 to 99999)	15.64 (13.77 to 19.42)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.3446
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.79
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.53
upper limit	1.19

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8031
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.94
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.64
upper limit	1.38

Statistical analysis title	Stratified analysis:Ipatasertib 400 mg v Placebo
Statistical analysis description:	
Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2159
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.72
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.47
upper limit	1.11

Statistical analysis title	Stratified analysis: Ipatasertib 200 mg v Placebo
-----------------------------------	---

Statistical analysis description:

Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).

Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8747
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.96
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.65
upper limit	1.43

Secondary: Phase II: Percentage of Participants Who Died in Participants With ICR PTEN Loss

End point title	Phase II: Percentage of Participants Who Died in Participants With ICR PTEN Loss ^[18]
-----------------	--

End point description:

Percentage of participants who died due to any cause. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to death (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[18] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: percentage of participants				
number (not applicable)	36	32	57.1	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II: Overall Survival in Participants With ICR IHC PTEN Loss

End point title	Phase II: Overall Survival in Participants With ICR IHC PTEN Loss ^[19]
-----------------	---

End point description:

Overall survival was defined as the interval between the date of randomization and death from any cause. Overall survival was estimated using Kaplan Meier method. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss. Here, "99999" indicates that the data was not evaluable due to insufficient number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to death (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[19] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: months				
median (confidence interval 90%)	19.15 (12.09 to 19.15)	99999 (15.74 to 99999)	14.75 (11.3 to 18.43)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2993
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.62
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.29
upper limit	1.33

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.3329
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.64
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.3
upper limit	1.37

Secondary: Phase II: Percentage of Participants With PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) - ITT Population

End point title	Phase II: Percentage of Participants With PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) - ITT Population ^[20]
-----------------	--

End point description:

PSA progression was defined as per Prostate Cancer Working Group 2 criteria. PSA progression was defined as a $\geq 25\%$ increase and an absolute increase of ≥ 2 nanogram per milliliter (ng/mL) from the baseline value, or a $\geq 25\%$ increase and an absolute increase of ≥ 2 ng/mL from the nadir if PSA decreases from baseline after treatment, which was confirmed by a second value obtained ≥ 3 weeks later. Analysis was performed on ITT population.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to PSA progression (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[20] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: percentage of participants				
number (not applicable)	57.1	69.8	72.3	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II: Time to PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) - ITT Population

End point title	Phase II: Time to PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) - ITT Population ^[21]
-----------------	--

End point description:

Time to PSA progression: Time from screening to the first occurrence of PSA progression, as determined by investigator. PSA progression was defined as per Prostate Cancer Working Group 2 criteria. PSA progression is defined as a $\geq 25\%$ increase and an absolute increase of ≥ 2 ng/mL from the baseline value, or a $\geq 25\%$ increase and an absolute increase of ≥ 2 ng/mL from the nadir if PSA decreases from baseline after treatment, which was confirmed by a second value obtained ≥ 3 weeks later. Analysis was performed on ITT population.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to PSA progression (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[21] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: months				
median (confidence interval 90%)	5.55 (4.17 to 7.39)	3.78 (2.79 to 5.49)	3.71 (2.79 to 4.67)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0665
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.7
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.51
upper limit	0.97

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Placebo + Abiraterone v Phase II: Ipatasertib 200 mg + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.9319
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.99
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.73
upper limit	1.33

Statistical analysis title	Stratified analysis:Ipatasertib 400 mg v Placebo
Statistical analysis description:	
Strata were: prior Enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.071
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.7
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.5
upper limit	0.97

Statistical analysis title	Stratified analysis:Ipatasertib 200 mg v Placebo
Statistical analysis description: Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.789
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.95
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.7
upper limit	1.31

Secondary: Phase II: Percentage of Participants With PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) in Participants With ICR PTEN Loss

End point title	Phase II: Percentage of Participants With PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) in Participants With ICR PTEN Loss ^[22]
-----------------	--

End point description:

PSA progression was defined as per Prostate Cancer Working Group 2 criteria. PSA progression was defined as a $\geq 25\%$ increase and an absolute increase of ≥ 2 ng/mL from the baseline value, or a $\geq 25\%$ increase and an absolute increase of ≥ 2 ng/mL from the nadir if PSA decreases from baseline after treatment, which was confirmed by a second value obtained ≥ 3 weeks later. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to PSA progression (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[22] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: percentage of participants				
number (not applicable)	72	64	66.7	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II: Time to PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) in Participants With ICR PTEN Loss

End point title	Phase II: Time to PSA Progression (as Assessed by Prostate Cancer Working Group 2 Criteria) in Participants With ICR PTEN Loss ^[23]
-----------------	--

End point description:

Time to PSA progression: Time from screening to the first occurrence of PSA progression, as determined by investigator. PSA progression was defined as per Prostate Cancer Working Group 2 criteria. PSA progression was defined as a $\geq 25\%$ increase and an absolute increase of ≥ 2 ng/mL from the baseline value, or a $\geq 25\%$ increase and an absolute increase of ≥ 2 ng/mL from the nadir if PSA decreases from baseline after treatment, which was confirmed by a second value obtained ≥ 3 weeks later. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to PSA progression (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[23] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: months				
median (confidence interval 90%)	3.71 (2.99 to 8.18)	2.92 (2.07 to 7.39)	2.79 (1.02 to 4.67)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2906
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.68
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.37
upper limit	1.25

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2716
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.65
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.35
upper limit	1.22

Secondary: Phase II: Percentage of Participants With PSA Response - ITT Population

End point title	Phase II: Percentage of Participants With PSA Response - ITT Population ^[24]
-----------------	---

End point description:

PSA response was defined as a > 50% decrease in PSA from baseline, which was to be confirmed after ≥ 4 weeks by a confirmatory PSA measurement. Analysis was performed on ITT population.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to 30 days after last dose (assessed at baseline, Day 1 of every cycle [starting from Cycle 2] till 30 days after last dose [up to overall 3.6 years]) (cycle length = 28 days)

Notes:

[24] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: percentage of participants				
number (confidence interval 90%)	36.9 (28.11 to 46.4)	33.7 (25.29 to 42.93)	34.9 (26.25 to 43.75)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7913
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	1.97
Confidence interval	
level	90 %
sides	2-sided
lower limit	-10.25
upper limit	14.18

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8675
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	-1.22
Confidence interval	
level	90 %
sides	2-sided
lower limit	-13.24
upper limit	10.8

Secondary: Phase II: Percentage of Participants With PSA Response in Participants With ICR PTEN Loss

End point title	Phase II: Percentage of Participants With PSA Response in Participants With ICR PTEN Loss ^[25]
-----------------	---

End point description:

PSA response was defined as a > 50% decrease in PSA from baseline, which was to be confirmed after ≥ 4 weeks by a confirmatory PSA measurement. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to 30 days after last dose (assessed at baseline, Day 1 of every cycle [starting from Cycle 2] till 30 days after last dose [up to overall 3.6 years]) (cycle length = 28 days)

Notes:

[25] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: percentage of participants				
number (confidence interval 90%)	40 (24.57 to 58.32)	44 (26.99 to 61.06)	28.6 (13.24 to 46.41)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.4176
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	11.43
Confidence interval	
level	90 %
sides	2-sided
lower limit	-11.43
upper limit	58.32

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2802
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	15.43
Confidence interval	
level	90 %
sides	2-sided
lower limit	-7.58
upper limit	38.44

Secondary: Phase II: Percentage of Participants With Objective Response as Assessed by RECIST 1.1 - ITT Population

End point title	Phase II: Percentage of Participants With Objective Response as Assessed by RECIST 1.1 - ITT Population ^[26]
-----------------	---

End point description:

Objective response was defined as having a confirm response (CR) or partial response (PR) according to a RECIST 1.1. CR was defined as disappearance of all target and non-target lesions and no new lesions, all pathological lymph nodes must have decreased to <10 mm in short axis. PR was defined as at least a 30% decrease in the sum of diameters of target lesions (taking as reference the baseline sum diameters), no progression in non-target lesions, and no new lesions. Analysis was performed on ITT population. Number of participants analysed=participants who were evaluable for this endpoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to radiographic progression or death, whichever occurred first (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[26] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	37	39	35	
Units: percentage of participants				
number (confidence interval 90%)	32.4 (20.56 to 46.41)	23.1 (13.69 to 35.63)	22.9 (11.91 to 36.46)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	72
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.3646
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	9.58
Confidence interval	
level	90 %
sides	2-sided
lower limit	-7.65
upper limit	26.8

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	74
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.9821
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	0.22
Confidence interval	
level	90 %
sides	2-sided
lower limit	-15.89
upper limit	16.33

Secondary: Phase II: Percentage of Participants With Objective Response (as Assessed by RECIST 1.1) in Participants With ICR PTEN Loss

End point title	Phase II: Percentage of Participants With Objective Response (as Assessed by RECIST 1.1) in Participants With ICR PTEN Loss ^[27]
-----------------	---

End point description:

Objective response was defined as having a CR or PR according to a RECIST 1.1. CR was defined as disappearance of all target and non-target lesions and no new lesions, all pathological lymph nodes must have decreased to <10 mm in short axis. PR was defined as at least a 30% decrease in the sum of diameters of target lesions (taking as reference the baseline sum diameters), no progression in non-target lesions, and no new lesions. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss and evaluable for this endpoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to radiographic progression or death, whichever occurred first (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[27] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	9	15	7	
Units: percentage of participants				
number (confidence interval 90%)	11.1 (1.16 to 39.09)	26.7 (12.18 to 50)	14.3 (1.49 to 50)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	16
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8489
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	-3.17
Confidence interval	
level	90 %
sides	2-sided
lower limit	-30.93
upper limit	24.58

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	22
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.5186
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	12.38
Confidence interval	
level	90 %
sides	2-sided
lower limit	-16.36
upper limit	41.12

Secondary: Phase II: Duration of Tumor Response - ITT Population

End point title	Phase II: Duration of Tumor Response - ITT Population ^[28]
-----------------	---

End point description:

Duration of tumor response: time period from 1st documentation of objective response (CR/PR) (whichever status was recorded first) to date of 1st occurrence of investigator documented disease progression or death. CR: disappearance of all target and non-target lesions and no new lesions, all pathological lymph nodes must have decreased to <10 mm in short axis. PR: at least 30% decrease in sum of diameters of target lesions (taking as reference baseline sum diameters), no progression in non-target lesions, and no new lesions. Progression: increase by at least 20% in sum of longest diameters of each target lesion, taking as reference smallest sum of longest diameters or appearance of one or more new lesions. Duration of tumor response was estimated using Kaplan Meier method. Number of participants analysed=participants from ITT population who had achieved an objective response. Here, 99999 reflects that data was not evaluable due to insufficient number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening up to radiographic progression or death, whichever occurred first (assessed at screening, after Cycles 3, 5, 7, 9, every three cycles [12 weeks] thereafter up to end of treatment [up to 3.6 years overall]) (cycle length = 28 days)

Notes:

[28] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	12	9	8	
Units: months				
median (confidence interval 90%)	8.77 (3.75 to 99999)	99999 (6.47 to 99999)	99999 (2.76 to 99999)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Placebo + Abiraterone v Phase II: Ipatasertib 400 mg + Abiraterone
Number of subjects included in analysis	20
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.678
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	1.31

Confidence interval	
level	90 %
sides	2-sided
lower limit	0.45
upper limit	3.8

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	17
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8372
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.85
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.24
upper limit	3.02

Statistical analysis title	Stratified analysis:Ipatasertib 400 mg v Placebo
Statistical analysis description:	
Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	20
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7733
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.77
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.17
upper limit	3.5

Statistical analysis title	Stratified analysis:Ipatasertib 200 mg v Placebo
Statistical analysis description:	
Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	

Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	17
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.4227
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	2.46
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.36
upper limit	16.59

Secondary: Phase II: Percentage of Participants With Circulating Tumor Cells (CTC) Reduction Response - ITT Population

End point title	Phase II: Percentage of Participants With Circulating Tumor Cells (CTC) Reduction Response - ITT Population ^[29]
End point description:	CTC reduction response was defined as participants with a reduction in CTCs of $\geq 30\%$ compared to baseline. Analysis was performed on ITT population. Number of participants analysed=participants with CTC > 0 cells/7.5 milliliters (mL) at baseline.
End point type	Secondary
End point timeframe:	Baseline, on Day 1 of Cycle 2, on Day 1 of Cycle 3, and at the treatment completion visit (up to overall 3.6 years) (cycle length=28 days)

Notes:

[29] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	58	62	63	
Units: percentage of participants				
number (confidence interval 90%)	67.2 (56.44 to 77.37)	71 (61.07 to 80.32)	63.5 (52.41 to 73.6)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	121
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.6652
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	3.75
Confidence interval	
level	90 %
sides	2-sided
lower limit	-10.47
upper limit	17.97

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	125
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.3734
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	7.48
Confidence interval	
level	90 %
sides	2-sided
lower limit	-6.29
upper limit	21.24

Secondary: Phase II: Percentage of Participants With CTC Reduction Response in Participants With ICR PTEN Loss

End point title	Phase II: Percentage of Participants With CTC Reduction Response in Participants With ICR PTEN Loss ^[30]
-----------------	---

End point description:

CTC reduction response was defined as participants with a reduction in CTCs of $\geq 30\%$ compared to baseline. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss and CTC > 0 cells/7.5 mL at baseline.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, on Day 1 of Cycle 2, on Day 1 of Cycle 3, and at the treatment completion visit (up to overall 3.6 years) (cycle length=28 days)

Notes:

[30] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	20	20	17	
Units: percentage of participants				
number (confidence interval 90%)	75 (55.8 to 87.42)	75 (55.8 to 87.42)	70.6 (50 to 85.95)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	37
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7633
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	4.41
Confidence interval	
level	90 %
sides	2-sided
lower limit	-19.76
upper limit	28.58

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	37
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7633
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	4.41
Confidence interval	
level	90 %
sides	2-sided
lower limit	-19.76
upper limit	28.58

Secondary: Phase II: Percentage of Participants With CTC Conversion - ITT Population

End point title	Phase II: Percentage of Participants With CTC Conversion - ITT Population ^[31]
End point description:	
CTC conversion was defined as a decline to < 5 cells/7.5 milliliter (mL) post baseline among participants with CTC ≥ 5 cells/7.5 mL at baseline. Analysis was performed on ITT population. Number of participants analysed=participants with CTC ≥ 5 cells/7.5 mL at baseline.	
End point type	Secondary
End point timeframe:	
Baseline, on Day 1 of Cycle 2, on Day 1 of Cycle 3, and at the treatment completion visit (up to overall 3.6 years) (cycle length=28 days)	

Notes:

[31] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	41	47	48	
Units: percentage of participants				
number (confidence interval 90%)	43.9 (31.18 to 57.87)	46.8 (34.48 to 59.72)	41.7 (29.59 to 54.55)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	89
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8317
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	2.24
Confidence interval	
level	90 %
sides	2-sided
lower limit	-15.07
upper limit	19.54

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	95
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.6139
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	5.14
Confidence interval	
level	90 %
sides	2-sided
lower limit	-11.6
upper limit	21.88

Secondary: Phase II: Percentage of Participants With CTC Conversion in Participants With ICR PTEN Loss

End point title	Phase II: Percentage of Participants With CTC Conversion in Participants With ICR PTEN Loss ^[32]
-----------------	---

End point description:

CTC conversion was defined as a decline to < 5 cells/7.5 milliliter (mL) post baseline among participants with CTC ≥ 5 cells/7.5 mL at baseline. PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss and CTC ≥ 5 cells/7.5 mL at baseline.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, on Day 1 of Cycle 2, on Day 1 of Cycle 3, and at the treatment completion visit (up to overall 3.6 years) (cycle length=28 days)

Notes:

[32] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	12	18	22	
Units: percentage of participants				
number (confidence interval 90%)	66.7 (39.84 to 84.58)	22.2 (10.6 to 41.88)	31.8 (18.11 to 50)	

Statistical analyses

Statistical analysis title	Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	34
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0505
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	34.85
Confidence interval	
level	90 %
sides	2-sided
lower limit	7.14
upper limit	62.56

Statistical analysis title	Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	40
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.4989
Method	Chi-squared
Parameter estimate	Difference in response rates
Point estimate	-9.6
Confidence interval	
level	90 %
sides	2-sided
lower limit	-32.54
upper limit	13.35

Secondary: Phase II: Percentage of Participants With Pain Progression - ITT Population

End point title	Phase II: Percentage of Participants With Pain Progression - ITT Population ^[33]
-----------------	---

End point description:

Pain progression was defined as ≥ 2 point-increase from baseline on the modified Brief Pain Inventory – short form (mBPI-sf). The mBPI-sf consists of four questions that assess pain intensity (worst, least, average, right now) and seven items within one question that assess the impact of pain on daily functions (general activity, mood, walking ability, normal work, relations with other people, sleep, enjoyment of life). Each item was assessed on 0-10 scale with 0 being no pain to 10 being worst imaginable pain. Total score was the average of individual item (range 0-10). Analysis was performed on ITT population.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening, Day 1 of each cycle (cycle length=28 days) up to treatment completion (up to 3.6 years)

Notes:

[33] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: percentage of participants				
number (not applicable)	33.3	34.9	34.9	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II: Time to Pain Progression - ITT Population

End point title	Phase II: Time to Pain Progression - ITT Population ^[34]
-----------------	---

End point description:

Time to pain progression was defined as the time from screening till first occurrence of pain progression. Pain progression was defined as ≥ 2 point-increase from baseline on the modified Brief Pain Inventory – short form (mBPI-sf). The mBPI-sf consists of four questions that assess pain intensity (worst, least, average, right now) and seven items within one question that assess the impact of pain on daily functions (general activity, mood, walking ability, normal work, relations with other people, sleep, enjoyment of life). Each item was assessed on 0-10 scale with 0 being no pain to 10 being worst imaginable pain. Total score was the average of individual item (range 0-10). Analysis was performed on ITT population. Here, "99999" reflects that the data was not evaluable because of insufficient number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening, Day 1 of each cycle (cycle length=28 days) up to treatment completion (up to 3.6 years)

Notes:

[34] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	84	86	83	
Units: months				
median (confidence interval 90%)	13.9 (8.61 to 99999)	16.16 (8.54 to 99999)	15.15 (11.07 to 99999)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8483
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.95
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.61
upper limit	1.47

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.785
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	1.08
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.7
upper limit	1.65

Statistical analysis title	Stratified analysis:Ipatasertib 400 mg v Placebo
Statistical analysis description:	
Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	167
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8847
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	1.04
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.66
upper limit	1.65

Statistical analysis title	Stratified analysis: Ipatasertib 200 mg v Placebo
Statistical analysis description:	
Strata were: prior enzalutamide (Yes vs. No), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one).	
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	169
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8647
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	1.05
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.67
upper limit	1.63

Secondary: Phase II: Percentage of Participants With Pain Progression in Participants With ICR PTEN Loss

End point title	Phase II: Percentage of Participants With Pain Progression in Participants With ICR PTEN Loss ^[35]
-----------------	---

End point description:

Pain progression was defined as ≥ 2 point-increase from baseline on the modified Brief Pain Inventory – short form (mBPI-sf). The mBPI-sf consists of four questions that assess pain intensity (worst, least, average, right now) and seven items within one question that assess the impact of pain on daily functions (general activity, mood, walking ability, normal work, relations with other people, sleep, enjoyment of life). Each item was assessed on 0-10 scale with 0 being no pain to 10 being worst imaginable pain. Total score was the average of individual item (range 0-10). PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening, Day 1 of each cycle (cycle length=28 days) up to treatment completion (up to 3.6 years)

Notes:

[35] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: percentage of participants				
number (not applicable)	40	28	33.3	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II: Time to Pain Progression in Participants With ICR PTEN Loss

End point title	Phase II: Time to Pain Progression in Participants With ICR PTEN Loss ^[36]
-----------------	---

End point description:

Time to pain progression was defined as the time from screening till first occurrence of pain progression. Pain progression: ≥ 2 point-increase from baseline on the modified Brief Pain Inventory – short form (mBPI-sf). The mBPI-sf consists of 4 questions that assess pain intensity (worst, least, average, right now) and seven items within one question that assess the impact of pain on daily functions (general activity, mood, walking ability, normal work, relations with other people, sleep, enjoyment of life). Each item was assessed on 0-10 scale with 0 being no pain to 10 being worst imaginable pain. Total score was the average of individual item (range 0-10). PTEN status was assessed by RUO IHC assay that was performed at ICR, UK. Samples with 100% of the tumor with no PTEN staining were classified as "ICR PTEN loss". Number of participants analyzed=participants with PTEN loss. Here, "99999" reflects that the data was not evaluable because of insufficient number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Screening, Day 1 of each cycle (cycle length=28 days) up to treatment completion (up to 3.6 years)

Notes:

[36] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase II stage of the study. Hence, arms of Phase Ib stage are not included.

End point values	Phase II: Ipatasertib 400 mg + Abiraterone	Phase II: Ipatasertib 200 mg + Abiraterone	Phase II: Placebo + Abiraterone	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	25	21	
Units: months				
median (confidence interval 90%)	16.49 (7.59 to 16.49)	99999 (8.28 to 99999)	6.93 (5.65 to 99999)	

Statistical analyses

Statistical analysis title	Unstratified analysis:Ipatasertib 400 mg v Placebo
Comparison groups	Phase II: Ipatasertib 400 mg + Abiraterone v Phase II: Placebo + Abiraterone

Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8271
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.89
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.39
upper limit	2.06

Statistical analysis title	Unstratified analysis:Ipatasertib 200 mg v Placebo
Comparison groups	Phase II: Ipatasertib 200 mg + Abiraterone v Phase II: Placebo + Abiraterone
Number of subjects included in analysis	46
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7383
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.84
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.35
upper limit	2.02

Secondary: Phase Ib: Maximum Plasma Concentration (Cmax) of Ipatasertib and GDC-0980 When Co-Administered With Abiraterone

End point title	Phase Ib: Maximum Plasma Concentration (Cmax) of Ipatasertib and GDC-0980 When Co-Administered With Abiraterone ^[37]
-----------------	---

End point description:

Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint. Results reported in "Phase Ib: Ipatasertib 400 mg" represents Cmax of Ipatasertib and results reported in arm "Phase Ib: GDC-0980 30 mg" represents Cmax of GDC-0980.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[37] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	14	6		
Units: ng/mL				
geometric mean (geometric coefficient of variation)				
Cycle 1, Day 1 (n=14,6)	269 (± 41.3)	190 (± 35.2)		
Cycle 1, Day 15 (n=12,2)	466 (± 36.2)	193 (± 16.2)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Time to Cmax (tmax) of Ipatasertib and GDC-0980 When Co-Administered With Abiraterone

End point title	Phase Ib: Time to Cmax (tmax) of Ipatasertib and GDC-0980 When Co-Administered With Abiraterone ^[38]
-----------------	---

End point description:

Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint. Results reported in "Phase Ib: Ipatasertib 400 mg" represents tmax of Ipatasertib and results reported in arm "Phase Ib: GDC-0980 30 mg" represents tmax of GDC-0980.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[38] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	14	6		
Units: hours				
median (full range (min-max))				
Cycle 1, Day 1 (n=14,6)	2 (0.97 to 4.05)	2.02 (1 to 4.1)		
Cycle 1, Day 15 (n=12,2)	2.02 (1 to 6)	2.04 (2 to 2.08)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Area Under The Concentration-Time Curve From Time 0 to 24

Hours (AUC0-24) of Ipatasertib and GDC-0980 When Co-Administered With Abiraterone

End point title	Phase Ib: Area Under The Concentration-Time Curve From Time 0 to 24 Hours (AUC0-24) of Ipatasertib and GDC-0980 When Co-Administered With Abiraterone ^[39]
-----------------	---

End point description:

Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint. Results reported in "Phase Ib: Ipatasertib 400 mg" represents AUC0-24 of Ipatasertib and results reported in arm "Phase Ib: GDC-0980 30 mg" represents AUC0-24 of GDC-0980.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[39] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	14	6		
Units: ng/mL*hours				
geometric mean (geometric coefficient of variation)				
Cycle 1, Day 1 (n=14,6)	1710 (\pm 54.7)	1600 (\pm 47.5)		
Cycle 1, Day 15 (n=12,2)	3290 (\pm 37)	1640 (\pm 5.63)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Total Body Clearance (CL/F) of Ipatasertib When Co-Administered With Abiraterone

End point title	Phase Ib: Total Body Clearance (CL/F) of Ipatasertib When Co-Administered With Abiraterone ^[40]
-----------------	--

End point description:

Clearance of a drug is a measure of the rate at which a drug is metabolized or eliminated by normal biological processes. Clearance obtained after oral dose is influenced by the fraction of the dose absorbed. Drug clearance is a quantitative measure of the rate at which a drug substance is removed from the blood. Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Numer of participants analyzed=participants with PK data available on Day 15 of Cycle 1.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 15 of Cycles 1

Notes:

[40] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg			
Subject group type	Reporting group			
Number of subjects analysed	12			
Units: milliliter per hour (mL/h)				
geometric mean (geometric coefficient of variation)	99600 (± 50.7)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Accumulation Ratio of Ipatasertib When Co-Administered With Abiraterone

End point title	Phase Ib: Accumulation Ratio of Ipatasertib When Co-Administered With Abiraterone ^[41]
-----------------	---

End point description:

Accumulation Ratio was calculated as AUC₀₋₂₄ (Cycle 1 Day 15) divided by AUC₀₋₂₄ (Cycle 1 Day 1). Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data Number of participants analyzed=participants with PK data available on both Day 1 and Day 15 of Cycle 1.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[41] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg			
Subject group type	Reporting group			
Number of subjects analysed	12			
Units: ratio				
geometric mean (geometric coefficient of variation)	1.82 (± 40.9)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Cmax of G-037720 (Metabolite of Ipatasertib)

End point title	Phase Ib: Cmax of G-037720 (Metabolite of Ipatasertib) ^[42]
-----------------	--

End point description:

G-037720 is N-dealkylated metabolite of ipatasertib and the main metabolite in circulation. Analysis was performed on Pharmacokinetic Analysis Population (only ipatasertib arm) which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint.

End point type	Secondary			
End point timeframe:				
Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1				
Notes:				
[42] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.				
Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.				
End point values	Phase Ib: Ipatasertib 400 mg			
	Subject group type	Reporting group		
	Number of subjects analysed	14		
	Units: ng/mL			
	geometric mean (geometric coefficient of variation)			
	Cycle 1, Day 1 (n=14) Cycle 1, Day 15 (n=12)	117 (± 64.4) 326 (± 42.4)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: tmax of G-037720 (Metabolite of Ipatasertib)

End point title	Phase Ib: tmax of G-037720 (Metabolite of Ipatasertib) ^[43]			
End point description:				
G-037220 is N-dealkylated metabolite of ipatasertib and the main metabolite in circulation. Analysis was performed on Pharmacokinetic Analysis Population (only ipatasertib arm) which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint.				
End point type	Secondary			
End point timeframe:				
Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1				
Notes:				
[43] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.				
Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.				

End point values	Phase Ib: Ipatasertib 400 mg			
Subject group type	Reporting group			
Number of subjects analysed	14			
Units: hours				
median (full range (min-max))				
Cycle 1, Day 1 (n=14)	2 (1 to 4.05)			
Cycle 1, Day 15 (n=12)	2.1 (1 to 6)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: AUC0-24 of G-037720 (Metabolite of Ipatasertib)

End point title	Phase Ib: AUC0-24 of G-037720 (Metabolite of Ipatasertib) ^[44]
-----------------	---

End point description:

G-037220 is N-dealkylated metabolite of ipatasertib and the main metabolite in circulation. Analysis was performed on Pharmacokinetic Analysis Population (only ipatasertib arm) which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[44] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg			
Subject group type	Reporting group			
Number of subjects analysed	14			
Units: ng/mL*hour				
geometric mean (geometric coefficient of variation)				
Cycle 1, Day 1 (n=14)	839 (± 70.3)			
Cycle 1, Day 15 (n=12)	2850 (± 71.7)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Accumulation Ratio of G-037720 (Metabolite of Ipatasertib)

End point title	Phase Ib: Accumulation Ratio of G-037720 (Metabolite of Ipatasertib) ^[45]
-----------------	--

End point description:

G-037220 is N-dealkylated metabolite of ipatasertib and the main metabolite in circulation. Accumulation Ratio was calculated as AUC0-24 (Cycle 1 Day 15) divided by AUC0-24 (Cycle 1 Day 1). Analysis was performed on Pharmacokinetic Analysis Population (only ipatasertib arm) which included all participants who had evaluable PK data. Number of participants analyzed=participants with PK data available on both Day 1 and Day 15 of Cycle 1.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[45] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg			
Subject group type	Reporting group			
Number of subjects analysed	12			
Units: ratio				
geometric mean (geometric coefficient of variation)	2.79 (\pm 54.4)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Cmax of Abiraterone When Co- Administered With Ipatasertib or GDC-0980

End point title	Phase Ib: Cmax of Abiraterone When Co- Administered With Ipatasertib or GDC-0980 ^[46]
-----------------	--

End point description:

Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[46] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	14	6		
Units: ng/mL				
geometric mean (geometric coefficient of variation)				
Cycle 1, Day 1 (n=14,6)	151 (\pm 123.6)	88.2 (\pm 218)		
Cycle 1, Day 15 (n=12,2)	140 (\pm 124.4)	52.7 (\pm 182)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: tmax of Abiraterone When Co- Administered With Ipatasertib or GDC-0980

End point title	Phase Ib: tmax of Abiraterone When Co- Administered With Ipatasertib or GDC-0980 ^[47]
-----------------	--

End point description:

Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint. Cycle 1, Day 1 (n=14,6)

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[47] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	14	6		
Units: hours				
median (full range (min-max))				
Cycle 1, Day 1 (n=14,6)	2.05 (2 to 6.45)	2.01 (1 to 4.03)		
Cycle 1, Day 15 (n=12,2)	2.17 (2 to 6)	3.04 (2 to 4.08)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: AUC0-24 of Abiraterone When Co- Administered With Ipatasertib or GDC-0980

End point title	Phase Ib: AUC0-24 of Abiraterone When Co- Administered With Ipatasertib or GDC-0980 ^[48]
-----------------	---

End point description:

Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint. Here, "99999" reflects that the data is not applicable because only 1 participant was evaluated at Day 15 Cycle 1 in GDC-0980 30 mg arm.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[48] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage

are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC- 0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	12	6		
Units: ng/mL*hr				
geometric mean (geometric coefficient of variation)				
Cycle 1, Day 1 (n=12,6)	749 (± 92)	475 (± 203.4)		
Cycle 1, Day 15 (n=6,1)	961 (± 87.2)	220 (± 99999)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Plasma Half-Life of Abiraterone When Co- Administered With Ipatasertib or GDC-0980

End point title	Phase Ib: Plasma Half-Life of Abiraterone When Co- Administered With Ipatasertib or GDC-0980 ^[49]
-----------------	---

End point description:

Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Here, "n" signifies the number of participants with PK data available for specified timepoint. Here, "99999" reflects that the data is not applicable because only 1 participant was evaluated at Day 15 Cycle 1 in GDC-0980 30 mg arm.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[49] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC- 0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	10	5		
Units: hours				
geometric mean (geometric coefficient of variation)				
Cycle 1, Day 1 (n=10,5)	5.25 (± 13)	7.68 (± 26.2)		
Cycle 1, Day 15 (n=6,1)	6.92 (± 23.5)	14.7 (± 99999)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Accumulation Ratio of Abiraterone When Co- Administered With Ipatasertib or GDC-0980

End point title	Phase Ib: Accumulation Ratio of Abiraterone When Co-Administered With Ipatasertib or GDC-0980 ^[50]
-----------------	---

End point description:

Accumulation Ratio was calculated as AUC₀₋₂₄ (Cycle 1 Day 15) divided by AUC₀₋₂₄ (Cycle 1 Day 1). Analysis was performed on Pharmacokinetic Analysis Population which included all participants who had evaluable PK data. Number of participants analyzed=participants with PK data available on both Day 1 and Day 15 of Cycle 1. Here, "99999" reflects that the data is not applicable because only 1 participant was evaluated in GDC--0980 30 mg arm.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hour), 1, 2, 4, 6, 24 hours post dose on Days 1, 15 of Cycles 1

Notes:

[50] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint was measured in Phase Ib stage of the study. Hence, arms of Phase II stage are not included.

End point values	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	12	1		
Units: ratio				
geometric mean (geometric coefficient of variation)	0.823 (± 70.7)	0.882 (± 99999)		

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

From first dose to 28 days after last dose (up to 3.6 years)

Adverse event reporting additional description:

Safety population

Assessment type	Non-systematic
-----------------	----------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	18.1
--------------------	------

Reporting groups

Reporting group title	Phase Ib: Ipatasertib 400 mg
-----------------------	------------------------------

Reporting group description:

Participants received ipatasertib 400 mg orally daily, abiraterone 1000 mg orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase Ib: GDC-0980 30 mg
-----------------------	--------------------------

Reporting group description:

Participants received GDC-0980 30 mg orally daily, abiraterone 1000 mg orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase II: Ipatasertib 400 mg + abiraterone
-----------------------	--

Reporting group description:

Participants received Ipatasertib 400 mg orally daily, abiraterone 1000 mg orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase II: Ipatasertib 200 mg + abiraterone
-----------------------	--

Reporting group description:

Participants received Ipatasertib 200 mg orally daily, abiraterone 1000 mg orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Reporting group title	Phase II: Placebo + abiraterone
-----------------------	---------------------------------

Reporting group description:

Participants received placebo (for Ipatasertib 400 mg or 200) orally daily, abiraterone 1000 mg orally daily, and prednisone/prednisolone 5 mg bid orally continuously in 28-day treatment cycles until disease progression or intolerable toxicity.

Serious adverse events	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg	Phase II: Ipatasertib 400 mg + abiraterone
Total subjects affected by serious adverse events			
subjects affected / exposed	9 / 14 (64.29%)	5 / 6 (83.33%)	36 / 84 (42.86%)
number of deaths (all causes)	2	1	4
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Prostate cancer			

subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	2 / 84 (2.38%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 1	0 / 1	0 / 2
Prostatic adenoma			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Small cell carcinoma			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Aortic aneurysm rupture			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Deep vein thrombosis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haematoma			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypotension			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Thrombosis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			

Asthenia	subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
	occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Death	subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
	occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Disease progression	subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
	occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Fatigue	subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
	occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General physical health deterioration	subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
	occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pain	subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	2 / 84 (2.38%)
	occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 3
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyrexia	subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
	occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders				
Pelvic pain	subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
	occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
	deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Prostatitis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Chronic obstructive pulmonary disease			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dyspnoea			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Interstitial lung disease			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonitis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Confusional state			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Aspartate aminotransferase increased			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haematocrit decreased			

subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haemoglobin decreased			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Radius fracture			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skull fracture			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Subdural haematoma			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 1	0 / 0	0 / 0
Ulna fracture			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Upper limb fracture			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders			
Atrial fibrillation			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Atrioventricular block			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bradycardia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Cardiac failure			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac failure acute			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac failure congestive			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Cerebral haemorrhage			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diplegia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dizziness			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ischaemic stroke			

subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorder			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Spinal cord compression			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Syncope			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Toxic encephalopathy			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular encephalopathy			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Coagulopathy			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haemolytic uraemic syndrome			

subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Abdominal pain upper			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ascites			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Colitis			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Constipation			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diarrhoea			
subjects affected / exposed	2 / 14 (14.29%)	0 / 6 (0.00%)	2 / 84 (2.38%)
occurrences causally related to treatment / all	2 / 2	0 / 0	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dyspepsia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dysphagia			

subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Enteritis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Enterocolitis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal haemorrhage			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haematochezia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Lower gastrointestinal haemorrhage			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nausea			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pharyngo-oesophageal diverticulum			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			

subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
Rash			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	2 / 84 (2.38%)
occurrences causally related to treatment / all	0 / 0	0 / 0	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash maculo-papular			
subjects affected / exposed	1 / 14 (7.14%)	2 / 6 (33.33%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Toxic skin eruption			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Acute kidney injury			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Calculus urinary			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haematuria			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	2 / 84 (2.38%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nephrolithiasis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nephropathy toxic			

subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal failure			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ureteric obstruction			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary retention			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bone pain			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Muscle haemorrhage			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Osteolysis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Spinal pain			

subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Cellulitis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Device related infection			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Osteomyelitis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			
subjects affected / exposed	2 / 14 (14.29%)	1 / 6 (16.67%)	3 / 84 (3.57%)
occurrences causally related to treatment / all	1 / 2	0 / 1	1 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyelonephritis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyelonephritis acute			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory tract infection			

subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Septic shock			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Spinal cord infection			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary tract infection			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	4 / 84 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urosepsis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Dehydration			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diabetes mellitus inadequate control			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Failure to thrive			

subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hyperglycaemia			
subjects affected / exposed	1 / 14 (7.14%)	2 / 6 (33.33%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	1 / 1	0 / 2	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypoglycaemia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypokalaemia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ketoacidosis			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Phase II: Ipatasertib 200 mg + abiraterone	Phase II: Placebo + abiraterone	
Total subjects affected by serious adverse events			
subjects affected / exposed	35 / 87 (40.23%)	16 / 82 (19.51%)	
number of deaths (all causes)	6	4	
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Prostate cancer			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
Prostatic adenoma			

subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Small cell carcinoma			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vascular disorders			
Aortic aneurysm rupture			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
Deep vein thrombosis			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haematoma			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypotension			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Thrombosis			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
General disorders and administration site conditions			
Asthenia			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Death			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 1	0 / 1	
Disease progression			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 1	0 / 1	
Fatigue			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
General physical health deterioration			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pain			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyrexia			
subjects affected / exposed	2 / 87 (2.30%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Reproductive system and breast disorders			
Pelvic pain			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Prostatitis			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Respiratory, thoracic and mediastinal disorders			
Chronic obstructive pulmonary disease			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Dyspnoea			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Interstitial lung disease			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonitis			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Psychiatric disorders			
Confusional state			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Investigations			
Aspartate aminotransferase increased			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haematocrit decreased			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haemoglobin decreased			

subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injury, poisoning and procedural complications			
Radius fracture			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skull fracture			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Subdural haematoma			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ulna fracture			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Upper limb fracture			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac disorders			
Atrial fibrillation			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Atrioventricular block			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Bradycardia			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac failure			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac failure acute			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Cardiac failure congestive			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
Cerebral haemorrhage			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Diplegia			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Dizziness			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ischaemic stroke			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorder			

subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Spinal cord compression			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Syncope			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Toxic encephalopathy			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vascular encephalopathy			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	2 / 87 (2.30%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 2	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Coagulopathy			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haemolytic uraemic syndrome			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			

Abdominal pain			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Abdominal pain upper			
subjects affected / exposed	0 / 87 (0.00%)	2 / 82 (2.44%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ascites			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Colitis			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Constipation			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diarrhoea			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Dyspepsia			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Dysphagia			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Enteritis			

subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Enterocolitis			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal haemorrhage			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haematochezia			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lower gastrointestinal haemorrhage			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nausea			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pharyngo-oesophageal diverticulum			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vomiting			
subjects affected / exposed	0 / 87 (0.00%)	2 / 82 (2.44%)	
occurrences causally related to treatment / all	0 / 0	0 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin and subcutaneous tissue disorders			
Rash			

subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Rash maculo-papular			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Toxic skin eruption			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal and urinary disorders			
Acute kidney injury			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Calculus urinary			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haematuria			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nephrolithiasis			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nephropathy toxic			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal failure			

subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	1 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ureteric obstruction			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary retention			
subjects affected / exposed	2 / 87 (2.30%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bone pain			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Muscle haemorrhage			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Osteolysis			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Spinal pain			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			

Cellulitis			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Device related infection			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Osteomyelitis			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia			
subjects affected / exposed	4 / 87 (4.60%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 5	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyelonephritis			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyelonephritis acute			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Respiratory tract infection			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Sepsis			

subjects affected / exposed	3 / 87 (3.45%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 3	0 / 0	
deaths causally related to treatment / all	0 / 2	0 / 0	
Septic shock			
subjects affected / exposed	2 / 87 (2.30%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Spinal cord infection			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract infection			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 2	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urosepsis			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Metabolism and nutrition disorders			
Dehydration			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diabetes mellitus inadequate control			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Failure to thrive			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hyperglycaemia			

subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypoglycaemia			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypokalaemia			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ketoacidosis			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 5 %

Non-serious adverse events	Phase Ib: Ipatasertib 400 mg	Phase Ib: GDC-0980 30 mg	Phase II: Ipatasertib 400 mg + abiraterone
Total subjects affected by non-serious adverse events			
subjects affected / exposed	14 / 14 (100.00%)	6 / 6 (100.00%)	82 / 84 (97.62%)
Vascular disorders			
Hot flush			
subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	9 / 84 (10.71%)
occurrences (all)	1	1	9
Hypertension			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	9 / 84 (10.71%)
occurrences (all)	1	0	13
Hypotension			
subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	3 / 84 (3.57%)
occurrences (all)	1	1	3
General disorders and administration site conditions			

Asthenia			
subjects affected / exposed	4 / 14 (28.57%)	1 / 6 (16.67%)	22 / 84 (26.19%)
occurrences (all)	4	1	32
Fatigue			
subjects affected / exposed	9 / 14 (64.29%)	1 / 6 (16.67%)	21 / 84 (25.00%)
occurrences (all)	11	1	33
Gait disturbance			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences (all)	1	0	1
Mucosal inflammation			
subjects affected / exposed	3 / 14 (21.43%)	0 / 6 (0.00%)	3 / 84 (3.57%)
occurrences (all)	3	0	4
Oedema			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences (all)	1	0	1
Oedema peripheral			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	8 / 84 (9.52%)
occurrences (all)	2	0	11
Pain			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	5 / 84 (5.95%)
occurrences (all)	0	1	5
Pyrexia			
subjects affected / exposed	1 / 14 (7.14%)	3 / 6 (50.00%)	9 / 84 (10.71%)
occurrences (all)	1	3	22
Thirst			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences (all)	0	1	0
Reproductive system and breast disorders			
Pelvic pain			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	7 / 84 (8.33%)
occurrences (all)	0	0	9
Penile pain			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0
Respiratory, thoracic and mediastinal disorders			

Cough			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	5 / 84 (5.95%)
occurrences (all)	1	0	9
Dry throat			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences (all)	0	1	0
Dysphonia			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	1 / 84 (1.19%)
occurrences (all)	0	1	1
Dyspnoea			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	8 / 84 (9.52%)
occurrences (all)	0	1	10
Dyspnoea exertional			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences (all)	1	0	1
Oropharyngeal pain			
subjects affected / exposed	2 / 14 (14.29%)	1 / 6 (16.67%)	3 / 84 (3.57%)
occurrences (all)	2	1	3
Pleural effusion			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0
Rhinorrhoea			
subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	1 / 84 (1.19%)
occurrences (all)	1	1	1
Sinus congestion			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	1 / 84 (1.19%)
occurrences (all)	0	1	1
Wheezing			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences (all)	0	1	0
Psychiatric disorders			
Agitation			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0
Anxiety			

subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	2 / 84 (2.38%)
occurrences (all)	1	0	2
Depression			
subjects affected / exposed	2 / 14 (14.29%)	0 / 6 (0.00%)	3 / 84 (3.57%)
occurrences (all)	2	0	3
Insomnia			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	8 / 84 (9.52%)
occurrences (all)	0	0	9
Restlessness			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0
Investigations			
Aspartate aminotransferase increased			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	5 / 84 (5.95%)
occurrences (all)	1	0	8
Blood alkaline phosphatase increased			
subjects affected / exposed	2 / 14 (14.29%)	0 / 6 (0.00%)	2 / 84 (2.38%)
occurrences (all)	2	0	2
Blood creatinine increased			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	7 / 84 (8.33%)
occurrences (all)	0	0	7
Blood urine present			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0
Injury, poisoning and procedural complications			
Meniscus injury			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0
Nervous system disorders			
Dizziness			
subjects affected / exposed	3 / 14 (21.43%)	0 / 6 (0.00%)	10 / 84 (11.90%)
occurrences (all)	4	0	14
Dysgeusia			
subjects affected / exposed	2 / 14 (14.29%)	0 / 6 (0.00%)	6 / 84 (7.14%)
occurrences (all)	2	0	8
Headache			

subjects affected / exposed occurrences (all)	2 / 14 (14.29%) 3	0 / 6 (0.00%) 0	8 / 84 (9.52%) 13
Hypoaesthesia subjects affected / exposed occurrences (all)	2 / 14 (14.29%) 2	0 / 6 (0.00%) 0	0 / 84 (0.00%) 0
Lethargy subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	1 / 6 (16.67%) 1	1 / 84 (1.19%) 1
Somnolence subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	3 / 84 (3.57%) 3
Blood and lymphatic system disorders Anaemia subjects affected / exposed occurrences (all)	2 / 14 (14.29%) 2	1 / 6 (16.67%) 1	13 / 84 (15.48%) 17
Pancytopenia subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	1 / 6 (16.67%) 1	0 / 84 (0.00%) 0
Eye disorders Vision blurred subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	0 / 84 (0.00%) 0
Gastrointestinal disorders Abdominal pain subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	7 / 84 (8.33%) 8
Abdominal pain lower subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	1 / 6 (16.67%) 1	2 / 84 (2.38%) 3
Abdominal pain upper subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	0 / 6 (0.00%) 0	8 / 84 (9.52%) 10
Cheilosis subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	0 / 84 (0.00%) 0
Constipation			

subjects affected / exposed	1 / 14 (7.14%)	2 / 6 (33.33%)	8 / 84 (9.52%)
occurrences (all)	1	2	8
Diarrhoea			
subjects affected / exposed	9 / 14 (64.29%)	4 / 6 (66.67%)	64 / 84 (76.19%)
occurrences (all)	14	6	181
Dry mouth			
subjects affected / exposed	2 / 14 (14.29%)	1 / 6 (16.67%)	3 / 84 (3.57%)
occurrences (all)	2	1	3
Dyspepsia			
subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	8 / 84 (9.52%)
occurrences (all)	1	1	8
Eructation			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences (all)	1	0	1
Nausea			
subjects affected / exposed	6 / 14 (42.86%)	4 / 6 (66.67%)	44 / 84 (52.38%)
occurrences (all)	11	4	68
Oral disorder			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences (all)	0	1	0
Rectal haemorrhage			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0
Retching			
subjects affected / exposed	2 / 14 (14.29%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences (all)	2	1	0
Tongue discolouration			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences (all)	0	1	0
Vomiting			
subjects affected / exposed	9 / 14 (64.29%)	3 / 6 (50.00%)	26 / 84 (30.95%)
occurrences (all)	14	3	40
Skin and subcutaneous tissue disorders			
Ecchymosis			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0

Haemorrhage subcutaneous subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	0 / 84 (0.00%) 0
Hyperhidrosis subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	1 / 6 (16.67%) 1	1 / 84 (1.19%) 2
Nail disorder subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	1 / 6 (16.67%) 1	0 / 84 (0.00%) 0
Night sweats subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	1 / 84 (1.19%) 2
Rash subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	3 / 6 (50.00%) 5	10 / 84 (11.90%) 12
Rash maculo-papular subjects affected / exposed occurrences (all)	2 / 14 (14.29%) 3	2 / 6 (33.33%) 3	4 / 84 (4.76%) 7
Renal and urinary disorders			
Acute kidney injury subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	1 / 6 (16.67%) 1	2 / 84 (2.38%) 4
Dysuria subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	7 / 84 (8.33%) 8
Haematuria subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	0 / 6 (0.00%) 0	8 / 84 (9.52%) 12
Hypertonic bladder subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	0 / 84 (0.00%) 0
Incontinence subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	1 / 84 (1.19%) 1
Pollakiuria			

subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	2 / 84 (2.38%)
occurrences (all)	1	1	2
Renal pain			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	1 / 84 (1.19%)
occurrences (all)	0	1	1
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	2 / 14 (14.29%)	0 / 6 (0.00%)	14 / 84 (16.67%)
occurrences (all)	3	0	21
Back pain			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	15 / 84 (17.86%)
occurrences (all)	1	0	21
Bone pain			
subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	10 / 84 (11.90%)
occurrences (all)	1	1	16
Flank pain			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	2 / 84 (2.38%)
occurrences (all)	2	0	2
Muscle spasms			
subjects affected / exposed	0 / 14 (0.00%)	0 / 6 (0.00%)	3 / 84 (3.57%)
occurrences (all)	0	0	5
Muscular weakness			
subjects affected / exposed	1 / 14 (7.14%)	1 / 6 (16.67%)	1 / 84 (1.19%)
occurrences (all)	1	1	1
Musculoskeletal chest pain			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	5 / 84 (5.95%)
occurrences (all)	1	0	13
Musculoskeletal pain			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	4 / 84 (4.76%)
occurrences (all)	1	0	5
Myalgia			
subjects affected / exposed	4 / 14 (28.57%)	1 / 6 (16.67%)	5 / 84 (5.95%)
occurrences (all)	5	1	7
Neck pain			

subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 2	0 / 6 (0.00%) 0	1 / 84 (1.19%) 1
Pain in extremity subjects affected / exposed occurrences (all)	2 / 14 (14.29%) 3	0 / 6 (0.00%) 0	11 / 84 (13.10%) 13
Pain in jaw subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 2	0 / 6 (0.00%) 0	2 / 84 (2.38%) 3
Infections and infestations Nasopharyngitis subjects affected / exposed occurrences (all)	2 / 14 (14.29%) 2	0 / 6 (0.00%) 0	2 / 84 (2.38%) 2
Urinary tract infection subjects affected / exposed occurrences (all)	2 / 14 (14.29%) 3	0 / 6 (0.00%) 0	12 / 84 (14.29%) 13
Urinary tract infection fungal subjects affected / exposed occurrences (all)	1 / 14 (7.14%) 1	0 / 6 (0.00%) 0	0 / 84 (0.00%) 0
Metabolism and nutrition disorders Decreased appetite subjects affected / exposed occurrences (all)	5 / 14 (35.71%) 5	1 / 6 (16.67%) 1	21 / 84 (25.00%) 28
Dehydration subjects affected / exposed occurrences (all)	3 / 14 (21.43%) 3	2 / 6 (33.33%) 2	3 / 84 (3.57%) 3
Failure to thrive subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	1 / 6 (16.67%) 1	0 / 84 (0.00%) 0
Hyperglycaemia subjects affected / exposed occurrences (all)	6 / 14 (42.86%) 7	2 / 6 (33.33%) 4	18 / 84 (21.43%) 26
Hyperkalaemia subjects affected / exposed occurrences (all)	0 / 14 (0.00%) 0	1 / 6 (16.67%) 1	1 / 84 (1.19%) 1
Hypertriglyceridaemia			

subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	1 / 84 (1.19%)
occurrences (all)	1	0	1
Hypocalcaemia			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	5 / 84 (5.95%)
occurrences (all)	1	0	6
Hypokalaemia			
subjects affected / exposed	0 / 14 (0.00%)	2 / 6 (33.33%)	8 / 84 (9.52%)
occurrences (all)	0	2	11
Hypovolaemia			
subjects affected / exposed	0 / 14 (0.00%)	1 / 6 (16.67%)	0 / 84 (0.00%)
occurrences (all)	0	1	0
Increased appetite			
subjects affected / exposed	1 / 14 (7.14%)	0 / 6 (0.00%)	0 / 84 (0.00%)
occurrences (all)	1	0	0

Non-serious adverse events	Phase II: Ipatasertib 200 mg + abiraterone	Phase II: Placebo + abiraterone	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	76 / 87 (87.36%)	76 / 82 (92.68%)	
Vascular disorders			
Hot flush			
subjects affected / exposed	3 / 87 (3.45%)	6 / 82 (7.32%)	
occurrences (all)	3	6	
Hypertension			
subjects affected / exposed	11 / 87 (12.64%)	8 / 82 (9.76%)	
occurrences (all)	12	10	
Hypotension			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences (all)	1	1	
General disorders and administration site conditions			
Asthenia			
subjects affected / exposed	17 / 87 (19.54%)	13 / 82 (15.85%)	
occurrences (all)	21	15	
Fatigue			
subjects affected / exposed	22 / 87 (25.29%)	24 / 82 (29.27%)	
occurrences (all)	30	32	
Gait disturbance			

subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences (all)	1	1	
Mucosal inflammation			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences (all)	0	1	
Oedema			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences (all)	1	0	
Oedema peripheral			
subjects affected / exposed	7 / 87 (8.05%)	7 / 82 (8.54%)	
occurrences (all)	8	7	
Pain			
subjects affected / exposed	8 / 87 (9.20%)	2 / 82 (2.44%)	
occurrences (all)	12	2	
Pyrexia			
subjects affected / exposed	11 / 87 (12.64%)	9 / 82 (10.98%)	
occurrences (all)	13	13	
Thirst			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Reproductive system and breast disorders			
Pelvic pain			
subjects affected / exposed	4 / 87 (4.60%)	0 / 82 (0.00%)	
occurrences (all)	6	0	
Penile pain			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Respiratory, thoracic and mediastinal disorders			
Cough			
subjects affected / exposed	3 / 87 (3.45%)	7 / 82 (8.54%)	
occurrences (all)	3	8	
Dry throat			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Dysphonia			

subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences (all)	1	0	
Dyspnoea			
subjects affected / exposed	8 / 87 (9.20%)	6 / 82 (7.32%)	
occurrences (all)	9	7	
Dyspnoea exertional			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Oropharyngeal pain			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Pleural effusion			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences (all)	1	0	
Rhinorrhoea			
subjects affected / exposed	3 / 87 (3.45%)	2 / 82 (2.44%)	
occurrences (all)	3	4	
Sinus congestion			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Wheezing			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Psychiatric disorders			
Agitation			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Anxiety			
subjects affected / exposed	5 / 87 (5.75%)	3 / 82 (3.66%)	
occurrences (all)	5	3	
Depression			
subjects affected / exposed	3 / 87 (3.45%)	2 / 82 (2.44%)	
occurrences (all)	3	2	
Insomnia			
subjects affected / exposed	10 / 87 (11.49%)	5 / 82 (6.10%)	
occurrences (all)	10	5	

Restlessness subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Investigations			
Aspartate aminotransferase increased subjects affected / exposed occurrences (all)	3 / 87 (3.45%) 3	1 / 82 (1.22%) 1	
Blood alkaline phosphatase increased subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	1 / 82 (1.22%) 1	
Blood creatinine increased subjects affected / exposed occurrences (all)	4 / 87 (4.60%) 5	1 / 82 (1.22%) 2	
Blood urine present subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Injury, poisoning and procedural complications			
Meniscus injury subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Nervous system disorders			
Dizziness subjects affected / exposed occurrences (all)	5 / 87 (5.75%) 7	5 / 82 (6.10%) 5	
Dysgeusia subjects affected / exposed occurrences (all)	1 / 87 (1.15%) 1	3 / 82 (3.66%) 3	
Headache subjects affected / exposed occurrences (all)	5 / 87 (5.75%) 5	6 / 82 (7.32%) 9	
Hypoaesthesia subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	2 / 82 (2.44%) 4	
Lethargy subjects affected / exposed occurrences (all)	1 / 87 (1.15%) 1	1 / 82 (1.22%) 1	

Somnolence subjects affected / exposed occurrences (all)	2 / 87 (2.30%) 2	0 / 82 (0.00%) 0	
Blood and lymphatic system disorders Anaemia subjects affected / exposed occurrences (all)	16 / 87 (18.39%) 29	8 / 82 (9.76%) 13	
Pancytopenia subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Eye disorders Vision blurred subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	3 / 82 (3.66%) 4	
Gastrointestinal disorders Abdominal pain subjects affected / exposed occurrences (all)	8 / 87 (9.20%) 9	3 / 82 (3.66%) 3	
Abdominal pain lower subjects affected / exposed occurrences (all)	1 / 87 (1.15%) 1	0 / 82 (0.00%) 0	
Abdominal pain upper subjects affected / exposed occurrences (all)	3 / 87 (3.45%) 4	5 / 82 (6.10%) 7	
Cheilosis subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Constipation subjects affected / exposed occurrences (all)	15 / 87 (17.24%) 20	17 / 82 (20.73%) 23	
Diarrhoea subjects affected / exposed occurrences (all)	39 / 87 (44.83%) 61	21 / 82 (25.61%) 37	
Dry mouth subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	4 / 82 (4.88%) 4	
Dyspepsia			

subjects affected / exposed	7 / 87 (8.05%)	3 / 82 (3.66%)	
occurrences (all)	8	3	
Eructation			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences (all)	1	0	
Nausea			
subjects affected / exposed	30 / 87 (34.48%)	20 / 82 (24.39%)	
occurrences (all)	43	29	
Oral disorder			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Rectal haemorrhage			
subjects affected / exposed	2 / 87 (2.30%)	0 / 82 (0.00%)	
occurrences (all)	3	0	
Retching			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Tongue discolouration			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Vomiting			
subjects affected / exposed	24 / 87 (27.59%)	12 / 82 (14.63%)	
occurrences (all)	69	20	
Skin and subcutaneous tissue disorders			
Ecchymosis			
subjects affected / exposed	0 / 87 (0.00%)	1 / 82 (1.22%)	
occurrences (all)	0	1	
Haemorrhage subcutaneous			
subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	
Hyperhidrosis			
subjects affected / exposed	0 / 87 (0.00%)	3 / 82 (3.66%)	
occurrences (all)	0	4	
Nail disorder			
subjects affected / exposed	1 / 87 (1.15%)	0 / 82 (0.00%)	
occurrences (all)	1	0	

Night sweats subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	1 / 82 (1.22%) 1	
Rash subjects affected / exposed occurrences (all)	4 / 87 (4.60%) 5	4 / 82 (4.88%) 4	
Rash maculo-papular subjects affected / exposed occurrences (all)	1 / 87 (1.15%) 2	1 / 82 (1.22%) 1	
Renal and urinary disorders Acute kidney injury subjects affected / exposed occurrences (all)	2 / 87 (2.30%) 2	0 / 82 (0.00%) 0	
Dysuria subjects affected / exposed occurrences (all)	2 / 87 (2.30%) 4	1 / 82 (1.22%) 1	
Haematuria subjects affected / exposed occurrences (all)	6 / 87 (6.90%) 10	5 / 82 (6.10%) 6	
Hypertonic bladder subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Incontinence subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Pollakiuria subjects affected / exposed occurrences (all)	6 / 87 (6.90%) 7	0 / 82 (0.00%) 0	
Renal pain subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Musculoskeletal and connective tissue disorders Arthralgia subjects affected / exposed occurrences (all)	12 / 87 (13.79%) 18	14 / 82 (17.07%) 16	
Back pain			

subjects affected / exposed	18 / 87 (20.69%)	21 / 82 (25.61%)	
occurrences (all)	31	30	
Bone pain			
subjects affected / exposed	11 / 87 (12.64%)	16 / 82 (19.51%)	
occurrences (all)	14	18	
Flank pain			
subjects affected / exposed	3 / 87 (3.45%)	0 / 82 (0.00%)	
occurrences (all)	3	0	
Muscle spasms			
subjects affected / exposed	6 / 87 (6.90%)	2 / 82 (2.44%)	
occurrences (all)	6	2	
Muscular weakness			
subjects affected / exposed	6 / 87 (6.90%)	3 / 82 (3.66%)	
occurrences (all)	7	3	
Musculoskeletal chest pain			
subjects affected / exposed	5 / 87 (5.75%)	6 / 82 (7.32%)	
occurrences (all)	8	9	
Musculoskeletal pain			
subjects affected / exposed	6 / 87 (6.90%)	11 / 82 (13.41%)	
occurrences (all)	7	13	
Myalgia			
subjects affected / exposed	3 / 87 (3.45%)	5 / 82 (6.10%)	
occurrences (all)	3	5	
Neck pain			
subjects affected / exposed	2 / 87 (2.30%)	3 / 82 (3.66%)	
occurrences (all)	2	3	
Pain in extremity			
subjects affected / exposed	10 / 87 (11.49%)	8 / 82 (9.76%)	
occurrences (all)	12	14	
Pain in jaw			
subjects affected / exposed	1 / 87 (1.15%)	1 / 82 (1.22%)	
occurrences (all)	1	1	
Infections and infestations			
Nasopharyngitis			
subjects affected / exposed	4 / 87 (4.60%)	4 / 82 (4.88%)	
occurrences (all)	5	5	

Urinary tract infection subjects affected / exposed occurrences (all)	6 / 87 (6.90%) 9	5 / 82 (6.10%) 7	
Urinary tract infection fungal subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Metabolism and nutrition disorders			
Decreased appetite subjects affected / exposed occurrences (all)	19 / 87 (21.84%) 23	12 / 82 (14.63%) 13	
Dehydration subjects affected / exposed occurrences (all)	2 / 87 (2.30%) 2	2 / 82 (2.44%) 2	
Failure to thrive subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Hyperglycaemia subjects affected / exposed occurrences (all)	7 / 87 (8.05%) 10	5 / 82 (6.10%) 7	
Hyperkalaemia subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	1 / 82 (1.22%) 1	
Hypertriglyceridaemia subjects affected / exposed occurrences (all)	2 / 87 (2.30%) 2	0 / 82 (0.00%) 0	
Hypocalcaemia subjects affected / exposed occurrences (all)	4 / 87 (4.60%) 5	4 / 82 (4.88%) 5	
Hypokalaemia subjects affected / exposed occurrences (all)	6 / 87 (6.90%) 10	7 / 82 (8.54%) 9	
Hypovolaemia subjects affected / exposed occurrences (all)	0 / 87 (0.00%) 0	0 / 82 (0.00%) 0	
Increased appetite			

subjects affected / exposed	0 / 87 (0.00%)	0 / 82 (0.00%)	
occurrences (all)	0	0	

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
31 January 2012	The following major changes were made in Amendment: 1) For Phase Ib, the exclusion criteria were changed: Participants who had received selected prior anti-cancer therapies (cytochrome P 17 inhibitors including abiraterone, inhibitors of Akt, PI3K/mTOR) were allowed. 2) For Phase Ib and II, participants who had received previous therapies with ketoconazole and/or MDV3100 (i.e., enzalutamide) were allowed to enroll. 3) The definition of DLT with respect to liver parameters was modified. 4) The number of sites participating in this study was increased from approximately 50 to approximately 70 sites. 5) Prednisolone was added as an alternative to prednisone. 6) Denosumab was added to the list of allowed concomitant therapies.
30 April 2012	The amendment was released to clarify the process for unblinding of treatment assignment by the investigator through the interactive voice response system/interactive web response system.
12 September 2012	The amendment was released to introduce the following main changes: 1) The study was amended to indicate the recommended dose of GDC-0068 (400 mg daily) in combination with abiraterone for the Phase II part of the study, on the basis of safety and pharmacokinetic data from the Phase Ib portion. In addition, because of adverse events observed with the combination GDC-0980 (30 mg daily) and abiraterone, further exploration of this combination in the study was discontinued. 2) The design of the Phase II portion of the study was amended accordingly. 3) Updated dose-modification guidelines for the management of hyperglycemia, skin toxicity, gastrointestinal toxicities, and pneumonitis were included.
07 March 2013	The following major changes were made in Amendment: 1) Eligibility criteria were modified to exclude patients with poor performance status (Eastern Cooperative Oncology Group [ECOG] 2). 2) Accordingly, the stratification factors in the Phase II portion of the study were changed: randomization was stratified by prior enzalutamide (yes vs. no), progression factor (PSA only vs. other), and number of prior chemotherapy regimens for metastatic disease (one vs. more than one). 3) Dose modification for prednisone was added. 4) Risk of abiraterone on the fetus was added. 5) Dose reduction of ipatasertib/placebo or GDC-0980 was added as an option for management of fasting Grade ≥ 3 hyperglycemia if fasting glucose levels recovered to Grade less than or equal to 2 within 3 days upon dose interruption. 6) Cases of potential drug-induced liver injury that included an elevated alanine aminotransferase or aspartate aminotransferase in combination with either an elevated bilirubin or clinical jaundice, and suspected transmission of an infectious agent by the study drug were added as protocol defined events of special interest.
28 November 2013	The disease inclusion criterion was updated to align with the study title and rationale that participants who progressed from docetaxel-based chemotherapy (including docetaxel and cabazitaxel, sharing the same cytotoxic mechanism) should be eligible.

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported