



## Clinical trial results:

### Assessment of Safety, Tolerability, and Pharmacodynamic Effects of LY2886721 in Patients with Mild Cognitive Impairment Due to Alzheimer's Disease or Mild Alzheimer's Disease

#### Summary

EudraCT number	2011-005217-37
Trial protocol	NL ES IT
Global end of trial date	06 August 2013

#### Results information

Result version number	v1 (current)
This version publication date	09 April 2018
First version publication date	09 April 2018

#### Trial information

##### Trial identification

Sponsor protocol code	I4O-MC-BACC
-----------------------	-------------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT01561430
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	Eli Lilly and Company
Sponsor organisation address	Lilly Corporate Center, Indianapolis, IN, United States, 46285
Public contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877-285-4559,
Scientific contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877-CTLILLY,

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

---

**Results analysis stage**

Analysis stage	Final
Date of interim/final analysis	06 August 2013
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	06 August 2013
Was the trial ended prematurely?	No

Notes:

---

**General information about the trial**

Main objective of the trial:

The purpose of this Phase 1/Phase 2 study is to evaluate how the body handles the drug and the drug's effect on the body of participants with mild cognitive impairment (MCI) due to Alzheimer's Disease (AD) or mild AD and who test positive for amyloid plaque.

Protection of trial subjects:

This study was conducted in accordance with International Conference on Harmonization (ICH) Good Clinical Practice, and the principles of the Declaration of Helsinki, in addition to following the laws and regulations of the country or countries in which a study is conducted.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	22 March 2012
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

---

**Population of trial subjects****Subjects enrolled per country**

Country: Number of subjects enrolled	United States: 60
Country: Number of subjects enrolled	Netherlands: 5
Country: Number of subjects enrolled	Japan: 5
Worldwide total number of subjects	70
EEA total number of subjects	5

Notes:

---

**Subjects enrolled per age group**

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	28
From 65 to 84 years	39

85 years and over	3
-------------------	---

## Subject disposition

### Recruitment

Recruitment details: -

### Pre-assignment

Screening details:

As a result of findings from the I4O-MC-BACJ study (NCT01534273), enrollment was discontinued to the 15 milligrams (mg) arm. The 9 participants already enrolled were allowed to continue study treatment at the 15 mg dose.

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator, Carer

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	15 mg LY2886721

Arm description:

LY2886721: 15 milligrams (mg), capsules, administered orally, once daily for 26 weeks.

Arm type	Experimental
Investigational medicinal product name	15 mg LY2886721
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

LY2886721: 15 milligrams (mg), capsules, administered orally, once daily for 26 weeks.

<b>Arm title</b>	35 mg LY2886721
------------------	-----------------

Arm description:

LY2886721: 35 mg, capsules, administered orally, once daily for 26 weeks.

Arm type	Experimental
Investigational medicinal product name	35 mg LY2886721
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

LY2886721: 35 mg, capsules, administered orally, once daily for 26 weeks.

<b>Arm title</b>	70 mg LY2886721
------------------	-----------------

Arm description:

LY2886721: 70 mg, capsules, administered orally, once daily for 26 weeks.

Arm type	Experimental
Investigational medicinal product name	70 mg LY2886721
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

LY2886721: 70 mg, capsules, administered orally, once daily for 26 weeks.

<b>Arm title</b>	Placebo
Arm description:	
Placebo: 1 placebo capsule, administered orally, once daily for 26 weeks.	
Arm type	Placebo
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

Placebo: 1 placebo capsule, administered orally, once daily for 26 weeks.

<b>Number of subjects in period 1</b>	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721
Started	9	23	18
Received at Least One Dose of Study Drug	9	23	18
Completed	8	8	1
Not completed	1	15	17
Physician decision	-	1	-
Consent withdrawn by subject	-	-	1
Adverse event, non-fatal	-	-	3
Sponsor Decision	1	14	13

<b>Number of subjects in period 1</b>	Placebo
Started	20
Received at Least One Dose of Study Drug	20
Completed	7
Not completed	13
Physician decision	-
Consent withdrawn by subject	-
Adverse event, non-fatal	1
Sponsor Decision	12

## Baseline characteristics

### Reporting groups

Reporting group title	Overall Study
-----------------------	---------------

Reporting group description: -

Reporting group values	Overall Study	Total	
Number of subjects	70	70	
Age categorical			
Units: Subjects			
In utero	0	0	
Preterm newborn infants (gestational age < 37 wks)	0	0	
Newborns (0-27 days)	0	0	
Infants and toddlers (28 days-23 months)	0	0	
Children (2-11 years)	0	0	
Adolescents (12-17 years)	0	0	
Adults (18-64 years)	28	28	
From 65-84 years	39	39	
85 years and over	3	3	
Gender, Male/Female			
Units:			
Female	33	33	
Male	37	37	
Race/Ethnicity, Customized			
Units: Subjects			
American Indian or Alaska Native	0	0	
Asian	6	6	
Black or African American	0	0	
Native Hawaiian or Other Pacific Islander	0	0	
White	64	64	
Region of Enrollment			
Units: Subjects			
United States	60	60	
Netherlands	5	5	
Japan	5	5	

## End points

### End points reporting groups

Reporting group title	15 mg LY2886721
Reporting group description: LY2886721: 15 milligrams (mg), capsules, administered orally, once daily for 26 weeks.	
Reporting group title	35 mg LY2886721
Reporting group description: LY2886721: 35 mg, capsules, administered orally, once daily for 26 weeks.	
Reporting group title	70 mg LY2886721
Reporting group description: LY2886721: 70 mg, capsules, administered orally, once daily for 26 weeks.	
Reporting group title	Placebo
Reporting group description: Placebo: 1 placebo capsule, administered orally, once daily for 26 weeks.	

### Primary: Change from Baseline to 12 Weeks in Cerebrospinal Fluid (CSF) Amyloid Beta (A $\beta$ )1-40 and A $\beta$ 1-42 Concentrations

End point title	Change from Baseline to 12 Weeks in Cerebrospinal Fluid (CSF) Amyloid Beta (A $\beta$ )1-40 and A $\beta$ 1-42 Concentrations
End point description: Percent change in lumbar CSF concentrations of A $\beta$ 1-40 and A $\beta$ 1-42 from baseline at 12 weeks post-dose was calculated. The units for CSF were picograms per milliliter (pg/mL). Least Squares (LS) means of percent change in concentration from baseline was calculated using analysis of covariance (ANCOVA) with baseline as a covariate and treatment as a fixed effect.  Analysis Population Description: All randomized participants with evaluable post-baseline CSF A $\beta$ 1-40 or A $\beta$ 1-42 data.	
End point type	Primary
End point timeframe: Baseline, 12 weeks	

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	7	9	5	12
Units: percent change in A $\beta$ 1-40 and A $\beta$ 1-42				
least squares mean (standard error)				
A $\beta$ 1-40 (n=7, 9, 5, 11)	-31.8 ( $\pm$ 7.35)	-57.7 ( $\pm$ 6.67)	-58.9 ( $\pm$ 8.73)	3.7 ( $\pm$ 5.96)
A $\beta$ 1-42 (n=7, 9, 5, 12)	-34.4 ( $\pm$ 7.27)	-52 ( $\pm$ 6.8)	-60.7 ( $\pm$ 8.42)	6.5 ( $\pm$ 6.06)

### Statistical analyses

Statistical analysis title	15 mg Change from Baseline to Week 12 CSF A $\beta$ 1-40
Comparison groups	15 mg LY2886721 v Placebo

Number of subjects included in analysis	19
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	ANCOVA
Parameter estimate	Least Squares Mean
Point estimate	-31.8
Confidence interval	
level	Other: 80 %
sides	2-sided
lower limit	-41.4
upper limit	-22.1
Variability estimate	Standard error of the mean

<b>Statistical analysis title</b>	35 mg Change from Baseline to Week 12 CSF A $\beta$ 1-40
Comparison groups	35 mg LY2886721 v Placebo
Number of subjects included in analysis	21
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	ANCOVA
Parameter estimate	Least Squire Means
Point estimate	-57.7
Confidence interval	
level	Other: 80 %
sides	2-sided
lower limit	-66.4
upper limit	-48.9
Variability estimate	Standard error of the mean

<b>Statistical analysis title</b>	70 mg Change from Baseline to Week 12 CSF A $\beta$ 1-40
Comparison groups	70 mg LY2886721 v Placebo
Number of subjects included in analysis	17
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	ANCOVA
Parameter estimate	Least Squares Mean
Point estimate	-58.9
Confidence interval	
level	Other: 80 %
sides	2-sided
lower limit	-70.4
upper limit	-47.4
Variability estimate	Standard error of the mean



<b>Statistical analysis title</b>	15mg Change from Baseline to 12 Weeks CSF A $\beta$ 1-42
Comparison groups	Placebo v 15 mg LY2886721
Number of subjects included in analysis	19
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	ANCOVA
Parameter estimate	Least Squares Mean
Point estimate	-34.4
Confidence interval	
level	Other: 80 %
sides	2-sided
lower limit	-44
upper limit	-24.9
Variability estimate	Standard error of the mean

<b>Statistical analysis title</b>	35 mg Change from Baseline to 12 Weeks CSF A $\beta$ 1-42
Comparison groups	35 mg LY2886721 v Placebo
Number of subjects included in analysis	21
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	ANCOVA
Parameter estimate	Least Squares Mean
Point estimate	-52
Confidence interval	
level	Other: 80 %
sides	2-sided
lower limit	-60.9
upper limit	-43.1
Variability estimate	Standard error of the mean

<b>Statistical analysis title</b>	70 mg Change from Baseline to 12 Weeks CSF A $\beta$ 1-42
Comparison groups	70 mg LY2886721 v Placebo
Number of subjects included in analysis	17
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	ANCOVA
Parameter estimate	Least Squares Mean
Point estimate	-60.7

Confidence interval	
level	Other: 80 %
sides	2-sided
lower limit	-71.7
upper limit	-49.6
Variability estimate	Standard error of the mean

### Primary: Change from Baseline to 26 Weeks in CSF A $\beta$ 1-40 and A $\beta$ 1-42 Concentrations

End point title	Change from Baseline to 26 Weeks in CSF A $\beta$ 1-40 and A $\beta$ 1-42 Concentrations <sup>[1]</sup>
-----------------	---

End point description:

Percent change in lumbar CSF concentrations of A $\beta$ 1-40 and A $\beta$ 1-42 from baseline at 26 weeks post-dose was to be calculated. The units for CSF were picograms per milliliter (pg/mL). LS means of percent change in concentration from baseline was calculated using ANCOVA with baseline as a covariate and treatment as a fixed effect.

9999=Data Not Available (N/A)

End point type	Primary
----------------	---------

End point timeframe:

Baseline, 26 weeks

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Zero total participants were analyzed for this End point; statistical analysis was not completed.

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	0 <sup>[2]</sup>	0 <sup>[3]</sup>	0 <sup>[4]</sup>	0 <sup>[5]</sup>
Units: percent change in A $\beta$ 1-40 and A $\beta$ 1-42				
least squares mean (standard error)	()	()	()	()

Notes:

[2] - No participants analyzed at 26 weeks due to the termination of the trial.

[3] - No participants analyzed at 26 weeks due to the termination of the trial.

[4] - No participants analyzed at 26 weeks due to the termination of the trial.

[5] - No participants analyzed at 26 weeks due to the termination of the trial.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in Plasma Amyloid Beta (A $\beta$ )1-40 and A $\beta$ 1-42 Concentrations

End point title	Change from Baseline in Plasma Amyloid Beta (A $\beta$ )1-40 and A $\beta$ 1-42 Concentrations
-----------------	--

End point description:

Percent change in plasma concentrations of A $\beta$ 1-40 and A $\beta$ 1-42 from baseline at 12 weeks and 26 weeks post-dose was calculated. The units for CSF were picograms per milliliter (pg/mL).

Analysis Population Description: All randomized participants with evaluable post-baseline CSF A $\beta$ 1-40 or A $\beta$ 1-42 data.

9999=Data Not Available (N/A)

End point type	Secondary
End point timeframe:	
Baseline, 12 weeks, 26 weeks	

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	6	8	4 <sup>[6]</sup>	12
Units: percent change in Aβ1-40 and Aβ1-42				
arithmetic mean (standard deviation)				
Aβ1-40, Week 12 (n=6, 8, 4, 12)	-73.4 (± 6.22)	-80.8 (± 8.07)	-88.1 (± 2.66)	1.9 (± 15)
Aβ1-40, Week 26 (n=5, 6, 1, 6)	-72.3 (± 11.5)	-85 (± 3.89)	-89.1 (± 9999)	0.944 (± 22.6)
Aβ1-42, Week 12 (n=6, 8, 4, 12)	-63.4 (± 6.11)	-71.1 (± 8.38)	-78.2 (± 5.08)	0.541 (± 10.1)
Aβ1-42, Week 26 (n=5, 6, 1, 6)	-65.1 (± 8.85)	-75 (± 4.33)	-80.8 (± 9999)	1.82 (± 9.98)

Notes:

[6] - Standard deviation not reported due to not enough participants for analyses.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Change from Baseline to 26 Weeks in Neuropsychological Test Battery (NTB)

End point title	Change from Baseline to 26 Weeks in Neuropsychological Test Battery (NTB)
-----------------	---

End point description:

The NTB is a composite cognitive measure in clinical Alzheimer's disease studies and is a collection of several written and oral tests that examines verbal and nonverbal brain functions. Lower scores indicate greater cognitive impairment. LS means were calculated using ANCOVA with baseline as a covariate and treatment as a fixed effect.

Analysis Population Description: All randomized participants with evaluable NTB data.

End point type	Secondary
End point timeframe:	
Baseline, 26 weeks	

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	8	7	1	9
Units: units on a scale				
least squares mean (standard error)				
NTB z-score	-0.039 (± 0.1096)	-0.161 (± 0.1165)	0.424 (± 0.3156)	-0.262 (± 0.1028)
NTB total score	2 (± 4.77)	-5.9 (± 5.09)	21.2 (± 13.69)	-14.9 (± 4.49)

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline to 26 Weeks in Alzheimer's Disease Assessment Scale - Cognitive Subscale (ADAS-Cog)

End point title	Change from Baseline to 26 Weeks in Alzheimer's Disease Assessment Scale - Cognitive Subscale (ADAS-Cog)
End point description: ADAS-Cog11 is an 11-item instrument measuring impairment in memory (Items 1-4, 7, 11), praxis (Items 4 and 5), orientation (Item 6), and language (Items 8-10). Item 1 ranged 0 (all items recalled correctly)-10 (none recalled correctly); Items 2-5 and 8-11 ranged 0 (all items named, performed, drawn, spoken, remember correctly/clearly)-5 (none correct/not clearly spoken); Item 6 ranged 0 (no incorrect responses)-8 (all incorrect); and Item 7 ranged 0 (all words remembered correctly)-12 (no words remembered correctly) for a total ADAS-Cog11 score of 0-70 with higher scores indicating greater disease severity. A score of 0-10 for delayed free recall and a conversion code of 0-5 for digit cancellation and maze completion was added to the total ADAS-Cog11 score for a total ADAS-Cog14 score ranging 0-90 with higher scores indicating greater impairment. LS means were calculated using Mixed Model Repeated Measures (MMRM) with Treatment + Visit + Treatment*Visit + Baseline + Baseline*Visit.	
End point type	Secondary
End point timeframe: Baseline, 26 weeks	

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	7 <sup>[7]</sup>	9 <sup>[8]</sup>	1 <sup>[9]</sup>	9 <sup>[10]</sup>
Units: units on a scale				
least squares mean (standard error)	1.1 (± 2.77)	0.6 (± 2.48)	0.6 (± 7.41)	1.3 (± 2.44)

Notes:

[7] - All randomized participants with evaluable ADAS-Cog data.

[8] - All randomized participants with evaluable ADAS-Cog data.

[9] - All randomized participants with evaluable ADAS-Cog data.

[10] - All randomized participants with evaluable ADAS-Cog data.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline to 26 Weeks in the Clinical Dementia Rating Scale - Sum of Boxes (CDR-SB)

End point title	Change from Baseline to 26 Weeks in the Clinical Dementia Rating Scale - Sum of Boxes (CDR-SB)
End point description: The CDR-SB is a composite measure of 6 domains of cognitive and functional performance: memory, orientation, judgment and problem solving, community affairs, home and hobbies, and personal care.	

Scores ranged from 0 to 18, with higher scores indicating greater impairment. LS means were calculated using MMRM with Treatment + Visit + Treatment\*Visit + Baseline + Baseline\*Visit.

Analysis Population Description: All randomized participants with evaluable CDR-SB data.

End point type	Secondary
End point timeframe:	
Baseline, 26 weeks	

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	8	9	1	8
Units: units on a scale				
least squares mean (standard error)	0.8 (± 0.432)	0.52 (± 0.395)	1.5 (± 1.013)	1.19 (± 0.402)

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline to 26 Weeks in Mini Mental State Examination (MMSE)

End point title	Change from Baseline to 26 Weeks in Mini Mental State Examination (MMSE)
-----------------	--

End point description:

The MMSE (Folstein et al. 1975) is one of the most widely used screening instruments for cognitive impairment. The test consists of five sections (orientation, registration, attention-calculation, recall, and language) and provides a total score ranging from 0 to 30, with lower scores indicative of greater cognitive impairment. LS means were calculated using MMRM with Treatment + Visit + Treatment\*Visit + Baseline + Baseline\*Visit.

Analysis Population Description: All randomized participants with evaluable MMSE data.

End point type	Secondary
End point timeframe:	
Baseline, 26 weeks	

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	8	9	1	10
Units: units on a scale				
least squares mean (standard error)	-0.7 (± 0.87)	-1.8 (± 0.81)	0.2 (± 2.41)	-3 (± 0.77)

### Statistical analyses

No statistical analyses for this end point

## Secondary: Change from Baseline in Cerebrospinal Fluid (CSF) Tau and Phosphorylated Tau (Ptau)-181 Concentrations

End point title	Change from Baseline in Cerebrospinal Fluid (CSF) Tau and Phosphorylated Tau (Ptau)-181 Concentrations
-----------------	--

End point description:

Percent change in lumbar CSF tau and ptau-181 concentrations from baseline at 12 weeks post-dose and 26 weeks post-dose was calculated. The units for CSF were picograms per milliliter (pg/mL). Least Squares (LS) means of percent change in concentration from baseline was calculated using analysis of covariance (ANCOVA) with baseline as a covariate and treatment as a fixed effect.

Analysis Population Description: All randomized participants with evaluable post-baseline CSF tau or ptau data.

9999=Data Not Available (N/A)

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, 12 weeks, 26 weeks

End point values	15 mg LY2886721	35 mg LY2886721	70 mg LY2886721	Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	7 <sup>[11]</sup>	9 <sup>[12]</sup>	5 <sup>[13]</sup>	12 <sup>[14]</sup>
Units: percent change in tau and ptau-181				
least squares mean (standard error)				
CSF Tau, Week 12	14.3 (± 6.06)	2.6 (± 5.42)	6.8 (± 7.49)	-4.4 (± 4.63)
CSF Tau, Week 26	9999 (± 9999)	9999 (± 9999)	9999 (± 9999)	9999 (± 9999)
CSF Ptau, Week 12	0.7 (± 4.81)	1.7 (± 4.01)	4.3 (± 5.41)	-3.6 (± 3.4)
CSF Ptau, Week 26	9999 (± 9999)	9999 (± 9999)	9999 (± 9999)	9999 (± 9999)

Notes:

[11] - No participants analyzed at 26 weeks due to the termination of the trial.

[12] - No participants analyzed at 26 weeks due to the termination of the trial.

[13] - No participants analyzed at 26 weeks due to the termination of the trial.

[14] - No participants analyzed at 26 weeks due to the termination of the trial.

## Statistical analyses

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Entire Study

Adverse event reporting additional description:

I4O-MC-BACC

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	16.0
--------------------	------

### Reporting groups

Reporting group title	Placebo
-----------------------	---------

Reporting group description: -

Reporting group title	15 mg LY2886721
-----------------------	-----------------

Reporting group description: -

Reporting group title	35 mg LY2886721
-----------------------	-----------------

Reporting group description: -

Reporting group title	70 mg LY2886721
-----------------------	-----------------

Reporting group description: -

Serious adverse events	Placebo	15 mg LY2886721	35 mg LY2886721
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	2 / 23 (8.70%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events	0	0	0
Nervous system disorders			
syncope			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
crohn's disease			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
small intestinal obstruction			
alternative dictionary used:			

MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

<b>Serious adverse events</b>	70 mg LY2886721		
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 18 (0.00%)		
number of deaths (all causes)	0		
number of deaths resulting from adverse events	0		
Nervous system disorders			
syncope			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Gastrointestinal disorders			
crohn's disease			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
small intestinal obstruction			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		

Frequency threshold for reporting non-serious adverse events: 0 %

<b>Non-serious adverse events</b>	Placebo	15 mg LY2886721	35 mg LY2886721
Total subjects affected by non-serious adverse events			
subjects affected / exposed	15 / 20 (75.00%)	8 / 9 (88.89%)	13 / 23 (56.52%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			



basal cell carcinoma alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
squamous cell carcinoma of skin alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
Vascular disorders hypertension alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
General disorders and administration site conditions asthenia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
fatigue alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	1 / 23 (4.35%) 1
oedema peripheral alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
pyrexia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
thirst alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
Psychiatric disorders			

abnormal dreams			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
agitation			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
anxiety			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
confusional state			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
delusion			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	1	0	1
depression			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
disorientation			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
hallucination			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
hallucination, visual			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
hypnopompic hallucination			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
initial insomnia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	1
insomnia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	1 / 9 (11.11%)	1 / 23 (4.35%)
occurrences (all)	1	1	2
nightmare			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
obsessive thoughts			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
parasomnia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	1
restlessness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
sleep talking			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0

stereotypy alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	1 / 23 (4.35%) 1
terminal insomnia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
Investigations			
alanine aminotransferase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
amylase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	1 / 23 (4.35%) 1
blood alkaline phosphatase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
blood creatine phosphokinase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	2 / 20 (10.00%) 2	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
blood pressure increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	1 / 23 (4.35%) 1
colour vision tests abnormal red-green alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
electrocardiogram change			

alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
gamma-glutamyltransferase increased			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
lipase increased			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	1
liver function test abnormal			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	2 / 23 (8.70%)
occurrences (all)	0	1	2
weight increased			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
Injury, poisoning and procedural complications			
accidental overdose			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
alcohol poisoning			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
contusion			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
fall			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	1	0	1
feeding tube complication			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
foot fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
joint injury			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
laceration			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
tooth fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
Cardiac disorders			
atrial fibrillation			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
bradycardia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
Nervous system disorders			
apraxia			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
balance disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	1
cerebral microhaemorrhage			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
decreased vibratory sense			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
dizziness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	1	1	0
dysgeusia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
headache			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	2 / 9 (22.22%)	1 / 23 (4.35%)
occurrences (all)	0	2	1
sinus headache			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
Blood and lymphatic system disorders			
anaemia			
alternative dictionary used: MedDRA 16.0			

<p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>leukocytosis</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 20 (0.00%)</p> <p>0</p> <p>0 / 20 (0.00%)</p> <p>0</p>	<p>0 / 9 (0.00%)</p> <p>0</p> <p>1 / 9 (11.11%)</p> <p>1</p>	<p>1 / 23 (4.35%)</p> <p>1</p> <p>0 / 23 (0.00%)</p> <p>0</p>
<p>Ear and labyrinth disorders</p> <p>vertigo</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 20 (0.00%)</p> <p>0</p>	<p>0 / 9 (0.00%)</p> <p>0</p>	<p>1 / 23 (4.35%)</p> <p>1</p>
<p>Eye disorders</p> <p>cataract nuclear</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>diplopia</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>eye irritation</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>lacrimation increased</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>visual acuity reduced</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>1 / 20 (5.00%)</p> <p>1</p> <p>0 / 20 (0.00%)</p> <p>0</p> <p>1 / 20 (5.00%)</p> <p>1</p> <p>0 / 20 (0.00%)</p> <p>0</p> <p>0 / 20 (0.00%)</p> <p>0</p>	<p>0 / 9 (0.00%)</p> <p>0</p> <p>0 / 9 (0.00%)</p> <p>0</p> <p>0 / 9 (0.00%)</p> <p>0</p> <p>0 / 9 (0.00%)</p> <p>0</p>	<p>0 / 23 (0.00%)</p> <p>0</p> <p>1 / 23 (4.35%)</p> <p>1</p> <p>0 / 23 (0.00%)</p> <p>0</p> <p>1 / 23 (4.35%)</p> <p>1</p> <p>1 / 23 (4.35%)</p> <p>1</p>
<p>Gastrointestinal disorders</p> <p>abdominal discomfort</p> <p>alternative dictionary used: MedDRA 16.0</p>			



subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	2
abdominal distension			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	1
abdominal pain upper			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
abdominal tenderness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
dental caries			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
diarrhoea			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	1	0	1
diverticulum intestinal			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	2	0	0
dry mouth			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
flatulence			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	1

gastritis erosive			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
gastrointestinal disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	2	0	0
gastrooesophageal reflux disease			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
haematochezia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
haemorrhoidal haemorrhage			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	0	0	1
haemorrhoids			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
hiatus hernia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
lip swelling			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
nausea			
alternative dictionary used: MedDRA 16.0			

<p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>1 / 20 (5.00%)</p> <p>1</p>	<p>0 / 9 (0.00%)</p> <p>0</p>	<p>0 / 23 (0.00%)</p> <p>0</p>
<p>oesophageal disorder</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>1 / 20 (5.00%)</p> <p>1</p>	<p>0 / 9 (0.00%)</p> <p>0</p>	<p>0 / 23 (0.00%)</p> <p>0</p>
<p>oesophagitis</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>1 / 20 (5.00%)</p> <p>1</p>	<p>0 / 9 (0.00%)</p> <p>0</p>	<p>0 / 23 (0.00%)</p> <p>0</p>
<p>small intestinal obstruction</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 20 (0.00%)</p> <p>0</p>	<p>1 / 9 (11.11%)</p> <p>1</p>	<p>0 / 23 (0.00%)</p> <p>0</p>
<p>vomiting</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>2 / 20 (10.00%)</p> <p>2</p>	<p>1 / 9 (11.11%)</p> <p>1</p>	<p>0 / 23 (0.00%)</p> <p>0</p>
<p>Hepatobiliary disorders</p> <p>hepatic function abnormal</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 20 (0.00%)</p> <p>0</p>	<p>0 / 9 (0.00%)</p> <p>0</p>	<p>0 / 23 (0.00%)</p> <p>0</p>
<p>Skin and subcutaneous tissue disorders</p> <p>actinic keratosis</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>eczema</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>rash</p> <p>alternative dictionary used: MedDRA 16.0</p>	<p>0 / 20 (0.00%)</p> <p>0</p> <p>0 / 20 (0.00%)</p> <p>0</p>	<p>1 / 9 (11.11%)</p> <p>2</p> <p>0 / 9 (0.00%)</p> <p>0</p>	<p>0 / 23 (0.00%)</p> <p>0</p> <p>1 / 23 (4.35%)</p> <p>1</p>

subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
rash macular alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
rash vesicular alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
urticaria alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
Renal and urinary disorders cystitis interstitial alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
pollakiuria alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	0 / 23 (0.00%) 0
renal failure chronic alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
Endocrine disorders hypothyroidism alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
Musculoskeletal and connective tissue disorders			

arthralgia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
back pain			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
joint range of motion decreased			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	1	0	0
joint swelling			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
muscle spasms			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 20 (10.00%)	0 / 9 (0.00%)	1 / 23 (4.35%)
occurrences (all)	2	0	1
muscle tightness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	1 / 9 (11.11%)	0 / 23 (0.00%)
occurrences (all)	0	1	0
musculoskeletal chest pain			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 20 (5.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	2	0	0
musculoskeletal stiffness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
Infections and infestations			

herpes zoster alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
nasopharyngitis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	1 / 23 (4.35%) 1
sinusitis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
upper respiratory tract infection alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	1 / 9 (11.11%) 1	1 / 23 (4.35%) 1
urinary tract infection alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 9 (11.11%) 1	1 / 23 (4.35%) 1
Metabolism and nutrition disorders decreased appetite alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
dehydration alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 9 (0.00%) 0	0 / 23 (0.00%) 0
hypokalaemia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 9 (0.00%) 0	1 / 23 (4.35%) 1
increased appetite alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	0 / 20 (0.00%)	0 / 9 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0

<b>Non-serious adverse events</b>	70 mg LY2886721		
Total subjects affected by non-serious adverse events			
subjects affected / exposed	16 / 18 (88.89%)		
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
basal cell carcinoma			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
squamous cell carcinoma of skin			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
Vascular disorders			
hypertension			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
General disorders and administration site conditions			
asthenia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
fatigue			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
oedema peripheral			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
pyrexia			
alternative dictionary used: MedDRA 16.0			

<p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>thirst</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Psychiatric disorders</p> <p>abnormal dreams</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p> <p>agitation</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p> <p>anxiety</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p> <p>confusional state</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p> <p>delusion</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>depression</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>3 / 18 (16.67%)</p> <p>occurrences (all)</p> <p>3</p> <p>disorientation</p> <p>alternative dictionary used: MedDRA 16.0</p>			



subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
hallucination			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
hallucination, visual			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
hypnopompic hallucination			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
initial insomnia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
insomnia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
nightmare			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
obsessive thoughts			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
parasomnia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		

restlessness alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
sleep talking alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
stereotypy alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
terminal insomnia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
Investigations alanine aminotransferase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
amylase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
blood alkaline phosphatase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
blood creatine phosphokinase increased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
blood pressure increased alternative dictionary used:			

MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
colour vision tests abnormal red-green			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
electrocardiogram change			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
gamma-glutamyltransferase increased			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	3 / 18 (16.67%)		
occurrences (all)	3		
lipase increased			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	2		
liver function test abnormal			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
weight increased			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
Injury, poisoning and procedural complications			
accidental overdose			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
alcohol poisoning			
alternative dictionary used:			

MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
contusion			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
fall			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 18 (11.11%)		
occurrences (all)	3		
feeding tube complication			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
foot fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
joint injury			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
laceration			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	2 / 18 (11.11%)		
occurrences (all)	3		
tooth fracture			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
Cardiac disorders			
atrial fibrillation			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
bradycardia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
Nervous system disorders			
apraxia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
balance disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
cerebral microhaemorrhage			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
decreased vibratory sense			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
dizziness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
dysgeusia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
headache			
alternative dictionary used: MedDRA 16.0			

<p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p> <p>sinus headache</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>4</p>			
<p>Blood and lymphatic system disorders</p> <p>anaemia</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>leukocytosis</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Ear and labyrinth disorders</p> <p>vertigo</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Eye disorders</p> <p>cataract nuclear</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>diplopia</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>eye irritation</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>lacrimation increased</p> <p>alternative dictionary used: MedDRA 16.0</p>			

subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
visual acuity reduced			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
Gastrointestinal disorders			
abdominal discomfort			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
abdominal distension			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
abdominal pain upper			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
abdominal tenderness			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
dental caries			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
diarrhoea			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		
diverticulum intestinal			
alternative dictionary used: MedDRA 16.0			

subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
dry mouth			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
flatulence			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
gastritis erosive			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
gastrointestinal disorder			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
gastrooesophageal reflux disease			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
haematochezia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
haemorrhoidal haemorrhage			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
haemorrhoids			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		



hiatus hernia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
lip swelling alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
nausea alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
oesophageal disorder alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
oesophagitis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
small intestinal obstruction alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
vomiting alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
Hepatobiliary disorders hepatic function abnormal alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
Skin and subcutaneous tissue disorders			

actinic keratosis alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
eczema alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
rash alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
rash macular alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
rash vesicular alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
urticaria alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
Renal and urinary disorders cystitis interstitial alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
pollakiuria alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
renal failure chronic alternative dictionary used: MedDRA 16.0			

subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1		
Endocrine disorders hypothyroidism alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0		
Musculoskeletal and connective tissue disorders arthralgia alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  back pain alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  joint range of motion decreased alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  joint swelling alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  muscle spasms alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  muscle tightness alternative dictionary used: MedDRA 16.0 subjects affected / exposed occurrences (all)  musculoskeletal chest pain alternative dictionary used: MedDRA 16.0	0 / 18 (0.00%) 0  1 / 18 (5.56%) 2  0 / 18 (0.00%) 0  1 / 18 (5.56%) 1  1 / 18 (5.56%) 1  0 / 18 (0.00%) 0		

<p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>musculoskeletal stiffness</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p>			
<p>Infections and infestations</p> <p>herpes zoster</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p> <p>nasopharyngitis</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>sinusitis</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>upper respiratory tract infection</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>1 / 18 (5.56%)</p> <p>occurrences (all)</p> <p>1</p> <p>urinary tract infection</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Metabolism and nutrition disorders</p> <p>decreased appetite</p> <p>alternative dictionary used: MedDRA 16.0</p> <p>subjects affected / exposed</p> <p>0 / 18 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>dehydration</p> <p>alternative dictionary used: MedDRA 16.0</p>			

subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
hypokalaemia			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	0 / 18 (0.00%)		
occurrences (all)	0		
increased appetite			
alternative dictionary used: MedDRA 16.0			
subjects affected / exposed	1 / 18 (5.56%)		
occurrences (all)	1		

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
02 October 2012	Amendment B:  Added the 70 mg dose and dropped the 15 mg dose.  Added change in CSF tau and p-tau181 as secondary outcomes.
08 April 2013	Amendment C:  Clarified the purpose of the study as a Ph2a PK/PD/safety study.  Revised diagnostic criteria to align with contemporary clinical practice.  Removed florbetapir scan at Visit 11.  Alterations to inclusion/exclusion criteria.  Added discontinuation criteria for patients with abnormal liver tests.  Altered the randomization ratio to increase the number of subjects assigned to 70 mg dose.

Notes:

### Interruptions (globally)

Were there any global interruptions to the trial? Yes

Date	Interruption	Restart date
12 June 2013	The study was terminated early due to hepatic safety finding.	-

Notes:

### Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

Study was terminated early because of abnormal liver biochemical tests levels for 4 participants.

Notes: