



Clinical trial results:

A Two Year, Multicenter, Randomized, Double-blind, Placebo-controlled, Parallel Group Trial to Evaluate Efficacy, Safety, Tolerability, and Pharmacokinetics of Teriflunomide Administered Orally Once Daily in Pediatric Patients With Relapsing Forms of Multiple Sclerosis Followed by an Open-label Extension

Summary

EudraCT number	2011-005249-12
Trial protocol	EE BE ES GR PL GB FR LT Outside EU/EEA IE NL BG PT SI IT
Global end of trial date	29 July 2024

Results information

Result version number	v3 (current)
This version publication date	04 January 2025
First version publication date	29 January 2021
Version creation reason	

Trial information

Trial identification

Sponsor protocol code	EFC11759
-----------------------	----------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02201108
WHO universal trial number (UTN)	U1111-1124-0983

Notes:

Sponsors

Sponsor organisation name	Genzyme, a Sanofi Company
Sponsor organisation address	50 Binney Street, Cambridge, Massachusetts, United States, 02142
Public contact	Trial Transparency Team, Sanofi aventis recherche & développement, Contact-US@sanofi.com
Scientific contact	Trial Transparency Team, Sanofi aventis recherche & développement, Contact-US@sanofi.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	Yes
EMA paediatric investigation plan number(s)	EMA-001094-PIP01-10
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	12 September 2024
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	29 July 2024
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

To assess the effect of teriflunomide in comparison to placebo on disease activity as measured by time to first clinical relapse after randomisation in children and adolescents 10 to 17 years of age with relapsing forms of multiple sclerosis.

Protection of trial subjects:

The study was conducted by investigators experienced in the treatment of paediatric patients. The parent(s) or guardian(s) as well as the children were fully informed of all pertinent aspects of the clinical trial as well as the possibility to discontinue at any time. In addition to the consent form for the parent(s)/guardian(s), an assent form in child-appropriate language was provided and explained to the child. Repeated invasive procedures were minimised. The number of blood samples as well as the amount of blood drawn were adjusted according to age and weight. A topical anesthesia might have been used to minimise distress and discomfort.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	16 July 2014
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Spain: 1
Country: Number of subjects enrolled	Tunisia: 4
Country: Number of subjects enrolled	Türkiye: 32
Country: Number of subjects enrolled	Ukraine: 7
Country: Number of subjects enrolled	United Kingdom: 2
Country: Number of subjects enrolled	United States: 5
Country: Number of subjects enrolled	Belgium: 2
Country: Number of subjects enrolled	Bulgaria: 2
Country: Number of subjects enrolled	Canada: 2
Country: Number of subjects enrolled	China: 37
Country: Number of subjects enrolled	Estonia: 4
Country: Number of subjects enrolled	France: 17
Country: Number of subjects enrolled	Greece: 2
Country: Number of subjects enrolled	Israel: 9

Country: Number of subjects enrolled	Lebanon: 4
Country: Number of subjects enrolled	Lithuania: 1
Country: Number of subjects enrolled	Morocco: 2
Country: Number of subjects enrolled	Netherlands: 2
Country: Number of subjects enrolled	Portugal: 2
Country: Number of subjects enrolled	Russian Federation: 27
Country: Number of subjects enrolled	Serbia: 2
Worldwide total number of subjects	166
EEA total number of subjects	33

Notes:

Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	16
Adolescents (12-17 years)	150
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

Subject disposition

Recruitment

Recruitment details:

Study was conducted at 57 active centres in 21 countries. A total of 185 subjects were screened between 16 July 2014 and 27 December 2017, of which 166 subjects were enrolled and randomised. A total of 19 subjects failed screening mainly due to meeting exclusion criteria.

Pre-assignment

Screening details:

Subjects were randomly assigned to receive either teriflunomide or placebo in a 2:1 ratio via Interactive Voice Response System. Randomisation was stratified by the country and subject's pubertal status.

Period 1

Period 1 title	Double-blind Period (up to Week 96)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

Arms

Are arms mutually exclusive?	Yes
Arm title	Placebo

Arm description:

Subjects received placebo matching to teriflunomide tablet orally once daily (QD) for 96 weeks.

Arm type	Placebo
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

A single tablet of placebo (matched to teriflunomide) was administered orally QD with or without food in the morning preferably at the same time each day of the double-blind treatment period (i.e., 96 weeks).

Arm title	Teriflunomide
------------------	---------------

Arm description:

Subjects received 1 Teriflunomide tablet, 3.5 milligrams (mg) (in case of body weight [BW] up to 40 kilograms [kg]) or 7 mg (in case of BW greater than [$>$]40 kg) orally QD for 8 weeks. After 8 weeks, based on subjects predicted pharmacokinetic (PK) parameters, teriflunomide was administered in following manner up to 96 weeks: if predicted PK parameters less than or equal to (\leq)95th percentile of adult range after 7 mg QD: 1 tablet of 7 mg daily (for BW up to 40 kg) or 1 tablet of 14 mg daily (for BW $>$ 40 kg); or if predicted PK parameters $>$ 95th percentile of adult range: 1 tablet of 3.5 mg daily (for BW up to 40 kg) or 1 tablet of 7 mg daily (for BW $>$ 40 kg). The adult range (5th-95th percentile) of predicted steady state PK parameters for 7 mg dose was defined as maximum concentration observed (C_{max}) ranging from 8.03 to 49.10 micrograms per millilitre (mcg/mL) and area under the curve from time 0 hour to 24 hours (AUC_{0-24}) ranging from 184 to 1160 micrograms*hour per millilitre (mcg*h/mL).

Arm type	Experimental
Investigational medicinal product name	Teriflunomide
Investigational medicinal product code	AUBAGIO, HMR1726
Other name	
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

Teriflunomide (3.5 mg or 7 mg or 14 mg) film-coated tablet was administered orally QD with or without food in the morning preferably at the same time each day of the double-blind treatment period (i.e., 96

weeks).

Number of subjects in period 1	Placebo	Teriflunomide
Started	57	109
Completed	53	102
Not completed	4	7
Consent withdrawn by subject	2	-
Adverse Event	-	6
Lack of efficacy	2	1

Period 2

Period 2 title	Open Label Period (up to Week 192)
Is this the baseline period?	No
Allocation method	Randomised - controlled
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
Arm title	Placebo/Teriflunomide

Arm description:

Subjects previously treated with placebo during the double blind (DB) period received 1 teriflunomide tablet 3.5 mg (BW ≤ 40 kg) or 7 mg teriflunomide (BW > 40 kg) for first 8 weeks. After 8 weeks if predicted PK parameter was ≤ 95th percentile of adult range then subjects received 7 mg teriflunomide (BW ≤ 40 kg) and 14 mg teriflunomide (BW > 40 kg) in the open label (OL) period for 96 weeks (i.e., up to Week 192). The adult range (5th-95th percentile) of predicted steady state PK parameters for 7 mg dose was defined as C_{max} ranging from 8.03 to 49.10 mcg/mL and AUC₀₋₂₄ ranging from 184 to 1160 mcg*h/mL.

Arm type	Experimental
Investigational medicinal product name	Teriflunomide
Investigational medicinal product code	AUBAGIO, HMR1726
Other name	
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

Teriflunomide (3.5 mg or 7 mg or 14 mg) film-coated tablet was administered orally QD with or without food in the morning preferably at the same time each day.

Arm title	Teriflunomide / Teriflunomide
-----------	-------------------------------

Arm description:

Subjects previously treated with Teriflunomide during DB period continued receiving Teriflunomide in the

OL period for additional 96 weeks (i.e., up to Week 192).

Arm type	Experimental
Investigational medicinal product name	Teriflunomide
Investigational medicinal product code	AUBAGIO, HMR1726
Other name	
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

Teriflunomide (3.5 mg or 7 mg or 14 mg) film-coated tablet was administered orally QD with or without food in the morning preferably at the same time each day.

Number of subjects in period 2^[1]	Placebo/Teriflunomide	Teriflunomide / Teriflunomide
Started	52	100
Completed	31	73
Not completed	21	27
Adverse Event	7	5
Other reason	4	7
Poor compliance to protocol	-	1
Lack of efficacy	10	14

Notes:

[1] - The number of subjects starting the period is not consistent with the number completing the preceding period. It is expected the number of subjects starting the subsequent period will be the same as the number completing the preceding period.

Justification: Post completion of DB period, only eligible subjects entered OL period.

Period 3

Period 3 title	Extension Period (Up to Week 492)
Is this the baseline period?	No
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Arm title	Teriflunomide
------------------	---------------

Arm description:

Subjects who were not adults when they completed the 192-week OL period were offered an optional additional extension period to receive 1 teriflunomide tablet 14 mg daily (or every other day if BW was <40 kg) until they became adults or until they were able to switch to the commercial product, whichever came first.

Arm type	Experimental
Investigational medicinal product name	Teriflunomide
Investigational medicinal product code	
Other name	AUBAGIO, HMR1726
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

Teriflunomide (14 mg) film-coated tablet was administered orally QD (or every other day if BW was <40 kg) with or without food in the morning preferably at the same time each day/every other day during the extension period (i.e., approximately 300 weeks)

Number of subjects in period 3^[2]	Teriflunomide
Started	27
Completed	24
Not completed	3
Adverse event, non-fatal	2
Poor compliance to protocol	1

Notes:

[2] - The number of subjects starting the period is not consistent with the number completing the preceding period. It is expected the number of subjects starting the subsequent period will be the same as the number completing the preceding period.

Justification: Post completion of OL period, only eligible subjects entered the extension period.

Baseline characteristics

Reporting groups

Reporting group title	Placebo
-----------------------	---------

Reporting group description:

Subjects received placebo matching to teriflunomide tablet orally once daily (QD) for 96 weeks.

Reporting group title	Teriflunomide
-----------------------	---------------

Reporting group description:

Subjects received 1 Teriflunomide tablet, 3.5 milligrams (mg) (in case of body weight [BW] up to 40 kilograms [kg]) or 7 mg (in case of BW greater than [$>$]40 kg) orally QD for 8 weeks. After 8 weeks, based on subjects predicted pharmacokinetic (PK) parameters, teriflunomide was administered in following manner up to 96 weeks: if predicted PK parameters less than or equal to (\leq)95th percentile of adult range after 7 mg QD: 1 tablet of 7 mg daily (for BW up to 40 kg) or 1 tablet of 14 mg daily (for BW $>$ 40 kg); or if predicted PK parameters $>$ 95th percentile of adult range: 1 tablet of 3.5 mg daily (for BW up to 40 kg) or 1 tablet of 7 mg daily (for BW $>$ 40 kg). The adult range (5th-95th percentile) of predicted steady state PK parameters for 7 mg dose was defined as maximum concentration observed (C_{max}) ranging from 8.03 to 49.10 micrograms per millilitre (mcg/mL) and area under the curve from time 0 hour to 24 hours (AUC₀₋₂₄) ranging from 184 to 1160 micrograms*hour per millilitre (mcg*h/mL).

Reporting group values	Placebo	Teriflunomide	Total
Number of subjects	57	109	166
Age categorical			
Units: Subjects			

Age continuous			
Units: years			
arithmetic mean	14.7	14.6	
standard deviation	± 2.1	± 2.0	-
Gender categorical			
Units: Subjects			
Female	39	72	111
Male	18	37	55
Race			
Units: Subjects			
Caucasian/White	42	75	117
Black	1	4	5
Asian/Oriental	12	25	37
Other	2	5	7

End points

End points reporting groups

Reporting group title	Placebo
-----------------------	---------

Reporting group description:

Subjects received placebo matching to teriflunomide tablet orally once daily (QD) for 96 weeks.

Reporting group title	Teriflunomide
-----------------------	---------------

Reporting group description:

Subjects received 1 Teriflunomide tablet, 3.5 milligrams (mg) (in case of body weight [BW] up to 40 kilograms [kg]) or 7 mg (in case of BW greater than [$>$]40 kg) orally QD for 8 weeks. After 8 weeks, based on subjects predicted pharmacokinetic (PK) parameters, teriflunomide was administered in following manner up to 96 weeks: if predicted PK parameters less than or equal to (\leq)95th percentile of adult range after 7 mg QD: 1 tablet of 7 mg daily (for BW up to 40 kg) or 1 tablet of 14 mg daily (for BW $>$ 40 kg); or if predicted PK parameters $>$ 95th percentile of adult range: 1 tablet of 3.5 mg daily (for BW up to 40 kg) or 1 tablet of 7 mg daily (for BW $>$ 40 kg). The adult range (5th-95th percentile) of predicted steady state PK parameters for 7 mg dose was defined as maximum concentration observed (C_{max}) ranging from 8.03 to 49.10 micrograms per millilitre (mcg/mL) and area under the curve from time 0 hour to 24 hours (AUC₀₋₂₄) ranging from 184 to 1160 micrograms*hour per millilitre (mcg*h/mL).

Reporting group title	Placebo/Teriflunomide
-----------------------	-----------------------

Reporting group description:

Subjects previously treated with placebo during the double blind (DB) period received 1 teriflunomide tablet 3.5 mg (BW \leq 40 kg) or 7 mg teriflunomide (BW $>$ 40 kg) for first 8 weeks. After 8 weeks if predicted PK parameter was \leq 95th percentile of adult range then subjects received 7 mg teriflunomide (BW \leq 40 kg) and 14 mg teriflunomide (BW $>$ 40 kg) in the open label (OL) period for 96 weeks (i.e., up to Week 192). The adult range (5th-95th percentile) of predicted steady state PK parameters for 7 mg dose was defined as C_{max} ranging from 8.03 to 49.10 mcg/mL and AUC₀₋₂₄ ranging from 184 to 1160 mcg*h/mL.

Reporting group title	Teriflunomide / Teriflunomide
-----------------------	-------------------------------

Reporting group description:

Subjects previously treated with Teriflunomide during DB period continued receiving Teriflunomide in the OL period for additional 96 weeks (i.e., up to Week 192).

Reporting group title	Teriflunomide
-----------------------	---------------

Reporting group description:

Subjects who were not adults when they completed the 192-week OL period were offered an optional additional extension period to receive 1 teriflunomide tablet 14 mg daily (or every other day if BW was $<$ 40 kg) until they became adults or until they were able to switch to the commercial product, whichever came first.

Subject analysis set title	Teriflunomide 7 mg
----------------------------	--------------------

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

Subjects with BW $>$ 40 kg received 1 Teriflunomide tablet, 7 mg orally QD for 8 weeks. After 8 weeks, based on subject's predicted PK parameters, teriflunomide was administered in following manner up to 96 weeks: if predicted PK parameters \leq 95th percentile of adult range after 7 mg QD: 1 tablet of 7 mg daily (for BW up to 40 kg) or 1 tablet of 14 mg daily (for BW $>$ 40 kg); or if predicted PK parameters $>$ 95th percentile of adult range: 1 tablet of 3.5 mg daily (for BW up to 40 kg) or 1 tablet of 7 mg daily (for BW $>$ 40 kg).

Subject analysis set title	Teriflunomide 14 mg
----------------------------	---------------------

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

After 8 weeks, based on subject's predicted PK parameters, teriflunomide was administered in following manner up to 96 weeks: if predicted PK parameters \leq 95th percentile of adult range after 7 mg QD: 1 tablet of 7 mg daily (for BW up to 40 kg) or 1 tablet of 14 mg daily (for BW $>$ 40 kg); or if predicted PK parameters $>$ 95th percentile of adult range: 1 tablet of 3.5 mg daily (for BW up to 40 kg) or 1 tablet of 7 mg daily (for BW $>$ 40 kg).

Subject analysis set title	Placebo / Teriflunomide 7 mg
----------------------------	------------------------------

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

Subjects previously treated with placebo during the DB period received Teriflunomide 7 mg for 96 weeks

in the OL period (i.e., up to Week 192).

Subject analysis set title	Placebo / Teriflunomide 14 mg
Subject analysis set type	Intention-to-treat

Subject analysis set description:

Subjects previously treated with placebo during the DB period received Teriflunomide 14 mg for 96 weeks in OL period (i.e., up to Week 192).

Subject analysis set title	Teriflunomide / Teriflunomide 7 mg
Subject analysis set type	Intention-to-treat

Subject analysis set description:

Subjects previously treated with Teriflunomide 7 mg during DB period continued receiving Teriflunomide 7 mg in the OL period for additional 96 weeks (i.e. up to Week 192).

Subject analysis set title	Teriflunomide / Teriflunomide 14 mg
Subject analysis set type	Intention-to-treat

Subject analysis set description:

Subjects previously treated with Teriflunomide 14 mg during DB period continued receiving Teriflunomide 14 mg in the OL period for additional 96 weeks (i.e. up to Week 192).

Primary: Time to First Confirmed Clinical Relapse

End point title	Time to First Confirmed Clinical Relapse
-----------------	--

End point description:

Time to first clinical relapse:duration (in weeks) between randomisation & first confirmed clinical relapse. Clinical relapses:new/recurrent neurological symptoms not associated with fever/infection,lasted at least 24 hours & accompanied by new objective neurological findings upon neurological examination & documented by standardised, quantified functional system score (FSSs) which included 8 items:rated on different scales:brain stem, cerebellar & cerebral functions rated on scale of 0 to 5;visual, pyramidal, sensory & bowel/bladder rated on scale of 0 to 6 & ambulation on scale of 0 to 12 where higher score in each scale indicated worsened neurological function. Confirmed clinical relapse were reviewed & confirmed by independent Relapse Adjudication Panel. Subject without confirmed clinical relapse;considered as clinical relapse free until end of Week 96. Analysed on Intent-to-treat (ITT) population:all randomised subjects analysed according to treatment allocated by

End point type	Primary
----------------	---------

End point timeframe:

Baseline up to Week 96

End point values	Placebo	Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	57	109		
Units: weeks				
median (full range (min-max))	39.14 (0.1 to 98.0)	75.29 (0.1 to 98.7)		

Statistical analyses

Statistical analysis title	Teriflunomide versus Placebo
----------------------------	------------------------------

Statistical analysis description:

Hazard ratio (HR) was estimated using a Cox proportional-hazards model with factors for treatment group, region, pubertal status, age, and number of relapses in the year prior to randomisation as covariates and with robust variance estimation. Derived from log-rank test with stratification of region and pubertal status.

Comparison groups	Placebo v Teriflunomide
-------------------	-------------------------

Number of subjects included in analysis	166
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2949 ^[1]
Method	Stratified Log-Rank test
Parameter estimate	Hazard ratio (HR)
Point estimate	0.657
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.388
upper limit	1.113

Notes:

[1] - Threshold for significance was < 0.05.

Secondary: Probability of Subjects Who Were Clinical Relapse Free at Weeks 24, 48, 72, 96, 120, 144, 168 and 192

End point title	Probability of Subjects Who Were Clinical Relapse Free at Weeks 24, 48, 72, 96, 120, 144, 168 and 192
-----------------	---

End point description:

Subject was considered free of clinical relapse if the subject had no confirmed clinical relapse before treatment discontinuation/completion in 192 weeks treatment period. Clinical relapses: new/recurrent neurological symptoms not associated with fever/infection, lasted at least 24 hours, & accompanied by new objective neurological findings upon neurological examination & documented by standardised, quantified FSSs which included 8 items: rated on different scales: brain stem, cerebellar & cerebral functions rated on scale of 0 to 5; visual, pyramidal, sensory & bowel/bladder rated on scale of 0 to 6 & ambulation on scale of 0 to 12, where higher score in each scale indicated worsened neurological function. New/recurrent symptoms occurred less than 30 days following onset of relapse were considered part of same relapse. Probability of subjects who were clinical relapse free at specified weeks was estimated by Kaplan-Meier method and reported. Analysed on efficacy population.

End point type	Secondary
----------------	-----------

End point timeframe:

Weeks 24, 48, 72, 96, 120, 144, 168 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: probability of relapse free subjects				
number (confidence interval 95%)				
Week 24	0.750 (0.609 to 0.846)	0.820 (0.730 to 0.883)		
Week 48	0.596 (0.451 to 0.715)	0.700 (0.600 to 0.780)		
Week 72	0.519 (0.377 to 0.644)	0.630 (0.528 to 0.716)		
Week 96	0.442 (0.305 to 0.571)	0.600 (0.497 to 0.688)		
Week 120	0.404 (0.271 to 0.533)	0.570 (0.467 to 0.660)		
Week 144	0.365 (0.238 to 0.494)	0.540 (0.437 to 0.631)		

Week 168	0.365 (0.238 to 0.494)	0.518 (0.416 to 0.611)		
Week 192	0.365 (0.238 to 0.494)	0.518 (0.416 to 0.611)		

Statistical analyses

No statistical analyses for this end point

Secondary: Magnetic Resonance Imaging Assessment: Change From Baseline in Volume of T2 Lesions at Weeks 24, 36, 48, 72, 96, 144 and 192

End point title	Magnetic Resonance Imaging Assessment: Change From Baseline in Volume of T2 Lesions at Weeks 24, 36, 48, 72, 96, 144 and 192
-----------------	--

End point description:

Volume of T2 lesions was measured by MRI scan. Analysis was performed on efficacy population. Here, 'n' = number of subjects analysed for each specified category.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, DB period: Weeks 24, 36, 48, 72 and 96; OL period: Weeks 48, 96, 144 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: millilitres				
arithmetic mean (standard deviation)				
DB Week 24 (n=40,88)	3.0 (± 7.0)	0.5 (± 1.8)		
DB Week 36 (n=19,39)	4.6 (± 11.7)	4.4 (± 21.3)		
DB Week 48 (n=21,68)	0.5 (± 0.6)	-0.2 (± 3.4)		
DB Week 72 (n=16,59)	1.2 (± 1.6)	0.1 (± 1.9)		
DB Week 96 (n=14,51)	0.9 (± 1.4)	0.2 (± 1.9)		
OL Week 48 (n=51,88)	3.0 (± 9.7)	0.6 (± 8.8)		
OL Week 96 (n=37,74)	2.7 (± 8.1)	1.9 (± 4.0)		
OL Week 144 (n=20,23)	4.7 (± 10.7)	0.4 (± 17.1)		
OL Week 192 (n=5,8)	0.4 (± 9.4)	-3.6 (± 30.6)		

Statistical analyses

No statistical analyses for this end point

Secondary: Magnetic Resonance Imaging (MRI) Assessment: Number of New or Enlarged T2 Lesions Per MRI Scan

End point title	Magnetic Resonance Imaging (MRI) Assessment: Number of New or Enlarged T2 Lesions Per MRI Scan
-----------------	--

End point description:

Number of new or enlarged T2 lesions per scan was defined as the total number of new or enlarged T2 lesion that occurred during the 192 weeks treatment period divided by the total number of scans performed during 192 weeks. To account for the different numbers of scans performed among the subjects, a negative binomial regression model with robust variance estimation was used. The model included the total number of new or enlarged T2-lesions as the response variable, with treatment group, region, pubertal status and age as covariates and log-transformed number of scans as an offset variable. Analysis was performed on efficacy population which included all subjects enrolled and treated with at least 1 dose of teriflumomide in OL period analysed according to the treatment group allocated by randomisation in the DB period.

End point type	Secondary
End point timeframe:	
Baseline up to Week 192	

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: lesions per scan				
number (confidence interval 95%)	11.087 (6.586 to 18.662)	5.664 (3.417 to 9.389)		

Statistical analyses

Statistical analysis title	Plac/ Teri versus Teri/ Teri
Comparison groups	Placebo/Teriflunomide v Teriflunomide / Teriflunomide
Number of subjects included in analysis	152
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Relative risk ratio
Point estimate	0.511
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.343
upper limit	0.762

Secondary: Magnetic Resonance Imaging Assessment: Number of T1 Gadolinium (Gd)-Enhancing T1 Lesions Per MRI Scan

End point title	Magnetic Resonance Imaging Assessment: Number of T1 Gadolinium (Gd)-Enhancing T1 Lesions Per MRI Scan
-----------------	---

End point description:

The number of T1 Gd-Enhancing lesions per scan was defined as the total number of lesions that occurred during the 192 weeks treatment period divided by the total number of scans performed during 192 weeks. To account for the different number of scans performed among the subjects, a negative binomial regression model with robust variance estimation was used. The model included the total number of T1-lesions as the response variable, with treatment group, region, pubertal status and age as covariates and log-transformed number of scans as an offset variable. Analysis was performed on efficacy population.

End point type	Secondary
End point timeframe:	
Baseline up to Week 192	

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: lesions per scan				
number (confidence interval 95%)	2.686 (1.263 to 5.712)	1.532 (0.624 to 3.762)		

Statistical analyses

Statistical analysis title	Plac/ Teri versus Teri/ Teri
Comparison groups	Placebo/Teriflunomide v Teriflunomide / Teriflunomide
Number of subjects included in analysis	152
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Relative risk ratio
Point estimate	0.57
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.331
upper limit	0.983

Secondary: Magnetic Resonance Imaging Assessment: Change From Baseline in Volume of T1 Hypointense Lesions

End point title	Magnetic Resonance Imaging Assessment: Change From Baseline in Volume of T1 Hypointense Lesions
End point description:	Volume of T1 hypointense lesions was measured by MRI scan. Analysis was performed on efficacy population. Here, 'n' = number of subjects analysed for each specified category.
End point type	Secondary
End point timeframe:	
Baseline, DB period: Weeks 24, 36, 48, 72 and 96; OL period: Weeks 48, 96, 144 and 192	

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: millilitres				
arithmetic mean (standard deviation)				
DB Week 24 (n=40,88)	0.3 (± 1.3)	0.0 (± 0.7)		
DB Week 36 (n=19,39)	0.8 (± 1.8)	0.2 (± 0.8)		
DB Week 48 (n=20,68)	0.2 (± 0.4)	0.4 (± 2.9)		
DB Week 72 (n=16,58)	0.1 (± 0.7)	0.1 (± 0.6)		
DB Week 96 (n=14,50)	0.1 (± 0.6)	0.1 (± 0.7)		
OL Week 48 (n=51,86)	0.7 (± 1.9)	0.5 (± 1.4)		
OL Week 96 (n=37,73)	1.0 (± 2.0)	0.6 (± 1.9)		
OL Week 144 (n=20,22)	2.0 (± 3.0)	1.9 (± 3.4)		
OL Week 192 (n=5,8)	4.0 (± 5.8)	2.5 (± 5.7)		

Statistical analyses

No statistical analyses for this end point

Secondary: Magnetic Resonance Imaging Assessment: Percentage of Subjects Free of New or Enlarged MRI T2-Lesions

End point title	Magnetic Resonance Imaging Assessment: Percentage of Subjects Free of New or Enlarged MRI T2-Lesions
End point description:	Percentage of subjects who were free of new or enlarged T2 lesions at Weeks 24, 48, 72, 96, 144 and 192 were reported. Analysis was performed on efficacy population.
End point type	Secondary
End point timeframe:	Baseline, Weeks 24, 48, 72, 96, 144 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: percentage of subjects				
number (not applicable)				
Week 24	86.5	86.0		
Week 48	32.7	25.0		
Week 72	15.4	17.0		
Week 96	15.4	16.0		
Week 144	11.5	14.0		
Week 192	7.7	9.0		

Statistical analyses

Secondary: Magnetic Resonance Imaging Assessment: Number of New T1 Hypointense Lesions Per MRI Scan

End point title	Magnetic Resonance Imaging Assessment: Number of New T1 Hypointense Lesions Per MRI Scan
-----------------	--

End point description:

The number of new T1 hypointense lesions were obtained from MRI scans. Analysis was performed on efficacy population.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to Week 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: lesions				
number (not applicable)	1561	1910		

Statistical analyses

Statistical analysis title	Plac/ Teri versus Teri/ Teri
Comparison groups	Placebo/Teriflunomide v Teriflunomide / Teriflunomide
Number of subjects included in analysis	152
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Relative risk ratio
Point estimate	0.498
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.296
upper limit	0.836

Secondary: Magnetic Resonance Imaging Assessment: Percent Change From Baseline in Brain Volume at Weeks 24, 36, 48, 72, 96, 144 and 192

End point title	Magnetic Resonance Imaging Assessment: Percent Change From Baseline in Brain Volume at Weeks 24, 36, 48, 72, 96, 144 and 192
-----------------	--

End point description:

Percent change from baseline in brain volume (assessed using MRI scans of the Brain) at Weeks 24, 36, 48, 72, 96, 144 and 192 was reported. Analysis was performed on efficacy population. Here, 'n' = number of subjects analysed for each specified category.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, DB period: Weeks 24, 36, 48, 72 and 96; OL period: Weeks 48, 96, 144 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: percent change				
arithmetic mean (standard deviation)				
DB Week 24 (n=38,83)	-0.3 (± 0.7)	-0.2 (± 0.7)		
DB Week 36 (n=17,39)	-0.7 (± 0.7)	-0.4 (± 1.0)		
DB Week 48 (n=20,64)	-0.6 (± 1.1)	-0.5 (± 0.9)		
DB Week 72 (n=16,53)	-0.9 (± 1.3)	-0.6 (± 1.1)		
DB Week 96 (n=13,47)	-0.9 (± 1.3)	-0.8 (± 1.1)		
OL Week 48 (n=43,73)	-1.4 (± 1.6)	-1.1 (± 1.3)		
OL Week 96 (n=33,60)	-1.8 (± 1.9)	-1.4 (± 1.6)		
OL Week 144 (n=16,17)	-3.1 (± 2.8)	-2.0 (± 1.7)		
OL Week 192 (n=4,6)	-2.2 (± 1.2)	-3.0 (± 1.9)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Total Number of Correct Substitutions Measured by Symbol Digit Modalities Test (SDMT) at Weeks 24, 48, 72, 96, 120, 144, 168 and 192

End point title	Cognitive Assessment: Change From Baseline in Total Number of Correct Substitutions Measured by Symbol Digit Modalities Test (SDMT) at Weeks 24, 48, 72, 96, 120, 144, 168 and 192
-----------------	--

End point description:

SDMT measures the time to pair abstract symbols with specific numbers. It is a simple substitution task that gives the examinee 90 seconds to pair specific numbers with given geometric figures as a measure for screening cognitive impairment. The SDMT score is the number of correct substitution and ranged from 0 (worst outcome) to 110 (best outcome), where higher score indicated better cognitive function. Analysis was performed on efficacy population. Here, 'n' = number of subjects analysed for each specified category. Here, '99999' is used as a space filler which specifies that the standard deviation was not calculated, since only 1 subject was assessed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, DB period: Weeks 24, 48, 72 and 96; OL period: Weeks 24, 48, 72, 96, 120, 144, 168 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: score on a scale				
arithmetic mean (standard deviation)				
DB Week 24 (n=40,85)	5.1 (± 12.0)	4.6 (± 9.1)		
DB Week 48 (n=20,64)	7.3 (± 14.1)	5.7 (± 10.3)		
DB Week 72 (n=17,58)	6.4 (± 12.9)	5.6 (± 11.5)		
DB Week 96 (n=14,52)	8.8 (± 10.7)	8.1 (± 11.1)		
OL Week 24 (n=51,91)	6.3 (± 13.9)	8.3 (± 12.3)		
OL Week 48 (n=44,88)	6.7 (± 13.3)	7.6 (± 13.0)		
OL Week 72 (41,76)	8.0 (± 15.5)	9.2 (± 13.0)		
OL Week 96 (n=36,73)	7.6 (± 16.7)	8.0 (± 14.8)		
OL Week 120 (n=22,28)	3.7 (± 15.9)	7.2 (± 10.3)		
OL Week 144 (n=18,19)	6.6 (± 14.4)	5.2 (± 13.0)		
OL Week 168 (n=13,15)	3.5 (± 17.9)	1.7 (± 14.0)		
OL Week 192 (n=1,4)	12.0 (± 99999)	-0.3 (± 18.2)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Number of Completed Items Measured by Symbol Digit Modalities Test at Weeks 24, 48, 72, 96, 120, 144, 168 and 192

End point title	Cognitive Assessment: Change From Baseline in Number of Completed Items Measured by Symbol Digit Modalities Test at Weeks 24, 48, 72, 96, 120, 144, 168 and 192
-----------------	---

End point description:

SDMT measures the time to pair abstract symbols with specific numbers. It is a simple substitution task that gives the examinee 90 seconds to pair specific numbers with given geometric figures as a measure for screening cognitive impairment. The SDMT score is the number of completed items and ranged from 0 (worst outcome) to 110 (best outcome), where higher score indicated better cognitive function. Analysis was performed on efficacy population. Here, 'n' = number of subjects analysed for each specified category. Here, '99999' was used as a space filler which specifies that the standard deviation was not calculated, since only 1 subject was assessed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, DB period: Weeks 24, 48, 72 and 96; OL period: Weeks 24, 48, 72, 96, 120, 144, 168 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: score on a scale				
arithmetic mean (standard deviation)				
DB Week 24 (n=40,85)	3.8 (± 11.7)	3.6 (± 9.0)		
DB Week 48 (n=20,64)	6.3 (± 13.8)	4.7 (± 10.3)		

DB Week 72 (n=17,58)	5.1 (± 13.4)	4.5 (± 11.4)		
DB Week 96 (n=14,52)	7.1 (± 11.2)	6.9 (± 11.1)		
OL Week 24 (n=51,91)	4.8 (± 13.5)	7.1 (± 12.4)		
OL Week 48 (n=44,88)	5.5 (± 12.8)	6.4 (± 13.0)		
OL Week 72 (n=41,76)	6.4 (± 15.4)	8.1 (± 12.8)		
OL Week 96 (n=36,73)	6.3 (± 16.6)	7.6 (± 12.5)		
OL Week 120 (n=22,28)	3.7 (± 14.7)	6.3 (± 11.5)		
OL Week 144 (n=18,19)	4.8 (± 15.1)	3.7 (± 13.4)		
OL Week 168 (n=13,15)	2.7 (± 15.5)	-0.3 (± 15.6)		
OL Week 192 (n=1,4)	12.0 (± 99999)	-1.5 (± 18.3)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Trail Making Test- Part A (TMT-A) Test Scores (in Seconds) at Weeks 96 and 192

End point title	Cognitive Assessment: Change From Baseline in Trail Making Test- Part A (TMT-A) Test Scores (in Seconds) at Weeks 96 and 192
-----------------	--

End point description:

'Trail Making Test Part A' is a neuropsychological test of visual attention and task switching. The task requires a subject to 'connect-the-dots' of 25 consecutive numbers (1, 2, 3, etc.) in sequential order on a sheet of paper or computer screen. The goal of the subject is to finish the test as quickly as possible, and the time taken to complete the test used as the primary performance metric (in seconds). This is a timed test and the number of seconds to complete the task is recorded. Maximum time allowed is 300 seconds. A lower score indicated better cognitive function. Analysis was performed on efficacy population. Here, 'number of subjects analysed' = number of subjects evaluable for this endpoint and "n" = number of subjects evaluable for each specified category.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Weeks 96 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	8	21		
Units: seconds				
arithmetic mean (standard deviation)				
Week 96 (n=3,13)	8.4 (± 9.0)	-3.1 (± 19.5)		
Week 192 (n=8,21)	6.3 (± 20.7)	-6.6 (± 17.4)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Trail Making Test B

(TMT-B) Test Scores (in Seconds) at Weeks 96 and 192

End point title	Cognitive Assessment: Change From Baseline in Trail Making Test B (TMT-B) Test Scores (in Seconds) at Weeks 96 and 192
-----------------	--

End point description:

TMT-B is a cognitive test that gives a measure of various aspects of cognitive performance. It is used to measure cognitive fatigue. The test consists of 25 circles containing 13 sequential numbers (1-13) and 12 sequential letters (A-L) positioned. The test evaluates the time (in seconds) to correctly order letters and numbers in alternate order (1, A, 2, B etc.). Maximum time allowed is 300 seconds, where less time/lower score indicates better cognitive function/performance. Analysis was performed on efficacy population. Here, "number of subjects analysed"= subjects evaluable for this endpoint and 'n' = number of subjects evaluable for each specified category.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Weeks 96 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	8	21		
Units: seconds				
arithmetic mean (standard deviation)				
Week 96 (n=3,13)	-19.8 (± 33.7)	-29.5 (± 56.6)		
Week 192 (n=8,21)	-37.0 (± 83.3)	-18.8 (± 37.3)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Brief Visuospatial Memory Test-Revised (BVM-T-R) Scores at Weeks 96 and 192

End point title	Cognitive Assessment: Change From Baseline in Brief Visuospatial Memory Test-Revised (BVM-T-R) Scores at Weeks 96 and 192
-----------------	---

End point description:

The BVM-T consists of three trials in which subjects must recall shapes by drawing figures on a blank page (response booklet) after being given the opportunity to memorise the figures (given in BVM-T-R form) for 10 seconds. BVM-T-R form consists of six figures. Points are awarded based on the accuracy of the drawn figure and by correct placement on the blank page. A minimum of 0 to 12 points/scores are awarded per trial, so a subject can score between 0 and 36 points for all three trials (by adding the points/score from each trial), where higher score indicates better outcome. Analysis was performed on ITT population which included all randomised subjects analysed according to the treatment group allocated by randomization. Here, 'number of subject analysed' = number of subjects evaluable for this endpoint and "n"= subjects evaluable for each specified category.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Weeks 96 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	16	45		
Units: score on a scale				
arithmetic mean (standard deviation)				
Week 96 (n=5,35)	-0.8 (± 9.3)	1.6 (± 5.3)		
Week 192 (n=16,45)	1.0 (± 6.9)	1.2 (± 5.4)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Beery Visual-motor Integration (BVMI) Scores at Weeks 96 and 192

End point title	Cognitive Assessment: Change From Baseline in Beery Visual-motor Integration (BVMI) Scores at Weeks 96 and 192
-----------------	--

End point description:

The Beery VMI is a non-verbal assessment that assessed the extent to which individuals can integrate their visual and motor abilities. The subjects were provided with geometric designs ranging from simple line drawings to more complex figures and were asked to copy the designs. The test consisted of 24 figures. One point was scored for each successful copy of drawings and no scoring was given when the subject failed to copy the drawings properly. Each successful copying of drawings was summed up and the total was scored on a scale ranged from 0 to 24, where higher score indicated better visual construction skills/better visual and motor abilities and lower score indicated poor visual construction skills/poor visual and motor abilities. Analysis was performed on efficacy population. Here, "number of subjects analysed" = subjects evaluable for this endpoint and 'n' = number of subjects evaluable for each specified category.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Weeks 96 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	26	58		
Units: scores on a scale				
arithmetic mean (standard deviation)				
Week 96 (n=10,49)	0.6 (± 2.4)	-0.4 (± 4.8)		
Week 192 (n=26,58)	0.1 (± 5.4)	0.4 (± 2.9)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Wechsler Abbreviated Scale of Intelligence-II (WASI-II) Vocabulary Total Raw Scores at Weeks 96 and 192

End point title	Cognitive Assessment: Change From Baseline in Wechsler Abbreviated Scale of Intelligence-II (WASI-II) Vocabulary Total Raw Scores at Weeks 96 and 192
End point description:	
The WASI-II: Vocabulary test is a quick estimate of an individual's level of intellectual functioning which comprised of 31 total items that require the subject to orally define 3 images and 28 words presented both orally and visually. Items 1 to 3 rated on a score of 0 or 1, items 4 and 5 rated on a score of 0 or 2, items 6 to 31 rated on a scale of 0 to 2. Each item score was summed up to derive the total score which was ranged from 0 (minimum score) to 59 (maximum score), where higher score indicated better level of intellectual functioning/higher level of intelligence. Analysis was performed on efficacy population. Here, "number of subjects analysed" = subjects evaluable for this endpoint. Here, "99999" was used as a space filler which specifies that the standard deviation was not calculated, since only 1 subject was assessed.	
End point type	Secondary
End point timeframe:	
Baseline, Weeks 96 and 192	

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1	1		
Units: scores on a scale				
arithmetic mean (standard deviation)				
Week 96	5.0 (± 99999)	4.0 (± 99999)		
Week 192	3.0 (± 99999)	5.0 (± 99999)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Delis-Kaplan Executive Function System (D-KEFS) Letter Fluency Total Correct Raw Score at Weeks 96 and 192

End point title	Cognitive Assessment: Change From Baseline in Delis-Kaplan Executive Function System (D-KEFS) Letter Fluency Total Correct Raw Score at Weeks 96 and 192
End point description:	
Letter fluency is a condition measured in Delis-Kaplan Executive Function System (D-KEFS). Subjects are asked to name as many words as they can, starting with a specified letter for 60 seconds. The words cannot be names, places, numbers or grammatical variants of previous answers. Repeated answers are not scored as a correct response. There are 3 trials, with 3 different letters. The total number of correct responses is totaled for all 3 trials and a letter fluency score is given. A higher score is considered better. There is no set range as the score depends on how many correct words the subject relays in the given time period. Analysis was performed on efficacy population. Here, "number of subjects analysed" = number of subjects evaluable for this endpoint. Here, 99999 was used as a space filler which specifies that the standard deviation was not calculated, since only 1 subject was assessed.	
End point type	Secondary
End point timeframe:	
Baseline, Weeks 96 and 192	

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1	2		
Units: scores on a scale				
arithmetic mean (standard deviation)				
Week 96	-3.0 (± 99999)	4.0 (± 9.9)		
Week 192	5.0 (± 99999)	-6.5 (± 16.3)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment - Selective Reminding Test (SRT): Change From Baseline in Total Number of Words on Delayed Recall at Weeks 96 and 192

End point title	Cognitive Assessment - Selective Reminding Test (SRT): Change From Baseline in Total Number of Words on Delayed Recall at Weeks 96 and 192
-----------------	--

End point description:

SRT is a test to assess verbal learning and memory. During the administration of the SRT only the examiner and the subject should be in the testing room. A list of twelve words is read aloud by the examiner at a rate of one word per two seconds. The subject is asked to recall all twelve words after a 30 minute delay. Only the words that are missed on the preceding trial are given in the consecutive trial. The total score represents a sum score of total 6 trials, therefore the range is from 0-72. The lower the value the worse the outcome, higher value indicates better recall. Analysis was performed on efficacy population. Here, "number of subjects analysed"= subjects evaluable for this endpoint and "n"= number of subjects evaluable for each specified category. Here, "99999" was used as a space filler and specified that the standard deviation was not calculated, since only 1 subject was assessed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Weeks 96 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1	2		
Units: scores on a scale				
arithmetic mean (standard deviation)				
Week 96 (n=1,2)	1.0 (± 99999)	-0.5 (± 2.1)		
Week 192 (n=1,1)	0.0 (± 99999)	0.0 (± 99999)		

Statistical analyses

No statistical analyses for this end point

Secondary: Cognitive Assessment: Change From Baseline in Delis-Kaplan Executive Function System Category Fluency Total Correct Raw Score at Weeks 96 and 192

End point title	Cognitive Assessment: Change From Baseline in Delis-Kaplan
-----------------	--

End point description:

Category fluency is a condition measured in the D-KEFS. It measures subject's ability to generate words from three different categories (e.g., fruits, vegetables and animals), within a minute for each category. Total score is number of correct words for each category with no points for repetitions or non-words. Score range 0 to unlimited, where 0 = low score, higher score indicates better performance. Analysis was performed on efficacy population. Here, "number of subjects analysed" = number of subjects evaluable for this endpoint. Here, '99999' was used as a space filler and specified that the standard deviation was not calculated, since only 1 subject was assessed.

End point type Secondary

End point timeframe:

Baseline, Weeks 96 and 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1	2		
Units: score on a scale				
arithmetic mean (standard deviation)				
Week 96	6.0 (± 99999)	5.0 (± 5.7)		
Week 192	-2.0 (± 99999)	-13.5 (± 17.7)		

Statistical analyses

No statistical analyses for this end point

Secondary: OL: Pharmacokinetics: Steady-state Trough Concentration (C_{trough}) of TeriflunomideEnd point title OL: Pharmacokinetics: Steady-state Trough Concentration (C_{trough}) of Teriflunomide

End point description:

C_{trough} was defined as the concentration reached by the drug before the next dose is administered. Data for this endpoint was planned to be collected and analysed separately for each dose of teriflunomide. Analysis was performed on PK population which included all randomised subjects exposed to study medication, regardless of the amount of treatment administered who had at least one PK sample taken. Here, 'number of subjects analysed' = subjects with available data for this endpoint.

End point type Secondary

End point timeframe:

Pre-dose at Week 36

End point values	Placebo / Teriflunomide 7 mg	Placebo / Teriflunomide 14 mg	Teriflunomide / Teriflunomide 7 mg	Teriflunomide / Teriflunomide 14 mg
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	12	31	2	79
Units: micrograms per millilitre				
arithmetic mean (standard deviation)	45.8 (± 35.3)	50.4 (± 23.8)	33.7 (± 10.7)	63.6 (± 35.5)

Statistical analyses

No statistical analyses for this end point

Secondary: DB: Pharmacokinetics (PK): Steady-state Trough Concentration (Ctrough) of Teriflunomide

End point title	DB: Pharmacokinetics (PK): Steady-state Trough Concentration (Ctrough) of Teriflunomide
-----------------	---

End point description:

Ctrough was defined as the concentration reached by the drug before the next dose is administered. Data for this endpoint was planned to be collected and analysed separately for each dose of teriflunomide. Analysis was performed on PK population which included all randomised subjects exposed to double-blind study medication and had at least 1 PK sample taken. PK samples for teriflunomide 3.5 mg were collected during the first 8 weeks but all subjects were switched to teriflunomide 7 mg after Week 8. Hence, plasma concentration of teriflunomide 7 mg and 14 mg were reported. Here, 'number of subjects analysed'=subjects with available data for this endpoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose on Week 36

End point values	Teriflunomide 7 mg	Teriflunomide 14 mg		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	10	66		
Units: micrograms per millilitre				
arithmetic mean (standard deviation)	53.1 (± 25.3)	67.8 (± 41.7)		

Statistical analyses

No statistical analyses for this end point

Secondary: OL: Time to First Confirmed Clinical Relapse

End point title	OL: Time to First Confirmed Clinical Relapse
-----------------	--

End point description:

Time to first clinical relapse: duration (in weeks) after enrollment in OL period & first confirmed clinical relapse. Clinical relapses: new/recurrent neurological symptoms not associated with fever/infection, lasted at least 24hours & accompanied by new objective neurological findings upon neurological examination & documented by standardised, quantified FSSs which included 8 items: rated on different scales: brain stem, cerebellar & cerebral functions rated on scale of 0 to 5; visual, pyramidal, sensory & bowel/bladder rated on scale of 0 to 6 & ambulation on scale of 0 to 12 where higher score in each scale indicated worsened neurological function. Confirmed clinical relapse were reviewed & confirmed by independent Relapse Adjudication Panel. Subject without confirmed clinical relapse; considered as clinical relapse free until end of Week 192. Analysed on efficacy population.

End point type	Secondary
----------------	-----------

End point timeframe:
Baseline up to Week 192

End point values	Placebo/Teriflunomide	Teriflunomide / Teriflunomide		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	52	100		
Units: weeks				
median (full range (min-max))	95.86 (0.6 to 176.0)	96.00 (1.0 to 183.4)		

Statistical analyses

Statistical analysis title	Plac/ Teri versus Teri/ Teri
Comparison groups	Placebo/Teriflunomide v Teriflunomide / Teriflunomide
Number of subjects included in analysis	152
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Hazard ratio (HR)
Point estimate	0.693
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.37
upper limit	1.296

Adverse events

Adverse events information

Timeframe for reporting adverse events:

Treatment-emergent adverse events (TEAEs), serious TEAEs and deaths: From first (1st) dose of drug up to 96 weeks (DB period), 1st dose of drug in OL period up to Week 192 (OL period) and 1st dose of drug in extension period up to Week 492 (extension period)

Adverse event reporting additional description:

Safety population included all randomised subjects exposed to study medication, regardless of the amount of treatment administered.

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	27.0
--------------------	------

Reporting groups

Reporting group title	Double-Blind treatment period: Placebo
-----------------------	--

Reporting group description:

Subjects received placebo matching to teriflunomide tablet orally QD for 96 weeks.

Reporting group title	Double-Blind treatment period: Teriflunomide
-----------------------	--

Reporting group description:

Subjects received 1 Teriflunomide tablet, 3.5 mg (in case of BW up to 40 kg) or 7 mg (in case of BW >40 kg) orally QD for 8 weeks. After 8 weeks, based on subjects predicted PK parameters, teriflunomide was administered in following manner up to 96 weeks: if predicted PK parameters ≤95th percentile of adult range after 7 mg QD: 1 tablet of 7 mg daily (for BW up to 40 kg) or 1 tablet of 14 mg daily (for BW >40 kg); or if predicted PK parameters >95th percentile of adult range: 1 tablet of 3.5 mg daily (for BW up to 40 kg) or 1 tablet of 7 mg daily (for BW >40 kg). The adult range (5th-95th percentile) of predicted steady state PK parameters for 7 mg dose was defined as C_{max} ranging from 8.03 to 49.10 mcg/mL and AUC₀₋₂₄ ranging from 184 to 1160 mcg*h/mL.

Reporting group title	Open-Label treatment period: Placebo/Teriflunomide
-----------------------	--

Reporting group description:

Subjects previously treated with placebo during the DB period received 1 teriflunomide tablet 3.5 mg (BW ≤40 kg) or 7 mg teriflunomide (BW >40 kg) for first 8 weeks. After 8 weeks if predicted PK parameter was ≤95th percentile of adult range then subjects received 7 mg teriflunomide (BW ≤40 kg) and 14 mg teriflunomide (BW >40 kg) in the OL period for 96 weeks (i.e., up to Week 192). The adult range (5th-95th percentile) of predicted steady state PK parameters for 7 mg dose was defined as C_{max} ranging from 8.03 to 49.10 mcg/mL and AUC₀₋₂₄ ranging from 184 to 1160 mcg*h/mL.

Reporting group title	Open-Label treatment period: Teriflunomide/Teriflunomide
-----------------------	--

Reporting group description:

Subjects previously treated with Teriflunomide during DB period continued receiving Teriflunomide in the OL period for additional 96 weeks (i.e., up to Week 192).

Reporting group title	Extension period/Teriflunomide
-----------------------	--------------------------------

Reporting group description:

Subjects who were not adults when they completed the 192-week OL period were offered an optional additional extension period to receive 1 teriflunomide tablet 14 mg daily (or every other day if BW <40 kg) until they became adults or until they were able to switch to the commercial product, whichever came first.

Serious adverse events	Double-Blind treatment period: Placebo	Double-Blind treatment period: Teriflunomide	Open-Label treatment period: Placebo/Teriflunomide
Total subjects affected by serious adverse events subjects affected / exposed	6 / 57 (10.53%)	12 / 109 (11.01%)	15 / 52 (28.85%)

number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events	0	0	0
Pregnancy, puerperium and perinatal conditions			
Pregnancy			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gait Disturbance			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Asthenia			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Asthma			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	2 / 52 (3.85%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Depression Suicidal			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicide Attempt			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Adjustment Disorder			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Affective Disorder			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Somatic Symptom Disorder			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Blood Creatine Phosphokinase Increased			
subjects affected / exposed	0 / 57 (0.00%)	2 / 109 (1.83%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	1 / 2	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Alanine Aminotransferase Increased			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	2 / 52 (3.85%)
occurrences causally related to treatment / all	0 / 0	1 / 1	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Transaminases Increased			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Lumbar Vertebral Fracture			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Intentional Overdose			

subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Joint Dislocation			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin Laceration			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Peripheral Nerve Injury			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Congenital, familial and genetic disorders			
Familial Mediterranean Fever			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders			
Cardiomyopathy			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bradycardia			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Dizziness			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Multiple Sclerosis			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypoaesthesia			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Headache			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Epilepsy			
subjects affected / exposed	1 / 57 (1.75%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Syncope			
subjects affected / exposed	1 / 57 (1.75%)	2 / 109 (1.83%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 3	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Multiple Sclerosis Pseudo Relapse			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Multiple Sclerosis Relapse			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Partial Seizures			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Transient Ischaemic Attack			

subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Neutropenia			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ear and labyrinth disorders			
Sudden Hearing Loss			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
Diplopia			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Pancreatic Disorder			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pancreatitis Acute			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Food Poisoning			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Constipation			

subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anal Fistula			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anal Fissure			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Hepatic Function Abnormal			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
Drug Eruption			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Renal Failure			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Pneumonia			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pulmonary Tuberculosis			

subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tonsillitis			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Upper Respiratory Tract Infection			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Coronavirus Infection			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Appendicitis			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acute Sinusitis			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Encephalitis Viral			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bronchitis			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Salpingo-Oophoritis			

subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Complicated Appendicitis			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Open-Label treatment period: Teriflunomide/Teriflu nomide	Extension period/Teriflunomide	
Total subjects affected by serious adverse events			
subjects affected / exposed	14 / 100 (14.00%)	8 / 27 (29.63%)	
number of deaths (all causes)	0	0	
number of deaths resulting from adverse events	0	0	
Pregnancy, puerperium and perinatal conditions			
Pregnancy			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gait Disturbance			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Asthenia			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Respiratory, thoracic and mediastinal disorders			

Asthma			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Psychiatric disorders			
Depression Suicidal			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Suicide Attempt			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Adjustment Disorder			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Affective Disorder			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Somatic Symptom Disorder			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Investigations			
Blood Creatine Phosphokinase Increased			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Alanine Aminotransferase Increased			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Transaminases Increased			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injury, poisoning and procedural complications			
Lumbar Vertebral Fracture			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Intentional Overdose			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Joint Dislocation			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin Laceration			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Peripheral Nerve Injury			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Congenital, familial and genetic disorders			
Familial Mediterranean Fever			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac disorders			
Cardiomyopathy			

subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bradycardia			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
Dizziness			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Multiple Sclerosis			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypoaesthesia			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Headache			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Epilepsy			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Syncope			
subjects affected / exposed	1 / 100 (1.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Multiple Sclerosis Pseudo Relapse			

subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Multiple Sclerosis Relapse			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Partial Seizures			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Transient Ischaemic Attack			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood and lymphatic system disorders			
Neutropenia			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ear and labyrinth disorders			
Sudden Hearing Loss			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Eye disorders			
Diplopia			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			
Pancreatic Disorder			

subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pancreatitis Acute			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	1 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Food Poisoning			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Constipation			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Anal Fistula			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Anal Fissure			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatobiliary disorders			
Hepatic Function Abnormal			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin and subcutaneous tissue disorders			
Drug Eruption			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal and urinary disorders			

Renal Failure			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
Pneumonia			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pulmonary Tuberculosis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Tonsillitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Upper Respiratory Tract Infection			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Coronavirus Infection			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Appendicitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Acute Sinusitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Encephalitis Viral			

subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bronchitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Salpingo-Oophoritis			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Complicated Appendicitis			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 2 %

Non-serious adverse events	Double-Blind treatment period: Placebo	Double-Blind treatment period: Teriflunomide	Open-Label treatment period: Placebo/Teriflunomide
Total subjects affected by non-serious adverse events			
subjects affected / exposed	42 / 57 (73.68%)	89 / 109 (81.65%)	40 / 52 (76.92%)
Vascular disorders			
Orthostatic Hypotension			
subjects affected / exposed	2 / 57 (3.51%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	3	0	0
Hypertension			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	1	0	0
General disorders and administration site conditions			
Influenza Like Illness			
subjects affected / exposed	1 / 57 (1.75%)	3 / 109 (2.75%)	0 / 52 (0.00%)
occurrences (all)	1	3	0
Asthenia			

subjects affected / exposed	0 / 57 (0.00%)	2 / 109 (1.83%)	2 / 52 (3.85%)
occurrences (all)	0	2	3
Fatigue			
subjects affected / exposed	3 / 57 (5.26%)	4 / 109 (3.67%)	4 / 52 (7.69%)
occurrences (all)	5	6	7
Pyrexia			
subjects affected / exposed	2 / 57 (3.51%)	5 / 109 (4.59%)	0 / 52 (0.00%)
occurrences (all)	2	8	0
Non-Cardiac Chest Pain			
subjects affected / exposed	0 / 57 (0.00%)	4 / 109 (3.67%)	0 / 52 (0.00%)
occurrences (all)	0	4	0
Reproductive system and breast disorders			
Dysmenorrhoea			
subjects affected / exposed	3 / 57 (5.26%)	1 / 109 (0.92%)	1 / 52 (1.92%)
occurrences (all)	3	1	4
Menstrual Disorder			
subjects affected / exposed	2 / 57 (3.51%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences (all)	2	1	0
Respiratory, thoracic and mediastinal disorders			
Rhinitis Allergic			
subjects affected / exposed	0 / 57 (0.00%)	3 / 109 (2.75%)	1 / 52 (1.92%)
occurrences (all)	0	3	1
Respiratory Disorder			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	2 / 52 (3.85%)
occurrences (all)	0	0	3
Oropharyngeal Pain			
subjects affected / exposed	2 / 57 (3.51%)	7 / 109 (6.42%)	2 / 52 (3.85%)
occurrences (all)	2	8	2
Nasal Congestion			
subjects affected / exposed	1 / 57 (1.75%)	3 / 109 (2.75%)	0 / 52 (0.00%)
occurrences (all)	1	3	0
Dyspnoea			
subjects affected / exposed	0 / 57 (0.00%)	3 / 109 (2.75%)	1 / 52 (1.92%)
occurrences (all)	0	3	1
Cough			

subjects affected / exposed	3 / 57 (5.26%)	5 / 109 (4.59%)	2 / 52 (3.85%)
occurrences (all)	3	6	2
Epistaxis			
subjects affected / exposed	1 / 57 (1.75%)	2 / 109 (1.83%)	0 / 52 (0.00%)
occurrences (all)	1	4	0
Psychiatric disorders			
Insomnia			
subjects affected / exposed	3 / 57 (5.26%)	1 / 109 (0.92%)	1 / 52 (1.92%)
occurrences (all)	3	1	3
Depression			
subjects affected / exposed	2 / 57 (3.51%)	2 / 109 (1.83%)	4 / 52 (7.69%)
occurrences (all)	2	2	4
Anxiety			
subjects affected / exposed	3 / 57 (5.26%)	4 / 109 (3.67%)	1 / 52 (1.92%)
occurrences (all)	3	6	1
Mixed Anxiety And Depressive Disorder			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Investigations			
Alanine Aminotransferase Increased			
subjects affected / exposed	1 / 57 (1.75%)	2 / 109 (1.83%)	6 / 52 (11.54%)
occurrences (all)	1	3	6
White Blood Cell Count Decreased			
subjects affected / exposed	1 / 57 (1.75%)	5 / 109 (4.59%)	2 / 52 (3.85%)
occurrences (all)	1	8	5
Weight Increased			
subjects affected / exposed	0 / 57 (0.00%)	3 / 109 (2.75%)	0 / 52 (0.00%)
occurrences (all)	0	4	0
Weight Decreased			
subjects affected / exposed	3 / 57 (5.26%)	6 / 109 (5.50%)	2 / 52 (3.85%)
occurrences (all)	3	7	2
Protein Urine Present			
subjects affected / exposed	1 / 57 (1.75%)	4 / 109 (3.67%)	2 / 52 (3.85%)
occurrences (all)	1	5	2
Blood Creatine Phosphokinase Increased			

subjects affected / exposed	0 / 57 (0.00%)	4 / 109 (3.67%)	1 / 52 (1.92%)
occurrences (all)	0	6	1
Neutrophil Count Decreased			
subjects affected / exposed	0 / 57 (0.00%)	3 / 109 (2.75%)	1 / 52 (1.92%)
occurrences (all)	0	5	4
Blood Bilirubin Increased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Blood Albumin Increased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Aspartate Aminotransferase Increased			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences (all)	0	1	0
Blood Uric Acid Decreased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Electrocardiogram St-T Segment Abnormal			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
White Blood Cells Urine Positive			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Neutrophil/Lymphocyte Ratio Decreased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	1 / 52 (1.92%)
occurrences (all)	0	0	1
Neutrophil Percentage Decreased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Monocyte Percentage Increased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Monocyte Count Decreased			

subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	1 / 52 (1.92%)
occurrences (all)	0	2	1
Gamma-Glutamyltransferase Increased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Glomerular Filtration Rate Increased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Haematocrit Decreased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Haemoglobin Decreased			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Injury, poisoning and procedural complications			
Contusion			
subjects affected / exposed	0 / 57 (0.00%)	4 / 109 (3.67%)	1 / 52 (1.92%)
occurrences (all)	0	4	1
Fall			
subjects affected / exposed	4 / 57 (7.02%)	6 / 109 (5.50%)	2 / 52 (3.85%)
occurrences (all)	5	8	3
Ligament Sprain			
subjects affected / exposed	2 / 57 (3.51%)	1 / 109 (0.92%)	1 / 52 (1.92%)
occurrences (all)	2	1	3
Accidental Overdose			
subjects affected / exposed	4 / 57 (7.02%)	5 / 109 (4.59%)	2 / 52 (3.85%)
occurrences (all)	7	5	2
Thermal Burn			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	2 / 52 (3.85%)
occurrences (all)	0	1	2
Hip Fracture			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Cardiac disorders			

Palpitations			
subjects affected / exposed	0 / 57 (0.00%)	3 / 109 (2.75%)	0 / 52 (0.00%)
occurrences (all)	0	3	0
Sinus Tachycardia			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Sinus Bradycardia			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Nervous system disorders			
Paraesthesia			
subjects affected / exposed	2 / 57 (3.51%)	11 / 109 (10.09%)	1 / 52 (1.92%)
occurrences (all)	3	19	1
Hypoaesthesia			
subjects affected / exposed	3 / 57 (5.26%)	6 / 109 (5.50%)	3 / 52 (5.77%)
occurrences (all)	5	7	4
Headache			
subjects affected / exposed	13 / 57 (22.81%)	18 / 109 (16.51%)	7 / 52 (13.46%)
occurrences (all)	18	30	22
Disturbance In Attention			
subjects affected / exposed	2 / 57 (3.51%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences (all)	2	1	0
Dizziness			
subjects affected / exposed	4 / 57 (7.02%)	9 / 109 (8.26%)	6 / 52 (11.54%)
occurrences (all)	5	9	12
Presyncope			
subjects affected / exposed	0 / 57 (0.00%)	3 / 109 (2.75%)	0 / 52 (0.00%)
occurrences (all)	0	3	0
Tremor			
subjects affected / exposed	2 / 57 (3.51%)	1 / 109 (0.92%)	2 / 52 (3.85%)
occurrences (all)	3	1	2
Syncope			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	3 / 52 (5.77%)
occurrences (all)	0	1	3
Blood and lymphatic system disorders			

Anaemia subjects affected / exposed occurrences (all)	1 / 57 (1.75%) 1	3 / 109 (2.75%) 3	0 / 52 (0.00%) 0
Neutropenia subjects affected / exposed occurrences (all)	0 / 57 (0.00%) 0	3 / 109 (2.75%) 5	0 / 52 (0.00%) 0
Lymphopenia subjects affected / exposed occurrences (all)	0 / 57 (0.00%) 0	0 / 109 (0.00%) 0	0 / 52 (0.00%) 0
Ear and labyrinth disorders Ear Pain subjects affected / exposed occurrences (all)	2 / 57 (3.51%) 2	3 / 109 (2.75%) 3	1 / 52 (1.92%) 1
Tinnitus subjects affected / exposed occurrences (all)	1 / 57 (1.75%) 1	3 / 109 (2.75%) 3	1 / 52 (1.92%) 1
Eye disorders Visual Impairment subjects affected / exposed occurrences (all)	2 / 57 (3.51%) 3	0 / 109 (0.00%) 0	2 / 52 (3.85%) 2
Vision Blurred subjects affected / exposed occurrences (all)	2 / 57 (3.51%) 3	2 / 109 (1.83%) 3	0 / 52 (0.00%) 0
Eye Pain subjects affected / exposed occurrences (all)	4 / 57 (7.02%) 8	4 / 109 (3.67%) 4	0 / 52 (0.00%) 0
Gastrointestinal disorders Abdominal Pain subjects affected / exposed occurrences (all)	1 / 57 (1.75%) 1	11 / 109 (10.09%) 13	2 / 52 (3.85%) 2
Abdominal Pain Upper subjects affected / exposed occurrences (all)	2 / 57 (3.51%) 2	6 / 109 (5.50%) 9	0 / 52 (0.00%) 0
Aphthous Ulcer subjects affected / exposed occurrences (all)	0 / 57 (0.00%) 0	2 / 109 (1.83%) 2	2 / 52 (3.85%) 2
Diarrhoea			

subjects affected / exposed occurrences (all)	4 / 57 (7.02%) 4	8 / 109 (7.34%) 9	6 / 52 (11.54%) 8
Constipation subjects affected / exposed occurrences (all)	1 / 57 (1.75%) 1	1 / 109 (0.92%) 1	2 / 52 (3.85%) 2
Mouth Ulceration subjects affected / exposed occurrences (all)	0 / 57 (0.00%) 0	2 / 109 (1.83%) 2	2 / 52 (3.85%) 4
Nausea subjects affected / exposed occurrences (all)	4 / 57 (7.02%) 5	10 / 109 (9.17%) 12	3 / 52 (5.77%) 3
Vomiting subjects affected / exposed occurrences (all)	5 / 57 (8.77%) 5	6 / 109 (5.50%) 7	2 / 52 (3.85%) 2
Gastrointestinal Disorder subjects affected / exposed occurrences (all)	0 / 57 (0.00%) 0	0 / 109 (0.00%) 0	0 / 52 (0.00%) 0
Toothache subjects affected / exposed occurrences (all)	1 / 57 (1.75%) 1	2 / 109 (1.83%) 2	0 / 52 (0.00%) 0
Skin and subcutaneous tissue disorders			
Alopecia subjects affected / exposed occurrences (all)	7 / 57 (12.28%) 7	24 / 109 (22.02%) 26	9 / 52 (17.31%) 10
Dry Skin subjects affected / exposed occurrences (all)	0 / 57 (0.00%) 0	0 / 109 (0.00%) 0	2 / 52 (3.85%) 2
Acne subjects affected / exposed occurrences (all)	4 / 57 (7.02%) 4	5 / 109 (4.59%) 5	0 / 52 (0.00%) 0
Eczema subjects affected / exposed occurrences (all)	1 / 57 (1.75%) 1	1 / 109 (0.92%) 1	0 / 52 (0.00%) 0
Rash subjects affected / exposed occurrences (all)	2 / 57 (3.51%) 2	4 / 109 (3.67%) 4	2 / 52 (3.85%) 2

Renal and urinary disorders			
Pollakiuria			
subjects affected / exposed	0 / 57 (0.00%)	2 / 109 (1.83%)	2 / 52 (3.85%)
occurrences (all)	0	2	2
Micturition Urgency			
subjects affected / exposed	1 / 57 (1.75%)	2 / 109 (1.83%)	5 / 52 (9.62%)
occurrences (all)	1	2	6
Endocrine disorders			
Autoimmune Thyroiditis			
subjects affected / exposed	2 / 57 (3.51%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences (all)	2	1	0
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	3 / 57 (5.26%)	5 / 109 (4.59%)	2 / 52 (3.85%)
occurrences (all)	7	5	2
Muscle Spasms			
subjects affected / exposed	2 / 57 (3.51%)	2 / 109 (1.83%)	2 / 52 (3.85%)
occurrences (all)	2	2	2
Back Pain			
subjects affected / exposed	2 / 57 (3.51%)	5 / 109 (4.59%)	2 / 52 (3.85%)
occurrences (all)	3	6	3
Muscular Weakness			
subjects affected / exposed	2 / 57 (3.51%)	1 / 109 (0.92%)	3 / 52 (5.77%)
occurrences (all)	2	1	3
Pain In Extremity			
subjects affected / exposed	2 / 57 (3.51%)	3 / 109 (2.75%)	0 / 52 (0.00%)
occurrences (all)	2	3	0
Infections and infestations			
Acute Sinusitis			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	2 / 52 (3.85%)
occurrences (all)	0	0	2
Bronchitis			
subjects affected / exposed	1 / 57 (1.75%)	5 / 109 (4.59%)	2 / 52 (3.85%)
occurrences (all)	1	5	2
Cystitis			

subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	2 / 52 (3.85%)
occurrences (all)	0	2	2
Gastroenteritis			
subjects affected / exposed	3 / 57 (5.26%)	3 / 109 (2.75%)	2 / 52 (3.85%)
occurrences (all)	3	3	2
Influenza			
subjects affected / exposed	4 / 57 (7.02%)	10 / 109 (9.17%)	5 / 52 (9.62%)
occurrences (all)	4	16	6
Nasopharyngitis			
subjects affected / exposed	5 / 57 (8.77%)	28 / 109 (25.69%)	8 / 52 (15.38%)
occurrences (all)	6	50	13
Upper Respiratory Tract Infection			
subjects affected / exposed	6 / 57 (10.53%)	24 / 109 (22.02%)	7 / 52 (13.46%)
occurrences (all)	23	66	32
Tracheobronchitis			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	2 / 52 (3.85%)
occurrences (all)	1	0	2
Tracheitis			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	2 / 52 (3.85%)
occurrences (all)	0	0	2
Tonsillitis			
subjects affected / exposed	2 / 57 (3.51%)	4 / 109 (3.67%)	2 / 52 (3.85%)
occurrences (all)	2	4	4
Tinea Versicolour			
subjects affected / exposed	2 / 57 (3.51%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences (all)	2	1	0
Sinusitis			
subjects affected / exposed	2 / 57 (3.51%)	4 / 109 (3.67%)	2 / 52 (3.85%)
occurrences (all)	2	4	3
Rhinitis			
subjects affected / exposed	2 / 57 (3.51%)	4 / 109 (3.67%)	2 / 52 (3.85%)
occurrences (all)	3	8	4
Respiratory Tract Infection Viral			
subjects affected / exposed	0 / 57 (0.00%)	4 / 109 (3.67%)	0 / 52 (0.00%)
occurrences (all)	0	7	0
Respiratory Tract Infection			

subjects affected / exposed	1 / 57 (1.75%)	1 / 109 (0.92%)	1 / 52 (1.92%)
occurrences (all)	1	1	1
Pharyngitis			
subjects affected / exposed	1 / 57 (1.75%)	7 / 109 (6.42%)	1 / 52 (1.92%)
occurrences (all)	1	8	2
Urinary Tract Infection			
subjects affected / exposed	1 / 57 (1.75%)	3 / 109 (2.75%)	2 / 52 (3.85%)
occurrences (all)	1	3	2
Pulpitis Dental			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Paronychia			
subjects affected / exposed	0 / 57 (0.00%)	1 / 109 (0.92%)	0 / 52 (0.00%)
occurrences (all)	0	1	0
Covid-19			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Root Canal Infection			
subjects affected / exposed	0 / 57 (0.00%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	0	0	0
Metabolism and nutrition disorders			
Decreased Appetite			
subjects affected / exposed	0 / 57 (0.00%)	2 / 109 (1.83%)	0 / 52 (0.00%)
occurrences (all)	0	3	0
Vitamin D Deficiency			
subjects affected / exposed	2 / 57 (3.51%)	3 / 109 (2.75%)	1 / 52 (1.92%)
occurrences (all)	2	4	1
Iron Deficiency			
subjects affected / exposed	1 / 57 (1.75%)	0 / 109 (0.00%)	0 / 52 (0.00%)
occurrences (all)	1	0	0

Non-serious adverse events	Open-Label treatment period: Teriflunomide/Teriflu nomide	Extension period/Teriflunomide	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	72 / 100 (72.00%)	18 / 27 (66.67%)	
Vascular disorders			

Orthostatic Hypotension subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	0 / 27 (0.00%) 0	
Hypertension subjects affected / exposed occurrences (all)	2 / 100 (2.00%) 5	1 / 27 (3.70%) 1	
General disorders and administration site conditions			
Influenza Like Illness subjects affected / exposed occurrences (all)	2 / 100 (2.00%) 2	0 / 27 (0.00%) 0	
Asthenia subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	0 / 27 (0.00%) 0	
Fatigue subjects affected / exposed occurrences (all)	4 / 100 (4.00%) 4	0 / 27 (0.00%) 0	
Pyrexia subjects affected / exposed occurrences (all)	3 / 100 (3.00%) 3	2 / 27 (7.41%) 2	
Non-Cardiac Chest Pain subjects affected / exposed occurrences (all)	2 / 100 (2.00%) 2	0 / 27 (0.00%) 0	
Reproductive system and breast disorders			
Dysmenorrhoea subjects affected / exposed occurrences (all)	2 / 100 (2.00%) 5	1 / 27 (3.70%) 1	
Menstrual Disorder subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	0 / 27 (0.00%) 0	
Respiratory, thoracic and mediastinal disorders			
Rhinitis Allergic subjects affected / exposed occurrences (all)	0 / 100 (0.00%) 0	0 / 27 (0.00%) 0	
Respiratory Disorder			

subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Oropharyngeal Pain			
subjects affected / exposed	7 / 100 (7.00%)	1 / 27 (3.70%)	
occurrences (all)	9	1	
Nasal Congestion			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Dyspnoea			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Cough			
subjects affected / exposed	1 / 100 (1.00%)	1 / 27 (3.70%)	
occurrences (all)	1	1	
Epistaxis			
subjects affected / exposed	1 / 100 (1.00%)	1 / 27 (3.70%)	
occurrences (all)	3	1	
Psychiatric disorders			
Insomnia			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Depression			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences (all)	2	0	
Anxiety			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences (all)	3	0	
Mixed Anxiety And Depressive Disorder			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Investigations			
Alanine Aminotransferase Increased			
subjects affected / exposed	3 / 100 (3.00%)	2 / 27 (7.41%)	
occurrences (all)	3	2	
White Blood Cell Count Decreased			

subjects affected / exposed	5 / 100 (5.00%)	3 / 27 (11.11%)
occurrences (all)	9	7
Weight Increased		
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)
occurrences (all)	3	0
Weight Decreased		
subjects affected / exposed	3 / 100 (3.00%)	0 / 27 (0.00%)
occurrences (all)	5	0
Protein Urine Present		
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)
occurrences (all)	1	0
Blood Creatine Phosphokinase Increased		
subjects affected / exposed	0 / 100 (0.00%)	2 / 27 (7.41%)
occurrences (all)	0	2
Neutrophil Count Decreased		
subjects affected / exposed	2 / 100 (2.00%)	3 / 27 (11.11%)
occurrences (all)	6	5
Blood Bilirubin Increased		
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)
occurrences (all)	0	1
Blood Albumin Increased		
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)
occurrences (all)	0	1
Aspartate Aminotransferase Increased		
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)
occurrences (all)	0	1
Blood Uric Acid Decreased		
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)
occurrences (all)	0	1
Electrocardiogram St-T Segment Abnormal		
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)
occurrences (all)	0	1
White Blood Cells Urine Positive		

subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Neutrophil/Lymphocyte Ratio Decreased			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Neutrophil Percentage Decreased			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Monocyte Percentage Increased			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Monocyte Count Decreased			
subjects affected / exposed	3 / 100 (3.00%)	0 / 27 (0.00%)	
occurrences (all)	4	0	
Gamma-Glutamyltransferase Increased			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Glomerular Filtration Rate Increased			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Haematocrit Decreased			
subjects affected / exposed	1 / 100 (1.00%)	1 / 27 (3.70%)	
occurrences (all)	1	1	
Haemoglobin Decreased			
subjects affected / exposed	1 / 100 (1.00%)	1 / 27 (3.70%)	
occurrences (all)	1	1	
Injury, poisoning and procedural complications			
Contusion			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Fall			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Ligament Sprain			

subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences (all)	2	0	
Accidental Overdose			
subjects affected / exposed	10 / 100 (10.00%)	3 / 27 (11.11%)	
occurrences (all)	12	3	
Thermal Burn			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Hip Fracture			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Cardiac disorders			
Palpitations			
subjects affected / exposed	2 / 100 (2.00%)	1 / 27 (3.70%)	
occurrences (all)	2	1	
Sinus Tachycardia			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Sinus Bradycardia			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Nervous system disorders			
Paraesthesia			
subjects affected / exposed	4 / 100 (4.00%)	0 / 27 (0.00%)	
occurrences (all)	7	0	
Hypoaesthesia			
subjects affected / exposed	3 / 100 (3.00%)	1 / 27 (3.70%)	
occurrences (all)	4	1	
Headache			
subjects affected / exposed	14 / 100 (14.00%)	1 / 27 (3.70%)	
occurrences (all)	28	1	
Disturbance In Attention			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Dizziness			

subjects affected / exposed occurrences (all)	2 / 100 (2.00%) 3	0 / 27 (0.00%) 0	
Presyncope subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	0 / 27 (0.00%) 0	
Tremor subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	1 / 27 (3.70%) 1	
Syncope subjects affected / exposed occurrences (all)	0 / 100 (0.00%) 0	0 / 27 (0.00%) 0	
Blood and lymphatic system disorders Anaemia subjects affected / exposed occurrences (all)	6 / 100 (6.00%) 6	0 / 27 (0.00%) 0	
Neutropenia subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	0 / 27 (0.00%) 0	
Lymphopenia subjects affected / exposed occurrences (all)	0 / 100 (0.00%) 0	1 / 27 (3.70%) 1	
Ear and labyrinth disorders Ear Pain subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	0 / 27 (0.00%) 0	
Tinnitus subjects affected / exposed occurrences (all)	0 / 100 (0.00%) 0	0 / 27 (0.00%) 0	
Eye disorders Visual Impairment subjects affected / exposed occurrences (all)	1 / 100 (1.00%) 1	0 / 27 (0.00%) 0	
Vision Blurred subjects affected / exposed occurrences (all)	3 / 100 (3.00%) 3	0 / 27 (0.00%) 0	
Eye Pain			

subjects affected / exposed occurrences (all)	3 / 100 (3.00%) 3	0 / 27 (0.00%) 0	
Gastrointestinal disorders			
Abdominal Pain			
subjects affected / exposed	8 / 100 (8.00%)	0 / 27 (0.00%)	
occurrences (all)	11	0	
Abdominal Pain Upper			
subjects affected / exposed	3 / 100 (3.00%)	0 / 27 (0.00%)	
occurrences (all)	4	0	
Aphthous Ulcer			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Diarrhoea			
subjects affected / exposed	7 / 100 (7.00%)	1 / 27 (3.70%)	
occurrences (all)	9	1	
Constipation			
subjects affected / exposed	3 / 100 (3.00%)	0 / 27 (0.00%)	
occurrences (all)	3	0	
Mouth Ulceration			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences (all)	3	0	
Nausea			
subjects affected / exposed	4 / 100 (4.00%)	0 / 27 (0.00%)	
occurrences (all)	4	0	
Vomiting			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences (all)	2	0	
Gastrointestinal Disorder			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Toothache			
subjects affected / exposed	2 / 100 (2.00%)	1 / 27 (3.70%)	
occurrences (all)	3	1	
Skin and subcutaneous tissue disorders			
Alopecia			

subjects affected / exposed	10 / 100 (10.00%)	1 / 27 (3.70%)	
occurrences (all)	10	1	
Dry Skin			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Acne			
subjects affected / exposed	4 / 100 (4.00%)	0 / 27 (0.00%)	
occurrences (all)	5	0	
Eczema			
subjects affected / exposed	3 / 100 (3.00%)	0 / 27 (0.00%)	
occurrences (all)	4	0	
Rash			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences (all)	2	0	
Renal and urinary disorders			
Pollakiuria			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Micturition Urgency			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Endocrine disorders			
Autoimmune Thyroiditis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Muscle Spasms			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Back Pain			
subjects affected / exposed	3 / 100 (3.00%)	0 / 27 (0.00%)	
occurrences (all)	5	0	
Muscular Weakness			

subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Pain In Extremity			
subjects affected / exposed	3 / 100 (3.00%)	0 / 27 (0.00%)	
occurrences (all)	4	0	
Infections and infestations			
Acute Sinusitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Bronchitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Cystitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Gastroenteritis			
subjects affected / exposed	5 / 100 (5.00%)	1 / 27 (3.70%)	
occurrences (all)	6	1	
Influenza			
subjects affected / exposed	5 / 100 (5.00%)	2 / 27 (7.41%)	
occurrences (all)	6	3	
Nasopharyngitis			
subjects affected / exposed	20 / 100 (20.00%)	4 / 27 (14.81%)	
occurrences (all)	37	4	
Upper Respiratory Tract Infection			
subjects affected / exposed	22 / 100 (22.00%)	6 / 27 (22.22%)	
occurrences (all)	59	12	
Tracheobronchitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Tracheitis			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Tonsillitis			
subjects affected / exposed	4 / 100 (4.00%)	0 / 27 (0.00%)	
occurrences (all)	5	0	

Tinea Versicolour			
subjects affected / exposed	0 / 100 (0.00%)	0 / 27 (0.00%)	
occurrences (all)	0	0	
Sinusitis			
subjects affected / exposed	2 / 100 (2.00%)	0 / 27 (0.00%)	
occurrences (all)	2	0	
Rhinitis			
subjects affected / exposed	6 / 100 (6.00%)	1 / 27 (3.70%)	
occurrences (all)	7	1	
Respiratory Tract Infection Viral			
subjects affected / exposed	1 / 100 (1.00%)	0 / 27 (0.00%)	
occurrences (all)	1	0	
Respiratory Tract Infection			
subjects affected / exposed	5 / 100 (5.00%)	1 / 27 (3.70%)	
occurrences (all)	5	1	
Pharyngitis			
subjects affected / exposed	4 / 100 (4.00%)	1 / 27 (3.70%)	
occurrences (all)	8	1	
Urinary Tract Infection			
subjects affected / exposed	5 / 100 (5.00%)	0 / 27 (0.00%)	
occurrences (all)	5	0	
Pulpitis Dental			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Paronychia			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Covid-19			
subjects affected / exposed	0 / 100 (0.00%)	3 / 27 (11.11%)	
occurrences (all)	0	3	
Root Canal Infection			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	
Metabolism and nutrition disorders			
Decreased Appetite			

subjects affected / exposed	3 / 100 (3.00%)	1 / 27 (3.70%)	
occurrences (all)	3	1	
Vitamin D Deficiency			
subjects affected / exposed	4 / 100 (4.00%)	0 / 27 (0.00%)	
occurrences (all)	4	0	
Iron Deficiency			
subjects affected / exposed	0 / 100 (0.00%)	1 / 27 (3.70%)	
occurrences (all)	0	1	

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
18 December 2013	Following changes were made: 1 MRI added at Weeks 24, 48 and 96 to strengthen MRI data; 1 MRI added at Week 36 in case of at least 5 new/enlarged T2 lesions; change in volume of T2 lesions, change in volume of T1 hypointense lesions, number of new T1 hypointense lesions, and brain atrophy were added and grouped as MRI endpoints; addition of immunoglobulin (IgG, IgM, and IgA) measurements at Baseline and every 24 weeks, aimed to provide additional information on a potential, although not expected, effect of teriflunomide on the immune system in this age group; poisson regression model changed in negative binomial model for the analysis of the number of new or enlarged T2-lesions and the number of T1 Gd-enhancing lesions per MRI scan. Reduced the impact of potential outliers, ordinal logistic regression model including treatment group, region, pubertal status, and age were also used to analyse these endpoints. The Poisson regression model with a robust error variance was to be used in sensitivity analyses.
26 June 2014	Following changes were made: Extension of the open-label period up to 192 weeks after randomisation; one MRI timepoint added at Week 72; addition of a criterion for switch into open-label period taking this additional MRI scan into account as follows: at least 5 new/enlarged T2 lesions on each of the 2 consecutive MRI scans of Week 48 and Week 72; change in exclusion criteria: the required minimum washout period duration for previous multiple sclerosis (MS) treatment was modified; addition of exclusion criteria: previous treatment with alemtuzumab; addition of a note about the content of teriflunomide tablets (lactose) and that therefore, investigators were to consider whether history of lactose intolerance could affect treatment tolerability; addition of a note about contraception and the association of local additional requirements (United Kingdom) to be followed, e.g., spermicidal foam/gel/film/cream/suppository, as per medicines and healthcare products regulatory agency rules; addition of endpoints: proportion of subjects free of new or enlarged MRI T2-lesions at Weeks 48 and 96. It was specified that the main MRI endpoints were the number of new/newly enlarged T2 lesions and the number of T1 Gd-enhancing T1 lesions; proportion of disease-free subjects as an exploratory endpoint; addition of text to introduce the rationale for placebo-controlled design; addition of endocrine function evaluation consisting of measurement of thyroid stimulating hormone (TSH) every 24 weeks and at end of treatment (EOT); addition of text to specify that local anesthetic was to be offered for blood draws to minimise pain and discomfort; correction of text: Pre-defined adverse events and laboratory abnormalities for specific reporting to restore the instructions that were included in the initial protocol but had been inadvertently taken out in Amended protocol 1.
10 December 2020	Following changes were done: Added specifications about weight and electrocardiogram during remote visits and reference to appendix G. Added text in case of emergency and reference to appendix G. Added IMP return specification and reference to appendix G. Added text in case of emergency and reference to appendix G. Added text in case of missing assessment and reference to Appendix G. Added cross references to sections. Added text on permanently discontinuation for subjects close to the end of treatment may be unable to continue their final treatment as scheduled. Added monitoring details, techniques, and source data verification. Added summaries of the 3 memos to detail the procedures set up in case of emergency.

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported