

**Clinical trial results:****A Double-Blind Efficacy and Safety Study of the Phosphodiesterase Type 5 Inhibitor Tadalafil in Pediatric Patients with Pulmonary Arterial Hypertension****Summary**

EudraCT number	2012-002354-23
Trial protocol	GB DE BE IT AT NL PL ES RO FR
Global end of trial date	

**Results information**

Result version number	v2
This version publication date	15 September 2021
First version publication date	22 March 2020
Version creation reason	

**Trial information****Trial identification**

Sponsor protocol code	H6D-MC-LVHV
-----------------------	-------------

**Additional study identifiers**

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT01824290
WHO universal trial number (UTN)	-
Other trial identifiers	Trial Number: 10609

Notes:

**Sponsors**

Sponsor organisation name	Eli Lilly and Company
Sponsor organisation address	Lilly Corporate Center, Indianapolis, IN, United States, 46285
Public contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877CTLilly,
Scientific contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 8772854559,

Notes:

**Paediatric regulatory details**

Is trial part of an agreed paediatric investigation plan (PIP)	Yes
EMA paediatric investigation plan number(s)	EMA-000452-PIP02-10
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Interim
Date of interim/final analysis	18 March 2019
Is this the analysis of the primary completion data?	Yes
Primary completion date	18 March 2019
Global end of trial reached?	No

Notes:

## General information about the trial

Main objective of the trial:

The main purpose of this study is to evaluate the safety and efficacy of tadalafil in pediatric participants with pulmonary arterial hypertension. Participants will receive study treatment for 6 months in the double-blind period (Period 1), and then will be eligible to enroll into an open-label 2 year extension period (Period 2) during which participants will receive tadalafil.

Protection of trial subjects:

This study was conducted in accordance with International Conference on Harmonization (ICH) Good Clinical Practice, and the principles of the Declaration of Helsinki, in addition to following the laws and regulations of the country or countries in which a study is conducted.

Background therapy:

Some participants received endothelin receptor agonist (ERA) background therapy (bosentan or ambrisentan).

Evidence for comparator: -

Actual start date of recruitment	05 February 2014
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Japan: 2
Country: Number of subjects enrolled	Turkey: 3
Country: Number of subjects enrolled	Brazil: 15
Country: Number of subjects enrolled	Mexico: 5
Country: Number of subjects enrolled	Israel: 6
Country: Number of subjects enrolled	France: 2
Country: Number of subjects enrolled	Germany: 1
Country: Number of subjects enrolled	Poland: 1
Worldwide total number of subjects	35
EEA total number of subjects	4

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0

Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	13
Adolescents (12-17 years)	22
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

Participants who experienced clinical worsening during double-blind period were considered as completers.

### Pre-assignment

Screening details:

Per protocol and statistical analysis plan (SAP), the primary and secondary analysis from period 1 were performed to compare all tadalafil participants together versus all placebo participants together.

### Period 1

Period 1 title	Double Blind
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Placebo

Arm description:

Period 1: Participants received placebo orally by tablets once a day.

Arm type	Placebo
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants received placebo orally by tablets once a day.

<b>Arm title</b>	Tadalafil
------------------	-----------

Arm description:

Period 1: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered tadalafil orally by tablets once a day.

Arm type	Experimental
Investigational medicinal product name	Tadalafil
Investigational medicinal product code	
Other name	LY450190
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants received tadalafil orally by tablets once a day.

Number of subjects in period 1	Placebo	Tadalafil
Started	18	17
Received at least one dose of study drug	18	17
Received 20 mg	0 <sup>[1]</sup>	4 <sup>[2]</sup>
Received 40 mg	0 <sup>[3]</sup>	13 <sup>[4]</sup>
Completed	16	16
Not completed	2	1
Parent/Caregiver Decision	1	1
Entry Criteria Not Met	1	-

Notes:

[1] - The number of subjects at this milestone seems inconsistent with the number of subjects in the arm. It is expected that the number of subjects will be greater than, or equal to the number that completed, minus those who left.

Justification: The number of participants in the milestones represent participants who received drug in each drug category (20 mg tadalafil, or 40 mg tadalafil).

[2] - The number of subjects at this milestone seems inconsistent with the number of subjects in the arm. It is expected that the number of subjects will be greater than, or equal to the number that completed, minus those who left.

Justification: The number of participants in the milestones represent participants who received drug in each drug category (20 mg tadalafil, or 40 mg tadalafil).

[3] - The number of subjects at this milestone seems inconsistent with the number of subjects in the arm. It is expected that the number of subjects will be greater than, or equal to the number that completed, minus those who left.

Justification: The number of participants in the milestones represent participants who received drug in each drug category (20 mg tadalafil, or 40 mg tadalafil).

[4] - The number of subjects at this milestone seems inconsistent with the number of subjects in the arm. It is expected that the number of subjects will be greater than, or equal to the number that completed, minus those who left.

Justification: The number of participants in the milestones represent participants who received drug in each drug category (20 mg tadalafil, or 40 mg tadalafil).

## Period 2

Period 2 title	Open-Label Treatment
Is this the baseline period?	No
Allocation method	Non-randomised - controlled
Blinding used	Not blinded

## Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Tadalafil/Tadalafil

Arm description:

Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered orally by tablets once a day.

Participants had received tadalafil during period 1.

Arm type	Experimental
Investigational medicinal product name	Tadalafil
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants received tadalafil orally by tablets once a day.

<b>Arm title</b>	Placebo/Tadalafil
Arm description:	
Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered orally by tablets once a day.	
Participants had received placebo during period 1.	
Arm type	Experimental
Investigational medicinal product name	Tadalafil
Investigational medicinal product code	
Other name	LY450190
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants received tadalafil orally by tablets once a day.

<b>Number of subjects in period 2</b>	Tadalafil/Tadalafil	Placebo/Tadalafil
Started	16	16
Received 20 mg	3 <sup>[5]</sup>	3 <sup>[6]</sup>
Received 40 mg	13	13
Completed	13	13
Not completed	3	3
Consent withdrawn by subject	1	2
Adverse Event	1	-
Parent caregiver Decision	1	1

Notes:

[5] - The number of subjects at this milestone seems inconsistent with the number of subjects in the arm. It is expected that the number of subjects will be greater than, or equal to the number that completed, minus those who left.

Justification: The number of participants in the milestones represent participants who received drug in each drug category (20 mg tadalafil, or 40 mg tadalafil).

[6] - The number of subjects at this milestone seems inconsistent with the number of subjects in the arm. It is expected that the number of subjects will be greater than, or equal to the number that completed, minus those who left.

Justification: The number of participants in the milestones represent participants who received drug in each drug category (20 mg tadalafil, or 40 mg tadalafil).

## Baseline characteristics

### Reporting groups

Reporting group title	Placebo
Reporting group description:	
Period 1: Participants received placebo orally by tablets once a day.	
Reporting group title	Tadalafil
Reporting group description:	
Period 1: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered tadalafil orally by tablets once a day.	

Reporting group values	Placebo	Tadalafil	Total
Number of subjects	18	17	35
Age categorical			
Units: Subjects			

Age continuous			
All participants who received at least one dose. Per protocol and statistical analysis plan (SAP), the primary and secondary analysis were performed to compare all tadalafil participants together versus all placebo participants together.			
Units: years			
arithmetic mean	12.8	14.1	
standard deviation	± 3.39	± 3.49	-
Gender categorical			
Units: Subjects			
Female	9	10	19
Male	9	7	16
Ethnicity (NIH/OMB)			
Units: Subjects			
Hispanic or Latino	7	8	15
Not Hispanic or Latino	6	4	10
Unknown or Not Reported	5	5	10
Race (NIH/OMB)			
Units: Subjects			
American Indian or Alaska Native	1	3	4
Asian	1	1	2
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	2	1	3
White	14	12	26
More than one race	0	0	0
Unknown or Not Reported	0	0	0
Region of Enrollment			
Units: Subjects			
Japan	1	1	2
Turkey	2	1	3
Brazil	9	6	15
Mexico	1	4	5
Israel	2	4	6
France	1	1	2

Germany	1	0	1
Poland	1	0	1

6 Minute Walk Distance			
Units: Meters			
arithmetic mean	476.7	485.8	
standard deviation	± 105.11	± 160.231	-



## End points

### End points reporting groups

Reporting group title	Placebo
-----------------------	---------

Reporting group description:

Period 1: Participants received placebo orally by tablets once a day.

Reporting group title	Tadalafil
-----------------------	-----------

Reporting group description:

Period 1: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered tadalafil orally by tablets once a day.

Reporting group title	Tadalafil/Tadalafil
-----------------------	---------------------

Reporting group description:

Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered orally by tablets once a day.

Participants had received tadalafil during period 1.

Reporting group title	Placebo/Tadalafil
-----------------------	-------------------

Reporting group description:

Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered orally by tablets once a day.

Participants had received placebo during period 1.

Subject analysis set title	20 mg Tadalafil
----------------------------	-----------------

Subject analysis set type	Per protocol
---------------------------	--------------

Subject analysis set description:

Period 1: 20 mg tadalafil administered orally by tablets once a day with concomitant endothelin receptor antagonist (ERA).

Subject analysis set title	40 mg Tadalafil
----------------------------	-----------------

Subject analysis set type	Per protocol
---------------------------	--------------

Subject analysis set description:

Period 1: 40 mg tadalafil administered orally by tablets once a day with concomitant endothelin receptor antagonist (ERA).

Subject analysis set title	Placebo/Tadalafil - Open-Label Treatment
----------------------------	--

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

Period 1: Participants received placebo orally by tablets once a day.

Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered orally by tablets once a day.

Subject analysis set title	Tadalafil/Tadalafil - Open-Label Treatment
----------------------------	--

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

Period 1: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered orally by tablets once a day.

Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered orally by tablets once a day.

### Primary: Period 1: Change from Baseline to Week 24 in a 6 Minute Walk (MW) Distance in Meters

End point title	Period 1: Change from Baseline to Week 24 in a 6 Minute Walk (MW) Distance in Meters
-----------------	--

End point description:

6MWD in meters assessed in a subset of participants who are  $\geq 6$  to  $< 18$  years of age who are developmentally capable of performing a 6MW test. Change from baseline was derived using mixed model repeated measures (MMRM) with terms for treatment group, visit, baseline 6MWD, and

treatment-by-visit interaction.

Analysis Population Description: All participants who received at least one dose of study drug who were > = 6 to < 18 years of age and were capable of performing a 6MW test.

End point type	Primary
End point timeframe:	
Baseline, Week 24	

End point values	Placebo	Tadalafil		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	15	15		
Units: Meters				
least squares mean (standard error)	36.60 (± 20.776)	60.48 (± 20.410)		

## Statistical analyses

Statistical analysis title	6 Minute Walk (MW) Distance in Meters
Comparison groups	Tadalafil v Placebo
Number of subjects included in analysis	30
Analysis specification	Pre-specified
Analysis type	superiority
Parameter estimate	Mean difference (final values)
Point estimate	23.88
Confidence interval	
level	Other: 80 %
sides	2-sided
lower limit	-14.25
upper limit	62

## Secondary: Period 1: Time to Adjudicated Clinical Worsening (CW)

End point title	Period 1: Time to Adjudicated Clinical Worsening (CW)
End point description:	
Clinical worsening was defined as any of the following: death, lung or heart transplantation, atrial septostomy or Potts' shunt, hospitalization for Pulmonary Arterial Hypertension (PAH) progression, new onset syncope, initiation of new PAH therapy (including increase in the dose of existing PAH specific concomitant therapy, such as endothelin receptor agonist or beraprost medication), or increase of 1 or more in World Health Organization (WHO) Functional Class (except for participants already in Class IV; only for participants unable to perform the 6 minute walk (6MW) test; worsening of WHO functional class by 1 or more for participants who can perform a 6 minute walk (6MW) test and who have a decrease of ≥ 20% in the 6 minute walk distance (for those participants who are ≥ 6 years of age). Criteria for CW (from Period 1) were adjudicated by an independent, blinded study-specific Clinical Endpoint Committee (CEC). This adjudication was used for data analysis, and was not used to guide	
End point type	Secondary
End point timeframe:	
Baseline through Week 24	

Analysis Population Description: All participants who received at least one dose of study drug.

End point values	Placebo	Tadalafil		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	18 <sup>[1]</sup>	17 <sup>[2]</sup>		
Units: Weeks				
median (confidence interval 95%)	9999 (9999 to 9999)	9999 (9999 to 9999)		

Notes:

[1] - 9999=Data Not Available (NA). There was no confirmed adjudicated CW case to report.

[2] - 9999=Data Not Available (NA). There was no confirmed adjudicated CW case to report.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Period 1: Percentage of Participants Who Experience CW

End point title	Period 1: Percentage of Participants Who Experience CW
-----------------	--

End point description:

Clinical worsening was defined as any of the following: death, lung or heart transplantation, atrial septostomy or Potts' shunt, hospitalization for Pulmonary Arterial Hypertension (PAH) progression, new onset syncope, initiation of new PAH therapy (including increase in the dose of existing PAH specific concomitant therapy, such as endothelin receptor agonist or beraprost medication), or increase of 1 or more in World Health Organization (WHO) Functional Class (except for participants already in Class IV; only for participants unable to perform the 6 minute walk (6MW) test; worsening of WHO functional class by 1 or more for participants who can perform a 6 minute walk (6MW) test and who have a decrease of  $\geq 20\%$  in the 6 minute walk distance (for those participants who are  $\geq 6$  years of age). Criteria for CW (from Period 1) were adjudicated by an independent, blinded study-specific Clinical Endpoint Committee (CEC). This adjudication was used for data analysis, and was not used to guide subject

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline through Week 24

Analysis Population Description: All participants who received at least one dose of study drug.

End point values	Placebo	Tadalafil		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	18	17		
Units: percentage of participants				
number (not applicable)	0	0		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Period 1: Pharmacokinetics (PK): Apparent Clearance (CL/F) of tadalafil

End point title	Period 1: Pharmacokinetics (PK): Apparent Clearance (CL/F) of tadalafil
-----------------	---

End point description:

Period 1: Pharmacokinetics (PK): Apparent Clearance (CL/F) of Tadalafil at Steady-state.

Analysis Population Description: All participants who received at least one dose of study drug and had evaluable PK data.

9999= NA.

For 20 mg tadalafil with concomitant bosentan, n=3.

For 20 mg tadalafil no bosentan (ERA: macitentan), n = 1. For n = 1, geometric mean and geometric coefficient of variation could not be calculated. Individual value is 2.68.

For 40 mg tadalafil No bosentan (ERA: Macitentan), n=0.

End point type	Secondary
End point timeframe:	
Week 2, Week 4, Week16 and Week 24	

End point values	20 mg Tadalafil	40 mg Tadalafil		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	4 <sup>[3]</sup>	13 <sup>[4]</sup>		
Units: Liter Per Hour (L/hr)				
geometric mean (geometric coefficient of variation)				
With concomitant bosentan	3.63 (± 38.1)	4.49 (± 28.2)		
No bosentan (ERA: Macitentan)	9999 (± 9999)	9999 (± 9999)		

Notes:

[3] - 9999= NA.

For 20 mg tadalafil No bosentan (ERA: Macitentan), n = 1.

[4] - 9999= NA.

For 40 mg tadalafil No bosentan (ERA: Macitentan), n=0.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Period 2: Time to First Occurrence of CW

End point title	Period 2: Time to First Occurrence of CW
-----------------	--

End point description:

Clinical worsening was defined as any of the following: death, lung or heart transplantation, atrial septostomy or Potts' shunt, hospitalization for Pulmonary Arterial Hypertension (PAH) progression, new onset syncope, initiation of new PAH therapy (including increase in the dose of existing PAH specific concomitant therapy, such as endothelin receptor agonist or beraprost medication), or increase of 1 or more in World Health Organization(WHO) Functional Class (except for participants already in Class IV; only for participants unable to perform the 6 minute walk (6MW) test; worsening of WHO functional class by 1 or more for participants who can perform a 6 minute walk (6MW) test and who have a decrease of ≥ 20% in the 6 minute walk distance (for those participants who are ≥6 years of age).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline through Study Completion (Up to 24 Months)

Analysis Population Description: Period 2: All participants who received at least one dose of study drug.

End point values	Placebo/Tadalafil - Open-Label Treatment	Tadalafil/Tadalafil - Open-Label Treatment		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	2 <sup>[5]</sup>	3 <sup>[6]</sup>		
Units: Months				
median (confidence interval 95%)	9999 (9999 to 9999)	9999 (9999 to 9999)		

Notes:

[5] - 9999=NA. Number of participants with CW were inadequate to perform the statistical analysis.

[6] - 9999=NA. Number of participants with CW were inadequate to perform the statistical analysis.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Period 2: Percentage of Participants Who Experience CW

End point title	Period 2: Percentage of Participants Who Experience CW
-----------------	--

End point description:

Clinical worsening was defined as any of the following: death, lung or heart transplantation, atrial septostomy or Potts' shunt, hospitalization for Pulmonary Arterial Hypertension (PAH) progression, new onset syncope, initiation of new PAH therapy (including increase in the dose of existing PAH specific concomitant therapy, such as endothelin receptor agonist or beraprost medication), or increase of 1 or more in World Health Organization(WHO) Functional Class (except for participants already in Class IV; only for participants unable to perform the 6 minute walk (6MW) test; worsening of WHO functional class by 1 or more for participants who can perform a 6 minute walk (6MW) test and who have a decrease of  $\geq 20\%$  in the 6 minute walk distance (for those participants who are  $\geq 6$  years of age).

End point type	Secondary
----------------	-----------

End point timeframe:

Period 2 Baseline through Study Completion (Up to 24 Months)

Analysis Population Description: Period 2: All participants who received at least one dose of study drug.

End point values	Placebo/Tadalafil - Open-Label Treatment	Tadalafil/Tadalafil - Open-Label Treatment		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	16	16		
Units: percentage of participants				
number (not applicable)	12.5	18.8		

### Statistical analyses

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Baseline Up To 24 Months

Adverse event reporting additional description:

All participants who received at least one dose of study drug.

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	24.0
--------------------	------

### Reporting groups

Reporting group title	Placebo - Double Blind
-----------------------	------------------------

Reporting group description:

Period 1: Participants received placebo orally by tablets once a day.

Reporting group title	Tadalafil - Double Blind
-----------------------	--------------------------

Reporting group description:

Period 1: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered tadalafil orally by tablets once a day.

Reporting group title	Tadalafil/Tadalafil - Open-Label Treatment
-----------------------	--

Reporting group description:

Period 1: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered tadalafil orally by tablets once a day.

Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered tadalafil orally by tablets once a day.

Reporting group title	Placebo/Tadalafil - Open-Label Treatment
-----------------------	--

Reporting group description:

Period 1: Participants received placebo orally by tablets once a day.

Period 2: 20 mg middle weight cohort or 40 mg for heavy weight cohort administered tadalafil orally by tablets once a day.

Serious adverse events	Placebo - Double Blind	Tadalafil - Double Blind	Tadalafil/Tadalafil - Open-Label Treatment
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events	0	0	0
Cardiac disorders			
acute right ventricular failure			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Blood and lymphatic system disorders			
anaemia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
haemoptysis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
gastroenteritis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
pneumonia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

<b>Serious adverse events</b>	Placebo/Tadalafil - Open-Label Treatment		
Total subjects affected by serious adverse events			
subjects affected / exposed	4 / 16 (25.00%)		
number of deaths (all causes)	0		
number of deaths resulting from adverse events	0		
Cardiac disorders			
acute right ventricular failure			
alternative dictionary used: MedDRA 24.0			

subjects affected / exposed	0 / 16 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Blood and lymphatic system disorders			
anaemia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Respiratory, thoracic and mediastinal disorders			
haemoptysis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Infections and infestations			
gastroenteritis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
pneumonia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		

Frequency threshold for reporting non-serious adverse events: 5 %

Non-serious adverse events	Placebo - Double Blind	Tadalafil - Double Blind	Tadalafil/Tadalafil - Open-Label Treatment
Total subjects affected by non-serious adverse events			
subjects affected / exposed	8 / 18 (44.44%)	15 / 17 (88.24%)	12 / 16 (75.00%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			



pharyngeal neoplasm alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	1 / 16 (6.25%) 1
Vascular disorders deep vein thrombosis alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  flushing alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  hypotension alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0   0 / 18 (0.00%) 0  0 / 18 (0.00%) 0	0 / 17 (0.00%) 0  1 / 17 (5.88%) 1  1 / 17 (5.88%) 1	1 / 16 (6.25%) 1  0 / 16 (0.00%) 0  0 / 16 (0.00%) 0
General disorders and administration site conditions asthenia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  exercise tolerance decreased alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  pyrexia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0   0 / 18 (0.00%) 0  0 / 18 (0.00%) 0	0 / 17 (0.00%) 0  0 / 17 (0.00%) 0  1 / 17 (5.88%) 1	0 / 16 (0.00%) 0  1 / 16 (6.25%) 1  0 / 16 (0.00%) 0
Immune system disorders hypersensitivity alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	0 / 16 (0.00%) 0

seasonal allergy alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	1 / 16 (6.25%) 2
Social circumstances menarche alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[1]</sup> occurrences (all)	1 / 9 (11.11%) 1	0 / 10 (0.00%) 0	0 / 10 (0.00%) 0
Reproductive system and breast disorders menorrhagia alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[2]</sup> occurrences (all) metrorrhagia alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[3]</sup> occurrences (all) penis disorder alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[4]</sup> occurrences (all) polymenorrhoea alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[5]</sup> occurrences (all) spontaneous penile erection alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[6]</sup> occurrences (all) uterine haemorrhage alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[7]</sup> occurrences (all)	0 / 9 (0.00%) 0  0 / 9 (0.00%) 0  0 / 9 (0.00%) 0  0 / 9 (0.00%) 0  0 / 9 (0.00%) 0  0 / 9 (0.00%) 0  0 / 9 (0.00%) 0	1 / 10 (10.00%) 1  0 / 10 (0.00%) 0  1 / 7 (14.29%) 1  0 / 10 (0.00%) 0  1 / 7 (14.29%) 2  0 / 10 (0.00%) 0	1 / 10 (10.00%) 2  1 / 10 (10.00%) 1  0 / 6 (0.00%) 0  1 / 10 (10.00%) 1  0 / 6 (0.00%) 0  0 / 10 (0.00%) 0
Respiratory, thoracic and mediastinal			

disorders			
cough			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
dyspnoea			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
epistaxis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 18 (5.56%)	2 / 17 (11.76%)	1 / 16 (6.25%)
occurrences (all)	1	2	1
oropharyngeal pain			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	1 / 17 (5.88%)	1 / 16 (6.25%)
occurrences (all)	0	1	1
pulmonary arterial hypertension			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 18 (5.56%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	1	0	0
respiratory distress			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Psychiatric disorders			
anxiety			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
excessive masturbation			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
insomnia			

alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	1 / 16 (6.25%) 1
Investigations blood creatine phosphokinase increased alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  hepatic enzyme increased alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0   0 / 18 (0.00%) 0	0 / 17 (0.00%) 0   1 / 17 (5.88%) 1	1 / 16 (6.25%) 1   0 / 16 (0.00%) 0
Injury, poisoning and procedural complications bone contusion alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  penis injury alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[8]</sup> occurrences (all)	0 / 18 (0.00%) 0   0 / 9 (0.00%) 0	1 / 17 (5.88%) 1   0 / 7 (0.00%) 0	0 / 16 (0.00%) 0   0 / 6 (0.00%) 0
Cardiac disorders palpitations alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  supraventricular tachycardia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  tachycardia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0   0 / 18 (0.00%) 0   0 / 18 (0.00%) 0	1 / 17 (5.88%) 1   0 / 17 (0.00%) 0   1 / 17 (5.88%) 1	0 / 16 (0.00%) 0   1 / 16 (6.25%) 1   0 / 16 (0.00%) 0

ventricular extrasystoles alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	0 / 16 (0.00%) 0
Nervous system disorders dizziness alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  headache alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  presyncope alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  somnolence alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  syncope alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0  2 / 18 (11.11%) 6  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0	0 / 17 (0.00%) 0  5 / 17 (29.41%) 5  1 / 17 (5.88%) 1  1 / 17 (5.88%) 1  0 / 17 (0.00%) 0	3 / 16 (18.75%) 3  3 / 16 (18.75%) 3  1 / 16 (6.25%) 1  0 / 16 (0.00%) 0  1 / 16 (6.25%) 1
Blood and lymphatic system disorders anaemia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  eosinophilia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0  1 / 18 (5.56%) 1	0 / 17 (0.00%) 0  0 / 17 (0.00%) 0	0 / 16 (0.00%) 0  0 / 16 (0.00%) 0
Ear and labyrinth disorders			

ear pain alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	1 / 16 (6.25%) 1
Eye disorders asthenopia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  photopsia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0  0 / 18 (0.00%) 0	0 / 17 (0.00%) 0  0 / 17 (0.00%) 0	0 / 16 (0.00%) 0  1 / 16 (6.25%) 1
Gastrointestinal disorders abdominal pain upper alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  dental caries alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  enterocolitis alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  gastrooesophageal reflux disease alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  irritable bowel syndrome alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  nausea	0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0	0 / 17 (0.00%) 0  0 / 17 (0.00%) 0  0 / 17 (0.00%) 0  0 / 17 (0.00%) 0  0 / 17 (0.00%) 0	2 / 16 (12.50%) 2  1 / 16 (6.25%) 1  0 / 16 (0.00%) 0  0 / 16 (0.00%) 0  1 / 16 (6.25%) 1

<p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 18 (0.00%)</p> <p>0</p>	<p>0 / 17 (0.00%)</p> <p>0</p>	<p>1 / 16 (6.25%)</p> <p>1</p>
<p>plicated tongue</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 18 (0.00%)</p> <p>0</p>	<p>0 / 17 (0.00%)</p> <p>0</p>	<p>1 / 16 (6.25%)</p> <p>1</p>
<p>vomiting</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 18 (0.00%)</p> <p>0</p>	<p>0 / 17 (0.00%)</p> <p>0</p>	<p>3 / 16 (18.75%)</p> <p>3</p>
<p>Skin and subcutaneous tissue disorders</p> <p>dermatitis allergic</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 18 (0.00%)</p> <p>0</p>	<p>0 / 17 (0.00%)</p> <p>0</p>	<p>1 / 16 (6.25%)</p> <p>2</p>
<p>haemorrhage subcutaneous</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 18 (0.00%)</p> <p>0</p>	<p>1 / 17 (5.88%)</p> <p>3</p>	<p>0 / 16 (0.00%)</p> <p>0</p>
<p>livedo reticularis</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>1 / 18 (5.56%)</p> <p>1</p>	<p>0 / 17 (0.00%)</p> <p>0</p>	<p>0 / 16 (0.00%)</p> <p>0</p>
<p>photosensitivity reaction</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 18 (0.00%)</p> <p>0</p>	<p>0 / 17 (0.00%)</p> <p>0</p>	<p>1 / 16 (6.25%)</p> <p>1</p>
<p>pruritus</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 18 (0.00%)</p> <p>0</p>	<p>0 / 17 (0.00%)</p> <p>0</p>	<p>0 / 16 (0.00%)</p> <p>0</p>
<p>rash</p> <p>alternative dictionary used: MedDRA 24.0</p>			

subjects affected / exposed	0 / 18 (0.00%)	1 / 17 (5.88%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
swelling face			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	1 / 17 (5.88%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Musculoskeletal and connective tissue disorders			
arthralgia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 18 (5.56%)	2 / 17 (11.76%)	2 / 16 (12.50%)
occurrences (all)	1	2	2
back pain			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	1 / 17 (5.88%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
musculoskeletal chest pain			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
pain in extremity			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
patellofemoral pain syndrome			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Infections and infestations			
acute sinusitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
ascariasis			
alternative dictionary used: MedDRA 24.0			



subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
body tinea			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
bronchitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 18 (5.56%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	1	0	0
ear infection			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	2
gastroenteritis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
impetigo			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
influenza			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	3 / 17 (17.65%)	1 / 16 (6.25%)
occurrences (all)	0	3	1
nasopharyngitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	3 / 16 (18.75%)
occurrences (all)	0	0	4
otitis media acute			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1

paronychia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
pharyngitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	1 / 17 (5.88%)	2 / 16 (12.50%)
occurrences (all)	0	1	4
rhinitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	1 / 17 (5.88%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
sinusitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 18 (5.56%)	1 / 17 (5.88%)	1 / 16 (6.25%)
occurrences (all)	1	1	2
tonsillitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
tracheobronchitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 18 (0.00%)	0 / 17 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
upper respiratory tract infection			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 18 (5.56%)	3 / 17 (17.65%)	2 / 16 (12.50%)
occurrences (all)	1	3	2
urinary tract infection			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 18 (5.56%)	1 / 17 (5.88%)	1 / 16 (6.25%)
occurrences (all)	1	1	1
vaginal infection			
alternative dictionary used: MedDRA 24.0			

subjects affected / exposed <sup>[9]</sup> occurrences (all)	0 / 9 (0.00%) 0	1 / 10 (10.00%) 1	0 / 10 (0.00%) 0
viral infection alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	0 / 16 (0.00%) 0
viral tonsillitis alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	1 / 18 (5.56%) 1	0 / 17 (0.00%) 0	0 / 16 (0.00%) 0
viral upper respiratory tract infection alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	1 / 17 (5.88%) 1	0 / 16 (0.00%) 0
Metabolism and nutrition disorders decreased appetite alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	0 / 16 (0.00%) 0
dyslipidaemia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	1 / 16 (6.25%) 1
vitamin b12 deficiency alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	0 / 16 (0.00%) 0
vitamin d deficiency alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 18 (0.00%) 0	0 / 17 (0.00%) 0	0 / 16 (0.00%) 0
<b>Non-serious adverse events</b>	Placebo/Tadalafil - Open-Label Treatment		
Total subjects affected by non-serious adverse events			

subjects affected / exposed	11 / 16 (68.75%)		
Neoplasms benign, malignant and unspecified (incl cysts and polyps) pharyngeal neoplasm alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 16 (0.00%) 0		
Vascular disorders deep vein thrombosis alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  flushing alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  hypotension alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 16 (0.00%) 0  0 / 16 (0.00%) 0  0 / 16 (0.00%) 0		
General disorders and administration site conditions asthenia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  exercise tolerance decreased alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  pyrexia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	1 / 16 (6.25%) 1  0 / 16 (0.00%) 0  0 / 16 (0.00%) 0		
Immune system disorders hypersensitivity alternative dictionary used: MedDRA 24.0			

subjects affected / exposed occurrences (all)	1 / 16 (6.25%) 1		
seasonal allergy alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 16 (0.00%) 0		
Social circumstances menarche alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[1]</sup> occurrences (all)	0 / 8 (0.00%) 0		
Reproductive system and breast disorders menorrhagia alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[2]</sup> occurrences (all)	1 / 8 (12.50%) 1		
metrorrhagia alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[3]</sup> occurrences (all)	0 / 8 (0.00%) 0		
penis disorder alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[4]</sup> occurrences (all)	0 / 8 (0.00%) 0		
polymenorrhoea alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[5]</sup> occurrences (all)	0 / 8 (0.00%) 0		
spontaneous penile erection alternative dictionary used: MedDRA 24.0 subjects affected / exposed <sup>[6]</sup> occurrences (all)	1 / 8 (12.50%) 1		
uterine haemorrhage alternative dictionary used: MedDRA 24.0			

subjects affected / exposed <sup>[7]</sup>	1 / 8 (12.50%)		
occurrences (all)	1		
Respiratory, thoracic and mediastinal disorders			
cough			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
dyspnoea			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
epistaxis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	1		
oropharyngeal pain			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
pulmonary arterial hypertension			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
respiratory distress			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
Psychiatric disorders			
anxiety			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	1		
excessive masturbation			
alternative dictionary used: MedDRA 24.0			

<p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>insomnia</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Investigations</p> <p>blood creatine phosphokinase increased</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>hepatic enzyme increased</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Injury, poisoning and procedural complications</p> <p>bone contusion</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>penis injury</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed<sup>[8]</sup></p> <p>1 / 8 (12.50%)</p> <p>occurrences (all)</p> <p>1</p>			
<p>Cardiac disorders</p> <p>palpitations</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>supraventricular tachycardia</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>tachycardia</p> <p>alternative dictionary used:</p>			

MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
ventricular extrasystoles			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	1		
Nervous system disorders			
dizziness			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	1		
headache			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	2 / 16 (12.50%)		
occurrences (all)	2		
presyncope			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
somnolence			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
syncope			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
Blood and lymphatic system disorders			
anaemia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	2 / 16 (12.50%)		
occurrences (all)	2		
eosinophilia			
alternative dictionary used: MedDRA 24.0			



subjects affected / exposed occurrences (all)	0 / 16 (0.00%) 0		
Ear and labyrinth disorders ear pain alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	0 / 16 (0.00%) 0		
Eye disorders asthenopia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  photopsia alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)	1 / 16 (6.25%) 1  0 / 16 (0.00%) 0		
Gastrointestinal disorders abdominal pain upper alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  dental caries alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  enterocolitis alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  gastrooesophageal reflux disease alternative dictionary used: MedDRA 24.0 subjects affected / exposed occurrences (all)  irritable bowel syndrome alternative dictionary used: MedDRA 24.0	0 / 16 (0.00%) 0  0 / 16 (0.00%) 0  1 / 16 (6.25%) 1  1 / 16 (6.25%) 1		

<p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>nausea</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>plicated tongue</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>vomiting</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Skin and subcutaneous tissue disorders</p> <p>dermatitis allergic</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>haemorrhage subcutaneous</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>livedo reticularis</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>photosensitivity reaction</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>pruritus</p> <p>alternative dictionary used: MedDRA 24.0</p>			

<p>subjects affected / exposed</p> <p>1 / 16 (6.25%)</p> <p>occurrences (all)</p> <p>1</p> <p>rash</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>swelling face</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Musculoskeletal and connective tissue disorders</p> <p>arthralgia</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>back pain</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>musculoskeletal chest pain</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>1 / 16 (6.25%)</p> <p>occurrences (all)</p> <p>1</p> <p>pain in extremity</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>patellofemoral pain syndrome</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>1 / 16 (6.25%)</p> <p>occurrences (all)</p> <p>1</p>			
<p>Infections and infestations</p> <p>acute sinusitis</p> <p>alternative dictionary used: MedDRA 24.0</p>			

subjects affected / exposed	2 / 16 (12.50%)		
occurrences (all)	2		
ascariasis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
body tinea			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
bronchitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
ear infection			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
gastroenteritis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
impetigo			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	1		
influenza			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
nasopharyngitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	2		

otitis media acute			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
paronychia			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
pharyngitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
rhinitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	2 / 16 (12.50%)		
occurrences (all)	2		
sinusitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
tonsillitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
tracheobronchitis			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	1		
upper respiratory tract infection			
alternative dictionary used: MedDRA 24.0			
subjects affected / exposed	0 / 16 (0.00%)		
occurrences (all)	0		
urinary tract infection			
alternative dictionary used: MedDRA 24.0			

<p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>vaginal infection</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed<sup>[9]</sup></p> <p>0 / 8 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>viral infection</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>1 / 16 (6.25%)</p> <p>occurrences (all)</p> <p>1</p>			
<p>viral tonsillitis</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>viral upper respiratory tract infection</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Metabolism and nutrition disorders</p> <p>decreased appetite</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>1 / 16 (6.25%)</p> <p>occurrences (all)</p> <p>1</p> <p>dyslipidaemia</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>0 / 16 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>vitamin b12 deficiency</p> <p>alternative dictionary used: MedDRA 24.0</p> <p>subjects affected / exposed</p> <p>1 / 16 (6.25%)</p> <p>occurrences (all)</p> <p>1</p> <p>vitamin d deficiency</p> <p>alternative dictionary used: MedDRA 24.0</p>			

subjects affected / exposed	1 / 16 (6.25%)		
occurrences (all)	1		

---

Notes:

[1] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[2] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[3] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[4] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[5] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[6] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[7] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[8] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly

[9] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: This event is gender specific, only occurring in male or female subjects. The number of subjects exposed has been adjusted accordingly.

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
27 September 2012	Change of primary endpoint for US from right heart catheterization (RHC) to Cardiac Echo. Per requirement from FDA that RHC can no longer be used as primary endpoint in Pediatric PAH trials.
14 December 2012	Per FDA removal of Tricuspid Annular Plane Systolic Excursion (TAPSE) as primary endpoint for US. Change of primary endpoint for US to improving time of 6-minute walk (6MW) test from baseline to week 24.
13 December 2018	Change of primary endpoint and study sample size.

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

The study is mainly descriptive in a small number of children with PAH and there were no participants enrolled in the light weight cohort.
--

Notes: