



## Clinical trial results:

### A Randomized, Double-Blind, Placebo-Controlled, Phase 2 Trial to Evaluate the Efficacy and Safety of a Vaccine, ASP0113, in Cytomegalovirus (CMV)-Seronegative Kidney Transplant Recipients Receiving an Organ from a CMV-Seropositive Donor

#### Summary

EudraCT number	2013-000464-29
Trial protocol	DE ES
Global end of trial date	

#### Results information

Result version number	v1
This version publication date	29 April 2017
First version publication date	29 April 2017

#### Trial information

##### Trial identification

Sponsor protocol code	0113-CL-2001
-----------------------	--------------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT01974206
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	Astellas Pharma Global Development, Inc.
Sponsor organisation address	1 Astellas Way, Northbrook, United States, 60062
Public contact	Clinical Trial Disclosure, Astellas Pharma Global Development, Inc., <a href="mailto:astellas.resultsdisclosure@astellas.com">astellas.resultsdisclosure@astellas.com</a>
Scientific contact	Clinical Trial Disclosure, Astellas Pharma Global Development, Inc., <a href="mailto:astellas.resultsdisclosure@astellas.com">astellas.resultsdisclosure@astellas.com</a>

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Interim
Date of interim/final analysis	13 May 2016
Is this the analysis of the primary completion data?	Yes
Primary completion date	13 May 2016
Global end of trial reached?	No

Notes:

## General information about the trial

Main objective of the trial:

The main objective of the trial was to evaluate efficacy and safety of ASP0113 compared to placebo in reducing the incidence of Cytomegalovirus (CMV) viremia through 1 year post first study drug injection in CMV-seronegative patients who received a kidney from a CMV-seropositive donor.

Protection of trial subjects:

This clinical study was written, conducted and reported in accordance with the protocol, International Council for Harmonisation of Technical Requirements for Pharmaceuticals for Human Use (ICH) Good Clinical Practice (GCP) Guidelines, and applicable local regulations, including the European Directive 2001/20/EC, on the protection of human rights, and with the ethical principles that have their origin in the Declaration of Helsinki. Astellas ensures that the use and disclosure of protected health information (PHI) obtained during a research study complies with the federal, national and/or regional legislation related to the privacy and protection of personal information.

Background therapy:

In the first 10 days after the transplant, patients received prophylactic valganciclovir or ganciclovir (dose per package insert) from the day of randomization to prevent Cytomegalovirus (CMV) disease. After randomization, patients continued to receive valganciclovir or ganciclovir until 100 days post-transplant. Valganciclovir or ganciclovir could be interrupted, dose adjusted or replaced by other CMV-specific antiviral prophylaxis (AVP) per standard of care after the day of randomization.

Evidence for comparator: -

Actual start date of recruitment	20 November 2013
Long term follow-up planned	Yes
Long term follow-up rationale	Safety
Long term follow-up duration	4 Years
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	United States: 93
Country: Number of subjects enrolled	Germany: 16
Country: Number of subjects enrolled	Canada: 7
Country: Number of subjects enrolled	France: 15
Country: Number of subjects enrolled	Spain: 8
Country: Number of subjects enrolled	Australia: 11
Worldwide total number of subjects	150
EEA total number of subjects	39

Notes:

<b>Subjects enrolled per age group</b>	
In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	129
From 65 to 84 years	21
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

A total of 150 patients were enrolled across 6 countries over approximately 3.5 years. Eligible patients for the study were  $\geq 18$  years of age and CMV-seronegative at the time of the transplant. Furthermore they had a kidney allograft from a CMV-seropositive living or deceased donor.

### Pre-assignment

Screening details:

Screening assessments were performed from 14-30 days after the transplant. Patients were randomized at day  $30 \pm 3$ , in relation to the day of the transplant, in a 1:1 ratio to ASP0113 or placebo. Patients were stratified by the use of antithymocyte globulin (ATG) prior to randomization and by the receipt of a kidney from a living or deceased donor.

### Period 1

Period 1 title	Primary Study Period (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator, Assessor

Blinding implementation details:

This was a double-blind study. Patients were randomized to receive ASP0113 or placebo in a double-blind fashion such that the investigator, sponsor's study management team, clinical staff nor the patient knew which agent was being administered.

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Placebo

Arm description:

Patients received 1 mL of 5 mg/mL of placebo via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).

Arm type	Placebo
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Injection
Routes of administration	Intramuscular use

Dosage and administration details:

Patients received Placebo in 2-mL vials containing phosphate-buffered saline.

<b>Arm title</b>	ASP0113 5mg
------------------	-------------

Arm description:

Patients received 1 mL of 5 mg/mL of ASP0113 via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).

Arm type	Experimental
Investigational medicinal product name	ASP0113
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Injection
Routes of administration	Intramuscular use

Dosage and administration details:

Patients received ASP0113 in single dose 2-ml vials containing 1.3 ml of 5 mg/mL of ASP0113.

<b>Number of subjects in period 1</b>	Placebo	ASP0113 5mg
Started	74	76
Completed	68	75
Not completed	6	1
Consent withdrawn by subject	3	-
Physician decision	1	-
Patient did not take study drug	-	1
Death	1	-
Lost to follow-up	1	-

## Baseline characteristics

### Reporting groups

Reporting group title	Placebo
-----------------------	---------

Reporting group description:

Patients received 1 mL of 5 mg/mL of placebo via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).

Reporting group title	ASP0113 5mg
-----------------------	-------------

Reporting group description:

Patients received 1 mL of 5 mg/mL of ASP0113 via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).

Reporting group values	Placebo	ASP0113 5mg	Total
Number of subjects	74	76	150
Age categorical			
Units: Subjects			

Age continuous			
Units: years			
arithmetic mean	47.9	50.8	
standard deviation	± 13.3	± 13.6	-
Gender categorical			
Units:			
Male	55	55	110
Female	19	21	40

## End points

### End points reporting groups

Reporting group title	Placebo
Reporting group description:	
Patients received 1 mL of 5 mg/mL of placebo via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).	
Reporting group title	ASP0113 5mg
Reporting group description:	
Patients received 1 mL of 5 mg/mL of ASP0113 via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).	

### Primary: Incidence of Cytomegalovirus (CMV) Viremia Through One Year Post First Study Drug Injection

End point title	Incidence of Cytomegalovirus (CMV) Viremia Through One Year Post First Study Drug Injection
End point description:	
CMV viremia was defined as presence of cytomegalovirus as measured in plasma viral load of $\geq 1000$ IU/mL by central laboratory assay. A patient who discontinued the study without a positive CMV viral load was imputed as having a CMV viremia. A patient who had more than one viral load $\geq 1000$ IU/mL by central assay was counted once in this summary. CMV viral loads after first injection (Day 1) through Day 380 (scheduled or unscheduled) were included in the analysis. The analysis population was the Full Analysis Set (FAS) which consisted of all randomized patients who received at least 1 dose of randomized study drug and who had at least 1 post dose viral load assessment within 1 year post first injection by central laboratory.	
End point type	Primary
End point timeframe:	
From first study dose injection (Day 1) up to one year post study drug injection (up to Day 380)	

End point values	Placebo	ASP0113 5mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	73	73		
Units: Percentage of Patients				
number (not applicable)				
Known CMV Viremia	35.6	35.6		
Imputed CMV Viremia Due to Discontinuation	5.5	0		

### Statistical analyses

Statistical analysis title	Common Odds Ratio for Patients With CMV
Statistical analysis description:	
The Cochran-Mantel-Haenszel (CMH) estimate of the common odds ratio (ASP0113 vs placebo) stratified by randomization strata, and the 90% CIs of the CMH odds ratio stratified by randomization group.	
Comparison groups	Placebo v ASP0113 5mg

Number of subjects included in analysis	146
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.307 <sup>[1]</sup>
Method	Cochran-Mantel-Haenszel
Parameter estimate	Common Odds Ratio
Point estimate	0.79
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.43
upper limit	1.47

Notes:

[1] - P-value (1-sided) of the CMH adjusted odds ratio stratified by randomization group.

### Secondary: Proportion of Patients with Adjudicated CMV-Associated Disease, Including CMV Syndrome and CMV Tissue-Invasive Disease

End point title	Proportion of Patients with Adjudicated CMV-Associated Disease, Including CMV Syndrome and CMV Tissue-Invasive Disease
-----------------	--

End point description:

An independent panel of medical experts reviewed/adjudicated events of CMV-associated disease including CMV syndrome and tissue invasive disease, which were defined according to the American Society of Transplantation Recommendations for Screening, Monitoring and Reporting of Infectious Complications in Immunosuppression Trials in Recipients of Organ Transplantation 2006. The analysis population was the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

From first study dose injection (Day 1) up to one year post study drug injection (up to Day 380)

End point values	Placebo	ASP0113 5mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	73	73		
Units: Percentage of Patients				
number (not applicable)	19.18	19.18		

### Statistical analyses

Statistical analysis title	Common Odds Ratio for CMV-Associated Disease
----------------------------	--

Statistical analysis description:

The exact Cochran-Mantel-Haenszel (CMH) estimate of the common odds ratio (ASP0113 vs placebo) stratified by randomization group, and the 90% CI of the exact CMH odds ratio stratified by randomization group.

Comparison groups	Placebo v ASP0113 5mg
-------------------	-----------------------



Number of subjects included in analysis	146
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.576 <sup>[2]</sup>
Method	Cochran-Mantel-Haenszel
Parameter estimate	Common Odds Ratio
Point estimate	0.99
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.46
upper limit	2.15

Notes:

[2] - P-value (1-sided) of the exact CMH method adjusted odds ratio stratified by randomization group.

### Secondary: Proportion of Patients with CMV Viremia Defined as Plasma Viral Load $\geq$ the Lower Limit of Quantification (LLOQ) Assessed by Central Laboratory

End point title	Proportion of Patients with CMV Viremia Defined as Plasma Viral Load $\geq$ the Lower Limit of Quantification (LLOQ) Assessed by Central Laboratory
-----------------	---

End point description:

The central laboratory had the LLOQ level for CMV viral load assessment. When the viral load was below the LLOQ the actual reading was not possible and was denoted as  $\leq$ LLOQ. If the patient had any CMV viral load assessments greater than the LLOQ, set up by the central laboratory, patient was classified as viremic and was included in the analysis. The analysis population was the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

From first study dose injection (Day 1) up to one year post study drug injection (up to Day 380)

End point values	Placebo	ASP0113 5mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	73	73		
Units: Percentage of Patients				
number (not applicable)	49.32	46.58		

### Statistical analyses

Statistical analysis title	Common Odds Ratio for Patients with CMV Viremia
----------------------------	---

Statistical analysis description:

The exact Cochran-Mantel-Haenszel (CMH) estimate of the common odds ratio (ASP0113 vs placebo) stratified by randomization group, and the 90% CI of the exact CMH odds ratio stratified by randomization group.

Comparison groups	Placebo v ASP0113 5mg
-------------------	-----------------------

Number of subjects included in analysis	146
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.408 <sup>[3]</sup>
Method	Cochran-Mantel-Haenszel
Parameter estimate	Common Odds Ratio
Point estimate	0.88
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.48
upper limit	1.59

Notes:

[3] - P-value (1-sided) of the exact CMH method adjusted odds ratio stratified by randomization group.

### Secondary: Proportion of Patients Who Took Adjudicated CMV-specific antiviral therapy (AVT) for the Treatment of CMV Viremia or Disease

End point title	Proportion of Patients Who Took Adjudicated CMV-specific antiviral therapy (AVT) for the Treatment of CMV Viremia or Disease
-----------------	--

End point description:

An independent panel of medical experts reviewed/adjudicated events of CMV-specific AVT for treatment of CMV viremia or disease. The analysis population was the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

From first study dose injection (Day 1) up to one year post study drug injection (up to Day 380)

End point values	Placebo	ASP0113 5mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	73	73		
Units: Percentage of Patients				
number (not applicable)	45.21	42.47		

### Statistical analyses

Statistical analysis title	Common Odds Ratio for Patients With CMV-AVT
----------------------------	---

Statistical analysis description:

The exact Cochran-Mantel-Haenszel (CMH) estimate of the common odds ratio (ASP0113 vs placebo) stratified by randomization group, and the 90% CI of the exact CMH odds ratio stratified by randomization group.

Comparison groups	Placebo v ASP0113 5mg
-------------------	-----------------------

Number of subjects included in analysis	146
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.419 <sup>[4]</sup>
Method	Cochran-Mantel-Haenszel
Parameter estimate	Common Odds Ratio
Point estimate	0.88
Confidence interval	
level	90 %
sides	2-sided
lower limit	0.48
upper limit	1.61

Notes:

[4] - P-value (1-sided) of the exact CMH method adjusted odds ratio stratified by randomization group.

## Secondary: Proportion of Patients with Graft Survival

End point title	Proportion of Patients with Graft Survival
End point description:	
Graft survival was defined for any patient that did not fit the definition of graft loss. Graft loss was defined as patient death, re-transplant, nephrectomy, or return to permanent dialysis (i.e., for > 30 days). Missing values for graft survival were not included in the denominator when making the proportion. The analysis population was the FAS.	
End point type	Secondary
End point timeframe:	
From first study dose injection (Day 1) up to one year post study drug injection (up to Day 380)	

End point values	Placebo	ASP0113 5mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	68	71		
Units: Percentage of Patients				
number (not applicable)	98.53	100		

## Statistical analyses

Statistical analysis title	Common Odds Ratio of Patients with Graft Survival
Statistical analysis description:	
Exact Cochran-Mantel-Haenszel estimate of the common odds ratio (ASP0113 versus Placebo) could not be estimated and is denoted as "9999." Moreover, upper limit of 90% CI of odds ratio is an infinity value and is also denoted as "9999."	
Comparison groups	Placebo v ASP0113 5mg
Number of subjects included in analysis	139
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.5 <sup>[5]</sup>
Method	Cochran-Mantel-Haenszel
Parameter estimate	Common Odds Ratio
Point estimate	9999

Confidence interval	
level	90 %
sides	2-sided
lower limit	0.11
upper limit	9999

Notes:

[5] - P-value (1-sided) of the Exact Cochran-Mantel-Haenszel method adjusted odds ratio stratified by randomization group.

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Day 1- Day 380

Adverse event reporting additional description:

Treatment Emergent Adverse Event (TEAE) was defined as an adverse event observed after the first study drug injection Day 1 through Day 380. Drug-related Adverse Event (AE) was defined as any AE with at least possible relationship to study treatment as assessed by the investigator or with missing assessment of the causal relationship.

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	16.0
--------------------	------

### Reporting groups

Reporting group title	ASP0113 5mg
-----------------------	-------------

Reporting group description:

Patients received 1 mL of 5 mg/mL of ASP0113 via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).

Reporting group title	Placebo
-----------------------	---------

Reporting group description:

Patients received 1 mL of 5 mg/mL of placebo via injection in the deltoid muscle alternating sides with each dose on days 30, 60, 90, 120 and 180 in relation to the day of transplant (Day 0).

Serious adverse events	ASP0113 5mg	Placebo	
Total subjects affected by serious adverse events			
subjects affected / exposed	44 / 75 (58.67%)	36 / 74 (48.65%)	
number of deaths (all causes)	0	1	
number of deaths resulting from adverse events	0	0	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Basal cell carcinoma			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Colorectal cancer			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Invasive lobular breast carcinoma			

subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Metastases to peritoneum			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Prostate cancer			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Squamous cell carcinoma			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Squamous cell carcinoma of skin			
subjects affected / exposed	0 / 75 (0.00%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vascular disorders			
Aortic aneurysm			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Deep vein thrombosis			
subjects affected / exposed	0 / 75 (0.00%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypotension			
subjects affected / exposed	0 / 75 (0.00%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Jugular vein thrombosis			

subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lymphocele			
subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Thrombophlebitis superficial			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Thrombosis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Surgical and medical procedures			
Nephrectomy			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Transurethral prostatectomy			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ureteral stent removal			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urostomy			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
General disorders and administration site conditions			

Malaise			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyrexia			
subjects affected / exposed	3 / 75 (4.00%)	3 / 74 (4.05%)	
occurrences causally related to treatment / all	0 / 3	0 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Immune system disorders			
Drug hypersensitivity			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Kidney transplant rejection			
subjects affected / exposed	3 / 75 (4.00%)	4 / 74 (5.41%)	
occurrences causally related to treatment / all	0 / 3	0 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
Respiratory, thoracic and mediastinal disorders			
Dyspnoea			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pleural effusion			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia aspiration			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Psychiatric disorders			
Depression			



subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Mental status changes			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Investigations			
Blood creatinine increased			
subjects affected / exposed	5 / 75 (6.67%)	4 / 74 (5.41%)	
occurrences causally related to treatment / all	0 / 5	0 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood glucose increased			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HLA marker study positive			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Immunosuppressant drug level increased			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injury, poisoning and procedural complications			
Bone fissure			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Graft haemorrhage			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Graft thrombosis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lumbar vertebral fracture			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Perinephric collection			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Post procedural haematuria			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Seroma			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Spinal compression fracture			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Wound			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac disorders			
Acute myocardial infarction			
subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Atrial fibrillation			

subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bradycardia			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Coronary artery disease			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
Convulsion			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Neuropathy peripheral			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood and lymphatic system disorders			
Agranulocytosis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Anaemia			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Febrile neutropenia			
subjects affected / exposed	2 / 75 (2.67%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 2	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Leukopenia			

subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	1 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Neutropenia			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	1 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Eye disorders			
Angle closure glaucoma			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Aphthous stomatitis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Constipation			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diarrhoea			
subjects affected / exposed	2 / 75 (2.67%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 2	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal toxicity			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Subileus			

subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Umbilical hernia			
subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatobiliary disorders			
Bile duct stone			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cholelithiasis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin and subcutaneous tissue disorders			
Diabetic foot			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal and urinary disorders			
Calculus urinary			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haematuria			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hydronephrosis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Nephrolithiasis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nephropathy			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal artery stenosis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal cyst			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal failure acute			
subjects affected / exposed	10 / 75 (13.33%)	6 / 74 (8.11%)	
occurrences causally related to treatment / all	0 / 12	0 / 10	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal impairment			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal tubular necrosis			
subjects affected / exposed	0 / 75 (0.00%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ureteric stenosis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary bladder atrophy			

subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary retention			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract disorder			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinoma			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Endocrine disorders			
Hyperparathyroidism tertiary			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Musculoskeletal and connective tissue disorders			
Rhabdomyolysis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
Abscess soft tissue			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
BK virus infection			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Bacterial pyelonephritis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Clostridium difficile colitis			
subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 1	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cytomegalovirus colitis			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	1 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cytomegalovirus infection			
subjects affected / exposed	7 / 75 (9.33%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	1 / 7	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cytomegalovirus mucocutaneous ulcer			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cytomegalovirus oesophagitis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cytomegalovirus syndrome			
subjects affected / exposed	2 / 75 (2.67%)	5 / 74 (6.76%)	
occurrences causally related to treatment / all	2 / 2	3 / 5	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cytomegalovirus viraemia			
subjects affected / exposed	5 / 75 (6.67%)	3 / 74 (4.05%)	
occurrences causally related to treatment / all	3 / 7	1 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Enterococcal bacteraemia			



subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Enterocolitis bacterial			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Epstein-Barr viraemia			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Escherichia bacteraemia			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Escherichia sepsis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Escherichia urinary tract infection			
subjects affected / exposed	5 / 75 (6.67%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 6	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis clostridial			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis norovirus			

subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis viral			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal infection			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haematoma infection			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infected lymphocele			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nasopharyngitis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Orchitis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Peritonitis			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Pneumocystis jiroveci pneumonia			

subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia cytomegaloviral			
subjects affected / exposed	3 / 75 (4.00%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	1 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia necrotising			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Polyomavirus-associated nephropathy			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyelonephritis			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyelonephritis acute			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Staphylococcal bacteraemia			
subjects affected / exposed	0 / 75 (0.00%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Staphylococcal sepsis			

subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Streptococcal bacteraemia			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract infection			
subjects affected / exposed	3 / 75 (4.00%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 4	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract infection bacterial			
subjects affected / exposed	3 / 75 (4.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 4	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract infection enterococcal			
subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract infection pseudomonal			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urosepsis			
subjects affected / exposed	3 / 75 (4.00%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Viral infection			
subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Metabolism and nutrition disorders			
Dehydration			

subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diabetes mellitus			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diabetes mellitus inadequate control			
subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Failure to thrive			
subjects affected / exposed	0 / 75 (0.00%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Fluid overload			
subjects affected / exposed	1 / 75 (1.33%)	1 / 74 (1.35%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hyperglycaemia			
subjects affected / exposed	1 / 75 (1.33%)	2 / 74 (2.70%)	
occurrences causally related to treatment / all	0 / 1	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hyperkalaemia			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypokalaemia			
subjects affected / exposed	2 / 75 (2.67%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypovolaemia			

subjects affected / exposed	1 / 75 (1.33%)	0 / 74 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	ASP0113 5mg	Placebo	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	75 / 75 (100.00%)	72 / 74 (97.30%)	
Vascular disorders			
Arteriosclerosis			
subjects affected / exposed	4 / 75 (5.33%)	1 / 74 (1.35%)	
occurrences (all)	4	1	
Hypertension			
subjects affected / exposed	12 / 75 (16.00%)	9 / 74 (12.16%)	
occurrences (all)	12	14	
Hypotension			
subjects affected / exposed	11 / 75 (14.67%)	8 / 74 (10.81%)	
occurrences (all)	16	10	
Orthostatic hypotension			
subjects affected / exposed	0 / 75 (0.00%)	4 / 74 (5.41%)	
occurrences (all)	0	5	
Surgical and medical procedures			
Ureteral stent removal			
subjects affected / exposed	1 / 75 (1.33%)	6 / 74 (8.11%)	
occurrences (all)	1	6	
General disorders and administration site conditions			
Asthenia			
subjects affected / exposed	5 / 75 (6.67%)	8 / 74 (10.81%)	
occurrences (all)	5	8	
Chills			
subjects affected / exposed	2 / 75 (2.67%)	4 / 74 (5.41%)	
occurrences (all)	2	4	
Fatigue			
subjects affected / exposed	27 / 75 (36.00%)	24 / 74 (32.43%)	
occurrences (all)	59	38	

Injection site erythema subjects affected / exposed occurrences (all)	8 / 75 (10.67%) 14	2 / 74 (2.70%) 2	
Injection site pain subjects affected / exposed occurrences (all)	47 / 75 (62.67%) 277	18 / 74 (24.32%) 46	
Oedema subjects affected / exposed occurrences (all)	5 / 75 (6.67%) 5	3 / 74 (4.05%) 3	
Oedema peripheral subjects affected / exposed occurrences (all)	9 / 75 (12.00%) 13	5 / 74 (6.76%) 8	
Pyrexia subjects affected / exposed occurrences (all)	9 / 75 (12.00%) 11	13 / 74 (17.57%) 14	
Respiratory, thoracic and mediastinal disorders Cough subjects affected / exposed occurrences (all)	7 / 75 (9.33%) 8	4 / 74 (5.41%) 4	
Dyspnoea subjects affected / exposed occurrences (all)	5 / 75 (6.67%) 9	8 / 74 (10.81%) 10	
Oropharyngeal pain subjects affected / exposed occurrences (all)	4 / 75 (5.33%) 4	5 / 74 (6.76%) 5	
Psychiatric disorders Insomnia subjects affected / exposed occurrences (all)	8 / 75 (10.67%) 11	6 / 74 (8.11%) 8	
Investigations Blood creatinine increased subjects affected / exposed occurrences (all)	11 / 75 (14.67%) 17	16 / 74 (21.62%) 19	
Weight increased subjects affected / exposed occurrences (all)	4 / 75 (5.33%) 4	3 / 74 (4.05%) 3	

White blood cell count decreased subjects affected / exposed occurrences (all)	1 / 75 (1.33%) 1	4 / 74 (5.41%) 12	
Injury, poisoning and procedural complications			
Complications of transplant surgery subjects affected / exposed occurrences (all)	2 / 75 (2.67%) 2	4 / 74 (5.41%) 4	
Complications of transplanted kidney subjects affected / exposed occurrences (all)	4 / 75 (5.33%) 4	4 / 74 (5.41%) 4	
Procedural pain subjects affected / exposed occurrences (all)	6 / 75 (8.00%) 10	2 / 74 (2.70%) 2	
Cardiac disorders			
Bradycardia subjects affected / exposed occurrences (all)	5 / 75 (6.67%) 5	3 / 74 (4.05%) 3	
Tachycardia subjects affected / exposed occurrences (all)	5 / 75 (6.67%) 5	3 / 74 (4.05%) 3	
Nervous system disorders			
Dizziness subjects affected / exposed occurrences (all)	3 / 75 (4.00%) 3	9 / 74 (12.16%) 12	
Headache subjects affected / exposed occurrences (all)	13 / 75 (17.33%) 13	12 / 74 (16.22%) 14	
Tremor subjects affected / exposed occurrences (all)	10 / 75 (13.33%) 11	7 / 74 (9.46%) 9	
Blood and lymphatic system disorders			
Anaemia subjects affected / exposed occurrences (all)	7 / 75 (9.33%) 7	4 / 74 (5.41%) 4	
Leukopenia			



subjects affected / exposed	29 / 75 (38.67%)	19 / 74 (25.68%)	
occurrences (all)	32	23	
Neutropenia			
subjects affected / exposed	10 / 75 (13.33%)	6 / 74 (8.11%)	
occurrences (all)	11	6	
Thrombocytopenia			
subjects affected / exposed	1 / 75 (1.33%)	5 / 74 (6.76%)	
occurrences (all)	1	6	
Eye disorders			
Vision blurred			
subjects affected / exposed	5 / 75 (6.67%)	1 / 74 (1.35%)	
occurrences (all)	5	1	
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	8 / 75 (10.67%)	6 / 74 (8.11%)	
occurrences (all)	9	7	
Abdominal pain upper			
subjects affected / exposed	5 / 75 (6.67%)	3 / 74 (4.05%)	
occurrences (all)	5	3	
Constipation			
subjects affected / exposed	4 / 75 (5.33%)	6 / 74 (8.11%)	
occurrences (all)	4	7	
Diarrhoea			
subjects affected / exposed	23 / 75 (30.67%)	24 / 74 (32.43%)	
occurrences (all)	33	38	
Dyspepsia			
subjects affected / exposed	5 / 75 (6.67%)	3 / 74 (4.05%)	
occurrences (all)	6	3	
Gastrooesophageal reflux disease			
subjects affected / exposed	1 / 75 (1.33%)	6 / 74 (8.11%)	
occurrences (all)	1	7	
Nausea			
subjects affected / exposed	13 / 75 (17.33%)	14 / 74 (18.92%)	
occurrences (all)	22	19	
Vomiting			

subjects affected / exposed occurrences (all)	9 / 75 (12.00%) 12	9 / 74 (12.16%) 13	
Skin and subcutaneous tissue disorders			
Rash			
subjects affected / exposed	4 / 75 (5.33%)	2 / 74 (2.70%)	
occurrences (all)	4	2	
Alopecia			
subjects affected / exposed	3 / 75 (4.00%)	8 / 74 (10.81%)	
occurrences (all)	3	9	
Renal and urinary disorders			
Dysuria			
subjects affected / exposed	5 / 75 (6.67%)	6 / 74 (8.11%)	
occurrences (all)	5	6	
Haematuria			
subjects affected / exposed	6 / 75 (8.00%)	7 / 74 (9.46%)	
occurrences (all)	8	7	
Kidney fibrosis			
subjects affected / exposed	7 / 75 (9.33%)	4 / 74 (5.41%)	
occurrences (all)	7	4	
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	5 / 75 (6.67%)	8 / 74 (10.81%)	
occurrences (all)	7	10	
Back pain			
subjects affected / exposed	5 / 75 (6.67%)	8 / 74 (10.81%)	
occurrences (all)	6	9	
Muscle spasms			
subjects affected / exposed	3 / 75 (4.00%)	5 / 74 (6.76%)	
occurrences (all)	3	6	
Muscular weakness			
subjects affected / exposed	0 / 75 (0.00%)	4 / 74 (5.41%)	
occurrences (all)	0	4	
Musculoskeletal discomfort			
subjects affected / exposed	4 / 75 (5.33%)	6 / 74 (8.11%)	
occurrences (all)	11	7	
Musculoskeletal pain			

subjects affected / exposed occurrences (all)	4 / 75 (5.33%) 4	2 / 74 (2.70%) 2	
Myalgia subjects affected / exposed occurrences (all)	21 / 75 (28.00%) 38	15 / 74 (20.27%) 22	
Pain in extremity subjects affected / exposed occurrences (all)	4 / 75 (5.33%) 5	8 / 74 (10.81%) 9	
Infections and infestations			
BK virus infection subjects affected / exposed occurrences (all)	16 / 75 (21.33%) 17	16 / 74 (21.62%) 18	
Cytomegalovirus infection subjects affected / exposed occurrences (all)	3 / 75 (4.00%) 4	5 / 74 (6.76%) 7	
Cytomegalovirus syndrome subjects affected / exposed occurrences (all)	7 / 75 (9.33%) 7	6 / 74 (8.11%) 6	
Cytomegalovirus viraemia subjects affected / exposed occurrences (all)	22 / 75 (29.33%) 30	19 / 74 (25.68%) 22	
Escherichia urinary tract infection subjects affected / exposed occurrences (all)	6 / 75 (8.00%) 9	5 / 74 (6.76%) 5	
Nasopharyngitis subjects affected / exposed occurrences (all)	8 / 75 (10.67%) 10	8 / 74 (10.81%) 10	
Upper respiratory tract infection subjects affected / exposed occurrences (all)	8 / 75 (10.67%) 9	7 / 74 (9.46%) 7	
Urinary tract infection subjects affected / exposed occurrences (all)	10 / 75 (13.33%) 17	4 / 74 (5.41%) 5	
Urinary tract infection bacterial subjects affected / exposed occurrences (all)	8 / 75 (10.67%) 11	3 / 74 (4.05%) 5	

Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	3 / 75 (4.00%)	5 / 74 (6.76%)	
occurrences (all)	3	5	
Diabetes mellitus			
subjects affected / exposed	4 / 75 (5.33%)	1 / 74 (1.35%)	
occurrences (all)	4	1	
Hypercalcaemia			
subjects affected / exposed	4 / 75 (5.33%)	5 / 74 (6.76%)	
occurrences (all)	4	5	
Hyperglycaemia			
subjects affected / exposed	2 / 75 (2.67%)	5 / 74 (6.76%)	
occurrences (all)	2	6	
Hyperkalaemia			
subjects affected / exposed	12 / 75 (16.00%)	8 / 74 (10.81%)	
occurrences (all)	16	8	
Hyperlipidaemia			
subjects affected / exposed	4 / 75 (5.33%)	3 / 74 (4.05%)	
occurrences (all)	4	3	
Hypoglycaemia			
subjects affected / exposed	1 / 75 (1.33%)	6 / 74 (8.11%)	
occurrences (all)	1	7	
Hypomagnesaemia			
subjects affected / exposed	7 / 75 (9.33%)	6 / 74 (8.11%)	
occurrences (all)	13	6	
Hyponatraemia			
subjects affected / exposed	5 / 75 (6.67%)	1 / 74 (1.35%)	
occurrences (all)	7	1	
Hypophosphataemia			
subjects affected / exposed	7 / 75 (9.33%)	7 / 74 (9.46%)	
occurrences (all)	8	8	
Metabolic acidosis			
subjects affected / exposed	11 / 75 (14.67%)	4 / 74 (5.41%)	
occurrences (all)	12	4	
Vitamin D deficiency			

subjects affected / exposed	7 / 75 (9.33%)	4 / 74 (5.41%)	
occurrences (all)	7	4	

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
13 August 2013	<p>Key changes in Substantial Amendment 1, dated 13 Aug 2013, are summarized below.</p> <ul style="list-style-type: none"><li>• Patients with graft failure would be discontinued from treatment</li><li>• 1-sided P value (not 2-sided) would be used for testing the hypothesis</li><li>• Urinalysis with microscopic evaluation was added to the safety labs</li><li>• Vital signs for reactogenicity assessments were added</li><li>• Screening laboratory tests could be repeated once</li><li>• Collect creatinine values during the long-term follow-up period, if available</li></ul>
06 January 2014	<p>Key changes in Substantial Amendment 2, dated 06 Jan 2014, are summarized below.</p> <ul style="list-style-type: none"><li>• Screening period could begin -14 days from transplant to day of randomization</li><li>• Redefined CMV viremia as plasma viral load <math>\geq 1000</math> IU/mL</li><li>• Expanded stratification criteria for ATG to day of randomization</li><li>• Revised CMV AVP period of adjustment to after the day of randomization</li><li>• Added vital signs and evaluation of patients 15 minutes after study drug injection</li><li>• Inclusion Criterion 5 - clarified patients received valganciclovir or ganciclovir per regulatory label (package insert)</li><li>• Exclusion Criteria: Excluded patients who required dialysis on day of randomization, Deleted criterion that excluded patients who had an episode of hyperacute or acuterejection prior to Randomization, Excluded patients who received eculizumab, bortezomib, and intravenous immunoglobulin (IVIG) and/or plasmapheresis from day of transplantation through day of randomization, Clarified contraindications to prophylaxis of CMV viremia/disease with valganciclovir and ganciclovir, Clarified a contraindication to an intramuscular injection also included those who were expected to have a contraindication to the injection, Clarified aspartate aminotransferase (AST) or alanine aminotransferase (ALT) criteria was within 3 days prior to randomization, Concomitant medications – included several clarifications (dose adjustments, interruptions to therapy, time periods for prohibited medications, recording) and additional prohibited medications, Spontaneously reported SAEs during long-term follow-up that were possibly or probably related to study drug should be reported.</li><li>• Patients with a temperature <math>\geq 100.4^{\circ}\text{F}</math> should not receive study drug</li><li>• Transplant surgery on day 0 would not be considered an AE or SAE</li></ul>

Notes:

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported