



## Clinical trial results:

**MULTICENTER, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED, PHASE 3 TRIAL OF FULVESTRANT (FASLODEX) WITH OR WITHOUT PD-0332991 (PALBOCICLIB) ±GOSERELIN IN WOMEN WITH HORMONE RECEPTOR-POSITIVE, HER2-NEGATIVE METASTATIC BREAST CANCER WHOSE DISEASE PROGRESSED AFTER PRIOR ENDOCRINE THERAPY**

### Summary

EudraCT number	2013-002580-26
Trial protocol	BE NL IE IT GB DE PT FR
Global end of trial date	28 September 2022

### Results information

Result version number	v2 (current)
This version publication date	02 September 2023
First version publication date	17 March 2016
Version creation reason	

### Trial information

#### Trial identification

Sponsor protocol code	A5481023
-----------------------	----------

#### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT01942135
WHO universal trial number (UTN)	-

Notes:

### Sponsors

Sponsor organisation name	Pfizer Inc.
Sponsor organisation address	235 East 42nd Street,, New York, United States, 10017
Public contact	Pfizer ClinicalTrials.gov Call Center, Pfizer Inc., +1 800 718 1021, ClinicalTrials.gov_Inquiries@pfizer.com
Scientific contact	Pfizer ClinicalTrials.gov Call Center, Pfizer Inc., +1 800 718 1021, ClinicalTrials.gov_Inquiries@pfizer.com

Notes:

### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	28 September 2022
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	28 September 2022
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

To demonstrate the superiority of palbociclib in combination with fulvestrant (with or without goserelin) over fulvestrant (with or without goserelin) along with prolonging investigator-assessed PFS in women with HR-positive/ HER2-negative metastatic breast cancer whose disease had progressed on prior endocrine therapy.

Protection of trial subjects:

The study was conducted in accordance with legal and regulatory requirements, as well as the general principles set forth in the International Ethical Guidelines for Biomedical Research Involving Human Subjects (Council for International Organizations of Medical Sciences 2002), Guidelines for Good Clinical Practice (GCP) (International Conference on Harmonisation [ICH] 1996), and the Declaration of Helsinki (World Medical Association 1996 and 2008). In addition, the study was conducted in accordance with the protocol, the ICH guideline on GCP, and applicable local regulatory requirements and laws.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	26 September 2013
Long term follow-up planned	Yes
Long term follow-up rationale	Safety, Efficacy
Long term follow-up duration	6 Months
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Australia: 32
Country: Number of subjects enrolled	Belgium: 27
Country: Number of subjects enrolled	Canada: 39
Country: Number of subjects enrolled	Germany: 3
Country: Number of subjects enrolled	Ireland: 3
Country: Number of subjects enrolled	Italy: 36
Country: Number of subjects enrolled	Japan: 35
Country: Number of subjects enrolled	Korea, Republic of: 43
Country: Number of subjects enrolled	Netherlands: 9
Country: Number of subjects enrolled	Portugal: 6
Country: Number of subjects enrolled	Romania: 8
Country: Number of subjects enrolled	Russian Federation: 17
Country: Number of subjects enrolled	Taiwan: 4
Country: Number of subjects enrolled	Turkey: 1
Country: Number of subjects enrolled	Ukraine: 44

Country: Number of subjects enrolled	United Kingdom: 13
Country: Number of subjects enrolled	United States: 201
Worldwide total number of subjects	521
EEA total number of subjects	92

Notes:

<b>Subjects enrolled per age group</b>	
In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	392
From 65 to 84 years	126
85 years and over	3

## Subject disposition

### Recruitment

Recruitment details:

The study was conducted at 144 sites in 17 countries that randomized 521 subjects. Eligible subjects were to have histologically or cytologically proven diagnosis of adenocarcinoma of the breast with evidence of recurrent (local or metastatic) disease.

### Pre-assignment

Screening details:

The study consisted of a screening within 28 days before randomization, an active treatment phase, divided in cycles of 28 days each, and a posttreatment follow-up period during which survival and new anti-cancer therapy information was collected every 3 months for the first 9 months, then every 6 months from the last dose of study intervention.

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

Blinding implementation details:

The blinding details was either a manual or electronic process. Blinding codes were only broken in emergency for reasons of participants safety, or if the participant discontinued treatment due to disease progression, as determined by the study physician using Response Evaluation Criteria in Solid Tumors (RECIST) v.1.1 criteria.

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Palbociclib + Fulvestrant

Arm description:

Subjects were administered an initial dose of 125 mg per day orally continuously for 3 weeks followed by 1 week off that can be reduced to 100 mg or 75 mg in case of toxicity; repeated at each subsequent cycle and fulvestrant 500 mg intramuscularly on days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase. Maximum treatment duration was 105 cycles.

Arm type	Experimental
Investigational medicinal product name	Palbociclib
Investigational medicinal product code	PD-0332991
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

Initial dose of 125 mg per day continuously for 3 weeks followed by 1 week off that can be reduced to 100 mg or 75 mg in case of toxicity; repeated at each subsequent cycle and fulvestrant 500 mg intramuscularly on days 1 and 15 of cycle 1 and then every 28 days.

<b>Arm title</b>	Placebo + Fulvestrant
------------------	-----------------------

Arm description:

Subjects were administered placebo orally continuously dosed for 3 weeks followed by 1 week off; repeated at each subsequent cycle and Fulvestrant 500 mg intramuscularly on Days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase. Maximum treatment duration was 93 cycles.

Arm type	Active comparator
----------	-------------------

Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

Placebo orally dosed for 3 weeks continuously followed by 1 week off; repeated at each subsequent cycle and fulvestrant 500 mg intramuscularly on days 1 and 15 of cycle 1 and then every 28 days.

<b>Number of subjects in period 1</b>	<b>Palbociclib + Fulvestrant</b>	<b>Placebo + Fulvestrant</b>
Started	347	174
Treated	345	172
Completed	0	0
Not completed	347	174
Consent withdrawn by subject	4	3
Global deterioration of health status	9	6
Adverse Event	21	7
Randomized not treated	2	2
Death	2	1
Not specified	19	4
Objective progression + progressive disease	279	148
Protocol deviation	1	-
Subject refused to continue treatment	10	3

## Baseline characteristics

### Reporting groups

Reporting group title	Palbociclib + Fulvestrant
Reporting group description:	
Subjects were administered an initial dose of 125 mg per day orally continuously for 3 weeks followed by 1 week off that can be reduced to 100 mg or 75 mg in case of toxicity; repeated at each subsequent cycle and fulvestrant 500 mg intramuscularly on days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase. Maximum treatment duration was 105 cycles.	
Reporting group title	Placebo + Fulvestrant
Reporting group description:	
Subjects were administered placebo orally continuously dosed for 3 weeks followed by 1 week off; repeated at each subsequent cycle and Fulvestrant 500 mg intramuscularly on Days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase. Maximum treatment duration was 93 cycles.	

Reporting group values	Palbociclib + Fulvestrant	Placebo + Fulvestrant	Total
Number of subjects	347	174	521
Age categorical Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	0	0	0
Infants and toddlers (28 days-23 months)	0	0	0
Children (2-11 years)	0	0	0
Adolescents (12-17 years)	0	0	0
Adults (18-64 years)	261	131	392
From 65-84 years	83	43	126
85 years and over	3	0	3
Age Continuous   Units: Years			
arithmetic mean	56.9	56.8	-
standard deviation	± 11.7	± 10.4	-
Gender, Male/Female Units: Subjects			
Female	347	174	521
Male	0	0	0
Menopausal Status Units: Subjects			
Postmenopausal	275	138	413
Pre/perimenopausal	72	36	108
Race Units: Subjects			
White	252	133	385
Black	12	8	20
Asian	74	31	105

Other	8	1	9
Unspecified	1	1	2
Ethnicity			
Units: Subjects			
Hispanic/Latino	17	11	28
Not Hispanic/Latino	329	161	490
Unspecified	1	2	3
Height			
Units: cm			
arithmetic mean	161.1	161.3	
standard deviation	± 7.0	± 7.6	-
Weight			
Units: kg			
arithmetic mean	70.4	72.0	
standard deviation	± 17.5	± 17.6	-

## End points

### End points reporting groups

Reporting group title	Palbociclib + Fulvestrant
-----------------------	---------------------------

Reporting group description:

Subjects were administered an initial dose of 125 mg per day orally continuously for 3 weeks followed by 1 week off that can be reduced to 100 mg or 75 mg in case of toxicity; repeated at each subsequent cycle and fulvestrant 500 mg intramuscularly on days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase. Maximum treatment duration was 105 cycles.

Reporting group title	Placebo + Fulvestrant
-----------------------	-----------------------

Reporting group description:

Subjects were administered placebo orally continuously dosed for 3 weeks followed by 1 week off; repeated at each subsequent cycle and Fulvestrant 500 mg intramuscularly on Days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase. Maximum treatment duration was 93 cycles.

### Primary: Progression-Free Survival (PFS) as assessed by the investigator

End point title	Progression-Free Survival (PFS) as assessed by the investigator
-----------------	---

End point description:

PFS which was defined as the time from the date of randomization to the date of the first documentation of objective progression of disease (PD) or death due to any cause in the absence of documented PD. PFS data were censored on the date of the last tumor assessment on study for subjects who did not have objective tumor progression and who did not die while on study. Subjects lacking an evaluation of tumor response after randomization had their PFS time censored on the date of randomization with the duration of one day. Subjects with documentation of PD or death after a long interval (ie, 2 or more incomplete or non-evaluable assessments) since the last tumor assessment were censored at the time of last objective assessment that did not show PD. The length of PFS was calculated as PFS time (months) = [progression/death date (censor date) - randomization date + 1]/30.4.

End point type	Primary
----------------	---------

End point timeframe:

From randomization date to date of first documentation of progression or death (assessed up to 2 years)

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: months				
median (confidence interval 95%)	11.2 (9.5 to 12.9)	4.6 (3.5 to 5.6)		

### Statistical analyses

Statistical analysis title	Statistical analysis for PFS
----------------------------	------------------------------

Statistical analysis description:

The primary hypothesis to be tested was  $H_0: \lambda \geq 1$  versus.  $H_A: \lambda < 1$ , where  $\lambda$  was the palbociclib plus



fulvestrant: placebo plus fulvestrant hazard ratio (HR). A HR less than 1 indicates a reduction in hazard rate in favor of Palbociclib + Fulvestrant. The study was planned to have 90% power and control the type-I error rate at 0.025.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	521
Analysis specification	Pre-specified
Analysis type	superiority
Method	Stratified Log Rank Test (1-sided)
Parameter estimate	Hazard ratio (HR)
Point estimate	0.497
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.398
upper limit	0.62

## Secondary: Overall Survival (OS) - Number of Subjects Who Died

End point title	Overall Survival (OS) - Number of Subjects Who Died
End point description:	
OS is defined as the time from date of randomization to date of death due to any cause. In the absence of confirmation of death, survival time was censored to last date the subject was known to be alive. For participants lacking survival data beyond the date of their last follow-up, the OS time was censored on the last date they were known to be alive. Subjects lacking survival data beyond randomization were to have their OS times be censored at randomization. The length of OS was calculated as OS time (months) = [death date (censor date) – randomization date + 1]/30.4.	
End point type	Secondary
End point timeframe:	
From randomization until death (up to 4.5 years)	

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: Subjects	201	109		

## Statistical analyses

No statistical analyses for this end point

## Secondary: Overall Survival (OS)

End point title	Overall Survival (OS)
End point description:	
OS is defined as the time from date of randomization to date of death due to any cause. In the absence of confirmation of death, survival time was censored to last date the subject was known to be alive. For participants lacking survival data beyond the date of their last follow-up, the OS time was censored on the last date they were known to be alive. Subjects lacking survival data beyond randomization were to have their OS times be censored at randomization. The length of OS was calculated as OS time	

(months) = [death date (censor date) – randomization date + 1]/30.4.

End point type	Secondary
End point timeframe:	
From randomization until death (up to 4.5 years)	

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: months				
median (confidence interval 95%)	34.9 (28.8 to 40.0)	28.0 (23.6 to 34.6)		

## Statistical analyses

Statistical analysis title	Statistical analysis for OS
Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	521
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0429 <sup>[1]</sup>
Method	Stratified Log Rank Test (1-sided)
Parameter estimate	Hazard ratio (HR)
Point estimate	0.814
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.644
upper limit	1.029

Notes:

[1] - 1-sided p-value from the log-rank test stratified by the presence of visceral metastases and sensitivity to prior endocrine therapy per randomization.

## Secondary: Survival Probabilities at Year 1, Year 2, and Year 3

End point title	Survival Probabilities at Year 1, Year 2, and Year 3
End point description:	
One-, Two- or Three-year Survival Probability is defined as the probability of survival 1 year, 2 or 3 years after the date of randomization based on the Kaplan-Meier estimate. Survival time was censored to last date the subject is known to be alive.	
End point type	Secondary
End point timeframe:	
From randomization until death (assessed up to 36 months)	

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: Survival Probability				
number (confidence interval 95%)				
Survival Probability at Year 1	85.5 (81.3 to 88.9)	84.8 (78.3 to 89.4)		
Survival Probability at Year2	65.3 (59.9 to 70.2)	57.3 (49.2 to 64.6)		
Survival Probability at Year 3	49.6 (44.0 to 54.9)	40.8 (32.9 to 48.5)		

## Statistical analyses

No statistical analyses for this end point

## Secondary: Duration of Response (DR)

End point title	Duration of Response (DR)
End point description:	
DR is defined as the time from the first documentation of objective tumor response (CR or PR) to the first documentation of disease progression or to death due to any cause, whichever occurs first. If tumor progression data included more than 1 date, the first date was used. DR was calculated as [the date response ended (ie, date of PD or death) – first CR or PR date + 1)]/30.4. DR was only be calculated for the subgroup of participants with an objective tumor response. Kaplan-Meier estimate of median of the DR is provided below. No inferential statistical analysis were done for DR.	
End point type	Secondary
End point timeframe:	
From randomization until end of treatment (assessed up to 2 years)	

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: Months				
median (confidence interval 95%)	10.4 (8.3 to 999999)	9.0 (5.6 to 999999)		

## Statistical analyses

No statistical analyses for this end point

## Secondary: Objective Response (OR)

End point title	Objective Response (OR)
End point description:	
OR is defined as the overall complete response (CR) or partial response (PR) according to the Response Evaluation Criteria in Solid Tumors (RECIST version 1.1) Objective Response Rate (ORR) is defined as the proportion of participants with CR or PR relative to (1) all randomized participants and (2)	

randomized participants with measurable disease at baseline. Designation of best response of stable disease (SD) requires the criteria to be met at least 8 weeks after randomization. Participants who do not have on-study radiographic tumor re-evaluation, who received anti-tumor treatment other than the study medication prior to reaching a CR or PR, or who died, progressed, or dropped out for any reason prior to reaching a CR or PR were counted as non-responders in the assessment of ORR. Percentage of participants with confirmed objective tumor response is mentioned below.

End point type	Secondary
----------------	-----------

End point timeframe:

From randomization until end of treatment (assessed up to 2 years)

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: percentage of participants				
number (confidence interval 95%)	21.0 (16.9 to 25.7)	8.6 (4.9 to 13.8)		

## Statistical analyses

Statistical analysis title	Statistical analysis for OR
----------------------------	-----------------------------

Statistical analysis description:

The exact test is testing the null hypothesis that the odds ratio of objective response rate is less than or equal to 1 vs. the alternative hypothesis that the odds ratio of objective response rate is greater than 1. An Odds Ratio >1 means better response in favor of palbociclib plus fulvestrant arm.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	521
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0001 [2]
Method	Exact test (1-sided)
Parameter estimate	Odds ratio (OR)
Point estimate	2.783
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.563
upper limit	5.603

Notes:

[2] - The p-value was not adjusted for multiple comparisons. The priori threshold for statistical significance is 1-sided, alpha=0.025.

## Secondary: Clinical Benefit Response (CBR)

End point title	Clinical Benefit Response (CBR)
-----------------	---------------------------------

End point description:

CBR is defined as the overall complete response (CR), partial response (PR), or stable disease (SD)  $\geq 24$  weeks according to the Response Evaluation Criteria in Solid Tumors (RECIST version 1.1). Clinical Benefit Response Rate (CBRR) is defined as the proportion of participants with CR, PR, or SD  $\geq 24$  weeks relative to (1) all randomized participants and (2) randomized participants with measurable disease at baseline. Designation of best response of SD  $\geq 24$  weeks requires the criteria to be met at least 24 weeks after randomization. Participants who do not have on-study radiographic tumor re-evaluation,

who received anti-tumor treatment other than the study medication prior to reaching a CR or PR, a best response of SD  $\geq$ 24 weeks, or who died, progressed, or dropped out for any reason prior to reaching a CR or PR and a best response of SD  $\geq$ 24 weeks was counted as non-responders in the assessment of CBR.

End point type	Secondary
End point timeframe:	
From randomization until end of treatment (assessed up to 2 years)	

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: percentage of participants				
number (confidence interval 95%)	66.3 (61.0 to 71.2)	39.7 (32.3 to 47.3)		

## Statistical analyses

Statistical analysis title	Statistical analysis of CBR
----------------------------	-----------------------------

Statistical analysis description:

The exact test is testing the null hypothesis that the odds ratio of objective response rate is less than or equal to 1 vs. the alternative hypothesis that the odds ratio of objective response rate is greater than 1. An Odds Ratio  $>1$  means better response in favor of palbociclib plus fulvestrant arm.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	521
Analysis specification	Pre-specified
Analysis type	superiority
P-value	$< 0.0001$ <sup>[3]</sup>
Method	Exact test (1-sided)
Parameter estimate	Odds ratio (OR)
Point estimate	3.016
Confidence interval	
level	95 %
sides	2-sided
lower limit	2.046
upper limit	4.565

Notes:

[3] - The p-value was not adjusted for multiple comparisons. The priori threshold for statistical significance is 1-sided,  $\alpha=0.025$ .

## Secondary: Observed Plasma Trough Concentration (C<sub>trough</sub>) for Palbociclib

End point title	Observed Plasma Trough Concentration (C <sub>trough</sub> ) for Palbociclib <sup>[4]</sup>
-----------------	--

End point description:

C<sub>trough</sub> was defined as steady-state predose concentration. Observed directly from data. For palbociclib, a steady-state trough was to be defined as a predose plasma concentration following at least 8 consecutive days of 125 mg daily dose without dosing interruption and the time window for the PK collection was to be between 22 and 26 hours after the dose (the day prior to PK collection) and no more than 1 hour post-dose on the day of PK collection. The subjects who were treated with Palbociclib + fulvestrant (with or without goserelin) or placebo + fulvestrant (with or without goserelin) and have at least one measured plasma drug concentration. The geometric mean and coefficient of variation was not

estimable for Cycle 1/Day 15 and Cycle 2/Day 15 for the reporting arm placebo plus fulvestrant.

End point type	Secondary
End point timeframe:	
Cycle 1/Day 15 and Cycle 2/Day 15	

Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

End point values	Palbociclib + Fulvestrant			
Subject group type	Reporting group			
Number of subjects analysed	347			
Units: nanograms per millilitre (ng/mL)				
geometric mean (geometric coefficient of variation)				
Cycle 1/Day 15 (N= 165)	70.70 ( $\pm$ 44)			
Cycle 2/Day 15 (N= 160)	75.29 ( $\pm$ 44)			

## Statistical analyses

No statistical analyses for this end point

## Secondary: Ctrough for Fulvestrant

End point title	Ctrough for Fulvestrant
End point description:	
Ctrough was defined as steady-state predose concentration. Observed directly from data. For fulvestrant, a steady-state trough was to be defined when a patient had received all prior planned doses and the sample was collected predose. The 40 subjects who participated in the early safety review, who are treated with palbociclib + fulvestrant $\pm$ goserelin or placebo + fulvestrant $\pm$ goserelin and have at least one measured plasma drug concentration.	
End point type	Secondary
End point timeframe:	
Cycles 2/Day 1 and Cycle 3/Day 1	

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: ng/mL				
geometric mean (geometric coefficient of variation)				
Cycle 2/Day 1 (N= 35, 19)	11.75 ( $\pm$ 41)	9.31 ( $\pm$ 52)		
Cycle 3/Day 1 (N= 29, 14)	9.90 ( $\pm$ 42)	7.60 ( $\pm$ 72)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Ctrough for Goserelin

End point title	Ctrough for Goserelin
-----------------	-----------------------

End point description:

Ctrough was defined as steady-state predose concentration. Observed directly from data. For goserelin, a steady-state trough was to be defined when a patient had received all prior planned doses and the sample was collected predose. The 40 subjects who participated in the early safety review, who are treated with palbociclib + fulvestrant ± goserelin or placebo + fulvestrant ± goserelin and have at least one measured plasma drug concentration.

End point type	Secondary
----------------	-----------

End point timeframe:

Cycles 2/ Day 1 and Cycle 3/ Day 1

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	347	174		
Units: picograms per millilitre (pg/mL)				
geometric mean (geometric coefficient of variation)				
Cycle 2/Day 1 (N= 9, 5)	295.1 (± 153)	302.5 (± 74)		
Cycle 3/Day 1 (N= 7, 3)	344.8 (± 64)	288.5 (± 40)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC QLQ-C30) Functional Scale Scores

End point title	Change from Baseline in European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC QLQ-C30) Functional Scale Scores
-----------------	--

End point description:

The EORTC-QLQ-C30 is a 30-item questionnaire composed of five multi-item functional subscales (physical, role, emotional, cognitive, and social functioning), three multi-item symptom scales (fatigue, nausea/vomiting, and pain), a global quality of life (QOL) subscale, and six single item symptom scales assessing other cancer-related symptoms (dyspnea, sleep disturbance, appetite loss, constipation, diarrhea, and the financial impact of cancer). The questionnaire employs 28 4-point Likert scales with responses from "not at all" to "very much" and two 7-point Likert scales for global health and overall QOL. Responses to all items are then converted to a 0 to 100 scale. For functional and global QOL scales, higher scores represent a better level of functioning/QOL. For symptom-oriented scales, a higher score represents more severe symptoms. A 10-point or higher change in scores from baseline is considered clinically significant.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Day 1 of Cycles 2 to 4, Day 1 of every alternate cycle after that until the end of treatment

<b>End point values</b>	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	335	166		
Units: Units on a scale				
least squares mean (confidence interval 95%)				
Global health status / QoL	-0.9 (-2.5 to 0.7)	-4.0 (-6.3 to -1.7)		
Physical functioning	-0.7 (-2.1 to 0.7)	-1.7 (-3.7 to 0.2)		
Role functioning	-1.8 (-3.7 to 0.1)	-3.7 (-6.5 to -0.9)		
Emotional functioning	2.7 (1.1 to 4.3)	-1.9 (-4.2 to 0.5)		
Cognitive functioning	-1.7 (-3.1 to -0.2)	-2.9 (-5.0 to -0.7)		
Social functioning	-0.5 (-2.5 to 1.5)	-0.6 (-3.4 to 2.3)		

## Statistical analyses

<b>Statistical analysis title</b>	Statistical significance for Global health status
Statistical analysis description:	
Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.	
Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0313 <sup>[5]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	3.1
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.3
upper limit	6

Notes:

[5] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for physical functioning
Statistical analysis description:	
Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.	
Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant



Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.4 <sup>[6]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.4
upper limit	3.5

Notes:

[6] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for role functioning
-----------------------------------	---

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2615 <sup>[7]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	1.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.5
upper limit	5.3

Notes:

[7] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for emotional functioning
-----------------------------------	--

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0016 <sup>[8]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	4.6

Confidence interval	
level	95 %
sides	2-sided
lower limit	1.7
upper limit	7.4

Notes:

[8] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for cognitive functioning
-----------------------------------	--

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.365 <sup>[9]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	1.2

Confidence interval

level	95 %
sides	2-sided
lower limit	-1.4
upper limit	3.8

Notes:

[9] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for social functioning
-----------------------------------	---

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.9615 <sup>[10]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	0.1

Confidence interval

level	95 %
sides	2-sided
lower limit	-3.4
upper limit	3.5

Notes:

[10] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

## Secondary: Change from Baseline in EORTC QLQ-C30 Symptom Scale Scores

End point title	Change from Baseline in EORTC QLQ-C30 Symptom Scale Scores
-----------------	--

**End point description:**

The EORTC-QLQ-C30 is a 30-item questionnaire composed of five multi-item functional subscales (physical, role, emotional, cognitive, and social functioning), three multi-item symptom scales (fatigue, nausea/vomiting, and pain), a global quality of life (QOL) subscale, and six single item symptom scales assessing other cancer-related symptoms (dyspnea, sleep disturbance, appetite loss, constipation, diarrhea, and the financial impact of cancer). The questionnaire employs 28 4-point Likert scales with responses from "not at all" to "very much" and two 7-point Likert scales for global health and overall QOL. Responses to all items are then converted to a 0 to 100 scale. For functional and global QOL scales, higher scores represent a better level of functioning/QOL. For symptom-oriented scales, a higher score represents more severe symptoms. A 10-point or higher change in scores from baseline is considered clinically significant.

End point type	Secondary
----------------	-----------

**End point timeframe:**

Baseline, Day 1 of Cycles 2 to 4, Day 1 of every alternate cycle after that until the end of treatment

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	335	166		
Units: Units on a scale				
least squares mean (confidence interval 95%)				
Fatigue	1.8 (0.1 to 3.5)	3.3 (0.9 to 5.8)		
Nausea and vomiting	1.7 (0.4 to 3.0)	4.2 (2.3 to 6.1)		
Pain	-3.3 (-5.1 to -1.5)	2.0 (-0.6 to 4.6)		
Dyspnoea	2.8 (1.0 to 4.7)	3.3 (0.6 to 6.0)		
Insomnia	-2.4 (-4.4 to -0.4)	-0.4 (-3.3 to 2.5)		
Appetite loss	1.1 (-0.8 to 3.1)	1.7 (-1.1 to 4.6)		
Constipation	3.5 (1.7 to 5.3)	2.8 (0.1 to 5.4)		
Diarrhoea	1.9 (0.6 to 3.1)	2.4 (0.5 to 4.3)		
Financial difficulties	-3.7 (-5.6 to -1.9)	-4.0 (-6.7 to -1.3)		

**Statistical analyses**

Statistical analysis title	Statistical significance for fatigue
----------------------------	--------------------------------------

**Statistical analysis description:**

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.32 <sup>[11]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-1.5

Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.5
upper limit	1.5

Notes:

[11] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for nausea/vomiting
-----------------------------------	--

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0369 <sup>[12]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-2.5

Confidence interval

level	95 %
sides	2-sided
lower limit	-4.8
upper limit	-0.2

Notes:

[12] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for pain
-----------------------------------	-----------------------------------

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0011 <sup>[13]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-5.3

Confidence interval

level	95 %
sides	2-sided
lower limit	-8.5
upper limit	-2.1

Notes:

[13] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for dyspnoea
-----------------------------------	---------------------------------------

**Statistical analysis description:**

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7699 <sup>[14]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-0.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.7
upper limit	2.8

Notes:

[14] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for insomnia
-----------------------------------	---------------------------------------

**Statistical analysis description:**

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2721 <sup>[15]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-5.5
upper limit	1.6

Notes:

[15] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for appetite loss
-----------------------------------	--

**Statistical analysis description:**

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7334 <sup>[16]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-0.6

Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.1
upper limit	2.9

Notes:

[16] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for constipation
-----------------------------------	---

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.6491 <sup>[17]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	0.7

Confidence interval

level	95 %
sides	2-sided
lower limit	-2.5
upper limit	3.9

Notes:

[17] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for diarrhoea
-----------------------------------	--

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.6293 <sup>[18]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-0.6

Confidence interval

level	95 %
sides	2-sided
lower limit	-2.8
upper limit	1.7

Notes:

[18] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for financial difficulty
-----------------------------------	---

**Statistical analysis description:**

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.8812 <sup>[19]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	0.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.1
upper limit	3.6

**Notes:**

[19] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

### Secondary: Change from Baseline in European Organization for Research and Treatment of Cancer Breast Cancer Module (EORTC QLQ BR23) Functional Scale Scores

End point title	Change from Baseline in European Organization for Research and Treatment of Cancer Breast Cancer Module (EORTC QLQ BR23) Functional Scale Scores
-----------------	--

**End point description:**

The EORTC-QLQ-BR23 is a 23-item breast cancer-specific companion module to the EORTC-QLQ-C30 and consists of four functional scales (body image, sexual functioning, sexual enjoyment, future perspective) and four symptom scales (systemic side effects, breast symptoms, arm symptoms, upset by hair loss). QLQ-BR23 questionnaire employs 4-point scales with responses from 'not at all' to 'very much'. All scores are converted to a 0 to 100 scale. For functional scales, higher scores represent a better level of functioning.

End point type	Secondary
----------------	-----------

**End point timeframe:**

Baseline, Day 1 of Cycles 2 to 4, Day 1 of every alternate cycle after that until the end of treatment

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	335	166		
Units: Units on a scale				
least squares mean (confidence interval 95%)				
Body image	1.9 (0.2 to 3.6)	-0.3 (-2.8 to 2.1)		
Sexual functioning	-1.1 (-2.5 to 0.2)	-0.4 (-2.3 to 1.5)		
Sexual enjoyment	-5.2 (-8.3 to -2.1)	-6.6 (-11.6 to -1.7)		
Future perspective	8.1 (5.8 to 10.4)	4.5 (1.2 to 7.9)		

## Statistical analyses

Statistical analysis title	Statistical significance for body image
Statistical analysis description: Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.	
Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.1386 <sup>[20]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	2.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.7
upper limit	5.2

Notes:

[20] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

Statistical analysis title	Statistical significance for future perspective
Statistical analysis description: Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.	
Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0845 <sup>[21]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	3.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.5
upper limit	7.6

Notes:

[21] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

Statistical analysis title	Statistical significance for sexual enjoyment
Statistical analysis description: Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time,	



treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.6271 [22]
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	1.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.4
upper limit	7.3

Notes:

[22] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for sexual functioning
-----------------------------------	---

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.5235 [23]
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-0.8
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.1
upper limit	1.6

Notes:

[23] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

## **Secondary: Change from Baseline in EORTC QLQ BR23 Symptom Scale Scores**

End point title	Change from Baseline in EORTC QLQ BR23 Symptom Scale Scores
-----------------	---

End point description:

The EORTC-QLQ-BR23 is a 23-item breast cancer-specific companion module to the EORTC-QLQ-C30 and consists of four functional scales (body image, sexual functioning, sexual enjoyment, future perspective) and four symptom scales (systemic side effects, breast symptoms, arm symptoms, upset by hair loss). QLQ-BR23 questionnaire employs 4-point scales with responses from 'not at all' to 'very much'. All scores are converted to a 0 to 100 scale. For symptom-oriented scales, a higher score represent more severe symptoms.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Day 1 of Cycles 2 to 4, Day 1 of every alternate cycle after that until the end of treatment

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	335	166		
Units: Units on a scale				
least squares mean (confidence interval 95%)				
Systemic therapy side effects	3.8 (2.6 to 4.9)	3.4 (1.8 to 5.0)		
Breast symptoms	-2.2 (-3.2 to -1.3)	-1.3 (-2.7 to 0.0)		
Arm symptoms	-2.2 (-3.6 to -0.9)	-2.0 (-4.0 to -0.1)		
Upset by hair loss	2.9 (-1.7 to 7.4)	-6.0 (-12.3 to 0.3)		

## Statistical analyses

Statistical analysis title	Statistical significance for systemic therapy
Statistical analysis description:	
Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.	
Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.7273 <sup>[24]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	0.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.6
upper limit	2.3

Notes:

[24] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

Statistical analysis title	Statistical significance for arm symptoms
Statistical analysis description:	
Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.	
Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant

Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.875 <sup>[25]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-0.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.6
upper limit	2.2

Notes:

[25] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for upset by hair loss
-----------------------------------	---

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0255 <sup>[26]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	8.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.1
upper limit	16.6

Notes:

[26] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

<b>Statistical analysis title</b>	Statistical significance for breast symptoms
-----------------------------------	--

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.2671 <sup>[27]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	-0.9

Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.6
upper limit	0.7

Notes:

[27] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

## Secondary: Change from Baseline in EuroQoL 5D (EQ-5D)- Health Index scores

End point title	Change from Baseline in EuroQoL 5D (EQ-5D)- Health Index scores
-----------------	---

End point description:

The EuroQoL-5D (version 3L) is a brief self-administered, validated instrument consisting of 2 parts. The first part consists of 5 descriptors of current health state (mobility, self care, usual activities, pain/discomfort, and anxiety/ depression); a participant is asked to rate each state on a three level scale (1=no problem, 2=some problem, and 3=extreme problem) with higher levels indicating greater severity/ impairment. Published weights are available that allow for the creation of a single summary score called the EQ-5D index, which basically ranges from 0 to 1 with low scores representing a higher level of dysfunction and 1 as perfect health. The second part consists of the EQ-5D general health status as measured by a visual analog scale (EQ-5D VAS). EQ-5D VAS measures the participant's self-rated health status on a scale from 0 (worst imaginable health state) to 100 (best imaginable health state).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Day 1 of Cycles 2 to 4, Day 1 of every alternate cycle after that until the end of treatment

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	335	166		
Units: Units on a scale				
least squares mean (confidence interval 95%)	0.006 (-0.01 to 0.03)	-0.031 (-0.06 to 0.00)		

## Statistical analyses

Statistical analysis title	Statistical Analysis for EQ-5D-Health Index scores
----------------------------	--

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.0308 [28]
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	0.037

Confidence interval	
level	95 %
sides	2-sided
lower limit	0
upper limit	0.07

Notes:

[28] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

## Secondary: Change from Baseline in EQ-5D Visual Analog Scale (VAS) scores scale

End point title	Change from Baseline in EQ-5D Visual Analog Scale (VAS) scores scale
-----------------	--

End point description:

The EuroQol-5D (version 3L) is a brief self-administered, validated instrument consisting of 2 parts. The second part consists of the EQ-5D general health status as measured by a visual analog scale (EQ-5D VAS). EQ-5D VAS measures the participant's self-rated health status on a scale from 0 (worst imaginable health state) to 100 (best imaginable health state).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Day 1 of Cycles 2 to 4, Day 1 of every alternate cycle after that until the end of treatment

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	335	166		
Units: Units on a scale				
least squares mean (confidence interval 95%)	-1.8 (-3.3 to -0.3)	-2.6 (-4.8 to -0.4)		

## Statistical analyses

Statistical analysis title	Statistical Analysis for EQ-5D VAS scores scale
----------------------------	---

Statistical analysis description:

Analysis based on repeated measures mixed-effects model with an intercept term, treatment, time, treatment-by-time, and baseline as covariate.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
Number of subjects included in analysis	501
Analysis specification	Pre-specified
Analysis type	superiority
P-value	= 0.5523 <sup>[29]</sup>
Method	Mixed Model Analysis (2-sided)
Parameter estimate	Mean difference (final values)
Point estimate	0.8
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.9
upper limit	3.5

Notes:

[29] - The priori threshold for statistical significance is 2-sided alpha=0.05. The p-value was not adjusted for multiple comparisons.

## Secondary: Time to Deterioration (TTD)

End point title	Time to Deterioration (TTD)
-----------------	-----------------------------

End point description:

A time to event analysis was pre-specified for pain. An analysis of TTD in pain defined as time between baseline and first occurrence of increase of  $\geq 10$  points in pain. Deterioration will be defined increase in score of 10 points or greater from baseline. The Kaplan-Meier estimates of quartiles (time to deterioration) with 95% CI is mentioned below.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Day 1 of Cycles 2 to 4, Day 1 of every alternate cycle after that until the end of treatment

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	335	166		
Units: Units on a scale				
median (confidence interval 95%)				
25% quartile	1.9 (1.2 to 2.2)	1.0 (1.0 to 1.9)		
50% quartile	8.0 (5.6 to 999999)	2.8 (2.3 to 5.4)		

## Statistical analyses

Statistical analysis title	Statistical Analysis 1 for TTD
----------------------------	--------------------------------

Statistical analysis description:

Treatment with palbociclib plus fulvestrant significantly delayed TTD in pain symptom compared with placebo plus fulvestrant for unstratified analysis.

Comparison groups	Palbociclib + Fulvestrant v Placebo + Fulvestrant
-------------------	---

Number of subjects included in analysis	501
---	-----

Analysis specification	Pre-specified
------------------------	---------------

Analysis type	superiority
---------------	-------------

P-value	< 0.001 <sup>[30]</sup>
---------	-------------------------

Method	Unstratified log-rank test (1-sided)
--------	--------------------------------------

Parameter estimate	Hazard ratio (HR)
--------------------	-------------------

Point estimate	0.642
----------------	-------

Confidence interval

level	95 %
-------	------

sides	2-sided
-------	---------

lower limit	0.487
-------------	-------

upper limit	0.846
-------------	-------

Notes:

[30] - The priori threshold for statistical significance is 1-sided alpha=0.025. The p-value was not adjusted for multiple comparisons.

## Secondary: Percentage of Subjects With Treatment-Emergent Adverse Events

**(TEAEs; All Causalities)**

End point title	Percentage of Subjects With Treatment-Emergent Adverse Events (TEAEs; All Causalities)
-----------------	--

End point description:

An AE is any untoward medical occurrence in a clinical investigation patient administered a product or medical device; the event need not necessarily have a causal relationship with the treatment or usage. An SAE is any untoward medical occurrence at any dose that results in death; is life-threatening; requires hospitalization; results in persistent or significant disability or in congenital anomaly/birth defect. Severity will be graded by the National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE), Version 4.0.

End point type	Secondary
----------------	-----------

End point timeframe:

From the date of randomization up to 28 calendar days ( $\pm 7$  days) after last dose of study intervention (up to 8.4 years).

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	345	172		
Units: percentage of subjects				
number (not applicable)				
With AEs	98.8	93.6		
With SAEs	22.6	19.2		
With Grade 3 or 4 AEs	80.6	26.7		
With Grade 5 AEs	2.3	1.7		
Discontinued palbociclib/placebo due to AEs	7.5	4.7		

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Subjects With Shifts From CTCAE Grade  $\leq 2$  at Baseline to CTCAE Grade 3 or 4 Postbaseline for Hematology Laboratory Results**

End point title	Subjects With Shifts From CTCAE Grade $\leq 2$ at Baseline to CTCAE Grade 3 or 4 Postbaseline for Hematology Laboratory Results
-----------------	---

End point description:

Number of subjects with shifts from Grade  $\leq 2$  at baseline values to post-baseline values (shift to Grade 3 or 4) were reported as per NCI-CTCAE, V4.0 graded from Grade 1 to 5. Grade 1: Mild; asymptomatic/mild symptoms; clinical/diagnostic observations only; intervention not indicated. Grade 2: Moderate; minimal, local/noninvasive intervention indicated. Grade 3: Severe/medically significant but not immediately life-threatening; hospitalization/prolongation of hospitalization indicated. Grade 4: Life-threatening consequences; urgent intervention indicated. Shifts in lab parameter from Grade  $\leq 2$  at baseline to Grade 3 or 4 postbaseline (for parameters Anemia, Hemoglobin increased, Neutrophil count decreased, Platelet count decreased, and White blood cell count decreased) were reported.

End point type	Secondary
----------------	-----------

End point timeframe:

From baseline to end of treatment/withdrawal (up to 4.5 years)

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	343	172		
Units: Subjects				
Anemia	16	3		
Hemoglobin increased	1	0		
Neutrophil count decreased	239	1		
Platelet count decreased	10	0		
White blood cell count decreased	167	0		

## Statistical analyses

No statistical analyses for this end point

## Secondary: Subjects With Shifts From CTCAE Grade ≤2 at Baseline to CTCAE Grade 3 or 4 Postbaseline for Chemistry Laboratory Results

End point title	Subjects With Shifts From CTCAE Grade ≤2 at Baseline to CTCAE Grade 3 or 4 Postbaseline for Chemistry Laboratory Results
-----------------	--

End point description:

Number of subjects with shifts from Grade ≤2 at baseline values to post-baseline values (shift to Grade 3 or 4) were reported as per NCI-CTCAE, V4.0 graded from Grade 1 to 5. Grade 1: Mild; asymptomatic/mild symptoms; clinical/diagnostic observations only; intervention not indicated. Grade 2: Moderate; minimal, local/noninvasive intervention indicated. Grade 3: Severe/medically significant but not immediately life-threatening; hospitalization/prolongation of hospitalization indicated. Grade 4: Life-threatening consequences; urgent intervention indicated. Shifts in lab parameter from Grade ≤2 at baseline to Grade 3 or 4 postbaseline (for parameters ALT increased, ALP increased, AST increased, Blood bilirubin increased, Creatinine increased, Hypercalcemia, Hyperkalemia, Hypermagnesemia, Hyponatremia, Hypoalbuminemia, Hypocalcemia, Hypokalemia, Hypomagnesemia, and Hyponatremia) were reported.

End point type	Secondary
----------------	-----------

End point timeframe:

From baseline to end of treatment/withdrawal (up to 4.5 years)

End point values	Palbociclib + Fulvestrant	Placebo + Fulvestrant		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	343	172		
Units: Subjects				
ALT increased	10	1		
ALP increased	2	2		
AST increased	13	7		
Blood bilirubin increased	2	3		
Creatinine increased	5	0		
Hypercalcemia	1	0		



Hyperkalemia	2	2		
Hypermagnesemia	7	2		
Hypernatremia	0	1		
Hypoalbuminemia	0	1		
Hypocalcemia	2	2		
Hypokalemia	0	0		
Hypomagnesemia	0	0		
Hyponatremia	12	4		

## Statistical analyses

---

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

From the date of randomization up to 28 calendar days ( $\pm 7$  days) after last dose of study intervention (up to 8.4 years).

Adverse event reporting additional description:

The safety analysis set included all subjects who received at least 1 dose of study intervention, with treatment assignments designated according to actual study intervention received.

Assessment type	Non-systematic
-----------------	----------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	25.1
--------------------	------

### Reporting groups

Reporting group title	Placebo + Fulvestrant
-----------------------	-----------------------

Reporting group description:

Subjects were administered placebo orally continuously dosed for 3 weeks followed by 1 week off; repeated at each subsequent cycle and Fulvestrant 500 mg intramuscularly on Days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase.

Reporting group title	Palbociclib + Fulvestrant
-----------------------	---------------------------

Reporting group description:

Subjects were administered an initial dose of 125 mg per day orally continuously for 3 weeks followed by 1 week off that can be reduced to 100 mg or 75 mg in case of toxicity; repeated at each subsequent cycle and fulvestrant 500 mg intramuscularly on days 1 and 15 of cycle 1 and then every 28 days. Pre- and perimenopausal women received goserelin at least 4 weeks before study treatment start and continued receiving concurrent ovarian function suppression with goserelin administered subcutaneously every 28 days during the active treatment phase.

Serious adverse events	Placebo + Fulvestrant	Palbociclib + Fulvestrant	
Total subjects affected by serious adverse events			
subjects affected / exposed	33 / 172 (19.19%)	78 / 345 (22.61%)	
number of deaths (all causes)	5	9	
number of deaths resulting from adverse events	5	9	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Metastases to central			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
Adenocarcinoma gastric			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Breast cancer			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
Colorectal cancer			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Endometrial cancer			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Rectal cancer			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vascular disorders			
Deep vein thrombosis			
subjects affected / exposed	0 / 172 (0.00%)	3 / 345 (0.87%)	
occurrences causally related to treatment / all	0 / 0	2 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
General disorders and administration site conditions			
Disease progression			
subjects affected / exposed	0 / 172 (0.00%)	4 / 345 (1.16%)	
occurrences causally related to treatment / all	0 / 0	0 / 4	
deaths causally related to treatment / all	0 / 1	0 / 5	
General physical health deterioration			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Chest pain			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Pyrexia			
subjects affected / exposed	1 / 172 (0.58%)	5 / 345 (1.45%)	
occurrences causally related to treatment / all	0 / 1	0 / 7	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pain			
subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Oedema peripheral			
subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Non-cardiac chest pain			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Asthenia			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injection site fibrosis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Malaise			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Multiple organ dysfunction syndrome			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 1	
Immune system disorders			

Hypersensitivity			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Reproductive system and breast disorders			
Breast mass			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Respiratory, thoracic and mediastinal disorders			
Acute respiratory distress syndrome			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
Chronic obstructive pulmonary disease			
subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Dyspnoea			
subjects affected / exposed	1 / 172 (0.58%)	2 / 345 (0.58%)	
occurrences causally related to treatment / all	0 / 1	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pleural effusion			
subjects affected / exposed	3 / 172 (1.74%)	2 / 345 (0.58%)	
occurrences causally related to treatment / all	0 / 5	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pulmonary embolism			
subjects affected / exposed	0 / 172 (0.00%)	4 / 345 (1.16%)	
occurrences causally related to treatment / all	0 / 0	0 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pulmonary hypertension			

subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Psychiatric disorders			
Depression			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Psychotic disorder			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Suicide attempt			
subjects affected / exposed	0 / 172 (0.00%)	2 / 345 (0.58%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Product issues			
Device occlusion			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Investigations			
Electrocardiogram QT prolonged			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Neutrophil count decreased			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Troponin increased			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Alanine aminotransferase increased subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood creatinine increased subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injury, poisoning and procedural complications			
Humerus fracture subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Fracture subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Femur fracture subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Road traffic accident subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Sedation complication subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lower limb fracture subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Heat illness			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Fall			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ankle fracture			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Subdural haematoma			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac disorders			
Pericarditis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Atrial fibrillation			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiovascular insufficiency			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Nervous system disorders			
Sciatica			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	



Migraine			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Facial paralysis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Drug withdrawal convulsions			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cerebrovascular accident			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cerebral haemorrhage			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
Carotid artery stenosis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Somnolence			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vocal cord paresis			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Transient ischaemic attack			

subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Paraesthesia			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Facial spasm			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cauda equina syndrome			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood and lymphatic system disorders			
Neutropenia			
subjects affected / exposed	0 / 172 (0.00%)	4 / 345 (1.16%)	
occurrences causally related to treatment / all	0 / 0	4 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
Febrile neutropenia			
subjects affected / exposed	0 / 172 (0.00%)	3 / 345 (0.87%)	
occurrences causally related to treatment / all	0 / 0	3 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Disseminated intravascular coagulation			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Leukopenia			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Eye disorders			

Cataract			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diplopia			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			
Ascites			
subjects affected / exposed	2 / 172 (1.16%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Duodenal ulcer			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Intestinal obstruction			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nausea			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pancreatitis			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vomiting			
subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Abdominal pain			

subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Oesophageal stenosis			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrooesophageal reflux disease			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastric volvulus			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diarrhoea			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Crohn's disease			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Colitis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Small intestinal obstruction			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hiatus hernia, obstructive			

subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatobiliary disorders			
Hepatic failure			
subjects affected / exposed	0 / 172 (0.00%)	2 / 345 (0.58%)	
occurrences causally related to treatment / all	0 / 0	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 1	
Cholelithiasis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cholecystitis			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bile duct stone			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin and subcutaneous tissue disorders			
Rash maculo-papular			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin disorder			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Endocrine disorders			
Hyperthyroidism			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Hypopituitarism			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Hypothyroidism			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pathological fracture			
subjects affected / exposed	2 / 172 (1.16%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pain in extremity			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Osteonecrosis of jaw			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Muscular weakness			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
Erysipelas			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Cellulitis			
subjects affected / exposed	0 / 172 (0.00%)	2 / 345 (0.58%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal infection			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract infection			
subjects affected / exposed	1 / 172 (0.58%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Upper respiratory tract infection			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyelonephritis			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia			
subjects affected / exposed	2 / 172 (1.16%)	3 / 345 (0.87%)	
occurrences causally related to treatment / all	0 / 3	0 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pharyngitis			
subjects affected / exposed	0 / 172 (0.00%)	2 / 345 (0.58%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Otitis media acute			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
COVID-19			

subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Appendicitis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Atypical pneumonia			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bacteraemia			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Neutropenic sepsis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 1	
Nasopharyngitis			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Meningitis aseptic			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lower respiratory tract infection			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Influenza			



subjects affected / exposed	0 / 172 (0.00%)	2 / 345 (0.58%)	
occurrences causally related to treatment / all	0 / 0	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infectious pleural effusion			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatitis C			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Escherichia sepsis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Sepsis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Viral upper respiratory tract infection			
subjects affected / exposed	1 / 172 (0.58%)	0 / 345 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Viral infection			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urosepsis			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Sinusitis			

subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Metabolism and nutrition disorders</b>			
Dehydration			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypokalaemia			
subjects affected / exposed	0 / 172 (0.00%)	1 / 345 (0.29%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	Placebo + Fulvestrant	Palbociclib + Fulvestrant	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	161 / 172 (93.60%)	341 / 345 (98.84%)	
<b>Vascular disorders</b>			
Hot flush			
subjects affected / exposed	30 / 172 (17.44%)	57 / 345 (16.52%)	
occurrences (all)	32	85	
Hypertension			
subjects affected / exposed	6 / 172 (3.49%)	26 / 345 (7.54%)	
occurrences (all)	7	52	
<b>General disorders and administration site conditions</b>			
Asthenia			
subjects affected / exposed	14 / 172 (8.14%)	27 / 345 (7.83%)	
occurrences (all)	17	38	
Chest pain			
subjects affected / exposed	11 / 172 (6.40%)	16 / 345 (4.64%)	
occurrences (all)	16	19	
Fatigue			
subjects affected / exposed	56 / 172 (32.56%)	151 / 345 (43.77%)	
occurrences (all)	89	301	

Injection site pain subjects affected / exposed occurrences (all)	19 / 172 (11.05%) 25	26 / 345 (7.54%) 40	
Oedema peripheral subjects affected / exposed occurrences (all)	13 / 172 (7.56%) 13	36 / 345 (10.43%) 44	
Pain subjects affected / exposed occurrences (all)	14 / 172 (8.14%) 16	20 / 345 (5.80%) 28	
Pyrexia subjects affected / exposed occurrences (all)	9 / 172 (5.23%) 11	44 / 345 (12.75%) 64	
Influenza like illness subjects affected / exposed occurrences (all)	5 / 172 (2.91%) 8	34 / 345 (9.86%) 62	
Respiratory, thoracic and mediastinal disorders Cough subjects affected / exposed occurrences (all)	23 / 172 (13.37%) 39	79 / 345 (22.90%) 97	
Dyspnoea subjects affected / exposed occurrences (all)	15 / 172 (8.72%) 17	50 / 345 (14.49%) 67	
Epistaxis subjects affected / exposed occurrences (all)	4 / 172 (2.33%) 4	25 / 345 (7.25%) 27	
Oropharyngeal pain subjects affected / exposed occurrences (all)	14 / 172 (8.14%) 18	53 / 345 (15.36%) 63	
Psychiatric disorders Anxiety subjects affected / exposed occurrences (all)	10 / 172 (5.81%) 13	20 / 345 (5.80%) 25	
Depression subjects affected / exposed occurrences (all)	10 / 172 (5.81%) 11	27 / 345 (7.83%) 36	
Insomnia			

subjects affected / exposed occurrences (all)	17 / 172 (9.88%) 18	46 / 345 (13.33%) 78	
Investigations			
Neutrophil count decreased subjects affected / exposed occurrences (all)	2 / 172 (1.16%) 6	84 / 345 (24.35%) 667	
Platelet count decreased subjects affected / exposed occurrences (all)	0 / 172 (0.00%) 0	43 / 345 (12.46%) 126	
White blood cell count decreased subjects affected / exposed occurrences (all)	7 / 172 (4.07%) 10	107 / 345 (31.01%) 581	
Aspartate aminotransferase increased subjects affected / exposed occurrences (all)	13 / 172 (7.56%) 21	42 / 345 (12.17%) 66	
Alanine aminotransferase increased subjects affected / exposed occurrences (all)	9 / 172 (5.23%) 14	32 / 345 (9.28%) 55	
Blood alkaline phosphatase increased subjects affected / exposed occurrences (all)	11 / 172 (6.40%) 14	13 / 345 (3.77%) 14	
Blood creatinine increased subjects affected / exposed occurrences (all)	3 / 172 (1.74%) 4	20 / 345 (5.80%) 36	
Weight decreased subjects affected / exposed occurrences (all)	5 / 172 (2.91%) 7	18 / 345 (5.22%) 28	
Injury, poisoning and procedural complications			
Contusion subjects affected / exposed occurrences (all)	4 / 172 (2.33%) 4	20 / 345 (5.80%) 21	
Fall subjects affected / exposed occurrences (all)	13 / 172 (7.56%) 14	18 / 345 (5.22%) 23	
Nervous system disorders			

Dizziness subjects affected / exposed occurrences (all)	18 / 172 (10.47%) 21	59 / 345 (17.10%) 101	
Dysgeusia subjects affected / exposed occurrences (all)	5 / 172 (2.91%) 6	18 / 345 (5.22%) 24	
Headache subjects affected / exposed occurrences (all)	38 / 172 (22.09%) 57	102 / 345 (29.57%) 182	
Paraesthesia subjects affected / exposed occurrences (all)	5 / 172 (2.91%) 5	19 / 345 (5.51%) 26	
Blood and lymphatic system disorders			
Anaemia subjects affected / exposed occurrences (all)	23 / 172 (13.37%) 39	108 / 345 (31.30%) 325	
Leukopenia subjects affected / exposed occurrences (all)	2 / 172 (1.16%) 9	113 / 345 (32.75%) 487	
Neutropenia subjects affected / exposed occurrences (all)	4 / 172 (2.33%) 10	231 / 345 (66.96%) 1689	
Thrombocytopenia subjects affected / exposed occurrences (all)	0 / 172 (0.00%) 0	51 / 345 (14.78%) 210	
Eye disorders			
Lacrimation increased subjects affected / exposed occurrences (all)	2 / 172 (1.16%) 2	27 / 345 (7.83%) 44	
Vision blurred subjects affected / exposed occurrences (all)	3 / 172 (1.74%) 3	23 / 345 (6.67%) 31	
Gastrointestinal disorders			
Diarrhoea subjects affected / exposed occurrences (all)	35 / 172 (20.35%) 54	95 / 345 (27.54%) 242	
Abdominal pain			

subjects affected / exposed	13 / 172 (7.56%)	31 / 345 (8.99%)	
occurrences (all)	17	43	
Abdominal pain upper			
subjects affected / exposed	15 / 172 (8.72%)	24 / 345 (6.96%)	
occurrences (all)	19	33	
Constipation			
subjects affected / exposed	29 / 172 (16.86%)	77 / 345 (22.32%)	
occurrences (all)	34	107	
Dry mouth			
subjects affected / exposed	16 / 172 (9.30%)	27 / 345 (7.83%)	
occurrences (all)	18	28	
Dyspepsia			
subjects affected / exposed	9 / 172 (5.23%)	41 / 345 (11.88%)	
occurrences (all)	11	58	
Nausea			
subjects affected / exposed	51 / 172 (29.65%)	126 / 345 (36.52%)	
occurrences (all)	62	221	
Stomatitis			
subjects affected / exposed	5 / 172 (2.91%)	51 / 345 (14.78%)	
occurrences (all)	6	140	
Vomiting			
subjects affected / exposed	27 / 172 (15.70%)	76 / 345 (22.03%)	
occurrences (all)	40	150	
Abdominal distension			
subjects affected / exposed	9 / 172 (5.23%)	21 / 345 (6.09%)	
occurrences (all)	12	27	
Gastrooesophageal reflux disease			
subjects affected / exposed	3 / 172 (1.74%)	23 / 345 (6.67%)	
occurrences (all)	3	29	
Skin and subcutaneous tissue disorders			
Alopecia			
subjects affected / exposed	11 / 172 (6.40%)	68 / 345 (19.71%)	
occurrences (all)	11	73	
Pruritus			
subjects affected / exposed	14 / 172 (8.14%)	32 / 345 (9.28%)	
occurrences (all)	17	43	

Rash			
subjects affected / exposed	8 / 172 (4.65%)	46 / 345 (13.33%)	
occurrences (all)	9	61	
Dry skin			
subjects affected / exposed	3 / 172 (1.74%)	29 / 345 (8.41%)	
occurrences (all)	4	32	
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	42 / 172 (24.42%)	90 / 345 (26.09%)	
occurrences (all)	61	186	
Back pain			
subjects affected / exposed	32 / 172 (18.60%)	65 / 345 (18.84%)	
occurrences (all)	46	97	
Muscle spasms			
subjects affected / exposed	12 / 172 (6.98%)	35 / 345 (10.14%)	
occurrences (all)	15	56	
Musculoskeletal chest pain			
subjects affected / exposed	11 / 172 (6.40%)	21 / 345 (6.09%)	
occurrences (all)	14	27	
Myalgia			
subjects affected / exposed	15 / 172 (8.72%)	35 / 345 (10.14%)	
occurrences (all)	18	51	
Pain in extremity			
subjects affected / exposed	27 / 172 (15.70%)	62 / 345 (17.97%)	
occurrences (all)	35	89	
Bone pain			
subjects affected / exposed	9 / 172 (5.23%)	28 / 345 (8.12%)	
occurrences (all)	11	45	
Neck pain			
subjects affected / exposed	8 / 172 (4.65%)	18 / 345 (5.22%)	
occurrences (all)	8	20	
Infections and infestations			
Nasopharyngitis			
subjects affected / exposed	14 / 172 (8.14%)	50 / 345 (14.49%)	
occurrences (all)	24	81	
Upper respiratory tract infection			

subjects affected / exposed occurrences (all)	13 / 172 (7.56%) 23	46 / 345 (13.33%) 67	
Influenza subjects affected / exposed occurrences (all)	10 / 172 (5.81%) 12	12 / 345 (3.48%) 19	
Urinary tract infection subjects affected / exposed occurrences (all)	14 / 172 (8.14%) 19	38 / 345 (11.01%) 57	
Metabolism and nutrition disorders Decreased appetite subjects affected / exposed occurrences (all)	18 / 172 (10.47%) 22	61 / 345 (17.68%) 86	



## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
01 April 2014	In protocol amendment 1, schedule of assessments was updated with ocular safety assessments procedures to newly enrolled lens grading evaluable participants to assess the potential risk of palbociclib-associated lens changes. In human pharmacokinetic data, included preliminary results from two clinical pharmacology studies of palbociclib. Clarified inclusion criteria #6 and exclusion criteria #4, 5 and 6 to address frequent questions from investigational sites. Added prohibition to take proton-pump inhibitors while receiving study drug. Clarified that routine safety assessments must continue if patient continues study treatment despite progression of disease.
30 September 2014	In protocol amendment 2, In schedule of activities, laboratory safety assessments: added prospective monitoring of hemoglobin A1c to characterize whether or not palbociclib affects glucose metabolism. Added clarification to the schedule of administration of fulvestrant and goserelin. Editorial changes made to differentiate between strong and moderate CYP3A nducers/inhibitors. Added clarification of adverse events follow-up procedure (telephone contact) at 28 calendar days after treatment discontinuation. Additional editorial changes made to clarify follow-up procedures for patients who discontinue treatment for reasons other than disease progression and for patients who discontinue treatment due to disease progression.
20 October 2015	In protocol amendment 3, in schedule of activities, added clarification that tumor assessments will be conducted as per local practice. In study design section, added clarification that the third-party core imaging lab will not performed any longer blinded independent central review of PFS data. The investigators will stop sending the imaging studies to the central lab. Added clarification that survival follow-up visists will be conducted every 3 months after approval of amendment 3.

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported