

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
All Strata Combined			
RBC Transfusion Dependent Rate at Week 24			
Not-Dependent, n(%)	52 ( 50.0%)	19 ( 36.5%)	71 ( 45.5%)
Transfusion Requiring, n(%)	7 ( 6.7%)	8 ( 15.4%)	15 ( 9.6%)
Transfusion Independent, n(%)	45 ( 43.3%)	11 ( 21.2%)	56 ( 35.9%)
Dependent, n(%)	52 ( 50.0%)	33 ( 63.5%)	85 ( 54.5%)
95% Exact CI	0.4003, 0.5997	0.4896, 0.7638	0.4633, 0.6247
Proportion Difference - Stratified CMH Method (95% CI)	-0.13 (-0.29, 0.03)		
p-value	0.10		
Proportion Difference - Unstratified Method (95% CI)	-0.13 (-0.30, 0.03)		
p-value	0.11		
Proportion Difference - Unstratified Exact Method (95% CI)	-0.13 (-0.30, 0.04)		
p-value	0.13		
>=4 units transfused in the last 8 weeks	15 ( 14.4%)	17 ( 32.7%)	32 ( 20.5%)
Any Hgb assessment < 8g/dL in the last 8 weeks	21 ( 20.2%)	16 ( 30.8%)	37 ( 23.7%)
Last Participation date < Day 162 in RT phase	28 ( 26.9%)	10 ( 19.2%)	38 ( 24.4%)
Other	10 ( 9.6%)	10 ( 19.2%)	20 ( 12.8%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>RBC Transfusion Dependent Rate at Week 12</b>			
Not-Dependent, n(%)	51 ( 49.0%)	20 ( 38.5%)	71 ( 45.5%)
Transfusion Requiring, n(%)	11 ( 10.6%)	11 ( 21.2%)	22 ( 14.1%)
Transfusion Independent, n(%)	40 ( 38.5%)	9 ( 17.3%)	49 ( 31.4%)
Dependent, n(%)	53 ( 51.0%)	32 ( 61.5%)	85 ( 54.5%)
95% Exact CI	0.4097, 0.6090	0.4702, 0.7470	0.4633, 0.6247
Proportion Difference - Stratified CMH Method (95% CI)	-0.11 (-0.27, 0.04)		
p-value	0.15		
Proportion Difference - Unstratified Method (95% CI)	-0.11 (-0.27, 0.06)		
p-value	0.21		
Proportion Difference - Unstratified Exact Method (95% CI)	-0.11 (-0.27, 0.07)		
p-value	0.24		
>=4 units transfused in the last 8 weeks	28 ( 26.9%)	15 ( 28.8%)	43 ( 27.6%)
Any Hgb assessment < 8g/dL in the last 8 weeks	34 ( 32.7%)	18 ( 34.6%)	52 ( 33.3%)
Last Participation date < Day 78 in RT phase	14 ( 13.5%)	9 ( 17.3%)	23 ( 14.7%)
Other	14 ( 13.5%)	10 ( 19.2%)	24 ( 15.4%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>Stratum 1</b>			
Transfusion Dependence Yes and Baseline TSS <18	37	15	52
<b>RBC Transfusion Dependent Rate at Week 24</b>			
Not-Dependent, n(%)	18 ( 48.6%)	5 ( 33.3%)	23 ( 44.2%)
Transfusion Requiring, n(%)	4 ( 10.8%)	5 ( 33.3%)	9 ( 17.3%)
Transfusion Independent, n(%)	14 ( 37.8%)	0	14 ( 26.9%)
Dependent, n(%)	19 ( 51.4%)	10 ( 66.7%)	29 ( 55.8%)
95% Exact CI	0.3440, 0.6808	0.3838, 0.8818	0.4133, 0.6953
Proportion Difference (95% CI)	-0.15 (-0.45, 0.14)		
Proportion Difference using Exact Method (95% CI)	-0.15 (-0.43, 0.15)		
>=4 units transfused in the last 8 weeks	6 ( 16.2%)	5 ( 33.3%)	11 ( 21.2%)
Any Hgb assessment < 8g/dL in the last 8 weeks	12 ( 32.4%)	6 ( 40.0%)	18 ( 34.6%)
Last Participation date < Day 162 in RT phase	7 ( 18.9%)	3 ( 20.0%)	10 ( 19.2%)
Other	6 ( 16.2%)	5 ( 33.3%)	11 ( 21.2%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>RBC Transfusion Dependent Rate at Week 12</b>			
Not-Dependent, n(%)	14 ( 37.8%)	4 ( 26.7%)	18 ( 34.6%)
Transfusion Requiring, n(%)	7 ( 18.9%)	4 ( 26.7%)	11 ( 21.2%)
Transfusion Independent, n(%)	7 ( 18.9%)	0	7 ( 13.5%)
Dependent, n(%)	23 ( 62.2%)	11 ( 73.3%)	34 ( 65.4%)
95% Exact CI	0.4476, 0.7754	0.4490, 0.9221	0.5091, 0.7803
Proportion Difference (95% CI)	-0.11(-0.40, 0.17)		
Proportion Difference using Exact Method (95% CI)	-0.11 (-0.40, 0.19)		
>=4 units transfused in the last 8 weeks	15 ( 40.5%)	6 ( 40.0%)	21 ( 40.4%)
Any Hgb assessment < 8g/dL in the last 8 weeks	21 ( 56.8%)	6 ( 40.0%)	27 ( 51.9%)
Last Participation date < Day 78 in RT phase	1 ( 2.7%)	3 ( 20.0%)	4 ( 7.7%)
Other	9 ( 24.3%)	5 ( 33.3%)	14 ( 26.9%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>Stratum 2</b>			
Transfusion Dependence Yes and Baseline TSS >=18	21	12	33
<b>RBC Transfusion Dependent Rate at Week 24</b>			
Not-Dependent, n(%)	7 ( 33.3%)	1 ( 8.3%)	8 ( 24.2%)
Transfusion Requiring, n(%)	2 ( 9.5%)	0	2 ( 6.1%)
Transfusion Independent, n(%)	5 ( 23.8%)	1 ( 8.3%)	6 ( 18.2%)
Dependent, n(%)	14 ( 66.7%)	11 ( 91.7%)	25 ( 75.8%)
95% Exact CI	0.4303, 0.8541	0.6152, 0.9979	0.5774, 0.8891
Proportion Difference (95% CI)	-0.25 (-0.53, 0.03)		
Proportion Difference using Exact Method (95% CI)	-0.25 (-0.57, 0.11)		
>=4 units transfused in the last 8 weeks	5 ( 23.8%)	7 ( 58.3%)	12 ( 36.4%)
Any Hgb assessment < 8g/dL in the last 8 weeks	5 ( 23.8%)	5 ( 41.7%)	10 ( 30.3%)
Last Participation date < Day 162 in RT phase	8 ( 38.1%)	3 ( 25.0%)	11 ( 33.3%)
Other	1 ( 4.8%)	2 ( 16.7%)	3 ( 9.1%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>RBC Transfusion Dependent Rate at Week 12</b>			
Not-Dependent, n(%)	5 ( 23.8%)	1 ( 8.3%)	6 ( 18.2%)
Transfusion Requiring, n(%)	0	1 ( 8.3%)	1 ( 3.0%)
Transfusion Independent, n(%)	5 ( 23.8%)	0	5 ( 15.2%)
Dependent, n(%)	16 ( 76.2%)	11 ( 91.7%)	27 ( 81.8%)
95% Exact CI	0.5283, 0.9178	0.6152, 0.9979	0.6454, 0.9302
Proportion Difference (95% CI)	-0.15(-0.42, 0.11)		
Proportion Difference using Exact Method (95% CI)	-0.15 (-0.49, 0.21)		
>=4 units transfused in the last 8 weeks	8 ( 38.1%)	5 ( 41.7%)	13 ( 39.4%)
Any Hgb assessment < 8g/dL in the last 8 weeks	9 ( 42.9%)	7 ( 58.3%)	16 ( 48.5%)
Last Participation date < Day 78 in RT phase	5 ( 23.8%)	3 ( 25.0%)	8 ( 24.2%)
Other	4 ( 19.0%)	3 ( 25.0%)	7 ( 21.2%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>Stratum 3</b>			
Transfusion Dependence No and Baseline TSS <18	24	13	37
<b>RBC Transfusion Dependent Rate at Week 24</b>			
Not-Dependent, n(%)	18 ( 75.0%)	7 ( 53.8%)	25 ( 67.6%)
Transfusion Requiring, n(%)	0	0	0
Transfusion Independent, n(%)	18 ( 75.0%)	7 ( 53.8%)	25 ( 67.6%)
Dependent, n(%)	6 ( 25.0%)	6 ( 46.2%)	12 ( 32.4%)
95% Exact CI	0.0977, 0.4671	0.1922, 0.7487	0.1801, 0.4979
Proportion Difference (95% CI)	-0.21 (-0.55, 0.12)		
Proportion Difference using Exact Method (95% CI)	-0.21 (-0.53, 0.12)		
>=4 units transfused in the last 8 weeks	0	3 ( 23.1%)	3 ( 8.1%)
Any Hgb assessment < 8g/dL in the last 8 weeks	1 ( 4.2%)	3 ( 23.1%)	4 ( 10.8%)
Last Participation date < Day 162 in RT phase	5 ( 20.8%)	1 ( 7.7%)	6 ( 16.2%)
Other	1 ( 4.2%)	3 ( 23.1%)	4 ( 10.8%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>RBC Transfusion Dependent Rate at Week 12</b>			
Not-Dependent, n(%)	18 ( 75.0%)	8 ( 61.5%)	26 ( 70.3%)
Transfusion Requiring, n(%)	2 ( 8.3%)	2 ( 15.4%)	4 ( 10.8%)
Transfusion Independent, n(%)	16 ( 66.7%)	6 ( 46.2%)	22 ( 59.5%)
Dependent, n(%)	6 ( 25.0%)	5 ( 38.5%)	11 ( 29.7%)
95% Exact CI	0.0977, 0.4671	0.1386, 0.6842	0.1587, 0.4698
Proportion Difference (95% CI)	-0.13(-0.46, 0.19)		
Proportion Difference using Exact Method (95% CI)	-0.13 (-0.46, 0.20)		
>=4 units transfused in the last 8 weeks	2 ( 8.3%)	2 ( 15.4%)	4 ( 10.8%)
Any Hgb assessment < 8g/dL in the last 8 weeks	2 ( 8.3%)	4 ( 30.8%)	6 ( 16.2%)
Last Participation date < Day 78 in RT phase	3 ( 12.5%)	0	3 ( 8.1%)
Other	1 ( 4.2%)	2 ( 15.4%)	3 ( 8.1%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>Stratum 4</b>			
Transfusion Dependence No and Baseline TSS >=18	22	12	34
<b>RBC Transfusion Dependent Rate at Week 24</b>			
Not-Dependent, n(%)	9 ( 40.9%)	6 ( 50.0%)	15 ( 44.1%)
Transfusion Requiring, n(%)	1 ( 4.5%)	3 ( 25.0%)	4 ( 11.8%)
Transfusion Independent, n(%)	8 ( 36.4%)	3 ( 25.0%)	11 ( 32.4%)
Dependent, n(%)	13 ( 59.1%)	6 ( 50.0%)	19 ( 55.9%)
95% Exact CI	0.3635, 0.7929	0.2109, 0.7891	0.3789, 0.7281
Proportion Difference (95% CI)	0.09 (-0.27, 0.45)		
Proportion Difference using Exact Method (95% CI)	0.09 (-0.26, 0.43)		
>=4 units transfused in the last 8 weeks	4 ( 18.2%)	2 ( 16.7%)	6 ( 17.6%)
Any Hgb assessment < 8g/dL in the last 8 weeks	3 ( 13.6%)	2 ( 16.7%)	5 ( 14.7%)
Last Participation date < Day 162 in RT phase	8 ( 36.4%)	3 ( 25.0%)	11 ( 32.4%)
Other	2 ( 9.1%)	0	2 ( 5.9%)

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46

Table 15.12.5.1: Analysis of RBC Transfusion Dependence Rate at Week 24 and at Week 12  
Randomized Treatment Phase  
Intent-to-Treat Analysis Set

	MMB (N=104)	BAT (N=52)	Total (N=156)
<b>RBC Transfusion Dependent Rate at Week 12</b>			
Not-Dependent, n(%)	14 ( 63.6%)	7 ( 58.3%)	21 ( 61.8%)
Transfusion Requiring, n(%)	2 ( 9.1%)	4 ( 33.3%)	6 ( 17.6%)
Transfusion Independent, n(%)	12 ( 54.5%)	3 ( 25.0%)	15 ( 44.1%)
Dependent, n(%)	8 ( 36.4%)	5 ( 41.7%)	13 ( 38.2%)
95% Exact CI	0.1720, 0.5934	0.1517, 0.7233	0.2217, 0.5644
Proportion Difference (95% CI)	-0.05(-0.41, 0.30)		
Proportion Difference using Exact Method (95% CI)	-0.05 (-0.40, 0.29)		
>=4 units transfused in the last 8 weeks	3 ( 13.6%)	2 ( 16.7%)	5 ( 14.7%)
Any Hgb assessment < 8g/dL in the last 8 weeks	2 ( 9.1%)	1 ( 8.3%)	3 ( 8.8%)
Last Participation date < Day 78 in RT phase	5 ( 22.7%)	3 ( 25.0%)	8 ( 23.5%)
Other	0	0	0

95% Exact CI is based on Clopper-Pearson method without stratification

Transfusion requiring is defined as being neither transfusion independent nor transfusion dependent

CMH = Cochran-Mantel-Haenszel;

RBC = Red Blood Cell

Source: Listing 16.3.3.2

Data Extracted: CRF data: 25JUN2019

Source: ...\\version4\prog\t-tf-td.sas v9.4 Output file: t-rbctd24.out 27AUG2020:20:46