

Proportion of patients with need for transfusions and number of units transfused

There was no significant difference between the treatment arms in the proportion of patients requiring transfusions and in the number of transfused units. In the FAS population, a median number of 6 units of PC and 7 units of pRBC were transfused within the first 6 months, see.

Number of units transfused within the first 6 months (FAS)

Parameter		Eltrombopag	Placebo	Total
Any	N	29	31	60
	Arithmetic mean / STD	15.5 / 17.4	17.9 / 16.9	16.7 / 17.1
	Median (IQR)	8.0 (5.0-17.0)	11.0 (4.5-32.0)	9.5 (5.0-25.0)
	Range	1 - 77	1 - 62	1 - 77
PC	N	25	20	45
	Arithmetic mean / STD	9.2 / 11.0	14.0 / 12.6	11.3 / 11.9
	Median (IQR)	4.0 (3.0-12.0)	12.0 (3.2-20.2)	6.0 (3.0-17.0)
	Range	1 - 43	1 - 43	1 - 43
pRBC	N	26	27	53
	Arithmetic mean / STD	8.5 / 8.6	10.2 / 6.9	9.3 / 7.7
	Median (IQR)	6.0 (2.0-10.8)	9.0 (5.5-15.0)	7.0 (2.0-14.0)
	Range	1 - 39	1 - 28	1 - 39