

# Time to best trileanage response

Time to... (months)		Eltrombopag	Placebo	Total
Best trilineage response	N	30	31	61
	Arithmetic mean / STD	9.35 / 8.21	10.62 / 6.88	10.00 / 7.53
	Median (IQR)	6.95 (2.89-11.20)	9.20 (8.28-11.15)	8.31 (5.49-11.24)
	Range	2.6 – 36.8	2.7 – 38.6	2.6 – 38.6
Erythroid response	N	26	32	58
	Arithmetic mean / STD	8.47 / 7.99	8.58 / 7.39	8.53 / 7.59
	Median (IQR)	5.59 (2.81-8.53)	8.28 (2.82-9.43)	7.08 (2.80-9.37)
	Range	2.8 – 35.6	2.6 – 38.6	2.6 – 38.6
Platelet response	N	28	33	61
	Arithmetic mean / STD	5.32 / 4.29	6.89 / 4.17	6.17 / 4.26
	Median (IQR)	2.92 (2.76-5.72)	5.68 (2.79-9.43)	5.52 (2.76-8.97)
	Range	2.7 – 22.1	2.5 – 21.8	2.5 – 22.1
Neutrophil response	N	32	34	66
	Arithmetic mean / STD	4.13 / 3.62	6.51 / 4.74	5.36 / 4.37
	Median (IQR)	2.79 (2.76-3.99)	5.52 (2.77-8.33)	2.91 (2.76-5.69)
	Range	2.6 – 22.4	2.5 – 21.8	2.5 – 22.4

The median time to best trilineage response for patients with response in the FAS population was 8.31 months. The median time to reach single lineage response was 7.08 months (erythroid), 5.52 months (platelet) and 2.91 months (neutrophil). A significant difference between the Placebo and Eltrombopag Arm was observed only for neutrophil response (p=0.025),