



Clinical trial results:

The exogenous progesterone free luteal phase after GnRHa trigger – a randomized controlled pilot study in high-responder IVF patients

Summary

| | |
|--------------------------|-----------------|
| EudraCT number | 2014-000448-13 |
| Trial protocol | DK |
| Global end of trial date | 15 October 2019 |

Results information

| | |
|-----------------------------------|--|
| Result version number | v1 (current) |
| This version publication date | 17 December 2020 |
| First version publication date | 17 December 2020 |
| Summary attachment (see zip file) | Summary of outcomes (Tables_Eudract_upload.docx) |

Trial information

Trial identification

| | |
|-----------------------|----------|
| Sponsor protocol code | Agonist6 |
|-----------------------|----------|

Additional study identifiers

| | |
|------------------------------------|---|
| ISRCTN number | - |
| ClinicalTrials.gov id (NCT number) | - |
| WHO universal trial number (UTN) | - |

Notes:

Sponsors

| | |
|------------------------------|--|
| Sponsor organisation name | The Fertility Clinic Skive |
| Sponsor organisation address | Reservevej 25, Skive, Denmark, 7800 |
| Public contact | Peter Humaidan, The Fertility Clinic, +45 78445773, peter.humaidan@midt.rm.dk |
| Scientific contact | Peter Humaidan, The Fertility Clinic, 78445760 78445773, peter.humaidan@midt.rm.dk |

Notes:

Paediatric regulatory details

| | |
|--|----|
| Is trial part of an agreed paediatric investigation plan (PIP) | No |
| Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial? | No |
| Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial? | No |

Notes:

Results analysis stage

| | |
|--|-----------------|
| Analysis stage | Final |
| Date of interim/final analysis | 30 October 2020 |
| Is this the analysis of the primary completion data? | Yes |
| Primary completion date | 15 October 2019 |
| Global end of trial reached? | Yes |
| Global end of trial date | 15 October 2019 |
| Was the trial ended prematurely? | No |

Notes:

General information about the trial

Main objective of the trial:

The objective of this pilot RCT is to explore the exogenous progesterone free luteal phase after GnRHa trigger in a population of IVF patients at risk of OHSS development, characterized by the development of 14-25 follicles ≥ 11 mm on the day of triggering of final follicular maturation.

Protection of trial subjects:

The study was approved by the scientific Ethics Committee of the Central Denmark Region – Project number: M201337713.

Written informed consent was obtained from all participants prior to inclusion.

Background therapy: -

Evidence for comparator: -

| | |
|---|---------------|
| Actual start date of recruitment | 03 March 2014 |
| Long term follow-up planned | No |
| Independent data monitoring committee (IDMC) involvement? | Yes |

Notes:

Population of trial subjects

Subjects enrolled per country

| | |
|--------------------------------------|--------------|
| Country: Number of subjects enrolled | Denmark: 120 |
| Worldwide total number of subjects | 120 |
| EEA total number of subjects | 120 |

Notes:

Subjects enrolled per age group

| | |
|---|-----|
| In utero | 0 |
| Preterm newborn - gestational age < 37 wk | 0 |
| Newborns (0-27 days) | 0 |
| Infants and toddlers (28 days-23 months) | 0 |
| Children (2-11 years) | 0 |
| Adolescents (12-17 years) | 0 |
| Adults (18-64 years) | 120 |
| From 65 to 84 years | 0 |
| 85 years and over | 0 |

Subject disposition

Recruitment

Recruitment details:

The first patient was enrolled in November 2014 and the last patient was enrolled in August 2019. All patients were from The fertility clinic, Skive Regional Hospital, Denmark.

Pre-assignment

Screening details:

A total of 275 IVF patients were assessed for eligibility and 250 patients were subsequently recruited for two studies, 2014-000448-13 and 2014-000447-32. 120 patients were randomized in 2014-000448-13 and 130 patients were randomized in 2014-000447-32. No patient was lost to follow-up.

Period 1

| | |
|------------------------------|---------------------------------|
| Period 1 title | Overall period (overall period) |
| Is this the baseline period? | Yes |
| Allocation method | Randomised - controlled |
| Blinding used | Not blinded |

Arms

| | |
|------------------------------|-------------|
| Are arms mutually exclusive? | Yes |
| Arm title | hCG trigger |

Arm description:

Ovulation trigger with 6500 IU hCG (Ovitrelle®), followed by 100 mg vaginal progesterone (Lutinus®) three times daily until the pregnancy test (14 days after OR), after which LPS was stopped.

| | |
|--|--|
| Arm type | Active comparator |
| Investigational medicinal product name | Ovitrelle |
| Investigational medicinal product code | |
| Other name | |
| Pharmaceutical forms | Concentrate for solution for injection |
| Routes of administration | Intradermal use |

Dosage and administration details:

Ovulation trigger with 6500 IU hCG (Ovitrelle®)

| | |
|------------------|---------------|
| Arm title | GnRHa trigger |
|------------------|---------------|

Arm description:

Ovulation trigger with a bolus of 0.5 mg Buserelin (Suprefact®), followed by a bolus of 1000 IU hCG (Pregnyl®) after OR and an additional bolus of 500 IU hCG (Pregnyl®) on OR + 4

| | |
|--|--|
| Arm type | Experimental |
| Investigational medicinal product name | Buserelin |
| Investigational medicinal product code | |
| Other name | |
| Pharmaceutical forms | Concentrate for cutaneous solution, Concentrate for solution for injection |
| Routes of administration | Subcutaneous use |

Dosage and administration details:

trigger with a bolus of 0.5 mg Buserelin (Suprefact®, Sanofi A/S, Copenhagen, Denmark)

| Number of subjects in period 1 | hCG trigger | GnRHa trigger |
|---------------------------------------|-------------|---------------|
| Started | 60 | 60 |
| Completed | 52 | 46 |
| Not completed | 8 | 14 |
| Protocol deviation | 8 | 14 |

Baseline characteristics

Reporting groups

| | |
|--------------------------------|----------------|
| Reporting group title | Overall period |
| Reporting group description: - | |

| Reporting group values | Overall period | Total | |
|--|----------------|-------|--|
| Number of subjects | 120 | 120 | |
| Age categorical | | | |
| Units: Subjects | | | |
| In utero | | 0 | |
| Preterm newborn infants (gestational age < 37 wks) | | 0 | |
| Newborns (0-27 days) | | 0 | |
| Infants and toddlers (28 days-23 months) | | 0 | |
| Children (2-11 years) | | 0 | |
| Adolescents (12-17 years) | | 0 | |
| Adults (18-64 years) | | 0 | |
| From 65-84 years | | 0 | |
| 85 years and over | | 0 | |
| Age continuous | | | |
| We adjusted for female age as continuous parameter | | | |
| Units: years | | | |
| arithmetic mean | 28.9 | | |
| standard deviation | ± 3.9 | - | |
| Gender categorical | | | |
| All were women | | | |
| Units: Subjects | | | |
| Female | 120 | 120 | |
| Male | 0 | 0 | |

Subject analysis sets

| | |
|----------------------------|--------------------------------------|
| Subject analysis set title | modified intention to treat analysis |
| Subject analysis set type | Modified intention-to-treat |

Subject analysis set description:

The primary analysis was the modified intention to treat analysis (White et al., 2011) of all randomized patients having an embryo transfer under the assumption that missing outcome data (patients not having an embryo transfer) was missing conditionally at random.

| | |
|----------------------------|--------------------|
| Subject analysis set title | Intention to treat |
| Subject analysis set type | Intention-to-treat |

Subject analysis set description:

Strict intention to treat for all patients randomized

| | |
|----------------------------|--------------|
| Subject analysis set title | Per protocol |
| Subject analysis set type | Per protocol |

Subject analysis set description:

Per protocol defined by those meeting all the eligibility criteria as well as complying with the protocol

| Reporting group values | modified intention to treat analysis | Intention to treat | Per protocol |
|---|--------------------------------------|--------------------|--------------|
| Number of subjects | 101 | 120 | 98 |
| Age categorical | | | |
| Units: Subjects | | | |
| In utero Preterm newborn infants (gestational age < 37 wks) Newborns (0-27 days) Infants and toddlers (28 days-23 months) Children (2-11 years) Adolescents (12-17 years) Adults (18-64 years) From 65-84 years 85 years and over | | | |
| Age continuous | | | |
| We adjusted for female age as continuous parameter | | | |
| Units: years | | | |
| arithmetic mean | 28.9 | 28.9 | 28.9 |
| standard deviation | ± 3.9 | ± 3.9 | ± 3.9 |
| Gender categorical | | | |
| All were women | | | |
| Units: Subjects | | | |
| Female | | | |
| Male | | | |

End points

End points reporting groups

| | |
|---|--------------------------------------|
| Reporting group title | hCG trigger |
| Reporting group description: Ovulation trigger with 6500 IU hCG (Ovitrelle®), followed by 100 mg vaginal progesterone (Lutinus®) three times daily until the pregnancy test (14 days after OR), after which LPS was stopped. | |
| Reporting group title | GnRHa trigger |
| Reporting group description: Ovulation trigger with a bolus of 0.5 mg Buserelin (Suprefact®), followed by a bolus of 1000 IU hCG (Pregnyl®) after OR and an additional bolus of 500 IU hCG (Pregnyl®) on OR + 4 | |
| Subject analysis set title | modified intention to treat analysis |
| Subject analysis set type | Modified intention-to-treat |
| Subject analysis set description: The primary analysis was the modified intention to treat analysis (White et al., 2011) of all randomized patients having an embryo transfer under the assumption that missing outcome data (patients not having an embryo transfer) was missing conditionally at random. | |
| Subject analysis set title | Intention to treat |
| Subject analysis set type | Intention-to-treat |
| Subject analysis set description: Strict intention to treat for all patients randomized | |
| Subject analysis set title | Per protocol |
| Subject analysis set type | Per protocol |
| Subject analysis set description: Per protocol defined by those meeting all the eligibility criteria as well as complying with the protocol | |

Primary: Ongoing pregnancy week 12

| | |
|----------------------------------|---------------------------|
| End point title | Ongoing pregnancy week 12 |
| End point description: | |
| End point type | Primary |
| End point timeframe: 12 weeks | |

| End point values | hCG trigger | GnRHa trigger | modified intention to treat analysis | |
|--------------------------------|-----------------|-----------------|--------------------------------------|--|
| Subject group type | Reporting group | Reporting group | Subject analysis set | |
| Number of subjects analysed | 52 | 49 | 101 | |
| Units: Fetal heartbeat at scan | 52 | 49 | 101 | |

Statistical analyses

| | |
|--|-------------------|
| Statistical analysis title | Binary regression |
| Statistical analysis description: A binary regression model was used to calculate the crude relative risks and relative differences (cRR, cRD) and adjusted relative risks and relative differences (aRR, aRD) for the primary outcome. | |

| | |
|---|--|
| Comparison groups | hCG trigger v GnRHa trigger v modified intention to treat analysis |
| Number of subjects included in analysis | 202 |
| Analysis specification | Pre-specified |
| Analysis type | equivalence |
| Method | Regression, Linear |
| Parameter estimate | Risk ratio (RR) |
| Point estimate | 0.86 |
| Confidence interval | |
| level | 95 % |
| sides | 2-sided |
| lower limit | 0.6 |
| upper limit | 1.22 |

Adverse events

Adverse events information^[1]

Timeframe for reporting adverse events:

Approximately 14 weeks.

Adverse event reporting additional description:

We did not investigate adverse events other than OHSS.

| | |
|-----------------|----------------|
| Assessment type | Non-systematic |
|-----------------|----------------|

Dictionary used

| | |
|-----------------|--------|
| Dictionary name | MedDRA |
|-----------------|--------|

| | |
|--------------------|----|
| Dictionary version | 23 |
|--------------------|----|

Frequency threshold for reporting non-serious adverse events: 5 %

Notes:

[1] - There are no non-serious adverse events recorded for these results. It is expected that there will be at least one non-serious adverse event reported.

Justification: We did not observe any adverse events. OHSS rates will be reported in the publication.

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? No

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

Sample size, lack of blinding.

Full manuscript has been submitted for publication.

Notes: