



Clinical trial results:

A Phase Ib/II, multicenter, open-label study of EGF816 in combination with INC280 in adult subjects with EGFR mutated non-small cell lung cancer

Summary

EudraCT number	2014-000726-37
Trial protocol	DE ES FR IT
Global end of trial date	11 May 2022

Results information

Result version number	v2 (current)
This version publication date	05 March 2023
First version publication date	26 January 2023
Version creation reason	<ul style="list-style-type: none">• Correction of full data set• Aligment with Clinicaltrials.gov information

Trial information

Trial identification

Sponsor protocol code	CINC280X2105C
-----------------------	---------------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02335944
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	Novartis Pharma AG
Sponsor organisation address	Novartis Campus, Basel, Switzerland,
Public contact	Clinical Disclosure Office, Novartis Pharma AG, 41 613241111, novartis.email@novartis.com
Scientific contact	Clinical Disclosure Office, Novartis Pharma AG, 41 613241111, novartis.email@novartis.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	10 November 2020
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	11 May 2022
Was the trial ended prematurely?	Yes

Notes:

General information about the trial

Main objective of the trial:

The purpose of this study was to determine the maximum tolerated dose (MTD) or recommended phase 2 dose (RP2D) of nazartinib in combination with capmatinib and to estimate the preliminary anti-tumor activity of nazartinib in combination with capmatinib in participants with advanced non-small cell lung cancer with documented EGFR mutation.

Protection of trial subjects:

The study was in compliance with the ethical principles derived from the Declaration of Helsinki and the International Conference on Harmonization (ICH) Good Clinical Practice (GCP) guidelines. All the local regulatory requirements pertinent to safety of trial subjects were also followed during the conduct of the trial.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	13 January 2015
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Australia: 8
Country: Number of subjects enrolled	Canada: 5
Country: Number of subjects enrolled	France: 8
Country: Number of subjects enrolled	Germany: 17
Country: Number of subjects enrolled	Italy: 21
Country: Number of subjects enrolled	Korea, Republic of: 14
Country: Number of subjects enrolled	Norway: 4
Country: Number of subjects enrolled	Singapore: 31
Country: Number of subjects enrolled	Spain: 38
Country: Number of subjects enrolled	Taiwan: 25
Country: Number of subjects enrolled	United States: 6
Worldwide total number of subjects	177
EEA total number of subjects	88

Notes:

Subjects enrolled per age group	
In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	112
From 65 to 84 years	65
85 years and over	0

Subject disposition

Recruitment

Recruitment details:

Participants took part in 19 investigative sites in 11 countries. Due to early study termination, Group 5 (Phase II part) was never opened

Pre-assignment

Screening details:

The screening period began once patients had signed the study informed consent. All screening/baseline evaluations were performed ≤ 28 days before Cycle 1 Day 1.

Period 1

Period 1 title	Treatment Phase (overall period)
Is this the baseline period?	Yes
Allocation method	Non-randomised - controlled
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
Arm title	Phase IB part- INC280 200mg BID/ EGF816 50mg QD

Arm description:

Combination of INC280 200mg twice daily (BID) and EGF816 50mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment

Arm type	Experimental
Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Capmatinib was administered orally, twice per day, at a dose of 200 mg, in fasted state.

Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use

Dosage and administration details:

Nazartinib was administered orally, once a day, at a dose of 50 mg, in fasted state

Arm title	Phase IB part- INC280 200mg BID/ EGF816 100mg QD
------------------	--

Arm description:

Combination of INC280 200mg twice daily (BID) and EGF816 100mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment

Arm type	Experimental
Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Capmatinib was administered orally, twice per day, at a dose of 200 mg, in fasted state.

Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use
Dosage and administration details:	
Nazartinib was administered orally, once a day, at a dose of 100 mg, in fasted state	
Arm title	Phase IB part- INC280 400mg BID/ EGF816 75mg QD
Arm description:	
Combination of INC280 400mg twice daily (BID) and EGF816 75mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Arm type	Experimental
Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use
Dosage and administration details:	
Nazartinib was administered orally, once a day, at a dose of 75 mg, in fasted state	
Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use
Dosage and administration details:	
Capmatinib was administered orally, twice per day, at a dose of 400 mg, in fasted state.	
Arm title	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Arm description:	
Combination of INC280 400mg twice daily (BID) and EGF816 100mg once daily (QD) in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Arm type	Experimental
Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use
Dosage and administration details:	
Capmatinib was administered orally, twice per day, at a dose of 400 mg, in fasted state.	
Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use
Dosage and administration details:	
Nazartinib was administered orally, once a day, at a dose of 100 mg, in fasted state	
Arm title	Phase IB part- INC280 400mg BID/ EGF816 150mg QD
Arm description:	
Combination of INC280 400mg twice daily (BID) and EGF816 150mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Arm type	Experimental

Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use
Dosage and administration details:	
Nazartinib was administered orally, once a day, at a dose of 150 mg, in fasted state	
Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use
Dosage and administration details:	
Capmatinib was administered orally, twice per day, at a dose of 400 mg, in fasted state.	
Arm title	Phase II- Group 1
Arm description:	
NSCLC participants with previously documented activating EGFR mutation, with any T790M and MET dysregulation status, who received 1 to 3 lines of systemic antineoplastic therapy prior to study entry including one line maximum of 1st or 2nd generation EGFR TKI. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Arm type	Experimental
Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use
Dosage and administration details:	
Nazartinib was administered orally, once a day, at a dose of 100 mg, in fasted state	
Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use
Dosage and administration details:	
Capmatinib was administered orally, twice per day, at a dose of 400 mg, in fasted state.	
Arm title	Phase II- Group 2
Arm description:	
NSCLC participants harboring T790M mutation in de novo setting, irrespective of the activating mutation status who received none to 2 lines of systemic antineoplastic therapy prior to study entry, but no therapy known to inhibit EGFR. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Arm type	Experimental
Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use
Dosage and administration details:	
Nazartinib was administered orally, once a day, at a dose of 100 mg, in fasted state	
Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet

Routes of administration	Oral use
--------------------------	----------

Dosage and administration details:

Capmatinib was administered orally, twice per day, at a dose of 400 mg, in fasted state.

Arm title	Phase II- Group 3
------------------	-------------------

Arm description:

NSCLC participants with previously documented EGFR activating mutation, T790M negative, and any MET status who never received any prior line of systemic antineoplastic therapy. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state

Arm type	Experimental
Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use

Dosage and administration details:

Nazartinib was administered orally, once a day, at a dose of 100 mg, in fasted state

Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Capmatinib was administered orally, twice per day, at a dose of 400 mg, in fasted state.

Arm title	Phase II- Group 4
------------------	-------------------

Arm description:

NSCLC participants with previously documented EGFR activating mutations and any T790M and MET status who received none to 2 prior lines of systemic antineoplastic therapy prior to study entry. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fed state.

Arm type	Experimental
Investigational medicinal product name	Nazartinib
Investigational medicinal product code	EGF816
Other name	
Pharmaceutical forms	Capsule, Tablet
Routes of administration	Oral use

Dosage and administration details:

Nazartinib was administered orally, once a day, at a dose of 100 mg, in fed state

Investigational medicinal product name	Capmatinib
Investigational medicinal product code	INC280
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Capmatinib was administered orally, twice per day, at a dose of 400 mg, in fed state.

Number of subjects in period 1	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD
Started	4	5	3
Completed	0	0	0
Not completed	4	5	3
Physician decision	-	-	-
Adverse event, non-fatal	2	-	-
Study Terminated By Sponsor	-	-	-
Progressive Disease	2	5	2
Subject/Guardian Decision	-	-	1

Number of subjects in period 1	Phase IB part- INC280 400mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 150mg QD	Phase II- Group 1
Started	16	5	52
Completed	1	0	2
Not completed	15	5	50
Physician decision	1	-	6
Adverse event, non-fatal	2	1	15
Study Terminated By Sponsor	1	-	3
Progressive Disease	11	4	25
Subject/Guardian Decision	-	-	1

Number of subjects in period 1	Phase II- Group 2	Phase II- Group 3	Phase II- Group 4
Started	3	47	42
Completed	0	5	6
Not completed	3	42	36
Physician decision	1	2	1
Adverse event, non-fatal	-	5	7
Study Terminated By Sponsor	-	1	1
Progressive Disease	2	30	27
Subject/Guardian Decision	-	4	-

Baseline characteristics

Reporting groups

Reporting group title	Phase IB part- INC280 200mg BID/ EGF816 50mg QD
Reporting group description: Combination of INC280 200mg twice daily (BID) and EGF816 50mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 200mg BID/ EGF816 100mg QD
Reporting group description: Combination of INC280 200mg twice daily (BID) and EGF816 100mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 75mg QD
Reporting group description: Combination of INC280 400mg twice daily (BID) and EGF816 75mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Reporting group description: Combination of INC280 400mg twice daily (BID) and EGF816 100mg once daily (QD) in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 150mg QD
Reporting group description: Combination of INC280 400mg twice daily (BID) and EGF816 150mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase II- Group 1
Reporting group description: NSCLC participants with previously documented activating EGFR mutation, with any T790M and MET dysregulation status, who received 1 to 3 lines of systemic antineoplastic therapy prior to study entry including one line maximum of 1st or 2nd generation EGFR TKI. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Reporting group title	Phase II- Group 2
Reporting group description: NSCLC participants harboring T790M mutation in de novo setting, irrespective of the activating mutation status who received none to 2 lines of systemic antineoplastic therapy prior to study entry, but no therapy known to inhibit EGFR. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Reporting group title	Phase II- Group 3
Reporting group description: NSCLC participants with previously documented EGFR activating mutation, T790M negative, and any MET status who never received any prior line of systemic antineoplastic therapy. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Reporting group title	Phase II- Group 4
Reporting group description: NSCLC participants with previously documented EGFR activating mutations and any T790M and MET status who received none to 2 prior lines of systemic antineoplastic therapy prior to study entry. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fed state.	

Reporting group values	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD
Number of subjects	4	5	3
Age Categorical Units: Participants			
<=18 years	0	0	0
Between 18 and 65 years	3	5	2

>=65 years	1	0	1
------------	---	---	---

Sex: Female, Male Units: Participants			
Female	4	3	1
Male	0	2	2
Race/Ethnicity, Customized Units: Subjects			
Asian	2	3	1
Caucasian	2	2	1
Missing	0	0	1
Black	0	0	0

Reporting group values	Phase IB part- INC280 400mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 150mg QD	Phase II- Group 1
Number of subjects	16	5	52
Age Categorical Units: Participants			
<=18 years	0	0	0
Between 18 and 65 years	11	3	33
>=65 years	5	2	19
Sex: Female, Male Units: Participants			
Female	9	4	38
Male	7	1	14
Race/Ethnicity, Customized Units: Subjects			
Asian	10	4	26
Caucasian	4	1	22
Missing	2	0	3
Black	0	0	1

Reporting group values	Phase II- Group 2	Phase II- Group 3	Phase II- Group 4
Number of subjects	3	47	42
Age Categorical Units: Participants			
<=18 years	0	0	0
Between 18 and 65 years	1	30	24
>=65 years	2	17	18
Sex: Female, Male Units: Participants			
Female	2	33	24
Male	1	14	18
Race/Ethnicity, Customized Units: Subjects			
Asian	2	30	20
Caucasian	1	16	21
Missing	0	1	1
Black	0	0	0

Reporting group values	Total		
Number of subjects	177		
Age Categorical Units: Participants			
<=18 years	0		
Between 18 and 65 years	112		
>=65 years	65		
Sex: Female, Male Units: Participants			
Female	118		
Male	59		
Race/Ethnicity, Customized Units: Subjects			
Asian	98		
Caucasian	70		
Missing	8		
Black	1		

End points

End points reporting groups

Reporting group title	Phase IB part- INC280 200mg BID/ EGF816 50mg QD
Reporting group description: Combination of INC280 200mg twice daily (BID) and EGF816 50mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 200mg BID/ EGF816 100mg QD
Reporting group description: Combination of INC280 200mg twice daily (BID) and EGF816 100mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 75mg QD
Reporting group description: Combination of INC280 400mg twice daily (BID) and EGF816 75mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Reporting group description: Combination of INC280 400mg twice daily (BID) and EGF816 100mg once daily (QD) in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 150mg QD
Reporting group description: Combination of INC280 400mg twice daily (BID) and EGF816 150mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment	
Reporting group title	Phase II- Group 1
Reporting group description: NSCLC participants with previously documented activating EGFR mutation, with any T790M and MET dysregulation status, who received 1 to 3 lines of systemic antineoplastic therapy prior to study entry including one line maximum of 1st or 2nd generation EGFR TKI. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Reporting group title	Phase II- Group 2
Reporting group description: NSCLC participants harboring T790M mutation in de novo setting, irrespective of the activating mutation status who received none to 2 lines of systemic antineoplastic therapy prior to study entry, but no therapy known to inhibit EGFR. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Reporting group title	Phase II- Group 3
Reporting group description: NSCLC participants with previously documented EGFR activating mutation, T790M negative, and any MET status who never received any prior line of systemic antineoplastic therapy. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	
Reporting group title	Phase II- Group 4
Reporting group description: NSCLC participants with previously documented EGFR activating mutations and any T790M and MET status who received none to 2 prior lines of systemic antineoplastic therapy prior to study entry. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fed state.	

Primary: Phase Ib: Number of participants with dose limiting toxicities (DLTs)

End point title	Phase Ib: Number of participants with dose limiting toxicities (DLTs) ^{[1][2]}
End point description: Number of participants with DLTs in the Phase Ib part. A DLT is defined as an AE or abnormal laboratory value assessed as unrelated to disease, disease progression, inter-current illness, or concomitant medications that occurs within the first 28 days of treatment with EGF816 in combination with INC280	

during the escalation part of the study (Phase Ib)

End point type	Primary
End point timeframe:	
Up to first 28 days of treatment	

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: No statistical analyses were planned for this endpoint

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	1	13
Units: Participants	1	0	0	1

End point values	Phase IB part- INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: Participants	2			

Statistical analyses

No statistical analyses for this end point

Primary: Phase II Group 1, 2 and 3: Overall Response Rate (ORR) by investigator's assessment per RECIST 1.1

End point title	Phase II Group 1, 2 and 3: Overall Response Rate (ORR) by investigator's assessment per RECIST 1.1 ^{[3][4]}
-----------------	--

End point description:

ORR is defined as the percentage of participants with a best overall response of complete response (CR) or partial response (PR) determined by investigator's assessment in accordance to Response Evaluation Criteria in Solid Tumors (RECIST 1.1). ORR was assessed in Group 1, 2 and 3 (Phase II part).

CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters.

End point type	Primary
End point timeframe:	
Up to approximately 4 years	

Notes:

[3] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: No statistical analyses were planned for this endpoint

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.
Justification: This endpoint is only applicable for Phase II Groups 1, 2 and 3

End point values	Phase II- Group 1	Phase II- Group 2	Phase II- Group 3	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	52	3	47	
Units: Percentage of participants				
number (confidence interval 95%)	28.8 (17.1 to 43.10)	33.3 (0.8 to 90.6)	61.7 (46.4 to 75.5)	

Statistical analyses

No statistical analyses for this end point

Primary: Phase II Group 4: Number of participants with Adverse Events (AEs) and Serious AEs (SAEs)

End point title	Phase II Group 4: Number of participants with Adverse Events (AEs) and Serious AEs (SAEs) ^{[5][6]}
-----------------	---

End point description:

Number of participants in Group 4 (Phase II part) with AEs and SAEs. An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of a study intervention, whether or not considered related to the study intervention. Any untoward event resulting in death, life threatening, requires inpatient hospitalization or prolongation of existing hospitalization, results in persistent disability/incapacity, congenital anomaly/birth defect or any other situation according to medical or scientific judgment is categorized as SAE.

End point type	Primary
----------------	---------

End point timeframe:

From start of treatment up to 30 days after last dose of study treatment, assessed up to 3.7 years

Notes:

[5] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: No statistical analyses were planned for this endpoint

[6] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Group 4

End point values	Phase II- Group 4			
Subject group type	Reporting group			
Number of subjects analysed	42			
Units: Participants				
AEs	42			
SAEs	26			

Statistical analyses

No statistical analyses for this end point

Primary: Phase II Group 4: Number of participants with dose reductions and dose interruptions of INC280 and EGF618

End point title	Phase II Group 4: Number of participants with dose reductions and dose interruptions of INC280 and EGF618 ^[7] ^[8]
-----------------	---

End point description:

Number of participants with at least one dose reduction of INC280, at least one dose interruption of INC280, at least one dose reduction of EGF816 and at least one dose interruption of EGF816 in the Group 4 (Phase II part).

End point type	Primary
----------------	---------

End point timeframe:

From start of treatment until end of treatment, assessed up to 3.6 years

Notes:

[7] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: No statistical analyses were planned for this endpoint

[8] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Group 4

End point values	Phase II- Group 4			
Subject group type	Reporting group			
Number of subjects analysed	42			
Units: Participants				
Dose Reduction of INC280	30			
Dose Interruption of INC280	31			
Dose Reduction of EGF816	12			
Dose Interruption of EGF816	35			

Statistical analyses

No statistical analyses for this end point

Primary: Phase II Group 4: Dose intensity

End point title	Phase II Group 4: Dose intensity ^[9] ^[10]
-----------------	---

End point description:

Dose intensity, defined as the ratio of total dose received and actual duration, for participants in Group 4 (Phase II part)

End point type	Primary
----------------	---------

End point timeframe:

From start of treatment until end of treatment, assessed up to 3.6 years

Notes:

[9] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: No statistical analyses were planned for this endpoint

[10] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Group 4

End point values	Phase II- Group 4			
Subject group type	Reporting group			
Number of subjects analysed	42			
Units: milligram/day				
arithmetic mean (standard deviation)				
INC280	666.2 (± 148.97)			
EGF816	87.4 (± 14.82)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Number of participants with dose reductions and dose interruptions of INC280 and EGF618

End point title	Phase Ib: Number of participants with dose reductions and dose interruptions of INC280 and EGF618 ^[11]
-----------------	---

End point description:

Number of participants in Phase Ib with at least one dose reduction of INC280, at least one dose interruption of INC280, at least one dose reduction of EGF816 and at least one dose interruption of EGF816 .

End point type	Secondary
----------------	-----------

End point timeframe:

From start of treatment until end of treatment, assessed up to approximately 5 years

Notes:

[11] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: Participants				
Dose Reduction of INC280	2	4	2	10
Dose Interruption of INC280	2	2	2	12
Dose Reduction of EGF816	2	1	1	5
Dose Interruption of EGF816	2	2	2	11

End point values	Phase IB part- INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: Participants				
Dose Reduction of INC280	5			

Dose Interruption of INC280	5			
Dose Reduction of EGF816	3			
Dose Interruption of EGF816	5			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Group 1, 2 and 3: Number of participants with dose reductions and dose interruptions of INC280 and EGF618

End point title	Phase II Group 1, 2 and 3: Number of participants with dose reductions and dose interruptions of INC280 and EGF618 ^[12]
-----------------	--

End point description:

Number of participants in Groups 1, 2 and 3 (Phase II part) with at least one dose reduction of INC280, at least one dose interruption of INC280, at least one dose reduction of EGF816 and at least one dose interruption of EGF816 .

End point type	Secondary
----------------	-----------

End point timeframe:

From start of treatment until end of treatment, assessed up to approximately 4 years

Notes:

[12] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 1, 2 and 3

End point values	Phase II-Group 1	Phase II-Group 2	Phase II-Group 3	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	52	3	47	
Units: Participants				
Dose Reduction of INC280	33	3	34	
Dose Interruption of INC280	31	3	39	
Dose Reduction of EGF816	17	2	23	
Dose Interruption of EGF816	34	3	40	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Dose intensity

End point title	Phase Ib: Dose intensity ^[13]
-----------------	--

End point description:

Dose intensity, defined as the ratio of total dose received and actual duration, in Phase Ib participants

End point type	Secondary
----------------	-----------

End point timeframe:

From start of treatment until end of treatment, assessed up to approximately 5 years

Notes:

[13] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: milligram/day				
arithmetic mean (standard deviation)				
INC280	319.1 (± 101.09)	356.9 (± 71.14)	596.3 (± 200.24)	658.2 (± 158.18)
EGF816	40.0 (± 12.60)	89.6 (± 17.93)	60.0 (± 13.36)	85.2 (± 17.28)

End point values	Phase IB part- INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: milligram/day				
arithmetic mean (standard deviation)				
INC280	463.4 (± 212.52)			
EGF816	97.7 (± 36.43)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Group 1, 2 and 3: Dose intensity

End point title	Phase II Group 1, 2 and 3: Dose intensity ^[14]
End point description:	
Dose intensity, defined as the ratio of total dose received and actual duration, in Group 1, 2 and 3 (Phase II)	
End point type	Secondary
End point timeframe:	
From start of treatment until end of treatment, assessed up to approximately 4 years	

Notes:

[14] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 1, 2 and 3

End point values	Phase II- Group 1	Phase II- Group 2	Phase II- Group 3	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	52	3	47	
Units: milligram/day				
arithmetic mean (standard deviation)				
INC280	662.9 (± 160.73)	562.9 (± 38.36)	634.7 (± 172.33)	
EGF816	85.2 (± 18.21)	76.3 (± 15.31)	84.8 (± 15.53)	

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Overall Response Rate (ORR) per RECIST 1.1 based on investigator's assessment

End point title	Phase Ib: Overall Response Rate (ORR) per RECIST 1.1 based on investigator's assessment ^[15]
-----------------	---

End point description:

ORR is defined as percentage of participants with best overall response of PR+CR determined by Investigator's assessment in accordance to RECIST 1.1. ORR was assessed in Phase Ib participants. CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to approximately 5 years

Notes:

[15] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: Percentage of participants				
number (confidence interval 95%)	0 (0.0 to 60.2)	40.0 (5.3 to 85.3)	33.3 (0.8 to 90.6)	50.0 (24.7 to 75.3)

End point values	Phase IB part- INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: Percentage of participants				
number (confidence interval 95%)	60.0 (14.7 to 94.7)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Group 4: Overall Response Rate (ORR) per RECIST 1.1 based on investigator's assessment

End point title	Phase II Group 4: Overall Response Rate (ORR) per RECIST 1.1 based on investigator's assessment ^[16]
-----------------	---

End point description:

ORR is defined as percentage of participants with best overall response of PR+CR determined by Investigator's assessment in accordance to RECIST 1.1. ORR was assessed in Group 4 (Phase II) CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to approximately 4 years

Notes:

[16] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Group 4

End point values	Phase II-Group 4			
Subject group type	Reporting group			
Number of subjects analysed	42			
Units: Percentage of participants				
number (confidence interval 95%)	42.9 (27.7 to 59.0)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Progression Free Survival (PFS) per RECIST 1.1 based on investigator's assessment

End point title	Phase Ib: Progression Free Survival (PFS) per RECIST 1.1 based on investigator's assessment ^[17]
-----------------	---

End point description:

PFS is defined as time from date of first dose of study treatment to date of first documented disease progression determined by Investigator assessment in accordance to RECIST 1.1.or death due to any cause. The PFS distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant has not had an event, PFS is censored at the date of last adequate tumor assessment. PFS was assessed in Phase Ib participants

End point type	Secondary
----------------	-----------

End point timeframe:

From date of first dose to first documented disease progression or death, assessed up to approximately 5 years

Notes:

[17] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: Months				
median (confidence interval 95%)	5.6 (3.5 to 999999)	7.4 (1.6 to 999999)	3.5 (0.8 to 999999)	5.7 (1.9 to 42.1)

End point values	Phase IB part-INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: Months				
median (confidence interval 95%)	14.5 (7.3 to 999999)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 1, 2, 3 and 4: Progression Free Survival (PFS) per RECIST 1.1 based on investigator's assessment

End point title	Phase II Groups 1, 2, 3 and 4: Progression Free Survival (PFS) per RECIST 1.1 based on investigator's assessment ^[18]
-----------------	--

End point description:

PFS is defined as time from date of first dose of study treatment to date of first documented disease progression determined by Investigator assessment in accordance to RECIST 1.1 or death due to any cause. The PFS distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant has not had an event, PFS is censored at the date of last adequate tumor assessment. PFS was assessed in Group 1, 2, 3 and 4 (Phase II)

End point type	Secondary
----------------	-----------

End point timeframe:

From date of first dose to first documented disease progression or death, assessed up to approximately 4 years

Notes:

[18] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 1, 2, 3 and 4

End point values	Phase II-Group 1	Phase II-Group 2	Phase II-Group 3	Phase II-Group 4
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	52	3	47	42
Units: Months				
median (confidence interval 95%)	5.6 (3.7 to 7.4)	3.8 (3.7 to 999999)	10.1 (7.6 to 13.8)	10.9 (5.6 to 19.2)

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Time to Response (TTR) per RECIST 1.1 based on investigator's assessment

End point title	Phase Ib: Time to Response (TTR) per RECIST 1.1 based on investigator's assessment ^[19]
-----------------	--

End point description:

TTR is defined as the time from the date of the first dose to the date of first documented response (CR or PR) determined by Investigator assessment in accordance to RECIST 1.1. The TTR distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant has not had an event, duration was censored at the date of last adequate tumor assessment. TTR was assessed in Phase Ib participants.

CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters.

End point type	Secondary
----------------	-----------

End point timeframe:

From the date of the first dose to the date of first documented response, up to approximately 5 years

Notes:

[19] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: Months				
median (confidence interval 95%)	999999 (999999 to 999999)	999999 (1.8 to 999999)	999999 (1.7 to 999999)	3.5 (1.6 to 999999)

End point values	Phase IB part-INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: Months				
median (confidence interval 95%)	4.5 (1.9 to			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 1, 2, 3 and 4: Time to Response (TTR) per RECIST 1.1 based on investigator's assessment

End point title	Phase II Groups 1, 2, 3 and 4: Time to Response (TTR) per RECIST 1.1 based on investigator's assessment ^[20]
-----------------	---

End point description:

TTR is defined as the time from the date of the first dose to the date of first documented response (CR or PR) determined by Investigator assessment in accordance to RECIST 1.1. The TTR distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant has not had an event, duration was censored at the date of last adequate tumor assessment. TTR was assessed in Group 1, 2, 3 and 4 (Phase II)

CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters.

End point type	Secondary
----------------	-----------

End point timeframe:

From the date of the first dose to the date of first documented response, up to approximately 4 years

Notes:

[20] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 1, 2, 3 and 4

End point values	Phase II-Group 1	Phase II-Group 2	Phase II-Group 3	Phase II-Group 4
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	52	3	47	42
Units: Months				
median (confidence interval 95%)	9999 (5.4 to 999999)	999999 (1.8 to 999999)	1.9 (1.8 to 5.9)	999999 (3.6 to 999999)

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Duration of Response (DOR) per RECIST 1.1 based on investigator's assessment

End point title	Phase Ib: Duration of Response (DOR) per RECIST 1.1 based on investigator's assessment ^[21]
-----------------	--

End point description:

DOR is defined as the time from first documented response (PR or CR) to the date of first documented disease progression determined by Investigator assessment in accordance to RECIST 1.1 or death due to underlying cancer. The DOR distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant has not had an event, duration was censored at the date of last adequate tumor assessment. DOR was assessed in Phase Ib participants

CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters.

End point type	Secondary
----------------	-----------

End point timeframe:

From date of first documented response to first documented disease progression or death, assessed up to approximately 5 years

Notes:

[21] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	0 ^[22]	2	1	8
Units: Months				
median (confidence interval 95%)	(to)	8.8 (5.6 to 999999)	14.8 (-999999 to 999999)	25.3 (3.6 to 47.9)

Notes:

[22] - No participants had a documented response (CR or PR)

End point values	Phase IB part-INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	3			
Units: Months				
median (confidence interval 95%)	8.0 (5.4 to 999999)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 1, 2, 3 and 4: Duration of Response (DOR) per RECIST 1.1 based on investigator's assessment

End point title	Phase II Groups 1, 2, 3 and 4: Duration of Response (DOR) per RECIST 1.1 based on investigator's assessment ^[23]
-----------------	---

End point description:

DOR is defined as the time from first documented response (PR or CR) to the date of first documented disease progression determined by Investigator assessment in accordance to RECIST 1.1 or death due to underlying cancer. The DOR distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant has not had an event, duration was censored at the date of last adequate tumor assessment. DOR was assessed in Group 1, 2, 3 and 4 (Phase II)

CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters.

End point type	Secondary
----------------	-----------

End point timeframe:

From date of first documented response to first documented disease progression or deaths, assessed up to approximately 4 years

Notes:

[23] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 1, 2, 3 and 4

End point values	Phase II-Group 1	Phase II-Group 2	Phase II-Group 3	Phase II-Group 4
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	15	1	29	18
Units: Months				
median (confidence interval 95%)	6.5 (3.7 to 10.8)	12.0 (-999999 to 999999)	11.6 (6.6 to 17.5)	14.5 (9.2 to 999999)

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Disease Control Rate (DCR) per RECIST 1.1 based on investigator's assessment

End point title	Phase Ib: Disease Control Rate (DCR) per RECIST 1.1 based on investigator's assessment ^[24]
-----------------	--

End point description:

DCR is defined as the percentage of participants with best overall response of CR, PR, or stable disease (SD) determined by Investigator assessment in accordance to RECIST 1.1. DCR was assessed in Phase Ib participants

CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters; SD= Neither sufficient shrinkage to qualify for PR or CR nor an increase in lesions which would qualify for progression.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to approximately 5 years

Notes:

[24] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase Ib part-INC280 200mg BID/ EGF816 50mg QD	Phase Ib part-INC280 200mg BID/ EGF816 100mg QD	Phase Ib part-INC280 400mg BID/ EGF816 75mg QD	Phase Ib part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: Percentage of participants				
number (confidence interval 95%)	100 (39.8 to 100)	60.0 (14.7 to 94.7)	33.3 (0.8 to 90.6)	62.5 (35.4 to 84.8)

End point values	Phase IB part- INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: Percentage of participants				
number (confidence interval 95%)	80.0 (28.4 to 99.5)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Group 1, 2 3 and 4: Disease Control Rate (DCR) per RECIST 1.1 based on investigator's assessment

End point title	Phase II Group 1, 2 3 and 4: Disease Control Rate (DCR) per RECIST 1.1 based on investigator's assessment ^[25]
-----------------	---

End point description:

DCR is defined as the percentage of participants with best overall response of CR, PR, or SD determined by Investigator assessment in accordance to RECIST 1.1. DCR was assessed in Group 1, 2, 3 and 4 (Phase II)

CR=Disappearance of all non-nodal target lesions. In addition, any pathological lymph nodes assigned as target lesions must have a reduction in short axis to < 10 mm; PR= At least a 30% decrease in the sum of diameters of all target lesions, taking as reference the baseline sum of diameters; SD= Neither sufficient shrinkage to qualify for PR or CR nor an increase in lesions which would qualify for progression.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to approximately 4 years

Notes:

[25] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 1, 2, 3 and 4

End point values	Phase II- Group 1	Phase II- Group 2	Phase II- Group 3	Phase II- Group 4
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	52	3	47	42
Units: Percentage of participants				
number (confidence interval 95%)	59.6 (45.1 to 73.0)	100 (29.2 to 100)	93.6 (82.5 to 98.7)	81.0 (65.9 to 91.4)

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Overall Survival (OS)

End point title	Phase Ib: Overall Survival (OS) ^[26]
-----------------	---

End point description:

OS is defined as the time from first dose of the study treatment to the date of death due to any cause. The OS distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant was not known to have died, survival was censored at the date of last contact. OS was assessed in Phase Ib participants

End point type	Secondary
----------------	-----------

End point timeframe:

From date of first dose to death, assessed up to approximately 5 years

Notes:

[26] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: Months				
median (confidence interval 95%)	10.1 (8.1 to 999999)	56.5 (6.5 to 999999)	7.0 (1.1 to 999999)	17.2 (5.7 to 58.7)

End point values	Phase IB part-INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	5			
Units: Months				
median (confidence interval 95%)	31.5 (16.6 to 999999)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 1, 2, 3 and 4: Overall Survival (OS)

End point title	Phase II Groups 1, 2, 3 and 4: Overall Survival (OS) ^[27]
-----------------	--

End point description:

OS is defined as the time from first dose of the study treatment to the date of death due to any cause. The OS distribution was estimated using the Kaplan-Meier method and associated 95% confidence intervals were calculated. If a participant was not known to have died, survival was censored at the date of last contact. OS was assessed in Group 1, 2, 3 and 4

End point type	Secondary
----------------	-----------

End point timeframe:

From date of first dose to death, assessed up to approximately 4 years

Notes:

[27] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 1, 2 3 and 4

End point values	Phase II-Group 1	Phase II-Group 2	Phase II-Group 3	Phase II-Group 4
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	52	3	47	42
Units: Months				
median (confidence interval 95%)	18.8 (14.9 to 26.0)	5.6 (3.7 to 999999)	25.6 (18.8 to 33.0)	28.9 (20.5 to 999999)

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Area under the plasma concentration versus time curve from time 0 to 12 hours (AUC0-12) of INC280

End point title	Phase Ib: Area under the plasma concentration versus time curve from time 0 to 12 hours (AUC0-12) of INC280 ^[28]
-----------------	---

End point description:

Pharmacokinetic (PK) parameters were calculated based on INC280 plasma concentrations by using non-compartmental methods.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr post-dose on Cycle 1 Day 1, Cycle 1 Day 15 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[28] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	1	12
Units: hours*nanogram/mililiter (hr*ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 3 / 5 / 1 / 12 / 3)	12300 (± 4710)	10500 (± 4320)	38300 (± 999999)	22200 (± 10700)
Cycle 1 Day 15 (n= 3 / 5 / 1 / 11 / 2)	11600 (± 3400)	11800 (± 3360)	48800 (± 999999)	28900 (± 12700)
Cycle 2 Day 1 (n= 3 / 5 / 0 / 9 / 2)	11300 (± 4050)	11100 (± 1770)	999999 (± 999999)	21600 (± 7370)

End point values	Phase IB part-			
------------------	----------------	--	--	--

	INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	3			
Units: hours*nanogram/mililiter (hr*ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 3 / 5 / 1 / 12 / 3)	23000 (± 11600)			
Cycle 1 Day 15 (n= 3 / 5 / 1 / 11 / 2)	23000 (± 3440)			
Cycle 2 Day 1 (n= 3 / 5 / 0 / 9 / 2)	24900 (± 5190)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Peak plasma concentration (Cmax) of INC280

End point title	Phase Ib: Peak plasma concentration (Cmax) of INC280 ^[29]
-----------------	--

End point description:

Pharmacokinetic (PK) parameters were calculated based on INC280 plasma concentrations by using non-compartmental methods.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr post-dose on Cycle 1 Day 1, Cycle 1 Day 15 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[29] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	1	12
Units: nanogram/mililiter (ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 4 / 5 / 1 / 12 / 3)	3630 (± 588)	2530 (± 1080)	8000 (± 999999)	5100 (± 2550)
Cycle 1 Day 15 (n= 3 / 5 / 1 / 11 / 2)	3070 (± 1120)	2840 (± 1350)	12000 (± 999999)	5780 (± 2640)
Cycle 2 Day 1 (N= 3 / 5 / 0 / 10 / 2)	3590 (± 1620)	2800 (± 797)	999999 (± 999999)	4830 (± 1390)

End point values	Phase IB part- INC280 400mg BID/ EGF816			
------------------	---	--	--	--

	150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	3			
Units: nanogram/mililiter (ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 4 / 5 / 1 / 12 / 3)	5930 (± 1330)			
Cycle 1 Day 15 (n= 3 / 5 / 1 / 11 / 2)	4350 (± 933)			
Cycle 2 Day 1 (N= 3 / 5 / 0 / 10 / 2)	6370 (± 1060)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Time to reach maximum concentration (Tmax) of INC280

End point title	Phase Ib: Time to reach maximum concentration (Tmax) of INC280 ^[30]
-----------------	--

End point description:

Pharmacokinetic (PK) parameters were calculated based on INC280 plasma concentrations by using non-compartmental methods. Actual time of sample collection was used (not the nominal time point as per scheduled assessment)

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr post-dose on Cycle 1 Day 1, Cycle 1 Day 15 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[30] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	1	12
Units: Hours				
median (full range (min-max))				
Cycle 1 Day 1 (n= 4 / 5 / 1 / 12 / 3)	1.07 (1.00 to 2.00)	2.00 (1.00 to 4.00)	2.00 (2.00 to 2.00)	2.01 (1.00 to 7.67)
Cycle 1 Day 15 (n= 3 / 5 / 1 / 11 / 2)	1.12 (1.03 to 2.05)	2.00 (0.883 to 7.98)	1.17 (1.17 to 1.17)	1.97 (0.950 to 4.00)
Cycle 2 Day 1 (n= 3 / 5 / 0 / 10 / 2)	1.00 (1.00 to 2.00)	2.00 (1.00 to 3.83)	999999 (999999 to 999999)	1.94 (0.983 to 4.00)

End point values	Phase IB part-INC280 400mg BID/ EGF816 150mg QD			
------------------	---	--	--	--

Subject group type	Reporting group			
Number of subjects analysed	3			
Units: Hours				
median (full range (min-max))				
Cycle 1 Day 1 (n= 4 / 5 / 1 / 12 / 3)	2.00 (1.02 to 2.13)			
Cycle 1 Day 15 (n= 3 / 5 / 1 / 11 / 2)	1.50 (1.00 to 2.00)			
Cycle 2 Day 1 (n= 3 / 5 / 0 / 10 / 2)	1.51 (1.02 to 2.00)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 3 and 4: Area under the plasma concentration versus time curve from time 0 to 12 hours (AUC0-12) of INC280

End point title	Phase II Groups 3 and 4: Area under the plasma concentration versus time curve from time 0 to 12 hours (AUC0-12) of INC280 ^[31]
-----------------	--

End point description:

Pharmacokinetic (PK) parameters were calculated based on INC280 plasma concentrations by using non-compartmental methods.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr post-dose on Cycle 1 Day 1 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[31] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 3 and 4

End point values	Phase II-Group 3	Phase II-Group 4		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	9	15		
Units: hours*nanogram/mililiter (hr*ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 8 / 12)	22000 (± 8000)	16900 (± 7360)		
Cycle 2 Day 1 (n= 8 / 15)	19700 (± 11200)	20500 (± 7440)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 3 and 4: Peak plasma concentration (Cmax) of INC280

End point title	Phase II Groups 3 and 4: Peak plasma concentration (Cmax) of
-----------------	--

End point description:

Pharmacokinetic (PK) parameters were calculated based on INC280 plasma concentrations by using non-compartmental methods.

End point type Secondary

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr post-dose on Cycle 1 Day 1 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[32] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 3 and 4

End point values	Phase II-Group 3	Phase II-Group 4		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	9	15		
Units: nanogram/mililiter (ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 8 / 12)	4770 (± 1460)	3420 (± 1550)		
Cycle 2 Day 1 (n= 8 / 15)	4360 (± 2060)	3940 (± 1660)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 3 and 4: Time to reach maximum concentration (Tmax) of INC280

End point title Phase II Groups 3 and 4: Time to reach maximum concentration (Tmax) of INC280^[33]

End point description:

Pharmacokinetic (PK) parameters were calculated based on INC280 plasma concentrations by using non-compartmental methods. Actual time of sample collection was used (not the nominal time point as per scheduled assessment)

End point type Secondary

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr post-dose on Cycle 1 Day 1 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[33] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 3 and 4

End point values	Phase II-Group 3	Phase II-Group 4		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	9	15		
Units: Hours				
median (full range (min-max))				
Cycle 1 Day 1 (n= 8 / 12)	1.99 (1.00 to 4.00)	2.08 (1.85 to 8.00)		

Cycle 2 Day 1 (n= 8 / 15)	1.47 (1.00 to 7.08)	3.82 (1.00 to 4.10)		
---------------------------	---------------------	---------------------	--	--

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Area under the plasma concentration versus time curve from time 0 to 24 hours (AUC0-24) of EGF816

End point title	Phase Ib: Area under the plasma concentration versus time curve from time 0 to 24 hours (AUC0-24) of EGF816 ^[34]
-----------------	---

End point description:

Pharmacokinetic (PK) parameters were calculated based on EGF816 plasma concentrations by using non-compartmental methods.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr and 24 hr post-dose on Cycle 1 Day 1, Cycle 1 Day 15 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[34] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	13
Units: hours*nanogram/mililiter (hr*ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 3 / 5 / 2 / 11 / 2)	3920 (± 1450)	5850 (± 4380)	4010 (± 534)	5930 (± 3600)
Cycle 1 Day 15 (n= 4 / 5 / 2 / 12 / 2)	5200 (± 2010)	9630 (± 4060)	7330 (± 108)	11100 (± 5340)
Cycle 2 Day 1 (n= 3 / 4 / 0 / 11 / 2)	4080 (± 1130)	7460 (± 2340)	999999 (± 999999)	10500 (± 7480)

End point values	Phase IB part-INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	3			
Units: hours*nanogram/mililiter (hr*ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 3 / 5 / 2 / 11 / 2)	10500 (± 3400)			

Cycle 1 Day 15 (n= 4 / 5 / 2 / 12 / 2)	16400 (± 5670)			
Cycle 2 Day 1 (n= 3 / 4 / 0 / 11 / 2)	11400 (± 2360)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Peak plasma concentration (Cmax) of EGF816

End point title	Phase Ib: Peak plasma concentration (Cmax) of EGF816 ^[35]
-----------------	--

End point description:

Pharmacokinetic (PK) parameters were calculated based on EGF816 plasma concentrations by using non-compartmental methods.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr and 24 hr post-dose on Cycle 1 Day 1, Cycle 1 Day 15 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[35] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase IB part-INC280 200mg BID/ EGF816 50mg QD	Phase IB part-INC280 200mg BID/ EGF816 100mg QD	Phase IB part-INC280 400mg BID/ EGF816 75mg QD	Phase IB part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	13
Units: nanogram/mililiter (ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 4 / 5 / 2 / 13 / 3)	258 (± 104)	465 (± 323)	302 (± 8.49)	381 (± 231)
Cycle 1 Day 15 (n= 3 / 5 / 3 / 12 / 2)	361 (± 165)	677 (± 321)	517 (± 180)	595 (± 261)
Cycle 2 Day 1 (n= 3 / 5 / 0 / 11 / 2)	269 (± 90.8)	583 (± 261)	999999 (± 999999)	604 (± 365)

End point values	Phase IB part-INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	3			
Units: nanogram/mililiter (ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 4 / 5 / 2 / 13 / 3)	777 (± 281)			
Cycle 1 Day 15 (n= 3 / 5 / 3 / 12 / 2)	1010 (± 354)			
Cycle 2 Day 1 (n= 3 / 5 / 0 / 11 / 2)	814 (± 137)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase Ib: Time to reach maximum concentration (Tmax) of EGF816

End point title	Phase Ib: Time to reach maximum concentration (Tmax) of EGF816 ^[36]
-----------------	--

End point description:

Pharmacokinetic (PK) parameters were calculated based on EGF816 plasma concentrations by using non-compartmental methods. Actual time of sample collection was used (not the nominal time point as per scheduled assessment)

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr and 24 hr post-dose on Cycle 1 Day 1, Cycle 1 Day 15 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[36] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase Ib arms

End point values	Phase Ib part-INC280 200mg BID/ EGF816 50mg QD	Phase Ib part-INC280 200mg BID/ EGF816 100mg QD	Phase Ib part-INC280 400mg BID/ EGF816 75mg QD	Phase Ib part-INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	13
Units: Hours (hr)				
median (full range (min-max))				
Cycle 1 Day 1 (n= 4 / 5 / 2 / 13 / 3)	3.03 (1.08 to 4.02)	3.90 (2.00 to 4.00)	3.00 (2.00 to 4.00)	4.00 (2.00 to 8.99)
Cycle 1 Day 15 (n= 3 / 5 / 3 / 12 / 2)	2.07 (2.00 to 4.05)	3.95 (2.00 to 4.00)	4.00 (2.17 to 4.05)	5.90 (2.17 to 11.9)
Cycle 2 Day 1 (n= 3 / 5 / 0 / 11 / 2)	4.00 (3.95 to 4.00)	4.00 (2.00 to 4.00)	999999 (999999 to 999999)	4.00 (1.98 to 7.00)

End point values	Phase Ib part-INC280 400mg BID/ EGF816 150mg QD			
Subject group type	Reporting group			
Number of subjects analysed	3			
Units: Hours (hr)				
median (full range (min-max))				
Cycle 1 Day 1 (n= 4 / 5 / 2 / 13 / 3)	4.00 (2.03 to 4.02)			

Cycle 1 Day 15 (n= 3 / 5 / 3 / 12 / 2)	3.93 (3.83 to 4.02)			
Cycle 2 Day 1 (n= 3 / 5 / 0 / 11 / 2)	3.05 (2.10 to 4.00)			

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 3 and 4: Area under the plasma concentration versus time curve from time 0 to 24 hours (AUC0-24) of EGF816

End point title	Phase II Groups 3 and 4: Area under the plasma concentration versus time curve from time 0 to 24 hours (AUC0-24) of EGF816 ^[37]
-----------------	--

End point description:

Pharmacokinetic (PK) parameters were calculated based on EGF816 plasma concentrations by using non-compartmental methods.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr and 24 hr post-dose on Cycle 1 Day 1 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[37] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 3 and 4

End point values	Phase II-Group 3	Phase II-Group 4		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	7	15		
Units: hours*nanogram/mililiter (hr*ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 6 / 12)	6390 (± 2460)	3420 (± 1700)		
Cycle 2 Day 1 (n= 5 / 15)	11100 (± 2000)	7670 (± 3330)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 3 and 4: Peak plasma concentration (Cmax) of EGF816

End point title	Phase II Groups 3 and 4: Peak plasma concentration (Cmax) of EGF816 ^[38]
-----------------	---

End point description:

Pharmacokinetic (PK) parameters were calculated based on EGF816 plasma concentrations by using non-compartmental methods.

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr and 24 hr post-dose on Cycle 1 Day 1 and

Notes:

[38] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 3 and 4

End point values	Phase II-Group 3	Phase II-Group 4		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	7	15		
Units: nanogram/mililiter (ng/mL)				
arithmetic mean (standard deviation)				
Cycle 1 Day 1 (n= 6 / 12)	448 (± 184)	239 (± 107)		
Cycle 2 Day 1 (n= 6 / 15)	658 (± 117)	447 (± 175)		

Statistical analyses

No statistical analyses for this end point

Secondary: Phase II Groups 3 and 4: Time to reach maximum concentration (Tmax) of EGF816

End point title	Phase II Groups 3 and 4: Time to reach maximum concentration (Tmax) of EGF816 ^[39]
-----------------	---

End point description:

Pharmacokinetic (PK) parameters were calculated based on EGF816 plasma concentrations by using non-compartmental methods. Actual time of sample collection was used (not the nominal time point as per scheduled assessment)

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose, 0.5 hours (hr) and 1 hr, 2 hr, 4 hr, 8 hr and 12 hr and 24 hr post-dose on Cycle 1 Day 1 and Cycle 2 Day 1 (Cycle=28 days)

Notes:

[39] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is only applicable for Phase II Groups 3 and 4

End point values	Phase II-Group 3	Phase II-Group 4		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	7	15		
Units: Hours (hr)				
median (full range (min-max))				
Cycle 1 Day 1 (n= 6 / 12)	4.00 (1.12 to 8.00)	4.00 (1.85 to 8.00)		
Cycle 2 Day 1 (n= 6 / 15)	4.00 (2.00 to 4.13)	4.00 (3.82 to 8.00)		

Statistical analyses

Post-hoc: All collected deaths

End point title	All collected deaths
-----------------	----------------------

End point description:

On-treatment deaths were collected from first dose of study medication to 30 days after the last dose of study medication for a maximum duration of approximately 5 years (Phase Ib) and 4 years (Phase II).

Post-treatment deaths were collected after 30 days post-treatment, for a maximum duration of approximately 5 years (Phase Ib) and 4 years (Phase II).

All deaths refer to the sum of on-treatment and post-treatment deaths

End point type	Post-hoc
----------------	----------

End point timeframe:

On-treatment: up to approximately 5 years (Phase Ib) and 4 years (Phase II). Post-treatment: Up to approximately 5 years (Phase Ib) and 4 years (Phase II).

End point values	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	4	5	3	16
Units: Participants				
On-treatment deaths	0	0	1	2
Post-treatment deaths	4	3	2	9
All deaths	4	3	3	11

End point values	Phase IB part- INC280 400mg BID/ EGF816 150mg QD	Phase II- Group 1	Phase II- Group 2	Phase II- Group 3
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	5	52	3	47
Units: Participants				
On-treatment deaths	0	9	2	5
Post-treatment deaths	3	26	1	25
All deaths	3	35	3	30

End point values	Phase II- Group 4			
Subject group type	Reporting group			
Number of subjects analysed	42			
Units: Participants				
On-treatment deaths	1			
Post-treatment deaths	22			
All deaths	23			

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

From first dose of study treatment until last dose of study treatment plus 30 days, up to approximately 5 years (Phase Ib) and 4 years (Phase II).

Adverse event reporting additional description:

Consistent with EudraCT disclosure specifications, Novartis has reported under the SAEs field "number of deaths resulting from adverse events" all those deaths, resulting from SAEs that are deemed to be causally related to treatment by the investigator.

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	25.0
--------------------	------

Reporting groups

Reporting group title	Phase IB part- INC280 200mg BID/ EGF816 50mg QD
-----------------------	---

Reporting group description:

Combination of INC280 200mg twice daily (BID) and EGF816 50mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment

Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 150mg QD
-----------------------	--

Reporting group description:

Combination of INC280 400mg twice daily (BID) and EGF816 150mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment

Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
-----------------------	--

Reporting group description:

Combination of INC280 400mg twice daily (BID) and EGF816 100mg once daily (QD) in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment

Reporting group title	Phase IB part- INC280 400mg BID/ EGF816 75mg QD
-----------------------	---

Reporting group description:

Combination of INC280 400mg twice daily (BID) and EGF816 75mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment

Reporting group title	Phase IB part- INC280 200mg BID/ EGF816 100mg QD
-----------------------	--

Reporting group description:

Combination of INC280 200mg twice daily (BID) and EGF816 100mg once daily (QD) in fasted state in NSCLC participants with previously documented EGFR mutation, who progressed on EGFR TKI treatment

Reporting group title	All Participants
-----------------------	------------------

Reporting group description:

All Participants

Reporting group title	Phase II- Group 4
-----------------------	-------------------

Reporting group description:

NSCLC participants with previously documented EGFR activating mutations and any T790M and MET status who received none to 2 prior lines of systemic antineoplastic therapy prior to study entry. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fed state.

Reporting group title	Phase II- Group 3
-----------------------	-------------------

Reporting group description:

NSCLC participants with previously documented EGFR activating mutation, T790M negative, and any MET status who never received any prior line of systemic antineoplastic therapy. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state

Reporting group title	Phase II- Group 2
-----------------------	-------------------

Reporting group description:

NSCLC participants harboring T790M mutation in de novo setting, irrespective of the activating mutation status who received none to 2 lines of systemic antineoplastic therapy prior to study entry, but no therapy known to inhibit EGFR. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state

Reporting group title	Phase II- Group 1
Reporting group description:	
NSCLC participants with previously documented activating EGFR mutation, with any T790M and MET dysregulation status, who received 1 to 3 lines of systemic antineoplastic therapy prior to study entry including one line maximum of 1st or 2nd generation EGFR TKI. Participants were treated with 400 mg twice daily for INC280 and 100 mg once daily for EGF816 in fasted state	

Serious adverse events	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 400mg BID/ EGF816 150mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Total subjects affected by serious adverse events			
subjects affected / exposed	3 / 4 (75.00%)	4 / 5 (80.00%)	11 / 16 (68.75%)
number of deaths (all causes)	0	0	2
number of deaths resulting from adverse events	0	0	0
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Malignant pleural effusion			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Invasive ductal breast carcinoma			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tumour associated fever			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Embolism			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypertensive crisis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypotension			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Fatigue			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Asthenia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General physical health deterioration			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Influenza like illness			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Non-cardiac chest pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Oedema			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyrexia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			

Anaphylactic shock			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypersensitivity			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Dyspnoea			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Aspiration			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypersensitivity pneumonitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haemoptysis			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pleural effusion			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Interstitial lung disease			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Pneumothorax			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonitis			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pulmonary embolism			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	2 / 16 (12.50%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory failure			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Confusional state			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Apathy			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Alanine aminotransferase increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Amylase increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Aspartate aminotransferase increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood bilirubin increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood creatinine increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ejection fraction decreased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Weight decreased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neutrophil count decreased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Fall			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Procedural pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Cardiac disorders			
Cardiac arrest			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardio-respiratory arrest			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pericardial effusion			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Myocardial ischaemia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Aphasia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dysarthria			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cerebral ischaemia			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Brain oedema			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Epilepsy			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Headache			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hydrocephalus			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Loss of consciousness			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypertensive hydrocephalus			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neurological decompensation			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Seizure			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Speech disorder			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tremor			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Spinal cord compression			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Wernicke's encephalopathy			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Disseminated intravascular coagulation			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Leukopenia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Thrombocytopenia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ear and labyrinth disorders			
Hypoacusis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
Visual impairment			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Gastrointestinal disorders			
Ascites			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Colitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Constipation			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diarrhoea			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastric ulcer			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hiatus hernia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Inguinal hernia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nausea			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 1	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Hepatic failure			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Skin and subcutaneous tissue disorders			
Dermatitis allergic			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Erythema multiforme			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Drug eruption			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Purpura			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash macular			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash maculo-papular			

subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	1 / 1	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urticaria			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Renal failure			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acute kidney injury			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Osteonecrosis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neck pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pain in extremity			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Infections and infestations			
Cellulitis			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	1 / 1	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Empyema			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Enterocolitis infectious			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Erysipelas			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Herpes zoster			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Herpes zoster meningitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Influenza			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Lower respiratory tract infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Meningitis			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Necrotising fasciitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Meningitis aseptic			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Pleural infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia aspiration			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia chlamydial			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory tract infection			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Soft tissue infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary tract infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urosepsis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Electrolyte imbalance			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypokalaemia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypoalbuminaemia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hyponatraemia			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	All Participants
Total subjects affected by serious adverse events			
subjects affected / exposed	3 / 3 (100.00%)	2 / 5 (40.00%)	113 / 177 (63.84%)
number of deaths (all causes)	1	0	20
number of deaths resulting from adverse events	0	0	2
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Malignant pleural effusion			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Invasive ductal breast carcinoma			
subjects affected / exposed	0 / 3 (0.00%)	1 / 5 (20.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tumour associated fever			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Embolism			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	1 / 1
Hypertensive crisis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypotension			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Fatigue			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences causally related to treatment / all	0 / 0	0 / 0	3 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Asthenia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General physical health deterioration			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Influenza like illness			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Non-cardiac chest pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Oedema			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyrexia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	8 / 177 (4.52%)
occurrences causally related to treatment / all	0 / 0	0 / 0	4 / 8
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			

Anaphylactic shock			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypersensitivity			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences causally related to treatment / all	0 / 0	0 / 0	3 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Dyspnoea			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	8 / 177 (4.52%)
occurrences causally related to treatment / all	0 / 1	0 / 0	1 / 9
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 2
Aspiration			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypersensitivity pneumonitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haemoptysis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pleural effusion			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	10 / 177 (5.65%)
occurrences causally related to treatment / all	0 / 1	0 / 0	3 / 12
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Interstitial lung disease			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences causally related to treatment / all	0 / 0	0 / 0	5 / 5
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Pneumothorax			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences causally related to treatment / all	0 / 0	0 / 0	3 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	1 / 1
Pulmonary embolism			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 7
deaths causally related to treatment / all	0 / 1	0 / 0	0 / 1
Respiratory failure			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	6 / 177 (3.39%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 7
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 3
Psychiatric disorders			
Confusional state			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Apathy			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Alanine aminotransferase increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Amylase increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Aspartate aminotransferase increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood bilirubin increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood creatinine increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ejection fraction decreased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Weight decreased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neutrophil count decreased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Fall			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Procedural pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Cardiac disorders			
Cardiac arrest			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Cardio-respiratory arrest			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Pericardial effusion			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Myocardial ischaemia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Nervous system disorders			
Aphasia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dysarthria			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cerebral ischaemia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Brain oedema			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Epilepsy			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Headache			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hydrocephalus			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Loss of consciousness			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypertensive hydrocephalus			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Neurological decompensation			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Seizure			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 8
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Speech disorder			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tremor			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Spinal cord compression			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Wernicke's encephalopathy			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Disseminated intravascular coagulation			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Leukopenia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	1 / 1	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Thrombocytopenia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ear and labyrinth disorders			
Hypoacusis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
Visual impairment			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Gastrointestinal disorders			
Ascites			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Colitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Constipation			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diarrhoea			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences causally related to treatment / all	0 / 0	0 / 0	3 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastric ulcer			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hiatus hernia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Inguinal hernia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nausea			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences causally related to treatment / all	0 / 0	0 / 0	3 / 5
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	2 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Hepatic failure			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 2
Skin and subcutaneous tissue disorders			
Dermatitis allergic			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Erythema multiforme			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Drug eruption			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Purpura			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash macular			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash maculo-papular			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences causally related to treatment / all	0 / 0	0 / 0	7 / 7
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urticaria			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Renal failure			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acute kidney injury			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Osteonecrosis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neck pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pain in extremity			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Infections and infestations Cellulitis subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	1 / 5 (20.00%) 1 / 2 0 / 0	9 / 177 (5.08%) 5 / 12 0 / 0
Empyema subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	0 / 5 (0.00%) 0 / 0 0 / 0	1 / 177 (0.56%) 0 / 1 0 / 0
Enterocolitis infectious subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	0 / 5 (0.00%) 0 / 0 0 / 0	1 / 177 (0.56%) 0 / 1 0 / 0
Erysipelas subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	0 / 5 (0.00%) 0 / 0 0 / 0	2 / 177 (1.13%) 3 / 3 0 / 0
Herpes zoster subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	0 / 5 (0.00%) 0 / 0 0 / 0	1 / 177 (0.56%) 1 / 1 0 / 0
Herpes zoster meningitis subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	0 / 5 (0.00%) 0 / 0 0 / 0	1 / 177 (0.56%) 0 / 1 0 / 0
Influenza subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	0 / 5 (0.00%) 0 / 0 0 / 0	2 / 177 (1.13%) 0 / 2 0 / 0
Lower respiratory tract infection subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 3 (0.00%) 0 / 0 0 / 0	0 / 5 (0.00%) 0 / 0 0 / 0	1 / 177 (0.56%) 0 / 1 0 / 0
Meningitis			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Necrotising fasciitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Meningitis aseptic			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	10 / 177 (5.65%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 10
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Pleural infection			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia aspiration			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia chlamydial			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 5
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Respiratory tract infection			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Soft tissue infection			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary tract infection			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urosepsis			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 1	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Electrolyte imbalance			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypokalaemia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypoalbuminaemia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hyponatraemia			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Phase II- Group 4	Phase II- Group 3	Phase II- Group 2
Total subjects affected by serious adverse events			
subjects affected / exposed	26 / 42 (61.90%)	33 / 47 (70.21%)	2 / 3 (66.67%)
number of deaths (all causes)	1	5	2
number of deaths resulting from adverse events	0	1	0
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Malignant pleural effusion			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Invasive ductal breast carcinoma			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tumour associated fever			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Embolism			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	1 / 1	0 / 0
Hypertensive crisis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypotension			

subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Fatigue			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Asthenia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General physical health deterioration			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Influenza like illness			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Non-cardiac chest pain			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Oedema			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyrexia			
subjects affected / exposed	2 / 42 (4.76%)	4 / 47 (8.51%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	2 / 2	1 / 4	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			

Anaphylactic shock			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypersensitivity			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Dyspnoea			
subjects affected / exposed	2 / 42 (4.76%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 2	1 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Aspiration			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypersensitivity pneumonitis			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haemoptysis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pleural effusion			
subjects affected / exposed	3 / 42 (7.14%)	4 / 47 (8.51%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 3	2 / 6	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Interstitial lung disease			
subjects affected / exposed	3 / 42 (7.14%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	3 / 3	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Pneumothorax			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonitis			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pulmonary embolism			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	1 / 3 (33.33%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory failure			
subjects affected / exposed	0 / 42 (0.00%)	3 / 47 (6.38%)	1 / 3 (33.33%)
occurrences causally related to treatment / all	0 / 0	0 / 3	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 1
Psychiatric disorders			
Confusional state			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Apathy			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Alanine aminotransferase increased			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Amylase increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Aspartate aminotransferase increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood bilirubin increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood creatinine increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ejection fraction decreased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Weight decreased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neutrophil count decreased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Fall			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Procedural pain			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Cardiac disorders			
Cardiac arrest			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardio-respiratory arrest			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pericardial effusion			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Myocardial ischaemia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Aphasia			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dysarthria			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cerebral ischaemia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Brain oedema			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Epilepsy			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Headache			
subjects affected / exposed	0 / 42 (0.00%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 3	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hydrocephalus			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Loss of consciousness			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypertensive hydrocephalus			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neurological decompensation			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Seizure			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Speech disorder			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tremor			

subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Spinal cord compression			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Wernicke's encephalopathy			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Disseminated intravascular coagulation			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Leukopenia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Thrombocytopenia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ear and labyrinth disorders			
Hypoacusis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
Visual impairment			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Gastrointestinal disorders			
Ascites			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Colitis			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Constipation			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diarrhoea			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastric ulcer			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hiatus hernia			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Inguinal hernia			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nausea			
subjects affected / exposed	1 / 42 (2.38%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	2 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			

subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 3	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Hepatic failure			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	1 / 3 (33.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Skin and subcutaneous tissue disorders			
Dermatitis allergic			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Erythema multiforme			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Drug eruption			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Purpura			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash macular			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash maculo-papular			

subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urticaria			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Renal failure			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acute kidney injury			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	0 / 42 (0.00%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 3	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Osteonecrosis			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neck pain			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pain in extremity			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Infections and infestations			
Cellulitis			
subjects affected / exposed	3 / 42 (7.14%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	2 / 4	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Empyema			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Enterocolitis infectious			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Erysipelas			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Herpes zoster			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Herpes zoster meningitis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Influenza			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Lower respiratory tract infection			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Meningitis			

subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Necrotising fasciitis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Meningitis aseptic			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			
subjects affected / exposed	4 / 42 (9.52%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 4	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pleural infection			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia aspiration			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia chlamydial			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 0
Respiratory tract infection			

subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Soft tissue infection			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary tract infection			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urosepsis			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Electrolyte imbalance			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypokalaemia			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypoalbuminaemia			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hyponatraemia			

subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Phase II- Group 1		
Total subjects affected by serious adverse events			
subjects affected / exposed	29 / 52 (55.77%)		
number of deaths (all causes)	9		
number of deaths resulting from adverse events	1		
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Malignant pleural effusion			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Invasive ductal breast carcinoma			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Tumour associated fever			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Vascular disorders			
Embolism			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypertensive crisis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypotension			

subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
General disorders and administration site conditions			
Fatigue			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	2 / 2		
deaths causally related to treatment / all	0 / 0		
Asthenia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
General physical health deterioration			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Influenza like illness			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Non-cardiac chest pain			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Oedema			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pyrexia			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	1 / 2		
deaths causally related to treatment / all	0 / 0		
Immune system disorders			

Anaphylactic shock			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypersensitivity			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	2 / 2		
deaths causally related to treatment / all	0 / 0		
Respiratory, thoracic and mediastinal disorders			
Dyspnoea			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences causally related to treatment / all	0 / 3		
deaths causally related to treatment / all	0 / 2		
Aspiration			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypersensitivity pneumonitis			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Haemoptysis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pleural effusion			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Interstitial lung disease			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	2 / 2		
deaths causally related to treatment / all	0 / 0		

Pneumothorax			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pneumonitis			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	2 / 2		
deaths causally related to treatment / all	1 / 1		
Pulmonary embolism			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Respiratory failure			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 1		
Psychiatric disorders			
Confusional state			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Apathy			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Investigations			
Alanine aminotransferase increased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Amylase increased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		

Aspartate aminotransferase increased				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Blood bilirubin increased				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Blood creatinine increased				
subjects affected / exposed	1 / 52 (1.92%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Ejection fraction decreased				
subjects affected / exposed	1 / 52 (1.92%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 1			
Weight decreased				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Neutrophil count decreased				
subjects affected / exposed	1 / 52 (1.92%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Injury, poisoning and procedural complications				
Fall				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Procedural pain				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			

Cardiac disorders			
Cardiac arrest			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 1		
Cardio-respiratory arrest			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 1		
Pericardial effusion			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Myocardial ischaemia			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 1		
Nervous system disorders			
Aphasia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Dysarthria			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Cerebral ischaemia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Brain oedema			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		

Epilepsy				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Headache				
subjects affected / exposed	1 / 52 (1.92%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Hydrocephalus				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Loss of consciousness				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Hypertensive hydrocephalus				
subjects affected / exposed	1 / 52 (1.92%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 1			
Neurological decompensation				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Seizure				
subjects affected / exposed	2 / 52 (3.85%)			
occurrences causally related to treatment / all	0 / 2			
deaths causally related to treatment / all	0 / 0			
Speech disorder				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Tremor				

subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Spinal cord compression			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Wernicke's encephalopathy			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Blood and lymphatic system disorders			
Disseminated intravascular coagulation			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Leukopenia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Thrombocytopenia			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Ear and labyrinth disorders			
Hypoacusis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Eye disorders			
Visual impairment			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		

Gastrointestinal disorders			
Ascites			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Colitis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Constipation			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Diarrhoea			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Gastric ulcer			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Hiatus hernia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Inguinal hernia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Nausea			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Vomiting			

subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Hepatobiliary disorders			
Hepatic failure			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Skin and subcutaneous tissue disorders			
Dermatitis allergic			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Erythema multiforme			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Drug eruption			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Rash			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Purpura			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Rash macular			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Rash maculo-papular			

subjects affected / exposed	4 / 52 (7.69%)		
occurrences causally related to treatment / all	4 / 4		
deaths causally related to treatment / all	0 / 0		
Urticaria			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	2 / 2		
deaths causally related to treatment / all	0 / 0		
Renal and urinary disorders			
Renal failure			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Acute kidney injury			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Osteonecrosis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Neck pain			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pain in extremity			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		

Infections and infestations Cellulitis subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 52 (1.92%) 0 / 2 0 / 0		
Empyema subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 52 (0.00%) 0 / 0 0 / 0		
Enterocolitis infectious subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 52 (0.00%) 0 / 0 0 / 0		
Erysipelas subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 52 (1.92%) 2 / 2 0 / 0		
Herpes zoster subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 52 (0.00%) 0 / 0 0 / 0		
Herpes zoster meningitis subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 52 (0.00%) 0 / 0 0 / 0		
Influenza subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 52 (0.00%) 0 / 0 0 / 0		
Lower respiratory tract infection subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 52 (0.00%) 0 / 0 0 / 0		
Meningitis			

subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Necrotising fasciitis				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Meningitis aseptic				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Pneumonia				
subjects affected / exposed	5 / 52 (9.62%)			
occurrences causally related to treatment / all	0 / 5			
deaths causally related to treatment / all	0 / 0			
Pleural infection				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Pneumonia aspiration				
subjects affected / exposed	1 / 52 (1.92%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Pneumonia chlamydial				
subjects affected / exposed	0 / 52 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Sepsis				
subjects affected / exposed	1 / 52 (1.92%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Respiratory tract infection				

subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Soft tissue infection			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Urinary tract infection			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Urosepsis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Electrolyte imbalance			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypokalaemia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypoalbuminaemia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hyponatraemia			

subjects affected / exposed	2 / 52 (3.85%)		
occurrences causally related to treatment / all	1 / 2		
deaths causally related to treatment / all	0 / 0		

Frequency threshold for reporting non-serious adverse events: 5 %

Non-serious adverse events	Phase IB part- INC280 200mg BID/ EGF816 50mg QD	Phase IB part- INC280 400mg BID/ EGF816 150mg QD	Phase IB part- INC280 400mg BID/ EGF816 100mg QD
Total subjects affected by non-serious adverse events			
subjects affected / exposed	4 / 4 (100.00%)	5 / 5 (100.00%)	16 / 16 (100.00%)
Vascular disorders			
Hot flush			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Hypotension			
subjects affected / exposed	1 / 4 (25.00%)	2 / 5 (40.00%)	0 / 16 (0.00%)
occurrences (all)	1	2	0
Pallor			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Thrombosis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
General disorders and administration site conditions			
Chest pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Asthenia			
subjects affected / exposed	2 / 4 (50.00%)	0 / 5 (0.00%)	4 / 16 (25.00%)
occurrences (all)	2	0	5
Chills			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	1	0	1
Fatigue			

subjects affected / exposed	2 / 4 (50.00%)	2 / 5 (40.00%)	6 / 16 (37.50%)
occurrences (all)	3	2	10
Face oedema			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	3 / 16 (18.75%)
occurrences (all)	0	1	3
Gait disturbance			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Influenza like illness			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Non-cardiac chest pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Oedema peripheral			
subjects affected / exposed	1 / 4 (25.00%)	4 / 5 (80.00%)	8 / 16 (50.00%)
occurrences (all)	3	8	17
Pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Peripheral swelling			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	2 / 16 (12.50%)
occurrences (all)	0	2	3
Pyrexia			
subjects affected / exposed	1 / 4 (25.00%)	3 / 5 (60.00%)	7 / 16 (43.75%)
occurrences (all)	1	3	11
Immune system disorders			
Anaphylactoid reaction			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Hypersensitivity			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Reproductive system and breast disorders			

Pelvic pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Perineal pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Vulvovaginal pruritus			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Prostatomegaly			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Respiratory, thoracic and mediastinal disorders			
Cough			
subjects affected / exposed	1 / 4 (25.00%)	1 / 5 (20.00%)	5 / 16 (31.25%)
occurrences (all)	1	1	10
Dry throat			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Dysphonia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Dyspnoea			
subjects affected / exposed	1 / 4 (25.00%)	1 / 5 (20.00%)	4 / 16 (25.00%)
occurrences (all)	2	1	8
Dyspnoea exertional			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences (all)	0	0	2
Epistaxis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Haemoptysis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Nasal discomfort			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	2
Hypoxia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Nasal dryness			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Oropharyngeal pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Pleural effusion			
subjects affected / exposed	2 / 4 (50.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	2	0	1
Pneumothorax			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Productive cough			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Pulmonary embolism			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Rhinorrhoea			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	3 / 16 (18.75%)
occurrences (all)	0	0	6
Throat irritation			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Psychiatric disorders			
Anxiety			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Confusional state			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1

Sleep disorder subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Depression subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Insomnia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 2	0 / 16 (0.00%) 0
Investigations			
Alanine aminotransferase increased subjects affected / exposed occurrences (all)	2 / 4 (50.00%) 4	2 / 5 (40.00%) 3	2 / 16 (12.50%) 2
Amylase increased subjects affected / exposed occurrences (all)	2 / 4 (50.00%) 4	2 / 5 (40.00%) 2	8 / 16 (50.00%) 13
Aspartate aminotransferase increased subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 2	1 / 5 (20.00%) 2	2 / 16 (12.50%) 2
Blood albumin decreased subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 1	1 / 5 (20.00%) 8	1 / 16 (6.25%) 1
Blood alkaline phosphatase increased subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 1	0 / 5 (0.00%) 0	1 / 16 (6.25%) 3
Blood bilirubin increased subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 1	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Blood creatine increased subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Blood creatine phosphokinase increased subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	4 / 16 (25.00%) 7
Blood creatinine increased			

subjects affected / exposed	2 / 4 (50.00%)	2 / 5 (40.00%)	8 / 16 (50.00%)
occurrences (all)	3	7	12
Blood glucose increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	2
Blood potassium increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Blood triglycerides increased			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	1	0	0
Blood urea increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Blood uric acid increased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
C-reactive protein increased			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	1	0	0
Electrocardiogram QT prolonged			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	1	0	0
Gamma-glutamyltransferase increased			
subjects affected / exposed	1 / 4 (25.00%)	1 / 5 (20.00%)	2 / 16 (12.50%)
occurrences (all)	1	2	3
Haemoglobin decreased			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	4	0	0
Lipase increased			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	4 / 16 (25.00%)
occurrences (all)	0	2	6
Lymphocyte count decreased			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	6

Protein total decreased subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Weight decreased subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	3 / 16 (18.75%) 3
Weight increased subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	2 / 5 (40.00%) 2	2 / 16 (12.50%) 2
White blood cell count decreased subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 2
Injury, poisoning and procedural complications Procedural pain subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 1	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Traumatic haematoma subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Cardiac disorders Angina pectoris subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Bradycardia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Nervous system disorders Amnesia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 2
Dizziness subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	4 / 16 (25.00%) 7
Dysaesthesia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0

Dysgeusia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Epilepsy			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Hypoaesthesia			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	1 / 16 (6.25%)
occurrences (all)	0	1	2
Lethargy			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences (all)	0	0	2
Headache			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	7 / 16 (43.75%)
occurrences (all)	0	0	12
Neuropathy peripheral			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Seizure			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Post herpetic neuralgia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Syncope			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	2 / 16 (12.50%)
occurrences (all)	0	2	2
Febrile neutropenia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Leukopenia			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Lymphopenia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Neutropenia			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	2	0
Thrombocytopenia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Ear and labyrinth disorders			
Deafness			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences (all)	0	0	4
Ear pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	2
Deafness transitory			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Hypoacusis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Tinnitus			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Eye disorders			
Eye pruritus			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Eye swelling			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Eyelid oedema			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Vision blurred			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Vitreous floaters			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Gastrointestinal disorders			
Abdominal distension			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Abdominal discomfort			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Abdominal pain			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	2 / 16 (12.50%)
occurrences (all)	0	1	2
Abdominal pain upper			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences (all)	1	0	3
Constipation			
subjects affected / exposed	1 / 4 (25.00%)	2 / 5 (40.00%)	5 / 16 (31.25%)
occurrences (all)	1	2	11
Diarrhoea			
subjects affected / exposed	1 / 4 (25.00%)	2 / 5 (40.00%)	6 / 16 (37.50%)
occurrences (all)	1	3	18
Dry mouth			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Dyspepsia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	4 / 16 (25.00%)
occurrences (all)	0	0	6
Dysphagia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	3 / 16 (18.75%)
occurrences (all)	0	0	3

Gastrooesophageal reflux disease subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Lip blister subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 2	0 / 16 (0.00%) 0
Lip dry subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Mouth ulceration subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Nausea subjects affected / exposed occurrences (all)	2 / 4 (50.00%) 2	5 / 5 (100.00%) 5	11 / 16 (68.75%) 27
Odynophagia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Toothache subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 2
Stomatitis subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	2 / 5 (40.00%) 3	3 / 16 (18.75%) 8
Vomiting subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	3 / 5 (60.00%) 6	6 / 16 (37.50%) 20
Hepatobiliary disorders Hypertransaminaemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Skin and subcutaneous tissue disorders Acne subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 1	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Dermatitis acneiform			

subjects affected / exposed	0 / 4 (0.00%)	2 / 5 (40.00%)	2 / 16 (12.50%)
occurrences (all)	0	4	2
Alopecia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences (all)	0	0	5
Dry skin			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	5 / 16 (31.25%)
occurrences (all)	0	4	7
Eczema			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Ingrowing nail			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Nail discolouration			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Night sweats			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Palmar-plantar erythrodysaesthesia syndrome			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Pruritus			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	5 / 16 (31.25%)
occurrences (all)	0	2	6
Rash			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Rash macular			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	1 / 16 (6.25%)
occurrences (all)	0	1	1
Rash maculo-papular			
subjects affected / exposed	0 / 4 (0.00%)	4 / 5 (80.00%)	3 / 16 (18.75%)
occurrences (all)	0	4	5

Rash papular subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Rash pruritic subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	2 / 16 (12.50%) 3
Skin hyperpigmentation subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0
Urticaria subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Renal and urinary disorders Acute kidney injury subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 1	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Chronic kidney disease subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Hypertonic bladder subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Renal failure subjects affected / exposed occurrences (all)	1 / 4 (25.00%) 1	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0
Endocrine disorders Hyperthyroidism subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Musculoskeletal and connective tissue disorders Arthralgia subjects affected / exposed occurrences (all)	2 / 4 (50.00%) 2	1 / 5 (20.00%) 1	2 / 16 (12.50%) 3
Back pain subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	0 / 16 (0.00%) 0

Bone pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Flank pain			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	1	0	1
Joint stiffness			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Muscular weakness			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Muscle spasms			
subjects affected / exposed	0 / 4 (0.00%)	2 / 5 (40.00%)	2 / 16 (12.50%)
occurrences (all)	0	2	3
Musculoskeletal pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Musculoskeletal chest pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Myalgia			
subjects affected / exposed	1 / 4 (25.00%)	1 / 5 (20.00%)	2 / 16 (12.50%)
occurrences (all)	2	1	2
Neck pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Pain in extremity			
subjects affected / exposed	0 / 4 (0.00%)	3 / 5 (60.00%)	2 / 16 (12.50%)
occurrences (all)	0	6	2
Tendon pain			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Infections and infestations			
Bronchitis			

subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	2	0
Conjunctivitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Cellulitis			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Cystitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Folliculitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Gastroenteritis			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	1	0	0
Herpes zoster			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences (all)	0	0	2
Nasopharyngitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Infection			
subjects affected / exposed	1 / 4 (25.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	1	0	0
Paronychia			
subjects affected / exposed	0 / 4 (0.00%)	2 / 5 (40.00%)	5 / 16 (31.25%)
occurrences (all)	0	3	6
Pharyngitis			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Pneumonia			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	2 / 16 (12.50%)
occurrences (all)	0	1	2
Respiratory tract infection			

subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Rhinitis			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Sinobronchitis			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Skin infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Tooth infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Upper respiratory tract infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	4 / 16 (25.00%)
occurrences (all)	0	0	5
Urinary tract infection			
subjects affected / exposed	0 / 4 (0.00%)	1 / 5 (20.00%)	0 / 16 (0.00%)
occurrences (all)	0	1	0
Viral infection			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	1
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	1 / 4 (25.00%)	1 / 5 (20.00%)	10 / 16 (62.50%)
occurrences (all)	1	2	17
Hyperglycaemia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	0 / 16 (0.00%)
occurrences (all)	0	0	0
Hyperkalaemia			
subjects affected / exposed	2 / 4 (50.00%)	0 / 5 (0.00%)	2 / 16 (12.50%)
occurrences (all)	3	0	2
Hyperlipasaemia			
subjects affected / exposed	0 / 4 (0.00%)	0 / 5 (0.00%)	1 / 16 (6.25%)
occurrences (all)	0	0	2

Hypoalbuminaemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	5 / 16 (31.25%) 8
Hypocalcaemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 2
Hypoglycaemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Hypokalaemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	2 / 16 (12.50%) 3
Hyponatraemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	3 / 16 (18.75%) 9
Hypomagnesaemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	3 / 16 (18.75%) 16
Hypophosphataemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	2 / 16 (12.50%) 15
Hypophagia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	0 / 5 (0.00%) 0	1 / 16 (6.25%) 1
Hypoproteinaemia subjects affected / exposed occurrences (all)	0 / 4 (0.00%) 0	1 / 5 (20.00%) 1	0 / 16 (0.00%) 0

Non-serious adverse events	Phase IB part- INC280 400mg BID/ EGF816 75mg QD	Phase IB part- INC280 200mg BID/ EGF816 100mg QD	All Participants
Total subjects affected by non-serious adverse events subjects affected / exposed	3 / 3 (100.00%)	5 / 5 (100.00%)	176 / 177 (99.44%)
Vascular disorders Hot flush subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	3 / 177 (1.69%) 3
Hypotension			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	10 / 177 (5.65%)
occurrences (all)	0	0	12
Pallor			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	2
Thrombosis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
General disorders and administration site conditions			
Chest pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Asthenia			
subjects affected / exposed	1 / 3 (33.33%)	2 / 5 (40.00%)	44 / 177 (24.86%)
occurrences (all)	2	3	62
Chills			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	0	0	8
Fatigue			
subjects affected / exposed	2 / 3 (66.67%)	0 / 5 (0.00%)	46 / 177 (25.99%)
occurrences (all)	3	0	57
Face oedema			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	9 / 177 (5.08%)
occurrences (all)	0	0	9
Gait disturbance			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Influenza like illness			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	10 / 177 (5.65%)
occurrences (all)	0	0	15
Non-cardiac chest pain			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	23 / 177 (12.99%)
occurrences (all)	1	0	26
Oedema peripheral			

subjects affected / exposed occurrences (all)	2 / 3 (66.67%) 2	1 / 5 (20.00%) 5	103 / 177 (58.19%) 226
Pain subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	6 / 177 (3.39%) 7
Peripheral swelling subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	7 / 177 (3.95%) 10
Pyrexia subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 5 (20.00%) 1	43 / 177 (24.29%) 55
Immune system disorders Anaphylactoid reaction subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Hypersensitivity subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 5 (20.00%) 2	3 / 177 (1.69%) 5
Reproductive system and breast disorders Pelvic pain subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Perineal pain subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Vulvovaginal pruritus subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Prostatomegaly subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Respiratory, thoracic and mediastinal disorders Cough subjects affected / exposed occurrences (all)	2 / 3 (66.67%) 2	2 / 5 (40.00%) 2	47 / 177 (26.55%) 75
Dry throat			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Dysphonia			
subjects affected / exposed	0 / 3 (0.00%)	1 / 5 (20.00%)	5 / 177 (2.82%)
occurrences (all)	0	1	5
Dyspnoea			
subjects affected / exposed	1 / 3 (33.33%)	1 / 5 (20.00%)	40 / 177 (22.60%)
occurrences (all)	1	1	48
Dyspnoea exertional			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Epistaxis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	0	0	5
Haemoptysis			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	1	0	5
Nasal discomfort			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	2
Hypoxia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	4
Nasal dryness			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Oropharyngeal pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	8 / 177 (4.52%)
occurrences (all)	0	0	8
Pleural effusion			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	9 / 177 (5.08%)
occurrences (all)	0	0	9
Pneumothorax			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	3
Productive cough			

subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	5 / 177 (2.82%) 7
Pulmonary embolism subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	3 / 177 (1.69%) 3
Rhinorrhoea subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 5 (0.00%) 0	13 / 177 (7.34%) 17
Throat irritation subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 5 (20.00%) 1	1 / 177 (0.56%) 1
Psychiatric disorders Anxiety subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	6 / 177 (3.39%) 6
Confusional state subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	3 / 177 (1.69%) 3
Sleep disorder subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	3 / 177 (1.69%) 3
Depression subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	2 / 177 (1.13%) 2
Insomnia subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	12 / 177 (6.78%) 14
Investigations Alanine aminotransferase increased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	41 / 177 (23.16%) 54
Amylase increased subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	2 / 5 (40.00%) 5	45 / 177 (25.42%) 103
Aspartate aminotransferase increased			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	34 / 177 (19.21%)
occurrences (all)	0	0	47
Blood albumin decreased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	0	0	12
Blood alkaline phosphatase increased			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	12 / 177 (6.78%)
occurrences (all)	1	0	18
Blood bilirubin increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	6 / 177 (3.39%)
occurrences (all)	0	0	10
Blood creatine increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	3
Blood creatine phosphokinase increased			
subjects affected / exposed	1 / 3 (33.33%)	1 / 5 (20.00%)	22 / 177 (12.43%)
occurrences (all)	1	6	42
Blood creatinine increased			
subjects affected / exposed	0 / 3 (0.00%)	2 / 5 (40.00%)	56 / 177 (31.64%)
occurrences (all)	0	3	108
Blood glucose increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	2
Blood potassium increased			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	1	0	1
Blood triglycerides increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Blood urea increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	4
Blood uric acid increased			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2

C-reactive protein increased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Electrocardiogram QT prolonged subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	7 / 177 (3.95%) 13
Gamma-glutamyltransferase increased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	16 / 177 (9.04%) 23
Haemoglobin decreased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 4
Lipase increased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 5 (20.00%) 3	33 / 177 (18.64%) 81
Lymphocyte count decreased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	2 / 177 (1.13%) 7
Protein total decreased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	8 / 177 (4.52%) 8
Weight decreased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	7 / 177 (3.95%) 7
Weight increased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	13 / 177 (7.34%) 13
White blood cell count decreased subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	3 / 177 (1.69%) 4
Injury, poisoning and procedural complications Procedural pain subjects affected / exposed occurrences (all) Traumatic haematoma	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	2 / 177 (1.13%) 2

subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	2 / 177 (1.13%) 2
Cardiac disorders			
Angina pectoris			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	3
Bradycardia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Nervous system disorders			
Amnesia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	3
Dizziness			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	36 / 177 (20.34%)
occurrences (all)	2	0	45
Dysaesthesia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	1	0	3
Dysgeusia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	6 / 177 (3.39%)
occurrences (all)	0	0	6
Epilepsy			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Hypoaesthesia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	8 / 177 (4.52%)
occurrences (all)	0	0	9
Lethargy			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	4
Headache			
subjects affected / exposed	0 / 3 (0.00%)	1 / 5 (20.00%)	45 / 177 (25.42%)
occurrences (all)	0	1	59
Neuropathy peripheral			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	3
Seizure			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	0	0	9
Post herpetic neuralgia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Syncope			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	1	0	1
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	2 / 3 (66.67%)	0 / 5 (0.00%)	30 / 177 (16.95%)
occurrences (all)	2	0	39
Febrile neutropenia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	1	0	2
Leukopenia			
subjects affected / exposed	1 / 3 (33.33%)	1 / 5 (20.00%)	4 / 177 (2.26%)
occurrences (all)	3	1	6
Lymphopenia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	1	0	11
Neutropenia			
subjects affected / exposed	1 / 3 (33.33%)	2 / 5 (40.00%)	6 / 177 (3.39%)
occurrences (all)	4	2	10
Thrombocytopenia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	10 / 177 (5.65%)
occurrences (all)	3	0	12
Ear and labyrinth disorders			
Deafness			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	1	0	7
Ear pain			

subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	2 / 177 (1.13%) 3
Deafness transitory subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Hypoacusis subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	9 / 177 (5.08%) 9
Tinnitus subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	7 / 177 (3.95%) 7
Eye disorders			
Eye pruritus subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	5 / 177 (2.82%) 5
Eye swelling subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Eyelid oedema subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 5 (0.00%) 0	3 / 177 (1.69%) 3
Vision blurred subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Vitreous floaters subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Gastrointestinal disorders			
Abdominal distension subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	13 / 177 (7.34%) 13
Abdominal discomfort subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	4 / 177 (2.26%) 4
Abdominal pain			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	14 / 177 (7.91%)
occurrences (all)	0	0	17
Abdominal pain upper			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	20 / 177 (11.30%)
occurrences (all)	0	0	22
Constipation			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	28 / 177 (15.82%)
occurrences (all)	3	0	42
Diarrhoea			
subjects affected / exposed	1 / 3 (33.33%)	1 / 5 (20.00%)	79 / 177 (44.63%)
occurrences (all)	1	1	132
Dry mouth			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	0	0	7
Dyspepsia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	30 / 177 (16.95%)
occurrences (all)	1	0	33
Dysphagia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	8 / 177 (4.52%)
occurrences (all)	0	0	8
Gastrooesophageal reflux disease			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	12 / 177 (6.78%)
occurrences (all)	0	0	13
Lip blister			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	2
Lip dry			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Mouth ulceration			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	9 / 177 (5.08%)
occurrences (all)	1	0	9
Nausea			
subjects affected / exposed	2 / 3 (66.67%)	0 / 5 (0.00%)	97 / 177 (54.80%)
occurrences (all)	3	0	150
Odynophagia			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	3
Toothache			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	5
Stomatitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	25 / 177 (14.12%)
occurrences (all)	0	0	49
Vomiting			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	64 / 177 (36.16%)
occurrences (all)	1	0	106
Hepatobiliary disorders			
Hypertransaminasaemia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	0	0	5
Skin and subcutaneous tissue disorders			
Acne			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	6
Dermatitis acneiform			
subjects affected / exposed	0 / 3 (0.00%)	1 / 5 (20.00%)	21 / 177 (11.86%)
occurrences (all)	0	1	26
Alopecia			
subjects affected / exposed	1 / 3 (33.33%)	1 / 5 (20.00%)	8 / 177 (4.52%)
occurrences (all)	1	1	11
Dry skin			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	23 / 177 (12.99%)
occurrences (all)	0	0	32
Eczema			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	9 / 177 (5.08%)
occurrences (all)	0	0	11
Ingrowing nail			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Nail discolouration			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Night sweats			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Palmar-plantar erythrodysaesthesia syndrome			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Pruritus			
subjects affected / exposed	0 / 3 (0.00%)	1 / 5 (20.00%)	36 / 177 (20.34%)
occurrences (all)	0	1	50
Rash			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	28 / 177 (15.82%)
occurrences (all)	0	0	32
Rash macular			
subjects affected / exposed	1 / 3 (33.33%)	1 / 5 (20.00%)	9 / 177 (5.08%)
occurrences (all)	1	1	10
Rash maculo-papular			
subjects affected / exposed	0 / 3 (0.00%)	2 / 5 (40.00%)	43 / 177 (24.29%)
occurrences (all)	0	2	56
Rash papular			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	0	0	6
Rash pruritic			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	0	0	9
Skin hyperpigmentation			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Urticaria			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	0	0	5
Renal and urinary disorders			
Acute kidney injury			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Chronic kidney disease			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Hypertonic bladder			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	1	0	1
Renal failure			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Endocrine disorders			
Hyperthyroidism			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	3
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	1 / 3 (33.33%)	1 / 5 (20.00%)	26 / 177 (14.69%)
occurrences (all)	2	2	32
Back pain			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	18 / 177 (10.17%)
occurrences (all)	1	0	20
Bone pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	6 / 177 (3.39%)
occurrences (all)	0	0	11
Flank pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	4
Joint stiffness			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Muscular weakness			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Muscle spasms			

subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	35 / 177 (19.77%)
occurrences (all)	2	0	45
Musculoskeletal pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	4
Musculoskeletal chest pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	0	0	8
Myalgia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	10 / 177 (5.65%)
occurrences (all)	1	0	12
Neck pain			
subjects affected / exposed	0 / 3 (0.00%)	1 / 5 (20.00%)	2 / 177 (1.13%)
occurrences (all)	0	1	2
Pain in extremity			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	23 / 177 (12.99%)
occurrences (all)	0	0	28
Tendon pain			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Infections and infestations			
Bronchitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	0	0	10
Conjunctivitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	4
Cellulitis			
subjects affected / exposed	0 / 3 (0.00%)	2 / 5 (40.00%)	12 / 177 (6.78%)
occurrences (all)	0	3	17
Cystitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	2 / 177 (1.13%)
occurrences (all)	0	0	2
Folliculitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	0	0	5

Gastroenteritis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	3
Herpes zoster			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	0	0	6
Nasopharyngitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	16 / 177 (9.04%)
occurrences (all)	0	0	19
Infection			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Paronychia			
subjects affected / exposed	0 / 3 (0.00%)	1 / 5 (20.00%)	25 / 177 (14.12%)
occurrences (all)	0	1	33
Pharyngitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	3 / 177 (1.69%)
occurrences (all)	0	0	3
Pneumonia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	9 / 177 (5.08%)
occurrences (all)	0	0	9
Respiratory tract infection			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	5
Rhinitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	0	0	10
Sinobronchitis			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Skin infection			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	4 / 177 (2.26%)
occurrences (all)	0	0	5
Tooth infection			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1

Upper respiratory tract infection subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	25 / 177 (14.12%) 34
Urinary tract infection subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	18 / 177 (10.17%) 31
Viral infection subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 1
Metabolism and nutrition disorders			
Decreased appetite subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	50 / 177 (28.25%) 63
Hyperglycaemia subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 2	0 / 5 (0.00%) 0	6 / 177 (3.39%) 8
Hyperkalaemia subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	8 / 177 (4.52%) 14
Hyperlipasaemia subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	1 / 177 (0.56%) 2
Hypoalbuminaemia subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	1 / 5 (20.00%) 1	34 / 177 (19.21%) 52
Hypocalcaemia subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 2	0 / 5 (0.00%) 0	7 / 177 (3.95%) 10
Hypoglycaemia subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	4 / 177 (2.26%) 5
Hypokalaemia subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 5 (0.00%) 0	15 / 177 (8.47%) 20
Hyponatraemia			

subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	10 / 177 (5.65%)
occurrences (all)	0	0	19
Hypomagnesaemia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	15 / 177 (8.47%)
occurrences (all)	0	0	33
Hypophosphataemia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	7 / 177 (3.95%)
occurrences (all)	3	0	22
Hypophagia			
subjects affected / exposed	0 / 3 (0.00%)	0 / 5 (0.00%)	1 / 177 (0.56%)
occurrences (all)	0	0	1
Hypoproteinaemia			
subjects affected / exposed	1 / 3 (33.33%)	0 / 5 (0.00%)	5 / 177 (2.82%)
occurrences (all)	3	0	7

Non-serious adverse events	Phase II- Group 4	Phase II- Group 3	Phase II- Group 2
Total subjects affected by non-serious adverse events			
subjects affected / exposed	42 / 42 (100.00%)	47 / 47 (100.00%)	3 / 3 (100.00%)
Vascular disorders			
Hot flush			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	0	2	0
Hypotension			
subjects affected / exposed	1 / 42 (2.38%)	4 / 47 (8.51%)	1 / 3 (33.33%)
occurrences (all)	2	5	1
Pallor			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	1 / 3 (33.33%)
occurrences (all)	0	0	2
Thrombosis			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
General disorders and administration site conditions			
Chest pain			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Asthenia			

subjects affected / exposed	10 / 42 (23.81%)	18 / 47 (38.30%)	1 / 3 (33.33%)
occurrences (all)	14	22	2
Chills			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	0	3	0
Fatigue			
subjects affected / exposed	9 / 42 (21.43%)	9 / 47 (19.15%)	2 / 3 (66.67%)
occurrences (all)	11	10	3
Face oedema			
subjects affected / exposed	1 / 42 (2.38%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	1	2	0
Gait disturbance			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	1	0	0
Influenza like illness			
subjects affected / exposed	5 / 42 (11.90%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	8	3	0
Non-cardiac chest pain			
subjects affected / exposed	6 / 42 (14.29%)	6 / 47 (12.77%)	1 / 3 (33.33%)
occurrences (all)	6	8	1
Oedema peripheral			
subjects affected / exposed	26 / 42 (61.90%)	31 / 47 (65.96%)	3 / 3 (100.00%)
occurrences (all)	42	93	3
Pain			
subjects affected / exposed	1 / 42 (2.38%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	1	3	0
Peripheral swelling			
subjects affected / exposed	3 / 42 (7.14%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	4	1	0
Pyrexia			
subjects affected / exposed	12 / 42 (28.57%)	11 / 47 (23.40%)	0 / 3 (0.00%)
occurrences (all)	14	13	0
Immune system disorders			
Anaphylactoid reaction			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0

Hypersensitivity subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Reproductive system and breast disorders			
Pelvic pain subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Perineal pain subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Vulvovaginal pruritus subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	1 / 3 (33.33%) 1
Prostatomegaly subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Respiratory, thoracic and mediastinal disorders			
Cough subjects affected / exposed occurrences (all)	14 / 42 (33.33%) 23	11 / 47 (23.40%) 13	1 / 3 (33.33%) 3
Dry throat subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Dysphonia subjects affected / exposed occurrences (all)	1 / 42 (2.38%) 1	1 / 47 (2.13%) 1	1 / 3 (33.33%) 1
Dyspnoea subjects affected / exposed occurrences (all)	7 / 42 (16.67%) 7	13 / 47 (27.66%) 13	1 / 3 (33.33%) 1
Dyspnoea exertional subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Epistaxis subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	4 / 47 (8.51%) 4	0 / 3 (0.00%) 0
Haemoptysis			

subjects affected / exposed	2 / 42 (4.76%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	3	0	0
Nasal discomfort			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Hypoxia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Nasal dryness			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Oropharyngeal pain			
subjects affected / exposed	5 / 42 (11.90%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	5	1	0
Pleural effusion			
subjects affected / exposed	3 / 42 (7.14%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	3	2	0
Pneumothorax			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	0	1	0
Productive cough			
subjects affected / exposed	0 / 42 (0.00%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	0	5	0
Pulmonary embolism			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	0	1	0
Rhinorrhoea			
subjects affected / exposed	5 / 42 (11.90%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	5	2	0
Throat irritation			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Psychiatric disorders			
Anxiety			
subjects affected / exposed	0 / 42 (0.00%)	4 / 47 (8.51%)	0 / 3 (0.00%)
occurrences (all)	0	4	0

Confusional state			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	0	2	0
Sleep disorder			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Depression			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	1	0	0
Insomnia			
subjects affected / exposed	4 / 42 (9.52%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	5	3	0
Investigations			
Alanine aminotransferase increased			
subjects affected / exposed	13 / 42 (30.95%)	13 / 47 (27.66%)	2 / 3 (66.67%)
occurrences (all)	17	19	2
Amylase increased			
subjects affected / exposed	7 / 42 (16.67%)	11 / 47 (23.40%)	1 / 3 (33.33%)
occurrences (all)	17	23	3
Aspartate aminotransferase increased			
subjects affected / exposed	11 / 42 (26.19%)	13 / 47 (27.66%)	2 / 3 (66.67%)
occurrences (all)	15	20	2
Blood albumin decreased			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	1	1	0
Blood alkaline phosphatase increased			
subjects affected / exposed	2 / 42 (4.76%)	6 / 47 (12.77%)	0 / 3 (0.00%)
occurrences (all)	2	10	0
Blood bilirubin increased			
subjects affected / exposed	1 / 42 (2.38%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	1	6	0
Blood creatine increased			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	2	0	0
Blood creatine phosphokinase increased			

subjects affected / exposed	6 / 42 (14.29%)	5 / 47 (10.64%)	0 / 3 (0.00%)
occurrences (all)	10	6	0
Blood creatinine increased			
subjects affected / exposed	15 / 42 (35.71%)	16 / 47 (34.04%)	1 / 3 (33.33%)
occurrences (all)	35	30	4
Blood glucose increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Blood potassium increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Blood triglycerides increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Blood urea increased			
subjects affected / exposed	2 / 42 (4.76%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	2	1	0
Blood uric acid increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	1 / 3 (33.33%)
occurrences (all)	0	0	1
C-reactive protein increased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Electrocardiogram QT prolonged			
subjects affected / exposed	3 / 42 (7.14%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	9	1	0
Gamma-glutamyltransferase increased			
subjects affected / exposed	5 / 42 (11.90%)	5 / 47 (10.64%)	0 / 3 (0.00%)
occurrences (all)	9	6	0
Haemoglobin decreased			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Lipase increased			
subjects affected / exposed	8 / 42 (19.05%)	7 / 47 (14.89%)	0 / 3 (0.00%)
occurrences (all)	11	19	0

Lymphocyte count decreased subjects affected / exposed occurrences (all)	1 / 42 (2.38%) 1	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Protein total decreased subjects affected / exposed occurrences (all)	3 / 42 (7.14%) 3	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Weight decreased subjects affected / exposed occurrences (all)	2 / 42 (4.76%) 2	1 / 47 (2.13%) 1	0 / 3 (0.00%) 0
Weight increased subjects affected / exposed occurrences (all)	2 / 42 (4.76%) 2	4 / 47 (8.51%) 4	0 / 3 (0.00%) 0
White blood cell count decreased subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	2 / 47 (4.26%) 2	0 / 3 (0.00%) 0
Injury, poisoning and procedural complications Procedural pain subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	1 / 47 (2.13%) 1	0 / 3 (0.00%) 0
Traumatic haematoma subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	1 / 47 (2.13%) 1	0 / 3 (0.00%) 0
Cardiac disorders Angina pectoris subjects affected / exposed occurrences (all)	1 / 42 (2.38%) 1	0 / 47 (0.00%) 0	1 / 3 (33.33%) 1
Bradycardia subjects affected / exposed occurrences (all)	1 / 42 (2.38%) 1	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Nervous system disorders Amnesia subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	1 / 47 (2.13%) 1	0 / 3 (0.00%) 0
Dizziness subjects affected / exposed occurrences (all)	10 / 42 (23.81%) 13	11 / 47 (23.40%) 12	0 / 3 (0.00%) 0

Dysaesthesia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Dysgeusia			
subjects affected / exposed	1 / 42 (2.38%)	4 / 47 (8.51%)	1 / 3 (33.33%)
occurrences (all)	1	4	1
Epilepsy			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	1 / 3 (33.33%)
occurrences (all)	0	0	1
Hypoaesthesia			
subjects affected / exposed	2 / 42 (4.76%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	2	3	0
Lethargy			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	1	1	0
Headache			
subjects affected / exposed	12 / 42 (28.57%)	17 / 47 (36.17%)	1 / 3 (33.33%)
occurrences (all)	14	23	1
Neuropathy peripheral			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Seizure			
subjects affected / exposed	1 / 42 (2.38%)	3 / 47 (6.38%)	1 / 3 (33.33%)
occurrences (all)	1	4	2
Post herpetic neuralgia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Syncope			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	4 / 42 (9.52%)	11 / 47 (23.40%)	1 / 3 (33.33%)
occurrences (all)	4	17	2
Febrile neutropenia			

subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Leukopenia			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	0	2	0
Lymphopenia			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	3	2	0
Neutropenia			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	0	1	0
Thrombocytopenia			
subjects affected / exposed	2 / 42 (4.76%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	2	3	0
Ear and labyrinth disorders			
Deafness			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	0	2	0
Ear pain			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Deafness transitory			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Hypoacusis			
subjects affected / exposed	3 / 42 (7.14%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	3	3	0
Tinnitus			
subjects affected / exposed	0 / 42 (0.00%)	6 / 47 (12.77%)	0 / 3 (0.00%)
occurrences (all)	0	6	0
Eye disorders			
Eye pruritus			
subjects affected / exposed	1 / 42 (2.38%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	1	3	0
Eye swelling			

subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Eyelid oedema subjects affected / exposed occurrences (all)	1 / 42 (2.38%) 1	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Vision blurred subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Vitreous floaters subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Gastrointestinal disorders			
Abdominal distension subjects affected / exposed occurrences (all)	4 / 42 (9.52%) 4	7 / 47 (14.89%) 7	0 / 3 (0.00%) 0
Abdominal discomfort subjects affected / exposed occurrences (all)	3 / 42 (7.14%) 3	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Abdominal pain subjects affected / exposed occurrences (all)	2 / 42 (4.76%) 3	5 / 47 (10.64%) 5	0 / 3 (0.00%) 0
Abdominal pain upper subjects affected / exposed occurrences (all)	7 / 42 (16.67%) 7	6 / 47 (12.77%) 7	0 / 3 (0.00%) 0
Constipation subjects affected / exposed occurrences (all)	8 / 42 (19.05%) 9	6 / 47 (12.77%) 9	1 / 3 (33.33%) 1
Diarrhoea subjects affected / exposed occurrences (all)	20 / 42 (47.62%) 30	26 / 47 (55.32%) 37	2 / 3 (66.67%) 2
Dry mouth subjects affected / exposed occurrences (all)	3 / 42 (7.14%) 3	1 / 47 (2.13%) 1	0 / 3 (0.00%) 0
Dyspepsia subjects affected / exposed occurrences (all)	6 / 42 (14.29%) 6	9 / 47 (19.15%) 9	0 / 3 (0.00%) 0

Dysphagia			
subjects affected / exposed	3 / 42 (7.14%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	3	2	0
Gastrooesophageal reflux disease			
subjects affected / exposed	6 / 42 (14.29%)	2 / 47 (4.26%)	1 / 3 (33.33%)
occurrences (all)	7	2	1
Lip blister			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Lip dry			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Mouth ulceration			
subjects affected / exposed	4 / 42 (9.52%)	4 / 47 (8.51%)	0 / 3 (0.00%)
occurrences (all)	4	4	0
Nausea			
subjects affected / exposed	19 / 42 (45.24%)	27 / 47 (57.45%)	3 / 3 (100.00%)
occurrences (all)	31	38	3
Odynophagia			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	1	1	0
Toothache			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	0	2	0
Stomatitis			
subjects affected / exposed	3 / 42 (7.14%)	12 / 47 (25.53%)	0 / 3 (0.00%)
occurrences (all)	13	20	0
Vomiting			
subjects affected / exposed	11 / 42 (26.19%)	22 / 47 (46.81%)	1 / 3 (33.33%)
occurrences (all)	12	33	1
Hepatobiliary disorders			
Hypertransaminasaemia			
subjects affected / exposed	1 / 42 (2.38%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	1	3	0
Skin and subcutaneous tissue disorders			

Acne			
subjects affected / exposed	0 / 42 (0.00%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	0	5	0
Dermatitis acneiform			
subjects affected / exposed	4 / 42 (9.52%)	10 / 47 (21.28%)	0 / 3 (0.00%)
occurrences (all)	5	11	0
Alopecia			
subjects affected / exposed	0 / 42 (0.00%)	4 / 47 (8.51%)	0 / 3 (0.00%)
occurrences (all)	0	4	0
Dry skin			
subjects affected / exposed	5 / 42 (11.90%)	7 / 47 (14.89%)	0 / 3 (0.00%)
occurrences (all)	5	11	0
Eczema			
subjects affected / exposed	6 / 42 (14.29%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	8	1	0
Ingrowing nail			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	0	1	0
Nail discolouration			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Night sweats			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	0	1	0
Palmar-plantar erythrodysaesthesia syndrome			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Pruritus			
subjects affected / exposed	14 / 42 (33.33%)	9 / 47 (19.15%)	0 / 3 (0.00%)
occurrences (all)	19	14	0
Rash			
subjects affected / exposed	9 / 42 (21.43%)	11 / 47 (23.40%)	0 / 3 (0.00%)
occurrences (all)	10	13	0
Rash macular			

subjects affected / exposed occurrences (all)	1 / 42 (2.38%) 1	3 / 47 (6.38%) 4	1 / 3 (33.33%) 1
Rash maculo-papular subjects affected / exposed occurrences (all)	9 / 42 (21.43%) 13	14 / 47 (29.79%) 21	1 / 3 (33.33%) 1
Rash papular subjects affected / exposed occurrences (all)	2 / 42 (4.76%) 3	2 / 47 (4.26%) 2	0 / 3 (0.00%) 0
Rash pruritic subjects affected / exposed occurrences (all)	1 / 42 (2.38%) 1	3 / 47 (6.38%) 4	0 / 3 (0.00%) 0
Skin hyperpigmentation subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	1 / 47 (2.13%) 1	0 / 3 (0.00%) 0
Urticaria subjects affected / exposed occurrences (all)	2 / 42 (4.76%) 2	2 / 47 (4.26%) 2	0 / 3 (0.00%) 0
Renal and urinary disorders Acute kidney injury subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Chronic kidney disease subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Hypertonic bladder subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Renal failure subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	0 / 47 (0.00%) 0	0 / 3 (0.00%) 0
Endocrine disorders Hyperthyroidism subjects affected / exposed occurrences (all)	0 / 42 (0.00%) 0	1 / 47 (2.13%) 1	0 / 3 (0.00%) 0
Musculoskeletal and connective tissue disorders			

Arthralgia			
subjects affected / exposed	6 / 42 (14.29%)	10 / 47 (21.28%)	0 / 3 (0.00%)
occurrences (all)	6	13	0
Back pain			
subjects affected / exposed	8 / 42 (19.05%)	7 / 47 (14.89%)	0 / 3 (0.00%)
occurrences (all)	9	8	0
Bone pain			
subjects affected / exposed	1 / 42 (2.38%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	1	8	0
Flank pain			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	0	1	0
Joint stiffness			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Muscular weakness			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Muscle spasms			
subjects affected / exposed	14 / 42 (33.33%)	12 / 47 (25.53%)	0 / 3 (0.00%)
occurrences (all)	19	15	0
Musculoskeletal pain			
subjects affected / exposed	0 / 42 (0.00%)	4 / 47 (8.51%)	0 / 3 (0.00%)
occurrences (all)	0	4	0
Musculoskeletal chest pain			
subjects affected / exposed	2 / 42 (4.76%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	2	4	0
Myalgia			
subjects affected / exposed	0 / 42 (0.00%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	0	3	0
Neck pain			
subjects affected / exposed	0 / 42 (0.00%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	0	1	0
Pain in extremity			
subjects affected / exposed	7 / 42 (16.67%)	7 / 47 (14.89%)	0 / 3 (0.00%)
occurrences (all)	8	8	0

Tendon pain			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	1 / 3 (33.33%)
occurrences (all)	0	0	1
Infections and infestations			
Bronchitis			
subjects affected / exposed	2 / 42 (4.76%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	2	4	0
Conjunctivitis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	1	0	0
Cellulitis			
subjects affected / exposed	5 / 42 (11.90%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	8	4	0
Cystitis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	1 / 3 (33.33%)
occurrences (all)	1	0	1
Folliculitis			
subjects affected / exposed	1 / 42 (2.38%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	1	3	0
Gastroenteritis			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	1	0	0
Herpes zoster			
subjects affected / exposed	3 / 42 (7.14%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	4	0	0
Nasopharyngitis			
subjects affected / exposed	5 / 42 (11.90%)	5 / 47 (10.64%)	0 / 3 (0.00%)
occurrences (all)	5	7	0
Infection			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Paronychia			
subjects affected / exposed	7 / 42 (16.67%)	7 / 47 (14.89%)	0 / 3 (0.00%)
occurrences (all)	8	8	0
Pharyngitis			

subjects affected / exposed	2 / 42 (4.76%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	2	0	0
Pneumonia			
subjects affected / exposed	2 / 42 (4.76%)	2 / 47 (4.26%)	0 / 3 (0.00%)
occurrences (all)	2	2	0
Respiratory tract infection			
subjects affected / exposed	0 / 42 (0.00%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	0	4	0
Rhinitis			
subjects affected / exposed	1 / 42 (2.38%)	2 / 47 (4.26%)	1 / 3 (33.33%)
occurrences (all)	1	2	1
Sinobronchitis			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Skin infection			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	1	2	0
Tooth infection			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Upper respiratory tract infection			
subjects affected / exposed	6 / 42 (14.29%)	6 / 47 (12.77%)	0 / 3 (0.00%)
occurrences (all)	10	8	0
Urinary tract infection			
subjects affected / exposed	8 / 42 (19.05%)	4 / 47 (8.51%)	0 / 3 (0.00%)
occurrences (all)	19	4	0
Viral infection			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	9 / 42 (21.43%)	14 / 47 (29.79%)	1 / 3 (33.33%)
occurrences (all)	11	15	1
Hyperglycaemia			
subjects affected / exposed	2 / 42 (4.76%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	2	4	0

Hyperkalaemia			
subjects affected / exposed	1 / 42 (2.38%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	1	0	0
Hyperlipasaemia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Hypoalbuminaemia			
subjects affected / exposed	7 / 42 (16.67%)	10 / 47 (21.28%)	0 / 3 (0.00%)
occurrences (all)	9	15	0
Hypocalcaemia			
subjects affected / exposed	0 / 42 (0.00%)	3 / 47 (6.38%)	0 / 3 (0.00%)
occurrences (all)	0	4	0
Hypoglycaemia			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	1	2	0
Hypokalaemia			
subjects affected / exposed	2 / 42 (4.76%)	4 / 47 (8.51%)	1 / 3 (33.33%)
occurrences (all)	2	5	1
Hyponatraemia			
subjects affected / exposed	2 / 42 (4.76%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	3	1	0
Hypomagnesaemia			
subjects affected / exposed	3 / 42 (7.14%)	3 / 47 (6.38%)	1 / 3 (33.33%)
occurrences (all)	3	5	1
Hypophosphataemia			
subjects affected / exposed	2 / 42 (4.76%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	2	1	0
Hypophagia			
subjects affected / exposed	0 / 42 (0.00%)	0 / 47 (0.00%)	0 / 3 (0.00%)
occurrences (all)	0	0	0
Hypoproteinaemia			
subjects affected / exposed	1 / 42 (2.38%)	1 / 47 (2.13%)	0 / 3 (0.00%)
occurrences (all)	1	1	0

Non-serious adverse events	Phase II- Group 1		
Total subjects affected by non-serious adverse events			

subjects affected / exposed	51 / 52 (98.08%)		
Vascular disorders			
Hot flush			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Hypotension			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Pallor			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Thrombosis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
General disorders and administration site conditions			
Chest pain			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Asthenia			
subjects affected / exposed	6 / 52 (11.54%)		
occurrences (all)	12		
Chills			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences (all)	3		
Fatigue			
subjects affected / exposed	14 / 52 (26.92%)		
occurrences (all)	15		
Face oedema			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Gait disturbance			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Influenza like illness			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences (all)	4		

Non-cardiac chest pain subjects affected / exposed occurrences (all)	8 / 52 (15.38%) 9		
Oedema peripheral subjects affected / exposed occurrences (all)	27 / 52 (51.92%) 53		
Pain subjects affected / exposed occurrences (all)	3 / 52 (5.77%) 3		
Peripheral swelling subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Pyrexia subjects affected / exposed occurrences (all)	8 / 52 (15.38%) 12		
Immune system disorders Anaphylactoid reaction subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Hypersensitivity subjects affected / exposed occurrences (all)	2 / 52 (3.85%) 3		
Reproductive system and breast disorders Pelvic pain subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Perineal pain subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Vulvovaginal pruritus subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Prostatomegaly subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Respiratory, thoracic and mediastinal disorders			

Cough			
subjects affected / exposed	10 / 52 (19.23%)		
occurrences (all)	20		
Dry throat			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Dysphonia			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Dyspnoea			
subjects affected / exposed	11 / 52 (21.15%)		
occurrences (all)	14		
Dyspnoea exertional			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Epistaxis			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Haemoptysis			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Nasal discomfort			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Hypoxia			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Nasal dryness			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Oropharyngeal pain			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Pleural effusion			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		

Pneumothorax			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Productive cough			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Pulmonary embolism			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Rhinorrhoea			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	3		
Throat irritation			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Psychiatric disorders			
Anxiety			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Confusional state			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Sleep disorder			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Depression			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Insomnia			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Investigations			
Alanine aminotransferase increased			
subjects affected / exposed	7 / 52 (13.46%)		
occurrences (all)	7		
Amylase increased			

subjects affected / exposed	11 / 52 (21.15%)		
occurrences (all)	35		
Aspartate aminotransferase increased			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Blood albumin decreased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Blood alkaline phosphatase increased			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Blood bilirubin increased			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	2		
Blood creatine increased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Blood creatine phosphokinase increased			
subjects affected / exposed	5 / 52 (9.62%)		
occurrences (all)	12		
Blood creatinine increased			
subjects affected / exposed	10 / 52 (19.23%)		
occurrences (all)	14		
Blood glucose increased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Blood potassium increased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Blood triglycerides increased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Blood urea increased			

subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Blood uric acid increased			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
C-reactive protein increased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Electrocardiogram QT prolonged			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Gamma-glutamyltransferase increased			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Haemoglobin decreased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Lipase increased			
subjects affected / exposed	12 / 52 (23.08%)		
occurrences (all)	40		
Lymphocyte count decreased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Protein total decreased			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Weight decreased			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Weight increased			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences (all)	3		
White blood cell count decreased			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		

Injury, poisoning and procedural complications Procedural pain subjects affected / exposed occurrences (all) Traumatic haematoma subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0 0 / 52 (0.00%) 0		
Cardiac disorders Angina pectoris subjects affected / exposed occurrences (all) Bradycardia subjects affected / exposed occurrences (all)	1 / 52 (1.92%) 1 0 / 52 (0.00%) 0		
Nervous system disorders Amnesia subjects affected / exposed occurrences (all) Dizziness subjects affected / exposed occurrences (all) Dysaesthesia subjects affected / exposed occurrences (all) Dysgeusia subjects affected / exposed occurrences (all) Epilepsy subjects affected / exposed occurrences (all) Hypoaesthesia subjects affected / exposed occurrences (all) Lethargy subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0 9 / 52 (17.31%) 10 1 / 52 (1.92%) 2 0 / 52 (0.00%) 0 0 / 52 (0.00%) 0 1 / 52 (1.92%) 1 0 / 52 (0.00%) 0		

Headache			
subjects affected / exposed	7 / 52 (13.46%)		
occurrences (all)	8		
Neuropathy peripheral			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Seizure			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Post herpetic neuralgia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Syncope			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	9 / 52 (17.31%)		
occurrences (all)	10		
Febrile neutropenia			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Leukopenia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Lymphopenia			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences (all)	4		
Neutropenia			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Thrombocytopenia			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Ear and labyrinth disorders			

Deafness			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Ear pain			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Deafness transitory			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Hypoacusis			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Tinnitus			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Eye disorders			
Eye pruritus			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Eye swelling			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Eyelid oedema			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Vision blurred			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Vitreous floaters			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Gastrointestinal disorders			
Abdominal distension			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Abdominal discomfort			

subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Abdominal pain			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	6		
Abdominal pain upper			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Constipation			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	6		
Diarrhoea			
subjects affected / exposed	20 / 52 (38.46%)		
occurrences (all)	39		
Dry mouth			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Dyspepsia			
subjects affected / exposed	10 / 52 (19.23%)		
occurrences (all)	11		
Dysphagia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Gastrooesophageal reflux disease			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Lip blister			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Lip dry			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Mouth ulceration			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Nausea			

subjects affected / exposed	28 / 52 (53.85%)		
occurrences (all)	41		
Odynophagia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Toothache			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Stomatitis			
subjects affected / exposed	5 / 52 (9.62%)		
occurrences (all)	5		
Vomiting			
subjects affected / exposed	20 / 52 (38.46%)		
occurrences (all)	33		
Hepatobiliary disorders			
Hypertransaminasaemia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Skin and subcutaneous tissue disorders			
Acne			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Dermatitis acneiform			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	3		
Alopecia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Dry skin			
subjects affected / exposed	5 / 52 (9.62%)		
occurrences (all)	5		
Eczema			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Ingrowing nail			

subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Nail discolouration			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Night sweats			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Palmar-plantar erythrodysesthesia syndrome			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Pruritus			
subjects affected / exposed	6 / 52 (11.54%)		
occurrences (all)	8		
Rash			
subjects affected / exposed	7 / 52 (13.46%)		
occurrences (all)	8		
Rash macular			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Rash maculo-papular			
subjects affected / exposed	10 / 52 (19.23%)		
occurrences (all)	10		
Rash papular			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Rash pruritic			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Skin hyperpigmentation			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Urticaria			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		

Renal and urinary disorders			
Acute kidney injury			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Chronic kidney disease			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Hypertonic bladder			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Renal failure			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Endocrine disorders			
Hyperthyroidism			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences (all)	3		
Back pain			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Bone pain			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Flank pain			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Joint stiffness			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Muscular weakness			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		

Muscle spasms			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Musculoskeletal pain			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Musculoskeletal chest pain			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Myalgia			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	3		
Neck pain			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Pain in extremity			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	4		
Tendon pain			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Infections and infestations			
Bronchitis			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Conjunctivitis			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences (all)	3		
Cellulitis			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Cystitis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Folliculitis			

subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Gastroenteritis			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Herpes zoster			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Nasopharyngitis			
subjects affected / exposed	5 / 52 (9.62%)		
occurrences (all)	6		
Infection			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Paronychia			
subjects affected / exposed	3 / 52 (5.77%)		
occurrences (all)	7		
Pharyngitis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Pneumonia			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	2		
Respiratory tract infection			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Rhinitis			
subjects affected / exposed	2 / 52 (3.85%)		
occurrences (all)	5		
Sinobronchitis			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Skin infection			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		
Tooth infection			

subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Upper respiratory tract infection subjects affected / exposed occurrences (all)	9 / 52 (17.31%) 11		
Urinary tract infection subjects affected / exposed occurrences (all)	5 / 52 (9.62%) 7		
Viral infection subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Metabolism and nutrition disorders			
Decreased appetite subjects affected / exposed occurrences (all)	14 / 52 (26.92%) 16		
Hyperglycaemia subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Hyperkalaemia subjects affected / exposed occurrences (all)	3 / 52 (5.77%) 8		
Hyperlipasaemia subjects affected / exposed occurrences (all)	0 / 52 (0.00%) 0		
Hypoalbuminaemia subjects affected / exposed occurrences (all)	9 / 52 (17.31%) 17		
Hypocalcaemia subjects affected / exposed occurrences (all)	2 / 52 (3.85%) 2		
Hypoglycaemia subjects affected / exposed occurrences (all)	1 / 52 (1.92%) 1		
Hypokalaemia subjects affected / exposed occurrences (all)	5 / 52 (9.62%) 8		

Hyponatraemia			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	6		
Hypomagnesaemia			
subjects affected / exposed	4 / 52 (7.69%)		
occurrences (all)	7		
Hypophosphataemia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Hypophagia			
subjects affected / exposed	0 / 52 (0.00%)		
occurrences (all)	0		
Hypoproteinaemia			
subjects affected / exposed	1 / 52 (1.92%)		
occurrences (all)	1		

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
28 November 2014	<p>Reduced nazartinib starting dose to 50 mg qd.</p> <p>Clarified the subsequent escalation of nazartinib and capmatinib doses.</p> <p>Revised the DLT definitions by adding Grade 2 pneumonitis as dose-limiting toxicity.</p> <p>Provided the latest available clinical safety and PK data for nazartinib and capmatinib tablet formulation.</p> <p>Provided updated hypothetical scenarios in reflection of the reduced starting dose of nazartinib and the updated priors based on the updated safety information.</p>
24 July 2015	<p>Introduced a formulation change for nazartinib from capsules to tablets (viable formulation for commercialization).</p> <p>Recommended guidelines for management and dose modification of rash/skin toxicities were provided for the nazartinib treatment.</p> <p>For capmatinib treatment, updated the dose modifications guideline for hepatotoxicity in regard to discontinuing study medication(s) with concurrent elevation of ALT and/or AST > 3×ULN and total bilirubin > 2×ULN with ALP < 2×ULN, in the absence of signs of cholestasis, hemolysis, and alternative causes of the liver injury (e.g., concomitant use of hepatotoxic drug(s), alcoholic hepatitis, etc.). Specific work-up for potential Hy's law cases had been added to the protocol.</p> <p>The dose modification rules and the dose limiting toxicity for liver toxicity as well as the follow-up evaluations for hepatic toxicities had also been updated accordingly.</p> <p>Included precautionary measures against ultraviolet exposure as capmatinib had photosensitization potential.</p> <p>Additionally, updated the exclusion criteria and few new exclusion criteria were added.</p> <p>Updated concomitant therapies, criteria for interruption and re-initiation of nazartinib and capmatinib.</p>
01 September 2015	<p>Provided detailed information about an urgent safety measure related to nazartinib.</p> <p>Excluded subjects with other malignancies, who underwent a bone marrow or solid organ transplant, with a known history of human immunodeficiency virus infection, or receiving immunosuppressive agents or chronic corticosteroids at study entry.</p> <p>Allowed subjects with either HBV surface antigen positive or HBV-DNA positive to enroll into the study if willing to take antiviral therapy 1-2 weeks prior to first dose of study treatment and continue antiviral therapy for at least 4 weeks after the last dose of study treatment.</p> <p>Allowed new subjects with negative hepatitis C antibody (HC Ab) or who were HC Ab positive but with an undetectable level of HCV-RNA into the study. Subjects with detectable HCV-RNA were not eligible to enroll into the study.</p>

02 November 2015	<p>Implemented a centralization of analysis for the resistance mutations EGFR T790M and MET oncogene amplification at a Novartis designated central laboratory for all subjects entering into Phase II.</p> <p>Introduced 50 mg dose strength of nazartinib tablets for better compliance.</p> <p>Optimized the subject safety and the toxicity monitoring as well as to align Novartis study protocols, the QTcF eligibility criterion and the management guidelines for pancreatic toxicity were revised.</p> <p>Updated the definition of the end of study to detail study continuation conditions after completion of the primary analysis until all subjects discontinued, or until another clinical trial was available for all ongoing subjects to be transferred to that clinical study and continue to receive capmatinib + nazartinib.</p> <p>Clarified the possibility of using bisphosphonates during the study.</p> <p>Detailed the use of an Interactive Response Technology system in the Phase II part of the study to specify the enrollment and the study drug dispensation tracking via the application.</p> <p>Modalities and conditions for subject re-screening after screen failure was introduced, together with screening assessments to be performed again during the new screening phase.</p>
29 July 2016	<p>Better defined the subject population to be enrolled in the Phase II part of the study (expansion phase). A new group was added (Group 3), and the sample size of Group 2 was revised.</p> <p>Implemented a requirement of ECOG performance status of 0-1 for all subjects to be enrolled in the study to ensure that subjects would be able to receive potential benefit from study treatment.</p> <p>Removed the exclusion criterion on fasting plasma glucose ≥ 160 mg/dL (≥ 8.9 mmol/L) in the absence of evidence of impact on the glycemia in light of the most recent available clinical data and according to the latest versions of capmatinib and nazartinib Investigator Brochures</p> <p>Introduced a new dose strength (150 mg) for capmatinib tablets.</p> <p>Introduced other rare EGFR activating sensitizing mutations that were also considered for enrollment onto the Phase II. These rare mutations, L861Q, G719X, and S768I, account for almost 10% of the cases whereas L858R point mutation and exon 19 deletions accounted for approximately 90% of the cases.</p> <p>Updated the overview of capmatinib and nazartinib according to last Investigator Brochure versions.</p> <p>Removed the restriction of proton pump inhibitors use during the course of the study.</p> <p>Clarified the conditions in which local laboratory assessments should be reported in the Case Report Form during the Phase 2 part of the trial.</p> <p>Clarified the time window for post-dose ECG during the visits with PK sampling.</p> <p>Clarified the definition of acquired resistance mechanisms and the definition of the 4 subgroups of subjects being enrolled in the Phase II Group 1.</p> <p>Clarified the biosample requirements for the optional biomarker assessments.</p> <p>Introduced the RP2D and the dose reduction steps starting from the RP2D.</p>
11 April 2017	<p>Added in Phase II a new group 4 of subjects (N=30) with EGFRmut, any T790M, any MET, in order to assess the safety and tolerability of the combination therapy when taken with food (unrestricted meal type).</p> <p>Proposed a new group 4 to administer study medication with food to improve the tolerability profile of the combination therapy, as dosing with food had been shown to improve the GI tolerability of some multi-kinase inhibitors such as imatinib (Gleevec®) and bosutinib (Bosulif®).</p> <p>Corrected the exclusion criterion for subjects with asymptomatic serum amylase and lipase > Grade 2, to clarify that subject now met exclusion criterion even if only one of the two parameters (asymptomatic serum amylase or lipase) was > Grade 2.</p> <p>Added the collection of an optional on-treatment biopsy with paired cfDNA sample to identify emergence of potential resistance markers during treatment as well as understand the correlation between tissue and plasma biomarker status.</p> <p>Clarified the exclusion criterion for subjects with brain metastases.</p> <p>Clarified the interpretation of study discontinuation after dose interruption.</p>

20 October 2021	<p>A new group 5 added per protocol amendment 7 will not be opened as Novartis made the decision to discontinue the study on 11-May-2022. Below are the key changes in amendment 7.</p> <p>As of 20-Oct-2021, a total of 177 participants have been enrolled in the study: 33 participants in Phase Ib and 144 in Phase II (Group 1 to Group 4). All participants from Phase Ib and Phase II Groups 1 to 4 had met one of the end of study completion criteria as considered under protocol amendment 6 and there were no participants receiving treatment.</p> <p>The main purpose of this amendment is to implement a new subject group, Phase II Group 5, to determine the efficacy and safety of capmatinib monotherapy. The rationale for this addition is based on the following: Based on evidence suggesting that MET gene amplification plays an important role as a mechanism of resistance to treatment with EGFR-TKIs in NSCLC patients carrying mutated EGFR. Furthermore, the fact that there are reports showing significant clinical antitumor activity in this subpopulation of patients when combining an EGFR-TKI with a specific cMET inhibitor (such as Capmatinib). Therefore, the contribution of capmatinib as a monodrug needs to be established. This information will pave the way for the optimal use of the combination in future trials.</p>
-----------------	--

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

Due to EudraCT system limitations, which EMA is aware of, data using 999999 as data points in this record are not an accurate representation of the clinical trial results. Please use <https://www.novctrd.com> for complete trial results.

Notes: