

Table 8: Patient treatment questionnaire

	A n=7	B n=7	OVERALL n=14
How did you find the application of the ointment?			
Easy			
Neither easy or difficult	6	6	12
Difficult	1	1	2
How long did it take to apply study treatment?	0	0	0
Less than 1 minute			
More than 1 minute but less than 5 minutes	0	2	2
More than 5 minutes but less than 10 minutes	4	3	7
More than 10 minutes	3	2	5
Overall, how satisfied were you with the study medication?	0	0	0
Very satisfied			
Satisfied	4	3	7
Neither satisfied or dissatisfied	1	2	3
Dissatisfied	0	1	1
Very dissatisfied	2	1	3
	0	0	0
If this ointment was available as a treatment would you use it?			
Yes	5	5	10
No	2	2	4

Table 8 indicates the acceptability of trial treatment at End Of Treatment (EOT) visit.

Table 9: Treatment compliance in cohort 2

Patients	A Left Active n=7	B Right Active n=7	Total n=14
Days on treatment			
Median	84	81	83
IQR, Range	(81,86) (81,86)	(80,85) (80,86)	(81,85) (80,86)
Patients missed treatment	n=5	n=5	n=10
Days of missed treatment			
Median	3	1	2
IQR, Range	(1,6) (1,8)	(1,9) (1,9)	(1,8) (1,9)
% protocol treatment received	n=7	n=7	n=14
Median			
IQR, Range	98.8 (92.9,100) (90.7,100)	98.8 (88.9,100) (88.8,100)	98.8 (92.9,100) (88.8,100)

14 participants completed the trial. 1 Participant withdrew consent during the trial. The table above provides information relating to median and IQR range participants complied and missed treatment per the trial protocol.

There were a total of 40 days of missed applications across 10 patients (4 patients were fully compliant). In 36 patient episodes of missed applications, all 10 tumours had been missed, in 1 patient episode the ointment had been applied to the wrong side, in 1 patient episode the reason was given as data had not been collected and in 2 patient episodes no reason was given.