



## Clinical trial results:

### Albiglutide Versus Placebo as Add-on to Intensified Basal-Bolus Insulin Therapy in Subjects With Type 2 Diabetes Mellitus

#### Summary

EudraCT number	2014-001969-27
Trial protocol	HU DE GB
Global end of trial date	09 June 2017

#### Results information

Result version number	v1 (current)
This version publication date	30 December 2018
First version publication date	30 December 2018
Summary attachment (see zip file)	Cancelled before Active Statement (Cancelled before Active Statement GLP111892.doc)

#### Trial information

##### Trial identification

Sponsor protocol code	GLP111892
-----------------------	-----------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	-
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	GlaxoSmithKline
Sponsor organisation address	980 Great West Road, Brentford, United Kingdom, Middlesex
Public contact	GSK Response Center, GlaxoSmithKline, 1 866-435-7343,
Scientific contact	GSK Response Center, GlaxoSmithKline, 1 866-435-7343,

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	09 June 2017
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	09 June 2017
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

To evaluate the efficacy of once-weekly albiglutide in providing similar (or better) glycemic control with less hypoglycemia when added to a regimen of intensified basal-bolus insulin therapy compared with intensified basal-bolus insulin therapy alone in subjects with T2DM.

Protection of trial subjects:

NA

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	09 June 2017
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Poland: 99999
Worldwide total number of subjects	99999
EEA total number of subjects	99999

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	99999
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

99999 is "Not applicable" value or 0 participants, this trial was discontinued with no participants enrolled in the trial

### Pre-assignment

Screening details:

NA

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Not applicable
Blinding used	Not blinded

### Arms

Arm title	Albiglutide
-----------	-------------

Arm description:

Subjects would have received Albiglutide 30 mg (with forced uptitration to albiglutide 50 mg at Week 4).

Arm type	Experimental
Investigational medicinal product name	Albiglutide
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Subjects were planned to receive once-weekly subcutaneous injections of albiglutide 30 mg (with forced uptitration to albiglutide 50 mg at Week 4) in addition to intensification of background basal-bolus insulin therapy (with or without metformin) according to predefined titration algorithms.

Number of subjects in period 1	Albiglutide
Started	99999
Completed	99999

## Baseline characteristics

### Reporting groups

Reporting group title	Overall Study
-----------------------	---------------

Reporting group description: -

Reporting group values	Overall Study	Total	
Number of subjects	99999	99999	
Age categorical			
Units: Subjects			
In utero		0	
Preterm newborn infants (gestational age < 37 wks)		0	
Newborns (0-27 days)		0	
Infants and toddlers (28 days-23 months)		0	
Children (2-11 years)		0	
Adolescents (12-17 years)		0	
Adults (18-64 years)		0	
From 65-84 years		0	
85 years and over		0	
Age continuous			
Units: years			
arithmetic mean	0		
standard deviation	± 0	-	
Gender categorical			
Units: Subjects			
Female	99999	99999	
Male	0	0	

## End points

### End points reporting groups

Reporting group title	Albiglutide
Reporting group description:	
Subjects would have received Albiglutide 30 mg (with forced uptitration to albiglutide 50 mg at Week 4).	

### Primary: 1.Percentage of subjects with severe or documented symptomatic hypoglycemia through Week 26

End point title	1.Percentage of subjects with severe or documented symptomatic hypoglycemia through Week 26 <sup>[1]</sup>
-----------------	--

End point description:

99999 is "Not applicable" value or 0 participants, this trial was discontinued with no participants enrolled in the trial.

End point type	Primary
----------------	---------

End point timeframe:

99999

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: No subjects were enrolled in the trial hence results are not available

End point values	Albiglutide			
Subject group type	Reporting group			
Number of subjects analysed	99999 <sup>[2]</sup>			
Units: Not available	99999			

Notes:

[2] - 99999 is "Not applicable" value or 0 participants, this trial was discontinued with no participants

### Statistical analyses

No statistical analyses for this end point

## Adverse events

---

### Adverse events information<sup>[1]</sup>

---

Timeframe for reporting adverse events:

All adverse events occurring during the course of the clinical trial were to be collected.

Assessment type	Systematic
-----------------	------------

---

### Dictionary used

---

Dictionary name	MedDRA
-----------------	--------

---

Dictionary version	0
--------------------	---

---

Frequency threshold for reporting non-serious adverse events: 5 %

---

### Notes:

[1] - There are no non-serious adverse events recorded for these results. It is expected that there will be at least one non-serious adverse event reported.

Justification: No subjects were enrolled in the trial hence results are not available

## **More information**

### **Substantial protocol amendments (globally)**

Were there any global substantial amendments to the protocol? No

---

### **Interruptions (globally)**

Were there any global interruptions to the trial? No

### **Limitations and caveats**

None reported