



## Clinical trial results:

### A Phase 2A Study Investigating the Safety, Pharmacokinetics, Immunogenicity, and Exploratory Efficacy of Dupilumab in Patients Aged 6 to <18 Years With Atopic Dermatitis

#### Summary

EudraCT number	2014-003263-37
Trial protocol	HU DE CZ PL GB Outside EU/EEA
Global end of trial date	14 March 2016

#### Results information

Result version number	v1
This version publication date	30 March 2017
First version publication date	30 March 2017

#### Trial information

##### Trial identification

Sponsor protocol code	R668-AD-1412
-----------------------	--------------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02407756
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	Regeneron Pharmaceuticals, Inc.
Sponsor organisation address	777 Old Saw Mill River Rd., Tarrytown, United States, 10591
Public contact	Clinical Trial Management, Regeneron Pharmaceuticals, Inc., clinicaltrials@regeneron.com
Scientific contact	Clinical Trial Management, Regeneron Pharmaceuticals, Inc., clinicaltrials@regeneron.com

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	Yes
EMA paediatric investigation plan number(s)	EMA-001501-PIP01-13
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	12 April 2016
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	14 March 2016
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

To characterize the safety and pharmacokinetics (PK) of Dupilumab in pediatric subjects with moderate-to-severe atopic dermatitis (AD) (for adolescents  $\geq 12$  to  $< 18$  years of age) or severe AD (for children  $\geq 6$  to  $< 12$  years of age).

Protection of trial subjects:

This study was conducted in accordance with the ethical principles that have their origin in the Declaration of Helsinki and that are consistent with the International Conference on Harmonisation (ICH) guidelines for Good Clinical Practice (GCP) and applicable regulatory requirements.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	17 March 2015
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Poland: 25
Country: Number of subjects enrolled	United Kingdom: 10
Country: Number of subjects enrolled	Czech Republic: 3
Country: Number of subjects enrolled	Germany: 18
Country: Number of subjects enrolled	Hungary: 10
Country: Number of subjects enrolled	Canada: 12
Worldwide total number of subjects	78
EEA total number of subjects	66

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age $< 37$ wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	38

Adolescents (12-17 years)	40
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

The study was conducted at 25 sites in 6 countries. A total of 88 subjects were screened between 17 Mar 2015 & 25 Sep 2015 of whom 78 subjects were randomized and 77 were treated. Ten subjects were screen failures as 5 were due to exclusion criteria met & inclusion criteria not met, 4 subjects withdrew consent and 1 subject was lost to follow-up.

### Pre-assignment

Screening details:

A total of 40 adolescents (aged  $\geq 12$  to  $< 18$  years) and 38 children (aged  $\geq 6$  to  $< 12$  years) were enrolled and randomized to 2 sequential ascending dose cohorts: Cohort 1 (2 mg/kg) and Cohort 2 (4 mg/kg). Dosing started with cohort 1. Proceeding to the next cohort occurred once all initial 8 patients enrolled had been observed for at least 2 weeks.

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Not applicable
Blinding used	Not blinded

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Dupilumab 2mg/kg: Adolescents

Arm description:

Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between  $\geq 12$  to  $< 18$  years.

Arm type	Experimental
Investigational medicinal product name	Dupilumab
Investigational medicinal product code	REGN668, SAR231893
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Subcutaneous injection in the different quadrants of the abdomen (avoiding navel and waist areas), upper thighs and upper arms.

<b>Arm title</b>	Dupilumab 2mg/kg: Children
------------------	----------------------------

Arm description:

Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between  $\geq 6$  to  $< 12$  years.

Arm type	Experimental
Investigational medicinal product name	Dupilumab
Investigational medicinal product code	REGN668, SAR231893
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Subcutaneous injection in the different quadrants of the abdomen (avoiding navel and waist areas), upper thighs and upper arms.

<b>Arm title</b>	Dupilumab 4mg/kg: Adolescents
------------------	-------------------------------

Arm description:

Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between  $\geq 12$  to  $< 18$  years.

Arm type	Experimental
----------	--------------

Investigational medicinal product name	Dupilumab
Investigational medicinal product code	REGN668, SAR231893
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Subcutaneous injection in the different quadrants of the abdomen (avoiding navel and waist areas), upper thighs and upper arms.

<b>Arm title</b>	Dupilumab 4mg/kg: Children
------------------	----------------------------

Arm description:

Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between  $\geq 6$  to  $<12$  years.

Arm type	Experimental
Investigational medicinal product name	Dupilumab
Investigational medicinal product code	REGN668, SAR231893
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Subcutaneous injection in the different quadrants of the abdomen (avoiding navel and waist areas), upper thighs and upper arms.

<b>Number of subjects in period 1<sup>[1]</sup></b>	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents
Started	20	18	20
Treated	20	18	20
Completed	20	18	20

<b>Number of subjects in period 1<sup>[1]</sup></b>	Dupilumab 4mg/kg: Children
Started	19
Treated	19
Completed	19

Notes:

[1] - The number of subjects reported to be in the baseline period are not the same as the worldwide number enrolled in the trial. It is expected that these numbers will be the same.

Justification: One subject in the 4 mg/kg (Cohort 2b) arm did not receive study drug and was withdrawn from the study on study day 1 during Part A period due to withdrawal of consent (fear of study drug injection) and was not included in the analysis. Analysis was performed on safety analysis set (SAF) that included all subjects who received any study drug.

## Baseline characteristics

### Reporting groups

Reporting group title	Dupilumab 2mg/kg: Adolescents
Reporting group description: Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 12$ to $< 18$ years.	
Reporting group title	Dupilumab 2mg/kg: Children
Reporting group description: Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 6$ to $< 12$ years.	
Reporting group title	Dupilumab 4mg/kg: Adolescents
Reporting group description: Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 12$ to $< 18$ years.	
Reporting group title	Dupilumab 4mg/kg: Children
Reporting group description: Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 6$ to $< 12$ years.	

Reporting group values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents
Number of subjects	20	18	20
Age categorical			
Units: Subjects			

Age continuous			
Units: years			
arithmetic mean	14.7	8.2	14.3
standard deviation	$\pm 2.01$	$\pm 1.62$	$\pm 1.66$
Gender categorical			
Units: Subjects			
Female	11	9	11
Male	9	9	9
Ethnicity			
Units: Subjects			
Hispanic or Latino	0	0	0
Not Hispanic or Latino	20	18	20
Race			
Units: Subjects			
White	17	17	15
Black or African American	0	0	1
Asian	2	0	3
Other	1	1	1
Number of Subjects with Investigator Global Assessment (IGA) score of 3 or 4			
IGA is an assessment scale used to determine severity of AD and clinical response to treatment on a 5-point scale (0 = clear; 1 = almost clear; 2 = mild; 3 = moderate; 4 = severe) based on erythema and papulation/ infiltration. Therapeutic response is an IGA score of 0 (clear) or 1 (almost clear). Analysis was performed on SAF.			
Units: Subjects			
IGA score of 3	8	1	11

IGA score of 4	12	17	9
----------------	----	----	---

Eczema Area and Severity Index (EASI) score			
The EASI score was used to measure the severity and extent of atopic dermatitis (AD) and measures erythema, infiltration, excoriation and lichenification on 4 anatomic regions of the body: head, trunk, upper and lower extremities. The total EASI score ranges from 0 to 72 points, with the higher scores reflecting the worse severity of AD. Analysis was performed on Safety Analysis set (SAF).			
Units: Score on a scale			
arithmetic mean	34.8	32.9	28.6
standard deviation	± 17	± 15.53	± 14.7
Pruritus Numerical Rating Scale (NRS)			
Pruritus NRS scale is an assessment tool that is used to report the intensity of subject's pruritus (itch), both maximum and average intensity, during a 24-hour recall period. Subjects were asked the following question: how would a subject rate his itch at the worst moment during the previous 24 hours (for maximum itch intensity on a scale of 0 – 10 [0 = no itch; 10 = worst itch imaginable]). Weekly average obtained in the 7-day period prior to the baseline visit. Analysis was performed on SAF.			
Units: Score on a scale			
arithmetic mean	6.1	6.4	6.9
standard deviation	± 2.47	± 2.23	± 2.21
Body Surface Area (BSA) Involvement with Atopic Dermatitis (AD)			
BSA affected by AD was assessed for each section of the body (the possible highest score for each region was: head and neck [9%], anterior trunk [18%], back [18%], upper limbs [18%], lower limbs [36%], and genitals [1%]). It was reported as a percentage of all major body sections combined. Analysis was performed on SAF.			
Units: Percentage of body surface area			
arithmetic mean	52.2	59	45.9
standard deviation	± 24.78	± 22.49	± 25.34
SCORing Atopic Dermatitis (SCORAD) Score			
SCORAD is a clinical tool for assessing the severity of atopic dermatitis developed by the European Task Force on Atopic Dermatitis ("Severity scoring of atopic dermatitis: the SCORAD index. Consensus Report of the European Task Force on Atopic Dermatitis". Dermatology (Basel) 186 (1): 23–31. 1993). Extent and intensity of eczema as well as subjective signs (insomnia, etc.) are assessed and scored. Total score ranges from 0 [absent disease] to 103 [severe disease]). Analysis was performed on SAF.			
Units: Score on a scale			
arithmetic mean	68	66.4	63
standard deviation	± 13.19	± 13.06	± 14.43

<b>Reporting group values</b>	Dupilumab 4mg/kg: Children	Total	
Number of subjects	19	77	
Age categorical			
Units: Subjects			

Age continuous			
Units: years			
arithmetic mean	8.2		
standard deviation	± 1.99	-	
Gender categorical			
Units: Subjects			
Female	8	39	
Male	11	38	

Ethnicity			
Units: Subjects			
Hispanic or Latino	1	1	
Not Hispanic or Latino	18	76	
Race			
Units: Subjects			
White	18	67	
Black or African American	1	2	
Asian	0	5	
Other	0	3	
Number of Subjects with Investigator Global Assessment (IGA) score of 3 or 4			
IGA is an assessment scale used to determine severity of AD and clinical response to treatment on a 5-point scale (0 = clear; 1 = almost clear; 2 = mild; 3 = moderate; 4 = severe) based on erythema and papulation/ infiltration. Therapeutic response is an IGA score of 0 (clear) or 1 (almost clear). Analysis was performed on SAF.			
Units: Subjects			
IGA score of 3	0	20	
IGA score of 4	19	57	
Eczema Area and Severity Index (EASI) score			
The EASI score was used to measure the severity and extent of atopic dermatitis (AD) and measures erythema, infiltration, excoriation and lichenification on 4 anatomic regions of the body: head, trunk, upper and lower extremities. The total EASI score ranges from 0 to 72 points, with the higher scores reflecting the worse severity of AD. Analysis was performed on Safety Analysis set (SAF).			
Units: Score on a scale			
arithmetic mean	38.8		
standard deviation	± 18.64	-	
Pruritus Numerical Rating Scale (NRS)			
Pruritus NRS scale is an assessment tool that is used to report the intensity of subject's pruritus (itch), both maximum and average intensity, during a 24-hour recall period. Subjects were asked the following question: how would a subject rate his itch at the worst moment during the previous 24 hours (for maximum itch intensity on a scale of 0 – 10 [0 = no itch; 10 = worst itch imaginable]). Weekly average obtained in the 7-day period prior to the baseline visit. Analysis was performed on SAF.			
Units: Score on a scale			
arithmetic mean	6.7		
standard deviation	± 2.35	-	
Body Surface Area (BSA) Involvement with Atopic Dermatitis (AD)			
BSA affected by AD was assessed for each section of the body (the possible highest score for each region was: head and neck [9%], anterior trunk [18%], back [18%], upper limbs [18%], lower limbs [36%], and genitals [1%]). It was reported as a percentage of all major body sections combined. Analysis was performed on SAF.			
Units: Percentage of body surface area			
arithmetic mean	62.3		
standard deviation	± 30.34	-	
SCORing Atopic Dermatitis (SCORAD) Score			
SCORAD is a clinical tool for assessing the severity of atopic dermatitis developed by the European Task Force on Atopic Dermatitis ("Severity scoring of atopic dermatitis: the SCORAD index. Consensus Report of the European Task Force on Atopic Dermatitis". Dermatology (Basel) 186 (1): 23–31. 1993). Extent and intensity of eczema as well as subjective signs (insomnia, etc.) are assessed and scored. Total score ranges from 0 [absent disease] to 103 [severe disease]). Analysis was performed on SAF.			
Units: Score on a scale			
arithmetic mean	72.7		
standard deviation	± 12.96	-	



## End points

### End points reporting groups

Reporting group title	Dupilumab 2mg/kg: Adolescents
Reporting group description: Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 12$ to $< 18$ years.	
Reporting group title	Dupilumab 2mg/kg: Children
Reporting group description: Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 6$ to $< 12$ years.	
Reporting group title	Dupilumab 4mg/kg: Adolescents
Reporting group description: Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 12$ to $< 18$ years.	
Reporting group title	Dupilumab 4mg/kg: Children
Reporting group description: Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period then 4 repeated doses weekly in subjects aged between $\geq 6$ to $< 12$ years.	

### Primary: Pharmacokinetics (PK) of dupilumab: Maximum plasma concentration observed (C<sub>max</sub>) after single administration

End point title	Pharmacokinetics (PK) of dupilumab: Maximum plasma concentration observed (C <sub>max</sub> ) after single administration <sup>[1]</sup>
End point description: Peak dupilumab concentration in serum following single dose administration. Analysis was performed on PK analysis set that included all treated subjects who received the study medication and had at least 1 quantified (non-missing) result for dupilumab concentration following the first dose of the study drug.	
End point type	Primary
End point timeframe: Day 2, 4, 8, 15, 22, 29, 36, 43, and 50	

#### Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Statistical analysis is descriptive only. No formal statistical comparison was performed.

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	20	18	20	19
Units: mg/L				
arithmetic mean (standard deviation)	9.91 ( $\pm$ 2.15)	14.3 ( $\pm$ 5.9)	23.1 ( $\pm$ 8.71)	32.4 ( $\pm$ 7.04)

### Statistical analyses

No statistical analyses for this end point

### Primary: PK of dupilumab: Area under the plasma concentration versus time curve

**(AUClast) after single administration**

End point title	PK of dupilumab: Area under the plasma concentration versus time curve (AUClast) after single administration <sup>[2]</sup>
-----------------	---

End point description:

Mean AUC estimates were calculated using mean concentration data at each time point, using a non-compartmental approach (NCA). Calculated AUClast (computed from time zero to the time of the last positive concentration) are presented. Analysis was performed on PK analysis set that included all treated subjects who received the study medication and had at least 1 quantified (non-missing) result for dupilumab concentration following the first dose of the study drug.

End point type	Primary
----------------	---------

End point timeframe:

Day 2, 4, 8, 15, 22, 29, 36, 43, and 50

Notes:

[2] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Statistical analysis is descriptive only. No formal statistical comparison was performed.

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	20	18	20	19
Units: Day*mg/L				
arithmetic mean (standard deviation)	104 (± 0)	160 (± 0)	362 (± 0)	330 (± 0)

**Statistical analyses**

No statistical analyses for this end point

**Primary: PK of dupilumab: Trough dupilumab concentration in serum (Ctrough) before 3rd and 4th repeated dose**

End point title	PK of dupilumab: Trough dupilumab concentration in serum (Ctrough) before 3rd and 4th repeated dose <sup>[3]</sup>
-----------------	--

End point description:

Analysis was performed on PK analysis set that included all treated subjects who received the study medication and had at least 1 quantified (non-missing) result for dupilumab concentration following the first dose of study drug.

End point type	Primary
----------------	---------

End point timeframe:

Pre-dose on Day 71 and Day 85

Notes:

[3] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Statistical analysis is descriptive only. No formal statistical comparison was performed.

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	20	18	20	19
Units: mg/L				
arithmetic mean (standard deviation)				
Day 71	10.4 (± 7.16)	17.2 (± 8.44)	32.8 (± 18.9)	42.1 (± 19.4)

Day 85	18.5 ( $\pm$ 12.4)	28 ( $\pm$ 12.9)	58.5 ( $\pm$ 24.4)	60.3 ( $\pm$ 36.3)
--------	--------------------	------------------	--------------------	--------------------

## Statistical analyses

No statistical analyses for this end point

## Secondary: Percent reduction from baseline in Eczema Area and Severity Index (EASI) at Week 12

End point title	Percent reduction from baseline in Eczema Area and Severity Index (EASI) at Week 12
-----------------	---

End point description:

The EASI score was used to measure the severity and extent of atopic dermatitis (AD) and measures erythema, infiltration, excoriation and lichenification on 4 anatomic regions of the body: head, trunk, upper and lower extremities. The total EASI score ranges from 0 to 72 points, with the higher scores reflecting the worse severity of AD. Analysis was performed on safety analysis set (SAF) that included all subjects who received any study drug. Data after rescue treatment use during the Part B period were set to missing, then missing values were imputed by last observation carried forward (LOCF).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline to Week 12 (one week after last dose)

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	20	18	20	19
Units: percent change				
arithmetic mean (standard deviation)	-66.4 ( $\pm$ 29.25)	-76.2 ( $\pm$ 25.48)	-69.7 ( $\pm$ 24.48)	-63.4 ( $\pm$ 25.37)

## Statistical analyses

No statistical analyses for this end point

## Secondary: Percent reduction from baseline in SCORing Atopic Dermatitis (SCORAD) Score at Week 12

End point title	Percent reduction from baseline in SCORing Atopic Dermatitis (SCORAD) Score at Week 12
-----------------	--

End point description:

SCORAD is a clinical tool for assessing the severity of atopic dermatitis developed by the European Task Force on Atopic Dermatitis ("Severity scoring of atopic dermatitis: the SCORAD index. Consensus Report of the European Task Force on Atopic Dermatitis". Dermatology (Basel) 186 (1): 23–31. 1993). Extent and intensity of eczema as well as subjective signs (insomnia, etc.) are assessed and scored. Total score ranges from 0 [absent disease] to 103 [severe disease]). Analysis was performed on SAF. Data after rescue treatment use during the Part B period were set to missing, then missing values were imputed by LOCF.

End point type	Secondary
End point timeframe:	
Baseline to Week 12 (one week after last dose)	

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	20	18	20	19
Units: percent change				
arithmetic mean (standard deviation)	-47.7 (± 27.27)	-57.5 (± 23.1)	-43.4 (± 25.38)	-46.9 (± 24.31)

### Statistical analyses

No statistical analyses for this end point

### Secondary: Percent reduction from baseline in Pruritus Numerical Rating Scale (NRS) at Week 12

End point title	Percent reduction from baseline in Pruritus Numerical Rating Scale (NRS) at Week 12
-----------------	---

End point description:

Pruritus NRS scale is an assessment tool that is used to report the intensity of subject's pruritus (itch), both maximum and average intensity, during a 24-hour recall period. Subjects were asked the following question: how would a subject rate his itch at the worst moment during the previous 24 hours (for maximum itch intensity on a scale of 0 – 10 [0 = no itch; 10 = worst itch imaginable]). Analysis was performed on SAF. Data after rescue treatment use during the Part B period were set to missing, then missing values were imputed by LOCF.

End point type	Secondary
End point timeframe:	
Baseline to Week 12 (one week after last dose)	

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	20	18	20	19
Units: Percent change				
arithmetic mean (standard deviation)	-30.8 (± 68.35)	-41.6 (± 35.32)	-37.6 (± 34.42)	-39.6 (± 40.88)

### Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of subjects with Investigator Global Assessment (IGA) score of "0" or "1" (clear or almost clear) at Week 12

End point title	Percentage of subjects with Investigator Global Assessment (IGA) score of "0" or "1" (clear or almost clear) at Week 12
-----------------	---

End point description:

IGA is an assessment scale used to determine severity of AD and clinical response to treatment on a 5-point scale (0 = clear; 1 = almost clear; 2 = mild; 3 = moderate; 4 = severe) based on erythema and papulation/infiltration. Therapeutic response is an IGA score of 0 (clear) or 1 (almost clear). Analysis was performed on SAF. Subjects with rescue treatment usage during the Part B period were specified as non-responders from the time the rescue was used.

End point type	Secondary
----------------	-----------

End point timeframe:

Week 12

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	20	18	20	19
Units: Percentage of subjects				
number (not applicable)	10	16.7	35	21.1

### Statistical analyses

No statistical analyses for this end point

### Secondary: Percent reduction from baseline in Body Surface Area (BSA) at week 12

End point title	Percent reduction from baseline in Body Surface Area (BSA) at week 12
-----------------	---

End point description:

Body surface area affected by AD was assessed for each section of the body (the possible highest score for each region was: head and neck [9%], anterior trunk [18%], back [18%], upper limbs [18%], lower limbs [36%], and genitals [1%]). It was reported as a percentage of all major body sections combined. Analysis was performed on SAF.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline to Week 12

End point values	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4mg/kg: Adolescents	Dupilumab 4mg/kg: Children
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	19	16	19	19
Units: Percentage of body surface area				
arithmetic mean (standard deviation)	-61 (± 31.08)	-70 (± 31.93)	-60.4 (±	-50 (± 30.8)

**Statistical analyses**

---

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

All Adverse Events (AEs) were collected from signature of the informed consent form up to the final visit (Week 20) regardless of seriousness or relationship to investigational product.

Adverse event reporting additional description:

Treatment Emergent Adverse Events (TEAEs) that developed/ worsened during the treatment and follow-up period (time period from the administration of first dose of study drug to End of Study (EOS) visit [8 weeks after the last dose]). Analysis was performed on SAF.

Assessment type	Non-systematic
-----------------	----------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	18.0
--------------------	------

### Reporting groups

Reporting group title	Dupilumab 2mg/kg: Adolescents
-----------------------	-------------------------------

Reporting group description:

Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period, then 4 repeated doses weekly in subjects aged between  $\geq 12$  to  $< 18$  years.

Reporting group title	Dupilumab 2mg/kg: Children
-----------------------	----------------------------

Reporting group description:

Dupilumab 2 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period, then 4 repeated doses weekly in subjects aged between  $\geq 6$  to  $< 12$  years.

Reporting group title	Dupilumab 4 mg/kg: Adolescents
-----------------------	--------------------------------

Reporting group description:

Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period, then 4 repeated doses weekly in subjects aged between  $\geq 12$  to  $< 18$  years.

Reporting group title	Dupilumab 4 mg/kg: Children
-----------------------	-----------------------------

Reporting group description:

Dupilumab 4 mg/kg as a single dose on Day 1 followed by 8-week PK sampling period), then 4 repeated doses weekly in subjects aged between  $\geq 6$  to  $< 12$  years.

Serious adverse events	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4 mg/kg: Adolescents
Total subjects affected by serious adverse events			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			
Cardiac disorders			
Palpitations			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			

Dermatitis atopic			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Staphylococcal skin infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Arthritis bacterial			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dermatitis infected			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

<b>Serious adverse events</b>	Dupilumab 4 mg/kg: Children		
Total subjects affected by serious adverse events			
subjects affected / exposed	2 / 19 (10.53%)		
number of deaths (all causes)	0		
number of deaths resulting from adverse events			
Cardiac disorders			
Palpitations			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Skin and subcutaneous tissue disorders			



Dermatitis atopic alternative assessment type: Systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	  1 / 19 (5.26%) 1 / 1 0 / 0		
Infections and infestations Staphylococcal skin infection alternative assessment type: Systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	  0 / 19 (0.00%) 0 / 0 0 / 0		
Arthritis bacterial alternative assessment type: Systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	  1 / 19 (5.26%) 1 / 1 0 / 0		
Dermatitis infected alternative assessment type: Systematic subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	  1 / 19 (5.26%) 1 / 1 0 / 0		

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	Dupilumab 2mg/kg: Adolescents	Dupilumab 2mg/kg: Children	Dupilumab 4 mg/kg: Adolescents
Total subjects affected by non-serious adverse events			
subjects affected / exposed	11 / 20 (55.00%)	11 / 18 (61.11%)	16 / 20 (80.00%)
General disorders and administration site conditions			
Fatigue			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	2
Chills			
alternative assessment type: Systematic			

subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	1	0	0
Chest pain			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	1	0	0
Pyrexia			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 20 (10.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	4	0	0
Injection site urticaria			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Injection site swelling			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Injection site irritation			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	2	0	0
Injection site erythema			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Feeling of body temperature change			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Immune system disorders			
Food allergy			
alternative assessment type: Systematic			

subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Hypersensitivity			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Allergic oedema			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Reproductive system and breast disorders			
Vaginal haemorrhage			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Respiratory, thoracic and mediastinal disorders			
Dysphonia			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Asthma			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	1	0	0
Cough			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	1 / 20 (5.00%)
occurrences (all)	0	2	2
Dyspnoea			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Epistaxis			
alternative assessment type: Systematic			

subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	2
Rhinitis allergic			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 20 (10.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	2	0	0
Oropharyngeal pain			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	1	0	0
Nasal disorder			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Rhinorrhoea			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	1	0	1
Psychiatric disorders			
Depressed mood			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Anxiety			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Investigations			
Protein urine present			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Injury, poisoning and procedural complications			

Animal bite alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 18 (0.00%) 0	0 / 20 (0.00%) 0
Ligament sprain alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 18 (0.00%) 0	0 / 20 (0.00%) 0
Contusion alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 18 (5.56%) 1	0 / 20 (0.00%) 0
Hand fracture alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 18 (0.00%) 0	1 / 20 (5.00%) 1
Bone contusion alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 18 (0.00%) 0	0 / 20 (0.00%) 0
Muscle strain alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	1 / 18 (5.56%) 1	0 / 20 (0.00%) 0
Post procedural inflammation alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 20 (5.00%) 1	0 / 18 (0.00%) 0	0 / 20 (0.00%) 0
Congenital, familial and genetic disorders Dermoid cyst alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 18 (0.00%) 0	1 / 20 (5.00%) 1
Cardiac disorders			

Cyanosis alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 18 (0.00%) 0	1 / 20 (5.00%) 1
Nervous system disorders Dizziness alternative assessment type: Systematic subjects affected / exposed occurrences (all) Akathisia alternative assessment type: Systematic subjects affected / exposed occurrences (all) Headache alternative assessment type: Systematic subjects affected / exposed occurrences (all) Lethargy alternative assessment type: Systematic subjects affected / exposed occurrences (all) Syncope alternative assessment type: Systematic subjects affected / exposed occurrences (all) Paraesthesia alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0  0 / 20 (0.00%) 0  1 / 20 (5.00%) 2  1 / 20 (5.00%) 2  1 / 20 (5.00%) 1  0 / 20 (0.00%) 0  1 / 20 (5.00%) 1  0 / 20 (0.00%) 0  0 / 20 (0.00%) 0	0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  1 / 18 (5.56%) 1  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0	0 / 20 (0.00%) 0  1 / 20 (5.00%) 2  1 / 20 (5.00%) 0  0 / 20 (0.00%) 0  0 / 20 (0.00%) 0  0 / 20 (0.00%) 0
Blood and lymphatic system disorders Iron deficiency anaemia alternative assessment type: Systematic subjects affected / exposed occurrences (all) Eosinophilia	1 / 20 (5.00%) 1	0 / 18 (0.00%) 0	1 / 20 (5.00%) 1

alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 18 (0.00%) 0	0 / 20 (0.00%) 0
Eye disorders Conjunctivitis allergic alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 20 (0.00%) 0	0 / 18 (0.00%) 0	0 / 20 (0.00%) 0
Gastrointestinal disorders Aphthous stomatitis alternative assessment type: Systematic subjects affected / exposed occurrences (all)  Cheilitis alternative assessment type: Systematic subjects affected / exposed occurrences (all)  Constipation alternative assessment type: Systematic subjects affected / exposed occurrences (all)  Diarrhoea alternative assessment type: Systematic subjects affected / exposed occurrences (all)  Abdominal pain upper alternative assessment type: Systematic subjects affected / exposed occurrences (all)  Vomiting alternative assessment type: Systematic subjects affected / exposed occurrences (all)  Mouth ulceration alternative assessment type: Systematic	0 / 20 (0.00%) 0  0 / 20 (0.00%) 0  0 / 20 (0.00%) 0  0 / 20 (0.00%) 0  0 / 20 (0.00%) 0  1 / 20 (5.00%) 1	0 / 18 (0.00%) 0  1 / 18 (5.56%) 1  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0  0 / 18 (0.00%) 0	0 / 20 (0.00%) 0  0 / 20 (0.00%) 0  1 / 20 (5.00%) 1  1 / 20 (5.00%) 1  2 / 20 (10.00%) 2  2 / 20 (10.00%) 2

subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Nausea			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Skin and subcutaneous tissue disorders			
Acne			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Dermatitis contact			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Alopecia areata			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Dermatitis atopic			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 20 (10.00%)	5 / 18 (27.78%)	3 / 20 (15.00%)
occurrences (all)	2	11	6
Alopecia			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Dry skin			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	2
Erythema			
alternative assessment type: Systematic			



subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Skin erosion			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Psoriasis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Rash			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	1	0	0
Pain of skin			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	1	0	0
Solar dermatitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Vitiligo			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Urticaria			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Renal and urinary disorders			
Dysuria			
alternative assessment type: Systematic			

subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Haematuria			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Musculoskeletal and connective tissue disorders			
Back pain			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 20 (5.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	1	0	0
Arthralgia			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Bone pain			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Epiphysiolysis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Pain in extremity			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Infections and infestations			
Bacterial disease carrier			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Acute tonsillitis			
alternative assessment type: Systematic			

subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Gastrointestinal infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Conjunctivitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Croup infectious			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Dermatitis infected			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	2 / 20 (10.00%)
occurrences (all)	0	1	4
Ear infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Folliculitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Gingivitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Herpes simplex			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	3

Molluscum contagiosum			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Influenza			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 20 (10.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	2	0	0
Laryngitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Herpes virus infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Nasopharyngitis			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 20 (10.00%)	5 / 18 (27.78%)	8 / 20 (40.00%)
occurrences (all)	3	9	11
Oral herpes			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Rhinitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	2	0
Pharyngitis bacterial			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	1 / 18 (5.56%)	0 / 20 (0.00%)
occurrences (all)	0	1	0
Pharyngitis streptococcal			
alternative assessment type: Systematic			

subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Paronychia			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Skin bacterial infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Skin candida			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Urinary tract infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1
Upper respiratory tract infection			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 20 (10.00%)	1 / 18 (5.56%)	1 / 20 (5.00%)
occurrences (all)	2	1	1
Tonsillitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Viral upper respiratory tract infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	0 / 20 (0.00%)
occurrences (all)	0	0	0
Metabolism and nutrition disorders			
Increased appetite			
alternative assessment type: Systematic			

subjects affected / exposed	0 / 20 (0.00%)	0 / 18 (0.00%)	1 / 20 (5.00%)
occurrences (all)	0	0	1

<b>Non-serious adverse events</b>	Dupilumab 4 mg/kg: Children		
Total subjects affected by non-serious adverse events			
subjects affected / exposed	18 / 19 (94.74%)		
General disorders and administration site conditions			
Fatigue			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Chills			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Chest pain			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Pyrexia			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 19 (10.53%)		
occurrences (all)	2		
Injection site urticaria			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	3		
Injection site swelling			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Injection site irritation			
alternative assessment type: Systematic			

<p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Injection site erythema</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p> <p>Feeling of body temperature change</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Immune system disorders</p> <p>Food allergy</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p> <p>Hypersensitivity</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p> <p>Allergic oedema</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Reproductive system and breast disorders</p> <p>Vaginal haemorrhage</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Respiratory, thoracic and mediastinal disorders</p> <p>Dysphonia</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p> <p>Asthma</p> <p>alternative assessment type:</p>			

Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Cough			
alternative assessment type: Systematic			
subjects affected / exposed	7 / 19 (36.84%)		
occurrences (all)	8		
Dyspnoea			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Epistaxis			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Rhinitis allergic			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Oropharyngeal pain			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	2		
Nasal disorder			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Rhinorrhoea			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Psychiatric disorders			
Depressed mood			
alternative assessment type: Systematic			



<p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Anxiety</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 19 (0.00%)</p> <p>0</p> <p>0 / 19 (0.00%)</p> <p>0</p>		
<p>Investigations</p> <p>Protein urine present</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 19 (0.00%)</p> <p>0</p>		
<p>Injury, poisoning and procedural complications</p> <p>Animal bite</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Ligament sprain</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Contusion</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Hand fracture</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Bone contusion</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Muscle strain</p> <p>alternative assessment type: Systematic</p>	<p>0 / 19 (0.00%)</p> <p>0</p> <p>1 / 19 (5.26%)</p> <p>1</p> <p>0 / 19 (0.00%)</p> <p>0</p> <p>0 / 19 (0.00%)</p> <p>0</p> <p>1 / 19 (5.26%)</p> <p>1</p>		

<p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Post procedural inflammation</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 19 (0.00%)</p> <p>0</p> <p>0 / 19 (0.00%)</p> <p>0</p>		
<p>Congenital, familial and genetic disorders</p> <p>Dermoid cyst</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 19 (0.00%)</p> <p>0</p>		
<p>Cardiac disorders</p> <p>Cyanosis</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 19 (0.00%)</p> <p>0</p>		
<p>Nervous system disorders</p> <p>Dizziness</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Akathisia</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Headache</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Lethargy</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>Syncope</p> <p>alternative assessment type: Systematic</p>	<p>2 / 19 (10.53%)</p> <p>3</p> <p>0 / 19 (0.00%)</p> <p>0</p> <p>2 / 19 (10.53%)</p> <p>3</p> <p>1 / 19 (5.26%)</p> <p>1</p>		

<p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Paraesthesia</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p>			
<p>Blood and lymphatic system disorders</p> <p>Iron deficiency anaemia</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Eosinophilia</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p>			
<p>Eye disorders</p> <p>Conjunctivitis allergic</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>3</p>			
<p>Gastrointestinal disorders</p> <p>Aphthous stomatitis</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p> <p>Cheilitis</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Constipation</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Diarrhoea</p> <p>alternative assessment type: Systematic</p>			

subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Abdominal pain upper			
alternative assessment type: Systematic			
subjects affected / exposed	3 / 19 (15.79%)		
occurrences (all)	3		
Vomiting			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Mouth ulceration			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Nausea			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 19 (10.53%)		
occurrences (all)	2		
Skin and subcutaneous tissue disorders			
Acne			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Dermatitis contact			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Alopecia areata			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Dermatitis atopic			
alternative assessment type: Systematic			

subjects affected / exposed	5 / 19 (26.32%)		
occurrences (all)	10		
Alopecia			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Dry skin			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Erythema			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Skin erosion			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Psoriasis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Rash			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Pain of skin			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Solar dermatitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		

<p>Vitiligo</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Urticaria</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>2</p>			
<p>Renal and urinary disorders</p> <p>Dysuria</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Haematuria</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p>			
<p>Musculoskeletal and connective tissue disorders</p> <p>Back pain</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Arthralgia</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p> <p>Bone pain</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>1 / 19 (5.26%)</p> <p>occurrences (all)</p> <p>1</p> <p>Epiphysiolysis</p> <p>alternative assessment type: Systematic</p> <p>subjects affected / exposed</p> <p>0 / 19 (0.00%)</p> <p>occurrences (all)</p> <p>0</p> <p>Pain in extremity</p>			

alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 19 (5.26%) 1		
Infections and infestations Bacterial disease carrier alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 19 (0.00%) 0		
Acute tonsillitis alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 19 (5.26%) 2		
Gastrointestinal infection alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 19 (5.26%) 1		
Conjunctivitis alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 19 (5.26%) 2		
Croup infectious alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 19 (0.00%) 0		
Dermatitis infected alternative assessment type: Systematic subjects affected / exposed occurrences (all)	2 / 19 (10.53%) 5		
Ear infection alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 19 (5.26%) 1		
Folliculitis alternative assessment type: Systematic			

subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Gingivitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Herpes simplex			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Molluscum contagiosum			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Influenza			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Laryngitis			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Herpes virus infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Nasopharyngitis			
alternative assessment type: Systematic			
subjects affected / exposed	10 / 19 (52.63%)		
occurrences (all)	11		
Oral herpes			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		



Rhinitis			
alternative assessment type: Systematic			
subjects affected / exposed	2 / 19 (10.53%)		
occurrences (all)	2		
Pharyngitis bacterial			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Pharyngitis streptococcal			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Paronychia			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Skin bacterial infection			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	2		
Skin candida			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Urinary tract infection			
alternative assessment type: Systematic			
subjects affected / exposed	0 / 19 (0.00%)		
occurrences (all)	0		
Upper respiratory tract infection			
alternative assessment type: Systematic			
subjects affected / exposed	1 / 19 (5.26%)		
occurrences (all)	1		
Tonsillitis			
alternative assessment type: Systematic			

subjects affected / exposed occurrences (all)  Viral upper respiratory tract infection alternative assessment type: Systematic subjects affected / exposed occurrences (all)	1 / 19 (5.26%)  1    1 / 19 (5.26%)  1		
Metabolism and nutrition disorders Increased appetite alternative assessment type: Systematic subjects affected / exposed occurrences (all)	0 / 19 (0.00%)  0		

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
24 June 2015	Following changes were made: •Changes were made regarding the use of topical corticosteroids (TCS) and topical calcineurin inhibitors (TCI) were permitted during the study to avoid subjects being unnecessarily deprived of topical medication. •Clarified the use of prescription moisturizers started prior to the study were permitted to continue during the study, but initiation of prescription moisturizers during the study was not allowed. •Clarified the definition of "childbearing potential" for the purpose of the study, and added a note regarding additional requirement for adequate contraceptive methods in certain countries to avoid ambiguity in the interpretation of the relevant exclusion criterion. •Added assessment of body surface area (BSA) affected by AD, at the same time points other efficacy assessments were being performed; added BSA related secondary endpoint. •Clarified that only subjects with a history of active infection with hepatitis B or C or evidence of active disease at screening were excluded from the study. •Clarified the duration of close monitoring required after study drug administration and assessments performed during the monitoring period. • Added text regarding opportunity for subjects who completed the study to enroll into an open-label extension study to continue Dupilumab treatment. •Reduced the amount of information gathered from subjects on pruritus based on NRS. •Clarified cheek swab sample taken for DNA extraction (removed the options of taking whole blood or saliva sample).

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported