



## Clinical trial results:

### **RAINBOW extension study: an extension study to evaluate the long-term efficacy and safety of RAnibizumab compared with laser therapy for the treatment of INfants BOrn prematurely With retinopathy of prematurity**

#### **Summary**

EudraCT number	2014-004048-36
Trial protocol	HU EE GB SK IT BE LT CZ DE AT SE GR DK FR HR PL
Global end of trial date	21 April 2022

#### **Results information**

Result version number	v1
This version publication date	05 November 2022
First version publication date	05 November 2022

#### **Trial information**

##### **Trial identification**

Sponsor protocol code	CRFB002H2301E1
-----------------------	----------------

##### **Additional study identifiers**

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02640664
WHO universal trial number (UTN)	-

Notes:

##### **Sponsors**

Sponsor organisation name	Novartis Pharma AG
Sponsor organisation address	CH-4002, Basel, Switzerland,
Public contact	Clinical Disclosure Office, Novartis Pharma AG, 41 613241111, Novartis.email@novartis.com
Scientific contact	Clinical Disclosure Office, Novartis Pharma AG, 41 613241111, Novartis.email@novartis.com

Notes:

##### **Paediatric regulatory details**

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	21 April 2022
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	21 April 2022
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

To evaluate the visual function of patients, by assessing the visual acuity (VA) in the better-seeing eye at the patients' 5th birthday.

Due to EudraCT system limitations, which EMA is aware of, data using 999 as data points in this record are not an accurate representation of the clinical trial results. Please use <https://www.novctrd.com/CtrdWeb/home.nov> for complete trial results.

Protection of trial subjects:

The study was in compliance with the ethical principles derived from the Declaration of Helsinki and the International Conference on Harmonization (ICH) Good Clinical Practice (GCP) guidelines. All the local regulatory requirements pertinent to safety of trial subjects were also followed during the conduct of the trial.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	16 June 2016
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Austria: 4
Country: Number of subjects enrolled	Belgium: 10
Country: Number of subjects enrolled	Croatia: 5
Country: Number of subjects enrolled	Czechia: 9
Country: Number of subjects enrolled	Denmark: 1
Country: Number of subjects enrolled	Egypt: 1
Country: Number of subjects enrolled	Estonia: 2
Country: Number of subjects enrolled	France: 2
Country: Number of subjects enrolled	Germany: 2
Country: Number of subjects enrolled	United Kingdom: 4
Country: Number of subjects enrolled	Greece: 10
Country: Number of subjects enrolled	Hungary: 2
Country: Number of subjects enrolled	India: 23
Country: Number of subjects enrolled	Italy: 12
Country: Number of subjects enrolled	Japan: 25
Country: Number of subjects enrolled	Lithuania: 1
Country: Number of subjects enrolled	Malaysia: 2

Country: Number of subjects enrolled	Romania: 13
Country: Number of subjects enrolled	Russian Federation: 18
Country: Number of subjects enrolled	Saudi Arabia: 1
Country: Number of subjects enrolled	Slovakia: 1
Country: Number of subjects enrolled	Taiwan: 7
Country: Number of subjects enrolled	Turkey: 10
Country: Number of subjects enrolled	United States: 15
Worldwide total number of subjects	180
EEA total number of subjects	74

Notes:

---

### **Subjects enrolled per age group**

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	3
Infants and toddlers (28 days-23 months)	177
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

Study was carried out in Austria (2), Belgium (2), Croatia (1), Czech Republic (3), Denmark (1), Egypt (1), Estonia (1), France (2), Germany (1), Greece (3), Hungary (2), India (6), Italy (4), Japan (16), Lithuania (1), Malaysia (2), Romania (3), Russian Federation (5), Saudi Arabia (1), Slovakia (1), Taiwan (2), Turkey (3), UK (2), US (9)

### Pre-assignment

Screening details:

This study was not randomized. In the core study, participants were randomized to 1 of the 3 treatment arms (ranibizumab 0.2 mg, ranibizumab 0.1 mg, and laser). Treatment arm assignment and participant identifier in the extension study remained the same as in the core study.

### Period 1

Period 1 title	Core study: Epoch 1
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Not blinded

Blinding implementation details:

This was an open-label study where the VA assessment at the child's 5th birthday visit was performed by an assessor who was masked to study treatment.

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Ranibizumab 0.2 mg

Arm description:

1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required

Arm type	Experimental
Investigational medicinal product name	Ranibizumab
Investigational medicinal product code	RFB002
Other name	
Pharmaceutical forms	Injection
Routes of administration	Intravitreal use

Dosage and administration details:

1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required

<b>Arm title</b>	Ranibizumab 0.1 mg
------------------	--------------------

Arm description:

1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required

Arm type	Experimental
Investigational medicinal product name	Ranibizumab
Investigational medicinal product code	RFB002
Other name	
Pharmaceutical forms	Injection
Routes of administration	Intravitreal use

Dosage and administration details:

1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required

<b>Arm title</b>	Laser therapy
------------------	---------------

Arm description:

Laser treatment to each eye on Day 1 (Baseline), with supplementary treatments allowed

Arm type | laser ablation therapy

No investigational medicinal product assigned in this arm

<b>Number of subjects in period 1</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy
Started	61	65	54
Completed	60	65	53
Not completed	1	0	1
Adverse event, serious fatal	-	-	1
Withdrawal of informed consent	1	-	-

## Period 2

Period 2 title | Extension study: Epoch 2

Is this the baseline period? | No

Allocation method | Randomised - controlled

Blinding used | Not blinded

Blinding implementation details:

This was an open-label study where the VA assessment at the child's 5th birthday visit was performed by an assessor who was masked to study treatment.

## Arms

Are arms mutually exclusive? | Yes

**Arm title** | Ranibizumab 0.2 mg

Arm description:

1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required

Arm type | Experimental

Investigational medicinal product name | Ranibizumab

Investigational medicinal product code | RFB002

Other name |

Pharmaceutical forms | Injection

Routes of administration | Intravitreal use

Dosage and administration details:

1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required

**Arm title** | Ranibizumab 0.1 mg

Arm description:

1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required

Arm type | Experimental

Investigational medicinal product name	Ranibizumab
Investigational medicinal product code	RFB002
Other name	
Pharmaceutical forms	Injection
Routes of administration	Intravitreal use
Dosage and administration details:	
1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required	
<b>Arm title</b>	Laser therapy
Arm description:	
Laser treatment to each eye on Day 1 (Baseline), with supplementary treatments allowed	
Arm type	laser ablation therapy
No investigational medicinal product assigned in this arm	

<b>Number of subjects in period 2</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy
Started	60	65	53
Completed	54	55	47
Not completed	6	10	6
Adverse event, serious fatal	-	-	1
Withdrawal of informed consent	3	4	2
Lost to follow-up	3	4	1
Subject/guardian decision	-	2	2

## Baseline characteristics

### Reporting groups

Reporting group title	Ranibizumab 0.2 mg
Reporting group description:	1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required
Reporting group title	Ranibizumab 0.1 mg
Reporting group description:	1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required
Reporting group title	Laser therapy
Reporting group description:	Laser treatment to each eye on Day 1 (Baseline), with supplementary treatments allowed

Reporting group values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy
Number of subjects	61	65	54
Age categorical			
Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	1	1	1
Infants and toddlers (28 days-23 months)	60	64	53
Children (2-11 years)	0	0	0
Adolescents (12-17 years)	0	0	0
Adults (18-64 years)	0	0	0
From 65-84 years	0	0	0
85 years and over	0	0	0
Age Continuous			
Age at core baseline visit			
Units: Weeks			
arithmetic mean	11.32	11.07	9.89
standard deviation	± 3.974	± 3.515	± 3.744
Sex: Female, Male			
Units: Participants			
Female	32	35	27
Male	29	30	27
Race/Ethnicity, Customized			
Units: Subjects			
Caucasian	38	40	32
Black	0	4	2
Asian	22	18	18
Native American	0	0	0
Pacific Islander	0	0	0
Unknown	0	0	0
Other	1	3	2

<b>Reporting group values</b>	Total		
-------------------------------	-------	--	--

Number of subjects	180		
Age categorical			
Units: Subjects			
In utero	0		
Preterm newborn infants (gestational age < 37 wks)	0		
Newborns (0-27 days)	3		
Infants and toddlers (28 days-23 months)	177		
Children (2-11 years)	0		
Adolescents (12-17 years)	0		
Adults (18-64 years)	0		
From 65-84 years	0		
85 years and over	0		
Age Continuous			
Age at core baseline visit			
Units: Weeks			
arithmetic mean			
standard deviation	-		
Sex: Female, Male			
Units: Participants			
Female	94		
Male	86		
Race/Ethnicity, Customized			
Units: Subjects			
Caucasian	110		
Black	6		
Asian	58		
Native American	0		
Pacific Islander	0		
Unknown	0		
Other	6		

## End points

### End points reporting groups

Reporting group title	Ranibizumab 0.2 mg
Reporting group description: 1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required	
Reporting group title	Ranibizumab 0.1 mg
Reporting group description: 1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required	
Reporting group title	Laser therapy
Reporting group description: Laser treatment to each eye on Day 1 (Baseline), with supplementary treatments allowed	
Reporting group title	Ranibizumab 0.2 mg
Reporting group description: 1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required	
Reporting group title	Ranibizumab 0.1 mg
Reporting group description: 1 intravitreal injection in both eyes on Day 1 (Baseline), with up to 2 re-treatments allowed for each eye if required	
Reporting group title	Laser therapy
Reporting group description: Laser treatment to each eye on Day 1 (Baseline), with supplementary treatments allowed	

### Primary: Visual acuity (letters) of the better-seeing eye at the participant's fifth birthday visit – comparison between treatment arms

End point title	Visual acuity (letters) of the better-seeing eye at the participant's fifth birthday visit – comparison between treatment arms
End point description: The VA assessment at the child's 5th birthday visit was performed using ETDRS methodology. VA measurements were taken in a sitting position at an initial test distance of 3 meters using Lea Symbols charts. Scoring was reported as a letter score representing the number of optotypes (Lea symbols) the participant identified. VA was tested in each eye, using the child's current refractive index. If the child had a score of '0' (could not identify any of the Lea symbols on the chart at 1 meter) on the VA assessment or the score was 'missing', they were tested with low VA testing. Low VA testing consisted of determining if the child could count the examiner's fingers, see hand motion, or perceive light. The better-seeing eye was defined as the eye with the higher ETDRS score at the 5th birthday visit. If both eyes had the same ETDRS letter score, then the right eye was assigned as the better-seeing eye.	
End point type	Primary
End point timeframe: at the participant's fifth birthday visit	

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	45	43	36	
Units: Score				
least squares mean (standard error)	66.8 (± 1.95)	64.6 (± 2.00)	62.1 (± 2.18)	

### Statistical analyses

<b>Statistical analysis title</b>	Ranibizumab 0.2 mg vs Laser
Comparison groups	Ranibizumab 0.2 mg v Laser therapy
Number of subjects included in analysis	81
Analysis specification	Pre-specified
Analysis type	superiority
Method	ANOVA
Parameter estimate	Mean difference (net)
Point estimate	4.7
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.1
upper limit	10.5
Variability estimate	Standard error of the mean
Dispersion value	2.93

<b>Statistical analysis title</b>	Ranibizumab 0.2 mg vs Ranibizumab 0.1 mg
Comparison groups	Ranibizumab 0.2 mg v Ranibizumab 0.1 mg
Number of subjects included in analysis	88
Analysis specification	Pre-specified
Analysis type	superiority
Method	ANOVA
Parameter estimate	Mean difference (net)
Point estimate	2.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.3
upper limit	7.8
Variability estimate	Standard error of the mean
Dispersion value	2.79

<b>Statistical analysis title</b>	Ranibizumab 0.1 mg vs Laser
Comparison groups	Ranibizumab 0.1 mg v Laser therapy

Number of subjects included in analysis	79
Analysis specification	Pre-specified
Analysis type	superiority
Method	ANOVA
Parameter estimate	Mean difference (net)
Point estimate	2.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.4
upper limit	8.3
Variability estimate	Standard error of the mean
Dispersion value	2.96

---

**Secondary: Number of participants with ocular adverse events regardless of study treatment or procedure relationship by preferred term**

End point title	Number of participants with ocular adverse events regardless of study treatment or procedure relationship by preferred term
End point description: Number of participants with ocular AEs starting during the core study and ongoing at extension baseline, or starting on/after extension baseline were reported.	
End point type	Secondary
End point timeframe: throughout the study, approximately 5 years	

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61	65	54	
Units: Participants	19	26	22	

**Statistical analyses**

No statistical analyses for this end point

---

**Secondary: Visual acuity (VA) of the worse-seeing eye at the participant's fifth birthday visit – comparison between treatment arms**

End point title	Visual acuity (VA) of the worse-seeing eye at the participant's fifth birthday visit – comparison between treatment arms
-----------------	--

End point description:

The VA assessment at the child's 5th birthday visit was performed using ETDRS methodology. VA measurements were taken in a sitting position at an initial test distance of 3 meters using Lea Symbols charts. Scoring was reported as a letter score representing the number of optotypes (Lea symbols) the patient identified. VA was tested in each eye, using the child's current refractive index. If the child had a score of '0' (could not identify any of the Lea symbols on the chart at 1 meter) on the VA assessment or the score was 'missing', they were tested with low VA testing. Low VA testing consisted of determining if the child could count the examiner's fingers, see hand motion, or perceive light. The worse-seeing eye

was the eye with a lower ETDRS letter score at the 5th birthday visit. If both eyes had the same ETDRS letter score, then the left eye is assigned as the worse-seeing eye.

End point type	Secondary
----------------	-----------

End point timeframe:

at the participant's fifth birthday visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	45	42	36	
Units: Score				
least squares mean (standard error)	60.2 (± 2.95)	53.8 (± 3.05)	52.2 (± 3.30)	

### Statistical analyses

<b>Statistical analysis title</b>	Ranibizumab 0.2 mg vs Laser
Comparison groups	Ranibizumab 0.2 mg v Laser therapy
Number of subjects included in analysis	81
Analysis specification	Pre-specified
Analysis type	superiority
Method	ANOVA
Parameter estimate	Mean difference (net)
Point estimate	8
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.8
upper limit	16.7
Variability estimate	Standard error of the mean
Dispersion value	4.42

<b>Statistical analysis title</b>	Ranibizumab 0.2 mg vs Ranibizumab 0.1 mg
Comparison groups	Ranibizumab 0.2 mg v Ranibizumab 0.1 mg
Number of subjects included in analysis	87
Analysis specification	Pre-specified
Analysis type	
Method	ANOVA
Parameter estimate	Mean difference (net)
Point estimate	6.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2
upper limit	14.8

Variability estimate	Standard error of the mean
Dispersion value	4.24

<b>Statistical analysis title</b>	Ranibizumab 0.1 mg vs Laser
Comparison groups	Ranibizumab 0.1 mg v Laser therapy
Number of subjects included in analysis	78
Analysis specification	Pre-specified
Analysis type	superiority
Method	ANOVA
Parameter estimate	Mean difference (net)
Point estimate	1.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-7.3
upper limit	10.5
Variability estimate	Standard error of the mean
Dispersion value	4.49

**Secondary: Number of participants with non-ocular adverse events regardless of study treatment or procedure relationship (greater than or equal to 3% in any arm) by preferred term**

End point title	Number of participants with non-ocular adverse events regardless of study treatment or procedure relationship (greater than or equal to 3% in any arm) by preferred term
-----------------	--

End point description:

Number of participants with non-ocular adverse events regardless of study treatment or procedure relationship (greater than or equal to 3% in any arm) by preferred term were reported.

End point type	Secondary
----------------	-----------

End point timeframe:

throughout the study, approximately 5 years

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61	65	54	
Units: Participants	46	53	46	

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Number of participants with absence of active Retinopathy of**

## Prematurity (ROP) at 40 weeks post core baseline visit

End point title	Number of participants with absence of active Retinopathy of Prematurity (ROP) at 40 weeks post core baseline visit
-----------------	---

End point description:

The absence of active ROP in both eyes is defined by the absence of all of the following features: (1) Vessel dilatation of plus disease in at least 2 quadrants (some persisting tortuosity is allowed), (2) Extra-retina vessels extending from the retina into the vitreous and judged to be a sign of active ROP disease.

End point type	Secondary
----------------	-----------

End point timeframe:

at 40 weeks post core baseline visit

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	55	58	46	
Units: Participants				
Absence of active ROP	55	58	46	

## Statistical analyses

No statistical analyses for this end point

## Secondary: Number of participants with absence of active Retinopathy of Prematurity (ROP) at 52 weeks post core baseline visit

End point title	Number of participants with absence of active Retinopathy of Prematurity (ROP) at 52 weeks post core baseline visit
-----------------	---

End point description:

The absence of active ROP in both eyes is defined by the absence of all of the following features: (1) Vessel dilatation of plus disease in at least 2 quadrants (some persisting tortuosity is allowed), (2) Extra-retina vessels extending from the retina into the vitreous and judged to be a sign of active ROP disease.

End point type	Secondary
----------------	-----------

End point timeframe:

at 52 weeks post core baseline visit

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	58	63	50	
Units: Participants	58	63	49	

## Statistical analyses

No statistical analyses for this end point

---

**Secondary: Number of participants with absence of all ocular structural abnormalities at or before 40 weeks post baseline visit**

---

End point title	Number of participants with absence of all ocular structural abnormalities at or before 40 weeks post baseline visit
-----------------	--

End point description:

The absence of ocular structural abnormalities is defined by the absence of all of the following fundus features in both eyes at or before the given time point: (1) Substantial temporal retinal vessel dragging causing abnormal structural features/macular Ectopia, (2) Retrolental membrane obscuring the view of the posterior pole, (3) Posterior retinal fold involving the macula, (4) Retinal detachment involving the macula

End point type	Secondary
----------------	-----------

End point timeframe:

at or before 40 weeks post baseline visit

---

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	60	65	53	
Units: Participants	59	61	47	

---

**Statistical analyses**

---

No statistical analyses for this end point

---

**Secondary: Number of participants with absence of all ocular structural abnormalities at or before the participant's fifth birthday visit**

---

End point title	Number of participants with absence of all ocular structural abnormalities at or before the participant's fifth birthday visit
-----------------	--

End point description:

The absence of ocular structural abnormalities is defined by the absence of all of the following fundus features in both eyes at or before the given time point: (1) Substantial temporal retinal vessel dragging causing abnormal structural features/macular Ectopia, (2) Retrolental membrane obscuring the view of the posterior pole, (3) Posterior retinal fold involving the macula, (4) Retinal detachment involving the macula

End point type	Secondary
----------------	-----------

End point timeframe:

at or before the participant's fifth birthday visit

---

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	60	65	53	
Units: Participants	59	61	47	

## Statistical analyses

No statistical analyses for this end point

### Secondary: Number of participants with absence of individual ocular structural abnormalities at or before 40 weeks post baseline visit

End point title	Number of participants with absence of individual ocular structural abnormalities at or before 40 weeks post baseline visit
-----------------	---

End point description:

The absence of ocular structural abnormalities is defined by the absence of all of the following fundus features in both eyes at or before the given time point: (1) Substantial temporal retinal vessel dragging causing abnormal structural features/macular Ectopia, (2) Retrolental membrane obscuring the view of the posterior pole, (3) Posterior retinal fold involving the macula, (4) Retinal detachment involving the macula

End point type	Secondary
----------------	-----------

End point timeframe:

at or before 40 weeks post baseline visit

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	60	65	53	
Units: Participants				
Absence of substantial temporal retinal vessel	59	63	49	
Absence of Retrolental membrane	60	65	53	
Absence of posterior retinal fold	59	65	51	
Absence of retinal detachment	60	63	51	
Abs of retinal detachment not involving the macula	60	62	50	
Absence of pre-retinal fibrosis	58	60	49	

## Statistical analyses

No statistical analyses for this end point

### Secondary: Number of participants with absence of individual ocular structural abnormalities at or before the participant's fifth birthday visit

End point title	Number of participants with absence of individual ocular structural abnormalities at or before the participant's fifth birthday visit
-----------------	---

End point description:

The absence of ocular structural abnormalities is defined by the absence of all of the following fundus features in both eyes at or before the given time point: (1) Substantial temporal retinal vessel dragging causing abnormal structural features/macular Ectopia, (2) Retrolental membrane obscuring the view of the posterior pole, (3) Posterior retinal fold involving the macula, (4) Retinal detachment involving the macula

End point type Secondary

End point timeframe:

at or before the participant's fifth birthday visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	60	65	53	
Units: Participants				
Absence of temporal retinal vessel dragging	59	63	49	
Absence of Retrolental membrane	60	65	52	
Absence of posterior retinal fold	59	65	51	
Absence of retinal detachment	60	63	50	
Abs of retinal detachment not involving macula	60	62	50	
Absence of pre-retinal fibrosis	58	59	48	
Absence of optic disc pallor	60	65	52	
Absence of optic disc swelling	60	65	53	
Absence of pigmentary disturbance in the macula	60	64	52	
Absence of atrophic changes in macula	60	65	51	

## Statistical analyses

No statistical analyses for this end point

## Secondary: Number of participants with absence of all ocular structural abnormalities at or before participant's 2 years corrected age visit

End point title Number of participants with absence of all ocular structural abnormalities at or before participant's 2 years corrected age visit

End point description:

The absence of ocular structural abnormalities is defined by the absence of all of the following fundus features in both eyes at or before the given time point: (1) Substantial temporal retinal vessel dragging causing abnormal structural features/macular Ectopia, (2) Retrolental membrane obscuring the view of the posterior pole, (3) Posterior retinal fold involving the macula, (4) Retinal detachment involving the macula

End point type Secondary

End point timeframe:

at or before participant's 2 years corrected age visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	60	65	53	
Units: Participants	59	61	47	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of participants with absence of individual ocular structural abnormalities at or before participant's 2 years corrected age visit

End point title	Number of participants with absence of individual ocular structural abnormalities at or before participant's 2 years corrected age visit
-----------------	--

End point description:

The absence of ocular structural abnormalities is defined by the absence of all of the following fundus features in both eyes at or before the given time point: (1) Substantial temporal retinal vessel dragging causing abnormal structural features/macular Ectopia, (2) Retrolental membrane obscuring the view of the posterior pole, (3) Posterior retinal fold involving the macula, (4) Retinal detachment involving the macula

End point type	Secondary
----------------	-----------

End point timeframe:

at or before participant's 2 years corrected age visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	60	65	53	
Units: Participants				
Abs of temporal retinal vessel dragging	59	63	49	
Absence of Retrolental membrane	60	65	52	
Absence of posterior retinal fold	59	65	51	
Abs of retinal detachment involving the macula	60	63	50	
Abs of retinal detachment not involving the macula	60	62	50	
Absence of pre-retinal fibrosis	58	59	48	
Absence of optic disc pallor	60	65	53	
Absence of optic disc swelling	60	65	53	
Absence of pigmentary disturbance in the macula	60	64	52	
Absence of atrophic changes in macula	60	65	52	

### Statistical analyses

No statistical analyses for this end point

**Secondary: Number of participants with recurrence of ROP up to 40 weeks post baseline visit in the core study**

End point title	Number of participants with recurrence of ROP up to 40 weeks post baseline visit in the core study
-----------------	--

End point description:

Recurrence of ROP was defined as ROP receiving any post-baseline intervention after the 1st study treatment in the core study. In the ranibizumab arms, post-baseline interventions were ranibizumab retreatment or switch to laser. In the laser arm, post-baseline interventions were supplementary laser treatments after 11 days post-baseline, or switch to ranibizumab; supplementary laser treatment within 11 days post-baseline was not counted as recurrence.

End point type	Secondary
----------------	-----------

End point timeframe:

up to 40 weeks post baseline visit in the core study

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61	65	54	
Units: Participants	19	22	11	

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Number of participants with recurrence of ROP up to 52 weeks post baseline visit in the core study**

End point title	Number of participants with recurrence of ROP up to 52 weeks post baseline visit in the core study
-----------------	--

End point description:

Recurrence of ROP was defined as ROP receiving any post-baseline intervention after the 1st study treatment in the core study. In the ranibizumab arms, post-baseline interventions were ranibizumab retreatment or switch to laser. In the laser arm, post-baseline interventions were supplementary laser treatments after 11 days post-baseline, or switch to ranibizumab; supplementary laser treatment within 11 days post-baseline was not counted as recurrence.

End point type	Secondary
----------------	-----------

End point timeframe:

up to 52 weeks post baseline visit in the core study

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61 <sup>[1]</sup>	65 <sup>[2]</sup>	54 <sup>[3]</sup>	
Units: Participants	999	999	999	

Notes:

[1] - analysis identical to 'recurrence of ROP upto 40 wks post-baseline' as recurrence remained unchanged

[2] - analysis identical to 'recurrence of ROP upto 40 wks post-baseline' as recurrence remained

unchanged

[3] - analysis identical to 'recurrence of ROP upto 40 wks post-baseline' as recurrence remained unchanged

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of ranibizumab injections received per participant over the whole safety observation period

End point title	Number of ranibizumab injections received per participant over the whole safety observation period
-----------------	--

End point description:

Number of ranibizumab injections received in the treatment of participants with ROP up to and including 40 weeks post baseline visit in the core study were reported.

End point type	Secondary
----------------	-----------

End point timeframe:

up to and including 40 weeks post baseline visit in the core study

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61	65	11	
Units: Number of injections				
arithmetic mean (standard deviation)	2.5 ( $\pm$ 0.96)	2.5 ( $\pm$ 1.06)	2.4 ( $\pm$ 0.92)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Refraction status: Summary of participants at participant's 2 years corrected age

End point title	Refraction status: Summary of participants at participant's 2 years corrected age
-----------------	---

End point description:

Summary of participants was reported to evaluate the refraction in each eye at the participant's 2 years corrected age

End point type	Secondary
----------------	-----------

End point timeframe:

at participant's 2 years corrected age

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	57	54	47	
Units: diopters				
arithmetic mean (standard deviation)				
Best eye	-0.697 (± 2.7032)	-0.713 (± 2.6100)	-1.793 (± 4.1570)	
Worst eye	-0.825 (± 2.6575)	-0.829 (± 2.8346)	-1.516 (± 3.5652)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Refraction status: Summary of participants at the participant's fifth birthday visit - Best eye

End point title	Refraction status: Summary of participants at the participant's fifth birthday visit - Best eye
End point description:	Summary of participants was reported to evaluate the refraction in each eye at the participant's fifth birthday visit
End point type	Secondary
End point timeframe:	at the participant's fifth birthday visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	52	55	45	
Units: diopters				
arithmetic mean (standard deviation)				
Best eye	-0.601 (± 2.8107)	-0.859 (± 2.7406)	-1.883 (± 4.4970)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Refraction status: Summary of participants at the participant's fifth birthday visit - Worst eye

End point title	Refraction status: Summary of participants at the participant's fifth birthday visit - Worst eye
End point description:	Summary of participants was reported to evaluate the refraction in each eye at the participant's fifth birthday visit
End point type	Secondary

End point timeframe:  
at the participant's fifth birthday visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	52	54	45	
Units: diopters				
arithmetic mean (standard deviation)	-0.904 ( $\pm$ 2.8501)	-1.074 ( $\pm$ 3.0405)	-1.706 ( $\pm$ 3.5740)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from baseline in weight at the participant's 2 years' corrected age

End point title	Change from baseline in weight at the participant's 2 years' corrected age
-----------------	--

End point description:

Subject's weight was reported to evaluate the physical development.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, at the participant's 2 years' corrected age

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	56	58	46	
Units: grams				
arithmetic mean (standard deviation)				
Year 2	8695.9 ( $\pm$ 1406.58)	8461.7 ( $\pm$ 1375.95)	8840.6 ( $\pm$ 1643.80)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from baseline in weight at the participant's fifth birthday

End point title	Change from baseline in weight at the participant's fifth birthday
-----------------	--

End point description:

Subject's weight was reported to evaluate the physical development.

End point type	Secondary
End point timeframe:	
Baseline, at the participant's fifth birthday	

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	52	54	45	
Units: grams				
arithmetic mean (standard deviation)	14611.9 ( $\pm$ 3018.65)	14324.7 ( $\pm$ 2616.96)	14689.4 ( $\pm$ 3300.05)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from baseline in Head Circumference

End point title	Change from baseline in Head Circumference
End point description:	
Subject's head circumference was reported to evaluate the physical development.	
End point type	Secondary
End point timeframe:	
Baseline and at the subject's 2 years' corrected age	

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	55	56	44	
Units: cm				
arithmetic mean (standard deviation)	16.5 ( $\pm$ 2.60)	16.2 ( $\pm$ 2.78)	17.2 ( $\pm$ 2.81)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from baseline in Sitting Diastolic Blood Pressure

End point title	Change from baseline in Sitting Diastolic Blood Pressure
End point description:	
Subject's Sitting Diastolic Blood Pressure was reported to evaluate the physical development.	
End point type	Secondary
End point timeframe:	
Baseline and at the subject's 2 years' corrected age	

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	38	33	31	
Units: mmHg				
arithmetic mean (standard deviation)	16.7 (± 14.92)	13.6 (± 14.10)	16.5 (± 12.14)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from baseline in Sitting Systolic Blood Pressure

End point title	Change from baseline in Sitting Systolic Blood Pressure
End point description:	Subject 's Sitting Systolic Blood Pressure was reported to evaluate the physical development.
End point type	Secondary
End point timeframe:	Baseline and at the subject's 2 years' corrected age

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	38	32	31	
Units: mmHg				
arithmetic mean (standard deviation)	20.6 (± 16.13)	16.2 (± 15.95)	17.6 (± 12.75)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of participants with the summary of respiratory function status

End point title	Number of participants with the summary of respiratory function status
End point description:	Number of participants with respiratory function status was reported
End point type	Secondary
End point timeframe:	at the participants' fifth birthday visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61	65	54	
Units: Participants				
participants with Wheezing/whistling status	6	4	2	
participants with attacks of wheezing	47	50	45	
participants with frequency of sleep disturbance	47	52	47	
participants with wheezing limiting speech ability	1	0	0	
participants with dry cough status at night	5	2	1	
participants with presence of smoker at home	15	8	4	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of participants with hearing impairment of any type

End point title	Number of participants with hearing impairment of any type
End point description:	Number of participants with hearing function status was reported
End point type	Secondary
End point timeframe:	at the subjects' fifth birthday visit

<b>End point values</b>	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61	65	54	
Units: Participants				
participants with hearing impairment of any type	2	2	4	
participants with use of hearing aid(s)	1	2	3	
participants with usual hearing	49	51	41	

### Statistical analyses

No statistical analyses for this end point

## Secondary: Weight at the time of first hospital discharge

End point title	Weight at the time of first hospital discharge
End point description:	Weight (gram) at the time of first hospital discharge was reported to evaluate the health status of the subject
End point type	Secondary
End point timeframe:	from birth to first hospital discharge home, up to 5 years

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	56	62	51	
Units: gram				
arithmetic mean (standard deviation)	2910.9 ( $\pm$ 1359.34)	2966.5 ( $\pm$ 1198.60)	2658.7 ( $\pm$ 926.92)	

## Statistical analyses

No statistical analyses for this end point

## Secondary: Duration of hospitalization

End point title	Duration of hospitalization
End point description:	Duration of hospitalization was reported to evaluate the health status of the subject
End point type	Secondary
End point timeframe:	from birth to first hospital discharge home, up to 5 years

End point values	Ranibizumab 0.2 mg	Ranibizumab 0.1 mg	Laser therapy	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	61	65	54	
Units: days				
arithmetic mean (standard deviation)	111.1 ( $\pm$ 62.27)	116.2 ( $\pm$ 78.10)	95.7 ( $\pm$ 57.07)	

## Statistical analyses

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Timeframe for AE

Adverse event reporting additional description:

AE additional description

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	25.0
--------------------	------

### Reporting groups

Reporting group title	Ranibizumab 0.2 mg
-----------------------	--------------------

Reporting group description:

Ranibizumab 0.2 mg

Reporting group title	Laser
-----------------------	-------

Reporting group description:

Laser

Reporting group title	Total
-----------------------	-------

Reporting group description:

Total

Reporting group title	Ranibizumab 0.1 mg
-----------------------	--------------------

Reporting group description:

Ranibizumab 0.1 mg

<b>Serious adverse events</b>	Ranibizumab 0.2 mg	Laser	Total
Total subjects affected by serious adverse events			
subjects affected / exposed	21 / 61 (34.43%)	26 / 54 (48.15%)	74 / 180 (41.11%)
number of deaths (all causes)	0	2	2
number of deaths resulting from adverse events	0	0	0
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Haemangioma			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
Perinatal brain damage			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	3 / 61 (4.92%)	2 / 54 (3.70%)	6 / 180 (3.33%)
occurrences causally related to treatment / all	0 / 3	0 / 3	0 / 7
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Inflammation			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Developmental delay			
subjects affected / exposed	2 / 61 (3.28%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Acute respiratory failure			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Asthma			
subjects affected / exposed	1 / 61 (1.64%)	2 / 54 (3.70%)	6 / 180 (3.33%)
occurrences causally related to treatment / all	0 / 2	0 / 3	0 / 9
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bronchial obstruction			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bronchopulmonary dysplasia			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bronchospasm			

subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Dyspnoea</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Cough</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 2	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Hypercapnia</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Hypoxia</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Nasal discharge discolouration</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Laryngospasm</b>			
subjects affected / exposed	2 / 61 (3.28%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Increased bronchial secretion</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Pulmonary vein stenosis</b>			

subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Pulmonary hypertension</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	1 / 1	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Pneumonitis</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Respiratory distress</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Respiratory failure</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Tonsillar hypertrophy</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Psychiatric disorders</b>			
<b>Psychomotor retardation</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Investigations</b>			
<b>Oxygen saturation decreased</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Injury, poisoning and procedural</b>			

complications			
Head injury			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Foreign body in respiratory tract			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Endotracheal intubation complication			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Periorbital haematoma			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Congenital, familial and genetic disorders			
Cerebral palsy			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Coarctation of the aorta			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cryptorchism			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders			
Cardiac arrest			

subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 1
<b>Bradycardia</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Nervous system disorders</b>			
<b>Cerebellar haemorrhage</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Febrile convulsion</b>			
subjects affected / exposed	2 / 61 (3.28%)	0 / 54 (0.00%)	5 / 180 (2.78%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 5
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Epilepsy</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Cognitive disorder</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Hydrocephalus</b>			
subjects affected / exposed	1 / 61 (1.64%)	1 / 54 (1.85%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Intracranial pressure increased</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Nystagmus</b>			

subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Myoclonus</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Motor dysfunction</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Quadripareisis</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Periventricular leukomalacia</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Ear and labyrinth disorders</b>			
<b>Deafness</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Eye disorders</b>			
<b>Conjunctival haemorrhage</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Eye disorder</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Retinal haemorrhage</b>			

subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Retinal detachment			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Retinopathy of prematurity			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Abdominal discomfort			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acetonaemic vomiting			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diarrhoea			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Constipation			
subjects affected / exposed	2 / 61 (3.28%)	1 / 54 (1.85%)	4 / 180 (2.22%)
occurrences causally related to treatment / all	0 / 2	0 / 1	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Flatulence			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Enterocolitis			

subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Gastrointestinal haemorrhage</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Haematemesis</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 2	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Gastrooesophageal reflux disease</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Haematochezia</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Ileus</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Inguinal hernia</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Necrotising colitis</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Mechanical ileus</b>			

subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			
subjects affected / exposed	1 / 61 (1.64%)	2 / 54 (3.70%)	6 / 180 (3.33%)
occurrences causally related to treatment / all	0 / 2	0 / 2	0 / 8
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Cholelithiasis			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatic failure			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Portal hypertension			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
Rash			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Erythema multiforme			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rash papular			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urticarial vasculitis			

subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Renal and urinary disorders</b>			
Acute kidney injury			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nephrolithiasis			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Endocrine disorders</b>			
Diabetes insipidus			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Musculoskeletal and connective tissue disorders</b>			
Osteopenia			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Infections and infestations</b>			
Adenovirus infection			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bacterial food poisoning			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bronchiolitis			

subjects affected / exposed	4 / 61 (6.56%)	0 / 54 (0.00%)	7 / 180 (3.89%)
occurrences causally related to treatment / all	0 / 4	0 / 0	0 / 7
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Bronchitis</b>			
subjects affected / exposed	3 / 61 (4.92%)	5 / 54 (9.26%)	11 / 180 (6.11%)
occurrences causally related to treatment / all	0 / 3	0 / 12	0 / 18
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Bronchitis viral</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Cellulitis</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Enterocolitis infectious</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Ear infection</b>			
subjects affected / exposed	1 / 61 (1.64%)	1 / 54 (1.85%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Croup infectious</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Gastroenteritis adenovirus</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Gastroenteritis</b>			

subjects affected / exposed	1 / 61 (1.64%)	2 / 54 (3.70%)	3 / 180 (1.67%)
occurrences causally related to treatment / all	0 / 1	0 / 2	0 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Epstein-Barr virus infection			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis norovirus			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis rotavirus			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Herpangina			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hand-foot-and-mouth disease			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis viral			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Influenza			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Laryngitis			

subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Leptospirosis</b>			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Lower respiratory tract infection</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 3	0 / 0	0 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Meningitis viral</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Otitis media acute</b>			
subjects affected / exposed	1 / 61 (1.64%)	1 / 54 (1.85%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Periorbital cellulitis</b>			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Parainfluenzae viral bronchitis</b>			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Pneumonia</b>			
subjects affected / exposed	2 / 61 (3.28%)	4 / 54 (7.41%)	10 / 180 (5.56%)
occurrences causally related to treatment / all	0 / 2	0 / 7	0 / 15
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
<b>Pharyngitis</b>			

subjects affected / exposed	1 / 61 (1.64%)	1 / 54 (1.85%)	4 / 180 (2.22%)
occurrences causally related to treatment / all	0 / 1	1 / 1	1 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pharyngitis streptococcal			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia aspiration			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia respiratory syncytial viral			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia viral			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyelonephritis			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory syncytial virus infection			
subjects affected / exposed	1 / 61 (1.64%)	1 / 54 (1.85%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory syncytial virus bronchitis			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	2 / 180 (1.11%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory syncytial virus bronchiolitis			

subjects affected / exposed	1 / 61 (1.64%)	2 / 54 (3.70%)	4 / 180 (2.22%)
occurrences causally related to treatment / all	0 / 1	0 / 2	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory tract infection viral			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Septic shock			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 1
Streptococcal infection			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Upper respiratory tract infection			
subjects affected / exposed	2 / 61 (3.28%)	1 / 54 (1.85%)	3 / 180 (1.67%)
occurrences causally related to treatment / all	0 / 2	0 / 1	0 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tonsillitis			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Viral infection			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Viral tonsillitis			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Viral pharyngitis			

subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Viral upper respiratory tract infection			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dehydration			
subjects affected / exposed	3 / 61 (4.92%)	1 / 54 (1.85%)	4 / 180 (2.22%)
occurrences causally related to treatment / all	0 / 3	0 / 1	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Electrolyte imbalance			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Failure to thrive			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypernatraemia			
subjects affected / exposed	0 / 61 (0.00%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hyperphosphatasaemia			
subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypophagia			

subjects affected / exposed	1 / 61 (1.64%)	0 / 54 (0.00%)	1 / 180 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

<b>Serious adverse events</b>	Ranibizumab 0.1 mg		
Total subjects affected by serious adverse events			
subjects affected / exposed	27 / 65 (41.54%)		
number of deaths (all causes)	0		
number of deaths resulting from adverse events	0		
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Haemangioma			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Pregnancy, puerperium and perinatal conditions			
Perinatal brain damage			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Inflammation			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Developmental delay			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Respiratory, thoracic and mediastinal disorders			

Acute respiratory failure				
subjects affected / exposed	2 / 65 (3.08%)			
occurrences causally related to treatment / all	0 / 2			
deaths causally related to treatment / all	0 / 0			
Asthma				
subjects affected / exposed	3 / 65 (4.62%)			
occurrences causally related to treatment / all	0 / 4			
deaths causally related to treatment / all	0 / 0			
Bronchial obstruction				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Bronchopulmonary dysplasia				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Bronchospasm				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Dyspnoea				
subjects affected / exposed	1 / 65 (1.54%)			
occurrences causally related to treatment / all	0 / 2			
deaths causally related to treatment / all	0 / 0			
Cough				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Hypercapnia				
subjects affected / exposed	1 / 65 (1.54%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Hypoxia				

subjects affected / exposed	2 / 65 (3.08%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Nasal discharge discolouration			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Laryngospasm			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Increased bronchial secretion			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Pulmonary vein stenosis			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Pulmonary hypertension			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pneumonitis			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Respiratory distress			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Respiratory failure			

subjects affected / exposed	2 / 65 (3.08%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Tonsillar hypertrophy			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Psychiatric disorders			
Psychomotor retardation			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Investigations			
Oxygen saturation decreased			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Injury, poisoning and procedural complications			
Head injury			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Foreign body in respiratory tract			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Endotracheal intubation complication			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Periorbital haematoma			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		

Congenital, familial and genetic disorders			
Cerebral palsy			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Coarctation of the aorta			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Cryptorchism			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Cardiac disorders			
Cardiac arrest			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Bradycardia			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Nervous system disorders			
Cerebellar haemorrhage			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Febrile convulsion			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences causally related to treatment / all	0 / 3		
deaths causally related to treatment / all	0 / 0		
Epilepsy			

subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Cognitive disorder</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Hydrocephalus</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Intracranial pressure increased</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Nystagmus</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Myoclonus</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Motor dysfunction</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Quadriparesis</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Periventricular leukomalacia</b>			

subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Ear and labyrinth disorders</b>			
Deafness			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Eye disorders</b>			
Conjunctival haemorrhage			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Eye disorder			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Retinal haemorrhage			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Retinal detachment			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Retinopathy of prematurity			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Gastrointestinal disorders</b>			
Abdominal discomfort			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		

Acetonaemic vomiting				
subjects affected / exposed	1 / 65 (1.54%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Diarrhoea				
subjects affected / exposed	1 / 65 (1.54%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Constipation				
subjects affected / exposed	1 / 65 (1.54%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Flatulence				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Enterocolitis				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Gastrointestinal haemorrhage				
subjects affected / exposed	1 / 65 (1.54%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Haematemesis				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Gastrooesophageal reflux disease				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Haematochezia				

subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Ileus</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Inguinal hernia</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Necrotising colitis</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Mechanical ileus</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Vomiting</b>			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences causally related to treatment / all	0 / 4		
deaths causally related to treatment / all	0 / 0		
<b>Hepatobiliary disorders</b>			
<b>Cholelithiasis</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Hepatic failure</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Portal hypertension</b>			

subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Skin and subcutaneous tissue disorders</b>			
<b>Rash</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Erythema multiforme</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Rash papular</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Urticarial vasculitis</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Renal and urinary disorders</b>			
<b>Acute kidney injury</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Nephrolithiasis</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Endocrine disorders</b>			
<b>Diabetes insipidus</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		

Musculoskeletal and connective tissue disorders			
Osteopenia			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Infections and infestations			
Adenovirus infection			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Bacterial food poisoning			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Bronchiolitis			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences causally related to treatment / all	0 / 3		
deaths causally related to treatment / all	0 / 0		
Bronchitis			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences causally related to treatment / all	0 / 3		
deaths causally related to treatment / all	0 / 0		
Bronchitis viral			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Cellulitis			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Enterocolitis infectious			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		

Ear infection				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Croup infectious				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Gastroenteritis adenovirus				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Gastroenteritis				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Epstein-Barr virus infection				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Gastroenteritis norovirus				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Gastroenteritis rotavirus				
subjects affected / exposed	1 / 65 (1.54%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Herpangina				
subjects affected / exposed	0 / 65 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Hand-foot-and-mouth disease				

subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Gastroenteritis viral</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Influenza</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Laryngitis</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Leptospirosis</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Lower respiratory tract infection</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Meningitis viral</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Otitis media acute</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Periorbital cellulitis</b>			

subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Parainfluenzae viral bronchitis			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pneumonia			
subjects affected / exposed	4 / 65 (6.15%)		
occurrences causally related to treatment / all	0 / 6		
deaths causally related to treatment / all	0 / 0		
Pharyngitis			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Pharyngitis streptococcal			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Pneumonia aspiration			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pneumonia respiratory syncytial viral			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Pneumonia viral			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pyelonephritis			

subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Respiratory syncytial virus infection			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Respiratory syncytial virus bronchitis			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Respiratory syncytial virus bronchiolitis			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Respiratory tract infection viral			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Septic shock			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Streptococcal infection			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Upper respiratory tract infection			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Tonsillitis			

subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Viral infection			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Viral tonsillitis			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Viral pharyngitis			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Viral upper respiratory tract infection			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Dehydration			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Electrolyte imbalance			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Failure to thrive			

subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Hypernatraemia</b>			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
<b>Hyperphosphatasaemia</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
<b>Hypophagia</b>			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		

Frequency threshold for reporting non-serious adverse events: 4 %

<b>Non-serious adverse events</b>	Ranibizumab 0.2 mg	Laser	Total
<b>Total subjects affected by non-serious adverse events</b>			
subjects affected / exposed	41 / 61 (67.21%)	40 / 54 (74.07%)	127 / 180 (70.56%)
<b>Nervous system disorders</b>			
<b>Febrile convulsion</b>			
subjects affected / exposed	3 / 61 (4.92%)	3 / 54 (5.56%)	8 / 180 (4.44%)
occurrences (all)	3	4	10
<b>Nystagmus</b>			
subjects affected / exposed	1 / 61 (1.64%)	1 / 54 (1.85%)	5 / 180 (2.78%)
occurrences (all)	1	1	6
<b>Seizure</b>			
subjects affected / exposed	3 / 61 (4.92%)	2 / 54 (3.70%)	5 / 180 (2.78%)
occurrences (all)	4	2	6
<b>General disorders and administration site conditions</b>			
Developmental delay			

subjects affected / exposed occurrences (all)	5 / 61 (8.20%) 5	3 / 54 (5.56%) 3	9 / 180 (5.00%) 9
Pyrexia subjects affected / exposed occurrences (all)	6 / 61 (9.84%) 6	10 / 54 (18.52%) 23	27 / 180 (15.00%) 59
Immune system disorders Seasonal allergy subjects affected / exposed occurrences (all)	3 / 61 (4.92%) 3	1 / 54 (1.85%) 1	6 / 180 (3.33%) 6
Eye disorders Astigmatism subjects affected / exposed occurrences (all)	3 / 61 (4.92%) 3	2 / 54 (3.70%) 2	7 / 180 (3.89%) 7
Myopia subjects affected / exposed occurrences (all)	6 / 61 (9.84%) 6	5 / 54 (9.26%) 6	14 / 180 (7.78%) 15
Strabismus subjects affected / exposed occurrences (all)	10 / 61 (16.39%) 11	12 / 54 (22.22%) 12	34 / 180 (18.89%) 38
Gastrointestinal disorders Vomiting subjects affected / exposed occurrences (all)	0 / 61 (0.00%) 0	2 / 54 (3.70%) 2	5 / 180 (2.78%) 13
Gastrooesophageal reflux disease subjects affected / exposed occurrences (all)	4 / 61 (6.56%) 4	5 / 54 (9.26%) 5	13 / 180 (7.22%) 13
Diarrhoea subjects affected / exposed occurrences (all)	3 / 61 (4.92%) 4	5 / 54 (9.26%) 6	13 / 180 (7.22%) 17
Constipation subjects affected / exposed occurrences (all)	4 / 61 (6.56%) 4	2 / 54 (3.70%) 2	14 / 180 (7.78%) 17
Respiratory, thoracic and mediastinal disorders Bronchopulmonary dysplasia subjects affected / exposed occurrences (all)	2 / 61 (3.28%) 2	4 / 54 (7.41%) 5	9 / 180 (5.00%) 10

Cough subjects affected / exposed occurrences (all)	5 / 61 (8.20%) 7	3 / 54 (5.56%) 6	14 / 180 (7.78%) 25
Dyspnoea subjects affected / exposed occurrences (all)	1 / 61 (1.64%) 1	0 / 54 (0.00%) 0	4 / 180 (2.22%) 10
Rhinorrhoea subjects affected / exposed occurrences (all)	2 / 61 (3.28%) 2	1 / 54 (1.85%) 2	6 / 180 (3.33%) 10
Skin and subcutaneous tissue disorders			
Erythema subjects affected / exposed occurrences (all)	0 / 61 (0.00%) 0	0 / 54 (0.00%) 0	3 / 180 (1.67%) 3
Dry skin subjects affected / exposed occurrences (all)	2 / 61 (3.28%) 2	3 / 54 (5.56%) 3	5 / 180 (2.78%) 5
Rash subjects affected / exposed occurrences (all)	0 / 61 (0.00%) 0	0 / 54 (0.00%) 0	3 / 180 (1.67%) 3
Psychiatric disorders			
Autism spectrum disorder subjects affected / exposed occurrences (all)	3 / 61 (4.92%) 3	1 / 54 (1.85%) 1	5 / 180 (2.78%) 5
Musculoskeletal and connective tissue disorders			
Rickets subjects affected / exposed occurrences (all)	0 / 61 (0.00%) 0	3 / 54 (5.56%) 3	3 / 180 (1.67%) 3
Infections and infestations			
Bronchiolitis subjects affected / exposed occurrences (all)	2 / 61 (3.28%) 4	0 / 54 (0.00%) 0	5 / 180 (2.78%) 7
Bronchitis subjects affected / exposed occurrences (all)	4 / 61 (6.56%) 11	6 / 54 (11.11%) 9	14 / 180 (7.78%) 24
COVID-19			

subjects affected / exposed	1 / 61 (1.64%)	3 / 54 (5.56%)	5 / 180 (2.78%)
occurrences (all)	1	3	6
Conjunctivitis			
subjects affected / exposed	2 / 61 (3.28%)	4 / 54 (7.41%)	10 / 180 (5.56%)
occurrences (all)	2	4	10
Ear infection			
subjects affected / exposed	4 / 61 (6.56%)	0 / 54 (0.00%)	7 / 180 (3.89%)
occurrences (all)	5	0	10
Nasopharyngitis			
subjects affected / exposed	11 / 61 (18.03%)	9 / 54 (16.67%)	31 / 180 (17.22%)
occurrences (all)	25	13	59
Influenza			
subjects affected / exposed	4 / 61 (6.56%)	1 / 54 (1.85%)	7 / 180 (3.89%)
occurrences (all)	4	1	9
Gastroenteritis viral			
subjects affected / exposed	1 / 61 (1.64%)	3 / 54 (5.56%)	5 / 180 (2.78%)
occurrences (all)	1	3	5
Exanthema subitum			
subjects affected / exposed	3 / 61 (4.92%)	1 / 54 (1.85%)	6 / 180 (3.33%)
occurrences (all)	3	1	6
Otitis media			
subjects affected / exposed	7 / 61 (11.48%)	3 / 54 (5.56%)	12 / 180 (6.67%)
occurrences (all)	9	3	14
Respiratory tract infection viral			
subjects affected / exposed	3 / 61 (4.92%)	1 / 54 (1.85%)	8 / 180 (4.44%)
occurrences (all)	5	3	16
Pneumonia			
subjects affected / exposed	5 / 61 (8.20%)	0 / 54 (0.00%)	6 / 180 (3.33%)
occurrences (all)	6	0	7
Pharyngitis			
subjects affected / exposed	2 / 61 (3.28%)	5 / 54 (9.26%)	10 / 180 (5.56%)
occurrences (all)	2	9	15
Upper respiratory tract infection			
subjects affected / exposed	3 / 61 (4.92%)	5 / 54 (9.26%)	11 / 180 (6.11%)
occurrences (all)	3	6	14
Viral upper respiratory tract infection			

subjects affected / exposed occurrences (all)	0 / 61 (0.00%) 0	0 / 54 (0.00%) 0	3 / 180 (1.67%) 5
Metabolism and nutrition disorders			
Iron deficiency			
subjects affected / exposed	4 / 61 (6.56%)	4 / 54 (7.41%)	8 / 180 (4.44%)
occurrences (all)	5	4	9
Malnutrition			
subjects affected / exposed	0 / 61 (0.00%)	1 / 54 (1.85%)	4 / 180 (2.22%)
occurrences (all)	0	1	4

<b>Non-serious adverse events</b>	Ranibizumab 0.1 mg		
Total subjects affected by non-serious adverse events			
subjects affected / exposed	46 / 65 (70.77%)		
Nervous system disorders			
Febrile convulsion			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences (all)	3		
Nystagmus			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences (all)	4		
Seizure			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences (all)	0		
General disorders and administration site conditions			
Developmental delay			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences (all)	1		
Pyrexia			
subjects affected / exposed	11 / 65 (16.92%)		
occurrences (all)	30		
Immune system disorders			
Seasonal allergy			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences (all)	2		
Eye disorders			
Astigmatism			

subjects affected / exposed occurrences (all)	2 / 65 (3.08%) 2		
Myopia subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 3		
Strabismus subjects affected / exposed occurrences (all)	12 / 65 (18.46%) 15		
Gastrointestinal disorders Vomiting subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 11		
Gastrooesophageal reflux disease subjects affected / exposed occurrences (all)	4 / 65 (6.15%) 4		
Diarrhoea subjects affected / exposed occurrences (all)	5 / 65 (7.69%) 7		
Constipation subjects affected / exposed occurrences (all)	8 / 65 (12.31%) 11		
Respiratory, thoracic and mediastinal disorders Bronchopulmonary dysplasia subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 3		
Cough subjects affected / exposed occurrences (all)	6 / 65 (9.23%) 12		
Dyspnoea subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 9		
Rhinorrhoea subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 6		
Skin and subcutaneous tissue disorders			

Erythema subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 3		
Dry skin subjects affected / exposed occurrences (all)	0 / 65 (0.00%) 0		
Rash subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 3		
Psychiatric disorders Autism spectrum disorder subjects affected / exposed occurrences (all)	1 / 65 (1.54%) 1		
Musculoskeletal and connective tissue disorders Rickets subjects affected / exposed occurrences (all)	0 / 65 (0.00%) 0		
Infections and infestations Bronchiolitis subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 3		
Bronchitis subjects affected / exposed occurrences (all)	4 / 65 (6.15%) 4		
COVID-19 subjects affected / exposed occurrences (all)	1 / 65 (1.54%) 2		
Conjunctivitis subjects affected / exposed occurrences (all)	4 / 65 (6.15%) 4		
Ear infection subjects affected / exposed occurrences (all)	3 / 65 (4.62%) 5		
Nasopharyngitis subjects affected / exposed occurrences (all)	11 / 65 (16.92%) 21		

Influenza			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences (all)	4		
Gastroenteritis viral			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences (all)	1		
Exanthema subitum			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences (all)	2		
Otitis media			
subjects affected / exposed	2 / 65 (3.08%)		
occurrences (all)	2		
Respiratory tract infection viral			
subjects affected / exposed	4 / 65 (6.15%)		
occurrences (all)	8		
Pneumonia			
subjects affected / exposed	1 / 65 (1.54%)		
occurrences (all)	1		
Pharyngitis			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences (all)	4		
Upper respiratory tract infection			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences (all)	5		
Viral upper respiratory tract infection			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences (all)	5		
Metabolism and nutrition disorders			
Iron deficiency			
subjects affected / exposed	0 / 65 (0.00%)		
occurrences (all)	0		
Malnutrition			
subjects affected / exposed	3 / 65 (4.62%)		
occurrences (all)	3		

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
31 July 2018	Introduced the collection of additional visual function data following EU national scientific advice meeting
10 July 2019	Included the requirement of masked VA assessments to be performed at the 5th birthday visit, following a request from the EMA.

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

Due to EudraCT system limitations, which EMA is aware of, data using 999 as data points in this record are not an accurate representation of the clinical trial results. Please use <https://www.novctrd.com/CtrdWeb/home.nov> for complete trial results.

Notes: