

SSAT 066: Statistical Analyses Results

Version 6.0: 22nd January 2018

Beta-2-microglobulin (urine)

ug/L	Total=60					
	Median (IQR)					
	Baseline	Week 4	Week 12	Week 24	Week 36	Week 48
	Arm 1: Truvada® (200mg/245mg film coated tablets) one tablet once daily + Raltegravir 400mg (one tablet) twice daily N=20					
Beta-2-microglobulin (urine)	N=16 101(63 to 399) Range: 40 to 653	N=16 137(74 to 218) Range: 40 to 386	N=11 98(40 to 310) Range: 40 to 467	N=14 131(40 to 302) Range: 40 to 1007	-	N=10 105(40 to 269) Range: 40 to 312
	Arm 2: Truvada® (200mg/245mg film coated tablets) one tablet once daily + Dolutegravir 50mg once daily N=20					
Beta-2-microglobulin (urine)	N=17 117(96 to 228) Range: 40 to 548	N=10 115(56 to 280) Range: 40 to 748	N=14 146(84 to 255) Range: 40 to 1618	N=12 114(46 to 242) Range: 40 to 3518	-	N=12 145(90 to 307) Range: 40 to 374
	Arm 3: Stribild®150mg/150mg/200mg/245mg tablet once daily N=20					
Beta-2-microglobulin (urine)	N=13 72(58 to 103) Range: 40 to 424	N=12 101(50 to 139) Range: 40 to 640	N=14 105(76 to 146) Range: 40 to 2747	N=11 172(40 to 561) Range: 40 to 5905	N=2 210(193 to 226) Range: 193 to 226	N=13 130(64 to 187) Range: 40 to 1188