

SSAT 066: Statistical Analyses Results

Version 6.0: 22nd January 2018

The changes in inflammatory markers

mg/L	High sensitivity CRP					
	Total=60					
	Median (IQR)					
	Baseline	Week 4	Week 12	Week 24	Week 36	Week 48
Arm 1: Truvada® (200mg/245mg film coated tablets) one tablet once daily + Raltegravir 400mg (one tablet) twice daily N=20						
hsCRP	N=20 1.2(0.8 to 1.2) Range: 0.4 to 13.4	N=19 1.4(0.8 to 2.8) Range: 0.3 to 6.5	N=19 1.0(0.7 to 2.4) Range: 0.3 to 52.4	N=19 0.9(0.6 to 3.1) Range: 0.3 to 31.1	-	N=18 0.9(0.6 to 1.4) Range: 0.3 to 8.3
Arm 2: Truvada® (200mg/245mg film coated tablets) one tablet once daily + Dolutegravir 50mg once daily N=20						
hsCRP	N=20 1.2(0.8 to 2.1) Range: 0.4 to 5.8	N=20 0.9(0.6 to 1.6) Range: 0.3 to 4.4	N=20 1.9(1.1 to 3.6) Range: 0.4 to 10.8	N=20 1.5(0.9 to 3.2) Range: 0.3 to 19.1	N=2 1.6, 3.6	N=19 1.2(0.8 to 2.6) Range: 0.4 to 12.5
Arm 3: Stribild®150mg/150mg/200mg/245mg tablet once daily N=20						
hsCRP	N=20 1.2(0.7 to 2.9) Range: 0.3 to 6.5	N=20 1.5(0.7 to 3.2) Range: 0.3 to 11.9	N=20 1.9(0.6 to 3.6) Range: 0.3 to 26.9	N=20 1.7(0.7 to 3.9) Range: 0.3 to 8.3	N=2 0.4, 0.5	N=20 1.4(0.7 to 1.1) Range: 0.3 to 42.3