



## Clinical trial results:

### Efficacy and safety of semaglutide versus dulaglutide as add-on to metformin in subjects with type 2 diabetes

#### Summary

EudraCT number	2014-005375-91
Trial protocol	IE FI DE GB SK LV LT GR ES PT HR
Global end of trial date	19 May 2017

#### Results information

Result version number	v1 (current)
This version publication date	30 May 2018
First version publication date	30 May 2018

#### Trial information

##### Trial identification

Sponsor protocol code	NN9535-4216
-----------------------	-------------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02648204
WHO universal trial number (UTN)	U1111-1164-8495

Notes:

#### Sponsors

Sponsor organisation name	Novo Nordisk A/S
Sponsor organisation address	Novo Allé, Bagsvaerd, Denmark, 2880
Public contact	Clinical Reporting Anchor and Disclosure (1452), Novo Nordisk A/S, clinicaltrials@novonordisk.com
Scientific contact	Clinical Reporting Anchor and Disclosure (1452), Novo Nordisk A/S, clinicaltrials@novonordisk.com

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:



## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	30 November 2017
Is this the analysis of the primary completion data?	Yes
Primary completion date	10 April 2017
Global end of trial reached?	Yes
Global end of trial date	19 May 2017
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

o compare the effect of once-weekly dosing of two dose levels of subcutaneous semaglutide (0.5 mg and 1.0 mg) versus once-weekly dosing of two dose levels of subcutaneous dulaglutide (0.75 mg and 1.5 mg) on glycaemic control in subjects with type 2 diabetes on a background treatment with metformin.

Protection of trial subjects:

The trial was conducted in accordance with the Declaration of Helsinki, ICH Good Clinical Practice, EN ISO 14155 Parts 1 and 2 and FDA 21 CFR 312.120.

Background therapy:

Subjects were to continue metformin at pre-trial dose and frequency during the whole treatment period unless rescue medication was needed.

Evidence for comparator:

Not applicable

Actual start date of recruitment	06 January 2016
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Bulgaria: 60
Country: Number of subjects enrolled	Germany: 58
Country: Number of subjects enrolled	Spain: 63
Country: Number of subjects enrolled	Finland: 31
Country: Number of subjects enrolled	United Kingdom: 70
Country: Number of subjects enrolled	Greece: 73
Country: Number of subjects enrolled	Hong Kong: 30
Country: Number of subjects enrolled	Croatia: 27
Country: Number of subjects enrolled	India: 130
Country: Number of subjects enrolled	Ireland: 22
Country: Number of subjects enrolled	Lithuania: 40
Country: Number of subjects enrolled	Latvia: 35
Country: Number of subjects enrolled	Portugal: 14
Country: Number of subjects enrolled	Romania: 65
Country: Number of subjects enrolled	Slovakia: 40
Country: Number of subjects enrolled	United States: 441



Worldwide total number of subjects	1199
EEA total number of subjects	598

Notes:

<b>Subjects enrolled per age group</b>	
In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	939
From 65 to 84 years	260
85 years and over	0



## Subject disposition

### Recruitment

Recruitment details:

A total of 210 sites were approved for recruiting subjects, of which 196 sites randomized the subjects: Bulgaria: 6; Croatia: 6; Finland: 6; Germany: 6; Greece: 8; Hong Kong: 1; India: 22; Ireland: 5; Latvia: 3; Lithuania: 5; Portugal: 4; Romania: 7; Slovakia: 6; Spain: 8; United Kingdom: 8; United States: 95.

### Pre-assignment

Screening details:

A total of 1201 subjects were randomized in the trial: Semaglutide 0.5 mg-301 subjects, Semaglutide 1.0 mg-300 subjects, Dulaglutide 0.75 mg-300 subjects, Dulaglutide 1.5 mg-300 subjects. Of the 1201 subjects, 1199 were exposed to trial products and 2 randomised subjects withdrew prior to exposure (1 subject in each Dulaglutide arm).

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Not blinded

Blinding implementation details:

Not applicable

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Semaglutide 0.5 mg

Arm description:

Subjects received subcutaneous (s.c., under the skin) injections of semaglutide once weekly (OW) for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for 36 weeks (weeks 5 to 40). Subjects were followed for 5 weeks after completion of treatment period.

Arm type	Experimental
Investigational medicinal product name	Semaglutide B 1.34mg/ml PDS290
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection in pre-filled pen
Routes of administration	Subcutaneous use

Dosage and administration details:

Semaglutide solution for injection in pre-filled PDS290 pen-injector, was administered by s.c. injection either in the thigh, abdomen or upper arm, at any time of day irrespective of meals. The injections were to be administered on the same day of the week during the trial.

<b>Arm title</b>	Semaglutide 1.0 mg
------------------	--------------------

Arm description:

Subjects received s.c. injections of semaglutide OW for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for another 4 weeks (weeks 5 to 8) and then semaglutide 1.0 mg for remaining 32 weeks (weeks 9-40). Subjects were followed for 5 weeks after completion of treatment period.

Arm type	Experimental
Investigational medicinal product name	Semaglutide B 1.34mg/ml PDS290
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection in pre-filled pen
Routes of administration	Subcutaneous use

Dosage and administration details:

Semaglutide solution for injection in pre-filled PDS290 pen-injector, was administered by s.c. injection



either in the thigh, abdomen or upper arm, at any time of day irrespective of meals. The injections were to be administered on the same day of the week during the trial.

<b>Arm title</b>	Dulaglutide 0.75 mg
------------------	---------------------

Arm description:

Subjects received s.c. injections of dulaglutide 0.75 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.

Arm type	Active comparator
Investigational medicinal product name	Dulaglutide
Investigational medicinal product code	
Other name	Trulicity®
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Dulaglutide solution for injection in a pre-filled pen was administered by s.c. injection either in the thigh, abdomen or upper arm, at any time of day irrespective of meals. The injections were to be administered on the same day of the week during the trial.

<b>Arm title</b>	Dulaglutide 1.5 mg
------------------	--------------------

Arm description:

Subjects received s.c. injections of dulaglutide 1.5 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.

Arm type	Active comparator
Investigational medicinal product name	Dulaglutide
Investigational medicinal product code	
Other name	Trulicity®
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Dulaglutide solution for injection in a pre-filled pen was administered by s.c. injection either in the thigh, abdomen or upper arm, at any time of day irrespective of meals. The injections were to be administered on the same day of the week during the trial.

<b>Number of subjects in period 1</b>	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg
Started	301	300	299
Completed	279	279	287
Not completed	22	21	12
Adverse event, serious fatal	1	1	2
Consent withdrawn by subject	12	8	1
Adverse event, non-fatal	-	-	1
Lost to follow-up	5	8	4
Missing follow-up information	4	4	4

<b>Number of subjects in period 1</b>	Dulaglutide 1.5 mg
---------------------------------------	--------------------



Started	299
Completed	284
Not completed	15
Adverse event, serious fatal	2
Consent withdrawn by subject	5
Adverse event, non-fatal	-
Lost to follow-up	6
Missing follow-up information	2



## Baseline characteristics

### Reporting groups

Reporting group title	Semaglutide 0.5 mg
-----------------------	--------------------

Reporting group description:

Subjects received subcutaneous (s.c., under the skin) injections of semaglutide once weekly (OW) for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for 36 weeks (weeks 5 to 40). Subjects were followed for 5 weeks after completion of treatment period.

Reporting group title	Semaglutide 1.0 mg
-----------------------	--------------------

Reporting group description:

Subjects received s.c. injections of semaglutide OW for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for another 4 weeks (weeks 5 to 8) and then semaglutide 1.0 mg for remaining 32 weeks (weeks 9-40). Subjects were followed for 5 weeks after completion of treatment period.

Reporting group title	Dulaglutide 0.75 mg
-----------------------	---------------------

Reporting group description:

Subjects received s.c. injections of dulaglutide 0.75 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.

Reporting group title	Dulaglutide 1.5 mg
-----------------------	--------------------

Reporting group description:

Subjects received s.c. injections of dulaglutide 1.5 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.

Reporting group values	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg
Number of subjects	301	300	299
Age Categorical Units: Subjects			
Adults (18-64 years)	222	247	238
From 65-74 years	67	45	53
75-84 years	12	8	8
Age Continuous Units: years			
arithmetic mean	56	55	55
standard deviation	± 10.9	± 10.6	± 10.4
Gender Categorical Units: Subjects			
Female	169	162	160
Male	132	138	139
Race Units: Subjects			
American Indian or Alaska Native	0	0	0
Asian	50	38	48
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	17	18	17
White	233	243	232
More than one race	0	0	0
Unknown or Not Reported	1	1	2
Ethnicity Units: Subjects			



Hispanic or Latino	29	35	31
Not Hispanic or Latino	272	265	268

Glycosylated haemoglobin (Hb1Ac) Units: percentage of HbA1c arithmetic mean standard deviation	8.3 ± 0.96	8.2 ± 0.92	8.2 ± 0.91
Body weight Units: kg arithmetic mean standard deviation	96.4 ± 24.38	95.5 ± 20.90	95.6 ± 23.01
Fasting plasma glucose			
Number of subject analysed=298, 299, 297, 297 for semaglutide 0.5 mg, semaglutide 1.0 mg, dulaglutide 0.75 mg and dulaglutide 1.5 mg, respectively.			
Units: mmol/L arithmetic mean standard deviation	9.8 ± 2.54	9.8 ± 2.58	9.7 ± 2.65

Reporting group values	Dulaglutide 1.5 mg	Total	
Number of subjects	299	1199	
Age Categorical Units: Subjects			
Adults (18-64 years)	232	939	
From 65-74 years	60	225	
75-84 years	7	35	
Age Continuous Units: years arithmetic mean standard deviation	56 ± 10.6	-	
Gender Categorical Units: Subjects			
Female	171	662	
Male	128	537	
Race Units: Subjects			
American Indian or Alaska Native	0	0	
Asian	55	191	
Native Hawaiian or Other Pacific Islander	0	0	
Black or African American	18	70	
White	220	928	
More than one race	0	0	
Unknown or Not Reported	6	10	
Ethnicity Units: Subjects			
Hispanic or Latino	43	138	
Not Hispanic or Latino	256	1061	
Glycosylated haemoglobin (Hb1Ac) Units: percentage of HbA1c arithmetic mean standard deviation	8.2 ± 0.89	-	
Body weight			



Units: kg			
arithmetic mean	93.4		
standard deviation	± 21.79	-	
Fasting plasma glucose			
Number of subject analysed=298, 299, 297, 297 for semaglutide 0.5 mg, semaglutide 1.0 mg, dulaglutide 0.75 mg and dulaglutide 1.5 mg, respectively.			
Units: mmol/L			
arithmetic mean	9.6		
standard deviation	± 2.29	-	



## End points

### End points reporting groups

Reporting group title	Semaglutide 0.5 mg
Reporting group description: Subjects received subcutaneous (s.c., under the skin) injections of semaglutide once weekly (OW) for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for 36 weeks (weeks 5 to 40). Subjects were followed for 5 weeks after completion of treatment period.	
Reporting group title	Semaglutide 1.0 mg
Reporting group description: Subjects received s.c. injections of semaglutide OW for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for another 4 weeks (weeks 5 to 8) and then semaglutide 1.0 mg for remaining 32 weeks (weeks 9-40). Subjects were followed for 5 weeks after completion of treatment period.	
Reporting group title	Dulaglutide 0.75 mg
Reporting group description: Subjects received s.c. injections of dulaglutide 0.75 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.	
Reporting group title	Dulaglutide 1.5 mg
Reporting group description: Subjects received s.c. injections of dulaglutide 1.5 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.	

### Primary: Change in HbA1c

End point title	Change in HbA1c
End point description: Results are based on HbA1c data from on-treatment without rescue medication observation period. The 'on-treatment' observation period was the period where the subject was considered to be exposed to trial product. The 'on-treatment without rescue medication' observation period was a subset of the 'on-treatment' observation period, where subjects did not receive any non-investigational antidiabetic medication (rescue medication). This period includes the observations recorded at, or after the date of first dose of trial product and not after the first occurrence of the following: the end-date of the 'on-treatment' observation period or initiation of rescue medication. The post-baseline responses are analysed using a mixed model for repeated measurements with treatment and country as fixed factors and baseline value as covariate, all nested within visit. Analysis was based on FAS (all randomised subjects exposed to at least one dose of trial product).	
End point type	Primary
End point timeframe: From baseline to week 40	

End point values	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg	Dulaglutide 1.5 mg
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	301	300	299	299
Units: percentage of HbA1c				
least squares mean (standard error)	-1.51 (± 0.06)	-1.78 (± 0.06)	-1.11 (± 0.05)	-1.37 (± 0.06)



## Statistical analyses

<b>Statistical analysis title</b>	Semaglutide 0.5 mg vs. Dulaglutide 0.75 mg
Comparison groups	Dulaglutide 0.75 mg v Semaglutide 0.5 mg
Number of subjects included in analysis	600
Analysis specification	Pre-specified
Analysis type	non-inferiority <sup>[1]</sup>
P-value	< 0.0001
Method	Mixed models analysis
Parameter estimate	Treatment difference
Point estimate	-0.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.55
upper limit	-0.25
Variability estimate	Standard error of the mean
Dispersion value	0.08

Notes:

[1] - Non-Inferiority margin: 0.4

<b>Statistical analysis title</b>	Semaglutide 0.5 mg vs. Dulaglutide 0.75 mg
Comparison groups	Semaglutide 0.5 mg v Dulaglutide 0.75 mg
Number of subjects included in analysis	600
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.0001
Method	Mixed models analysis
Parameter estimate	Treatment difference
Point estimate	-0.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.55
upper limit	-0.25
Variability estimate	Standard error of the mean
Dispersion value	0.08

<b>Statistical analysis title</b>	Semaglutide 1.0 mg vs. Dulaglutide 1.5 mg
Comparison groups	Semaglutide 1.0 mg v Dulaglutide 1.5 mg
Number of subjects included in analysis	599
Analysis specification	Pre-specified
Analysis type	non-inferiority <sup>[2]</sup>
P-value	< 0.0001
Method	Mixed models analysis
Parameter estimate	Treatment difference
Point estimate	-0.41



Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.57
upper limit	-0.25
Variability estimate	Standard error of the mean
Dispersion value	0.08

Notes:

[2] - Non-Inferiority margin: 0.04

<b>Statistical analysis title</b>	Semaglutide 1.0 mg vs. Dulaglutide 1.5 mg
Comparison groups	Semaglutide 1.0 mg v Dulaglutide 1.5 mg
Number of subjects included in analysis	599
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.0001
Method	Mixed models analysis
Parameter estimate	Treatment difference
Point estimate	-0.41
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.57
upper limit	-0.25
Variability estimate	Standard error of the mean
Dispersion value	0.08

## Secondary: Change in body weight (kg)

End point title	Change in body weight (kg)
End point description:	
Results are based on body weight data from on-treatment without rescue medication observation period. The 'on-treatment' observation period was the period where the subject was considered to be exposed to trial product. The 'on-treatment without rescue medication' observation period was a subset of the 'on-treatment' observation period, where subjects did not receive any non-investigational antidiabetic medication (rescue medication). This period includes the observations recorded at, or after the date of first dose of trial product and not after the first occurrence of the following: the end-date of the 'on-treatment' observation period or initiation of rescue medication. Analysis was based on FAS. Number of subjects analysed=number of subjects with available data for body weight.	
End point type	Secondary
End point timeframe:	
From baseline to week 40	



<b>End point values</b>	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg	Dulaglutide 1.5 mg
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	301	300	299	298
Units: kg				
least squares mean (standard error)	-4.56 ( $\pm$ 0.28)	-6.53 ( $\pm$ 0.28)	-2.30 ( $\pm$ 0.27)	-2.98 ( $\pm$ 0.27)

## Statistical analyses

<b>Statistical analysis title</b>	Semaglutide 1.0 mg vs. Dulaglutide 1.5 mg
Comparison groups	Semaglutide 1.0 mg v Dulaglutide 1.5 mg
Number of subjects included in analysis	598
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.0001
Method	Mixed models analysis
Parameter estimate	Treatment difference
Point estimate	-3.55
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.32
upper limit	-2.78
Variability estimate	Standard error of the mean
Dispersion value	0.39

<b>Statistical analysis title</b>	Semaglutide 0.5 mg vs. Dulaglutide 0.75 mg
Comparison groups	Semaglutide 0.5 mg v Dulaglutide 0.75 mg
Number of subjects included in analysis	600
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.0001
Method	Mixed models analysis
Parameter estimate	Treatment difference
Point estimate	-2.26
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.02
upper limit	-1.51
Variability estimate	Standard error of the mean
Dispersion value	0.39

## Secondary: Change in fasting plasma glucose



End point title	Change in fasting plasma glucose
End point description:	
Results are based on fasting plasma glucose data from on-treatment without rescue medication observation period. The 'on-treatment' observation period was the period where the subject was considered to be exposed to trial product. The 'on-treatment without rescue medication' observation period was a subset of the 'on -treatment' observation period, where subjects did not receive any non-investigational antidiabetic medication (rescue medication). This period includes the observations recorded at, or after the date of first dose of trial product and not after the first occurrence of the following: the end-date of the 'on-treatment' observation period or initiation of rescue medication. Analysis was based on FAS. Number of participants analysed=number of participants with available data for fasting plasma glucose.	
End point type	Secondary
End point timeframe:	
From baseline to week 40	

End point values	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg	Dulaglutide 1.5 mg
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	298	299	297	297
Units: mmol/L				
least squares mean (standard error)	-2.18 (± 0.12)	-2.83 (± 0.12)	-1.87 (± 0.12)	-2.25 (± 0.12)

## Statistical analyses

No statistical analyses for this end point

## Secondary: Change in systolic and diastolic blood pressure

End point title	Change in systolic and diastolic blood pressure
End point description:	
Results are based on systolic and diastolic blood pressure data from on-treatment without rescue medication observation period. The 'on-treatment' observation period was the period where the subject was considered to be exposed to trial product. The 'on-treatment without rescue medication' observation period was a subset of the 'on -treatment' observation period, where subjects did not receive any non-investigational antidiabetic medication (rescue medication). This period includes the observations recorded at, or after the date of first dose of trial product and not after the first occurrence of the following: the end-date of the 'on-treatment' observation period or initiation of rescue medication. Analysis was based on FAS.	
End point type	Secondary
End point timeframe:	
From baseline to week 40	

End point values	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg	Dulaglutide 1.5 mg
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	301	300	299	299
Units: mmHg				
least squares mean (standard error)				
Diastolic BP	-0.57 (± 0.48)	-2.05 (± 0.49)	-0.35 (± 0.47)	-0.03 (± 0.47)



Systolic BP	-2.44 (± 0.76)	-4.88 (± 0.77)	-2.16 (± 0.75)	-2.86 (± 0.75)
-------------	----------------	----------------	----------------	----------------

## Statistical analyses

No statistical analyses for this end point

## Secondary: Overall scores for patient reported outcomes: Diabetes Treatment Satisfaction Questionnaire

End point title	Overall scores for patient reported outcomes: Diabetes Treatment Satisfaction Questionnaire
End point description: The questionnaire contains 8 items and evaluates subjects' diabetes treatment in terms of convenience, flexibility and general feelings towards treatment. The result presented is 'Treatment Satisfaction' summary score (sum of 6 of the 8 items). Response options: 6 (best case) to 0 (worst case). Total scores range: 0-36. Higher scores=higher satisfaction. Results are based on data from on-treatment without rescue medication observation period. The 'on-treatment' observation period was period where subject was considered to be exposed to trial product. The 'on-treatment without rescue medication' observation period was a subset of the 'on -treatment' observation period, where subjects did not receive any non-investigational antidiabetic medication (rescue medication). This includes observations recorded at, or after the date of first dose of trial product and not after first occurrence of following: end-date of the 'on-treatment' observation period or initiation of rescue medication	
End point type	Secondary
End point timeframe: From baseline to week 40	

End point values	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg	Dulaglutide 1.5 mg
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	301 <sup>[3]</sup>	299 <sup>[4]</sup>	299 <sup>[5]</sup>	298 <sup>[6]</sup>
Units: units on a scale				
least squares mean (standard error)	4.60 (± 0.28)	4.55 (± 0.29)	4.52 (± 0.28)	4.65 (± 0.28)

Notes:

[3] - Number of participants with available data for diabetes treatment satisfaction questionnaire.

[4] - Number of participants with available data for diabetes treatment satisfaction questionnaire

[5] - Number of participants with available data for diabetes treatment satisfaction questionnaire

[6] - Number of participants with available data for diabetes treatment satisfaction questionnaire

## Statistical analyses

No statistical analyses for this end point

## Secondary: Subjects who achieve (yes/no) HbA1c ≤6.5% (48 mmol/mol) American Association of Clinical Endocrinologists target

End point title	Subjects who achieve (yes/no) HbA1c ≤6.5% (48 mmol/mol) American Association of Clinical Endocrinologists target
End point description: Percentage of subjects who achieved HbA1c target below or equal to 6.5% (48 mmol/mol) after 40 weeks of treatment. Results are based on data from on-treatment without rescue medication period. The 'on-treatment' observation period was the period where the subject was considered to be exposed	



to trial product. The 'on-treatment without rescue medication' observation period was a subset of the 'on-treatment' observation period, where subjects did not receive any non-investigational antidiabetic medication (rescue medication). This period includes the observations recorded at, or after the date of first dose of trial product and not after the first occurrence of the following: the end-date of the 'on-treatment' observation period or initiation of rescue medication. Missing data imputed from a mixed model for repeated measurements with treatment and country as fixed factors and baseline value as covariate, all nested within visit. Results are based on the FAS.

End point type	Secondary
End point timeframe:	
After 40 weeks treatment	

End point values	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg	Dulaglutide 1.5 mg
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	301	300	299	299
Units: percentage of subjects				
number (not applicable)				
Yes	49.2	66.7	34.1	47.2
No	50.8	33.3	65.9	52.8

### Statistical analyses

No statistical analyses for this end point



## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Baseline (week 0) to week 40 + follow up of 5 weeks (including a visit window of +7 days)

Adverse event reporting additional description:

A TEAE was defined as an AE with onset in the 'on-treatment' period (information collected while subjects were considered as exposed to trial product). This corresponded to information collected from first administration of trial product until the follow-up (5 weeks after the last treatment including a visit window of +7 days)

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	19
--------------------	----

### Reporting groups

Reporting group title	Semaglutide 0.5 mg
-----------------------	--------------------

Reporting group description:

Subjects received s.c. injections of semaglutide OW for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for 36 weeks (weeks 5 to 40).

Reporting group title	Semaglutide 1.0 mg
-----------------------	--------------------

Reporting group description:

Subjects received s.c. injections of semaglutide OW for treatment duration of 40 weeks. Subjects received semaglutide 0.25 mg for 4 weeks (weeks 1 to 4) followed by 0.5 mg for another 4 weeks (weeks 5 to 8) and then semaglutide 1.0 mg for remaining 32 weeks (weeks 9-40). Subjects were followed for 5 weeks after completion of treatment period.

Reporting group title	Dulaglutide 0.75 mg
-----------------------	---------------------

Reporting group description:

Subjects received s.c. injections of dulaglutide 0.75 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.

Reporting group title	Dulaglutide 1.5 mg
-----------------------	--------------------

Reporting group description:

Subjects received s.c. injections of dulaglutide 1.5 mg OW for treatment duration of 40 weeks. Subjects were followed for 5 weeks after completion of treatment period.

Serious adverse events	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg
Total subjects affected by serious adverse events			
subjects affected / exposed	17 / 301 (5.65%)	23 / 300 (7.67%)	24 / 299 (8.03%)
number of deaths (all causes)	0	1	2
number of deaths resulting from adverse events	0	0	0
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Bladder cancer			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Lipoma			



subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Malignant neoplasm of ampulla of Vater			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metastatic squamous cell carcinoma			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pancreatic carcinoma stage IV			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 0
Rectal adenocarcinoma			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rectal cancer			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Hypertensive crisis			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Surgical and medical procedures			
Cardioversion			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0



Gastrectomy			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hip arthroplasty			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hospitalisation			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hysterectomy			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Orchidectomy			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Drowning			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Multiple organ dysfunction syndrome			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Non-cardiac chest pain			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0



Immune system disorders			
Hypersensitivity			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sarcoidosis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders			
Cystocele			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Uterine prolapse			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Acute respiratory failure			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Chronic obstructive pulmonary disease			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Epistaxis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pulmonary embolism			



subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Anxiety			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Depression			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicide attempt			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Lipase increased			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Contusion			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Face injury			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Fall			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0



Femur fracture			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Meniscus injury			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Post procedural myocardial infarction			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Road traffic accident			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tibia fracture			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders			
Angina pectoris			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Angina unstable			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Aortic valve stenosis			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Atrial fibrillation			



subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Atrial flutter			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Atrioventricular block complete			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardio-respiratory arrest			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Congestive cardiomyopathy			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Coronary artery stenosis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Myocardial infarction			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ventricular extrasystoles			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Carpal tunnel syndrome			



subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	2 / 299 (0.67%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cerebral infarction			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diabetic neuropathy			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypoglycaemic unconsciousness			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Paraesthesia			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ear and labyrinth disorders			
Acute vestibular syndrome			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Colitis			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Crohn's disease			



subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroduodenitis			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Haemorrhoids			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Impaired gastric emptying			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Inguinal hernia			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nausea			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pancreatitis acute			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pancreatolithiasis			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Rectal prolapse			



subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Cholecystitis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cholecystitis acute			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cholelithiasis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gallbladder pain			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatic cirrhosis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Nephrolithiasis			
subjects affected / exposed	1 / 301 (0.33%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal colic			



subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ureterolithiasis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	2 / 299 (0.67%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Intervertebral disc degeneration			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Intervertebral disc protrusion			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal chest pain			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Spinal column stenosis			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Appendicitis			
subjects affected / exposed	2 / 301 (0.66%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0



Bronchitis			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cellulitis			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis viral			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
HIV infection			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Kidney infection			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Perirectal abscess			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Postoperative wound infection			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyelonephritis			



subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Upper respiratory tract infection			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary tract infection			
subjects affected / exposed	1 / 301 (0.33%)	0 / 300 (0.00%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Dehydration			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diabetic complication			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
Hypercalcaemia			
subjects affected / exposed	0 / 301 (0.00%)	0 / 300 (0.00%)	1 / 299 (0.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hypoglycaemia			
subjects affected / exposed	0 / 301 (0.00%)	1 / 300 (0.33%)	0 / 299 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

<b>Serious adverse events</b>	Dulaglutide 1.5 mg		
-------------------------------	--------------------	--	--



Total subjects affected by serious adverse events			
subjects affected / exposed	22 / 299 (7.36%)		
number of deaths (all causes)	2		
number of deaths resulting from adverse events	0		
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Bladder cancer			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Lipoma			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Malignant neoplasm of ampulla of Vater			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Metastatic squamous cell carcinoma			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pancreatic carcinoma stage IV			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Rectal adenocarcinoma			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Rectal cancer			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Vascular disorders			



Hypertensive crisis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Surgical and medical procedures			
Cardioversion			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Gastrectomy			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hip arthroplasty			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hospitalisation			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hysterectomy			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Orchidectomy			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
General disorders and administration site conditions			
Drowning			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 1		



Multiple organ dysfunction syndrome			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 1		
Non-cardiac chest pain			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Immune system disorders			
Hypersensitivity			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Sarcoidosis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Reproductive system and breast disorders			
Cystocele			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Uterine prolapse			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Respiratory, thoracic and mediastinal disorders			
Acute respiratory failure			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Chronic obstructive pulmonary disease			



subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Epistaxis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pulmonary embolism			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Psychiatric disorders			
Anxiety			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Depression			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Suicide attempt			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Investigations			
Lipase increased			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Injury, poisoning and procedural complications			
Contusion			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		



Face injury			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Fall			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Femur fracture			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Meniscus injury			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Post procedural myocardial infarction			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Road traffic accident			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Tibia fracture			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Cardiac disorders			
Angina pectoris			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Angina unstable			



subjects affected / exposed	2 / 299 (0.67%)			
occurrences causally related to treatment / all	0 / 2			
deaths causally related to treatment / all	0 / 0			
Aortic valve stenosis				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Atrial fibrillation				
subjects affected / exposed	2 / 299 (0.67%)			
occurrences causally related to treatment / all	0 / 2			
deaths causally related to treatment / all	0 / 0			
Atrial flutter				
subjects affected / exposed	1 / 299 (0.33%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Atrioventricular block complete				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Cardio-respiratory arrest				
subjects affected / exposed	1 / 299 (0.33%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 1			
Congestive cardiomyopathy				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Coronary artery stenosis				
subjects affected / exposed	1 / 299 (0.33%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Myocardial infarction				



subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Ventricular extrasystoles			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Nervous system disorders			
Carpal tunnel syndrome			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Cerebral infarction			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Diabetic neuropathy			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypoglycaemic unconsciousness			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Paraesthesia			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Ear and labyrinth disorders			
Acute vestibular syndrome			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Gastrointestinal disorders			



Abdominal pain				
subjects affected / exposed	1 / 299 (0.33%)			
occurrences causally related to treatment / all	1 / 1			
deaths causally related to treatment / all	0 / 0			
Colitis				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Crohn's disease				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Gastroduodenitis				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Haemorrhoids				
subjects affected / exposed	1 / 299 (0.33%)			
occurrences causally related to treatment / all	0 / 1			
deaths causally related to treatment / all	0 / 0			
Impaired gastric emptying				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Inguinal hernia				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Nausea				
subjects affected / exposed	0 / 299 (0.00%)			
occurrences causally related to treatment / all	0 / 0			
deaths causally related to treatment / all	0 / 0			
Pancreatitis acute				



subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pancreatolithiasis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Rectal prolapse			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Vomiting			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hepatobiliary disorders			
Cholecystitis			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Cholecystitis acute			
subjects affected / exposed	2 / 299 (0.67%)		
occurrences causally related to treatment / all	0 / 2		
deaths causally related to treatment / all	0 / 0		
Cholelithiasis			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Gallbladder pain			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Hepatic cirrhosis			



subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Renal and urinary disorders			
Nephrolithiasis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Renal colic			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Ureterolithiasis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Intervertebral disc degeneration			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Intervertebral disc protrusion			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Musculoskeletal chest pain			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		



Spinal column stenosis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Infections and infestations			
Appendicitis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Bronchitis			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Cellulitis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Gastroenteritis viral			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
HIV infection			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Kidney infection			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Perirectal abscess			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pneumonia			



subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 1		
Postoperative wound infection			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Pyelonephritis			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Sepsis			
subjects affected / exposed	1 / 299 (0.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 1		
Upper respiratory tract infection			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Urinary tract infection			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Metabolism and nutrition disorders			
Dehydration			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Diabetic complication			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypercalcaemia			



subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Hypoglycaemia			
subjects affected / exposed	0 / 299 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	Semaglutide 0.5 mg	Semaglutide 1.0 mg	Dulaglutide 0.75 mg
Total subjects affected by non-serious adverse events			
subjects affected / exposed	132 / 301 (43.85%)	134 / 300 (44.67%)	106 / 299 (35.45%)
Investigations			
Lipase increased			
subjects affected / exposed	19 / 301 (6.31%)	17 / 300 (5.67%)	16 / 299 (5.35%)
occurrences (all)	23	17	17
Nervous system disorders			
Headache			
subjects affected / exposed	25 / 301 (8.31%)	22 / 300 (7.33%)	12 / 299 (4.01%)
occurrences (all)	35	30	20
Gastrointestinal disorders			
Constipation			
subjects affected / exposed	16 / 301 (5.32%)	14 / 300 (4.67%)	10 / 299 (3.34%)
occurrences (all)	18	14	10
Diarrhoea			
subjects affected / exposed	43 / 301 (14.29%)	41 / 300 (13.67%)	23 / 299 (7.69%)
occurrences (all)	79	96	42
Nausea			
subjects affected / exposed	67 / 301 (22.26%)	63 / 300 (21.00%)	39 / 299 (13.04%)
occurrences (all)	144	192	66
Vomiting			
subjects affected / exposed	30 / 301 (9.97%)	31 / 300 (10.33%)	12 / 299 (4.01%)
occurrences (all)	50	48	16
Infections and infestations			



Nasopharyngitis subjects affected / exposed occurrences (all)	15 / 301 (4.98%) 16	14 / 300 (4.67%) 16	17 / 299 (5.69%) 20
Upper respiratory tract infection subjects affected / exposed occurrences (all)	13 / 301 (4.32%) 18	10 / 300 (3.33%) 11	21 / 299 (7.02%) 26
Metabolism and nutrition disorders Decreased appetite subjects affected / exposed occurrences (all)	25 / 301 (8.31%) 26	27 / 300 (9.00%) 27	9 / 299 (3.01%) 12

<b>Non-serious adverse events</b>	Dulaglutide 1.5 mg		
Total subjects affected by non-serious adverse events subjects affected / exposed	139 / 299 (46.49%)		
Investigations Lipase increased subjects affected / exposed occurrences (all)	17 / 299 (5.69%) 20		
Nervous system disorders Headache subjects affected / exposed occurrences (all)	19 / 299 (6.35%) 30		
Gastrointestinal disorders Constipation subjects affected / exposed occurrences (all)  Diarrhoea subjects affected / exposed occurrences (all)  Nausea subjects affected / exposed occurrences (all)  Vomiting subjects affected / exposed occurrences (all)	15 / 299 (5.02%) 18  53 / 299 (17.73%) 75  60 / 299 (20.07%) 108  29 / 299 (9.70%) 40		
Infections and infestations Nasopharyngitis			



subjects affected / exposed occurrences (all)	20 / 299 (6.69%) 24		
Upper respiratory tract infection subjects affected / exposed occurrences (all)	16 / 299 (5.35%) 21		
Metabolism and nutrition disorders Decreased appetite subjects affected / exposed occurrences (all)	31 / 299 (10.37%) 36		



## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
28 October 2016	The protocol was updated to include diabetic retinopathy complications in the risk-benefit assessment of semaglutide based on clinical findings from the phase 3a programme. Other minor language corrections and updates were also included for general clarification.

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported