

**Table 14.2.3.1.1 Summary statistics, changes from baseline and frequencies of abnormal values of CD3 [ $10^9/L$ ] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	1.490 (0.4225)		1.590 (0.4160)	
	Geometric mean	1.432		1.540	
	Median	1.460		1.540	
	IQR	0.420		0.330	
	Q1 / Q3	1.220 / 1.640		1.430 / 1.760	
	Min / Max	0.63 / 2.71		0.94 / 2.71	
	Abnormal <sup>1</sup> , n (%)	3 (5.8%)		1 (4.8%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	1.541 (0.4935)	0.052 (0.4891)	1.848 (0.5038)	0.258 (0.4453)
	Geometric mean	1.464		1.792	
	Median	1.550	0.060	1.650	0.170
	IQR	0.741	0.510	0.450	0.520
	Q1 / Q3	1.099 / 1.840	-0.300 / 0.210	1.530 / 1.980	0.020 / 0.540
	Min / Max	0.65 / 2.97	-1.11 / 1.50	1.22 / 3.10	-0.55 / 1.09
	Abnormal <sup>1</sup> , n (%)	4 (7.8%)		4 (19.0%)	
Visit 5	n / missing	45 / 2	45 / 2	19 / 0	19 / 0
	Mean (SD)	1.513 (0.4905)	-0.006 (0.5126)	1.850 (0.5322)	0.218 (0.4206)
	Geometric mean	1.438		1.787	
	Median	1.450	-0.003	1.690	0.130
	IQR	0.510	0.440	0.600	0.650
	Q1 / Q3	1.200 / 1.710	-0.270 / 0.170	1.540 / 2.140	-0.110 / 0.540
	Min / Max	0.67 / 2.73	-1.09 / 1.88	1.04 / 3.40	-0.42 / 1.14
	Abnormal <sup>1</sup> , n (%)	6 (13.3%)		2 (10.5%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.2 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD3 [%] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	77.38 (5.637)		78.63 (4.067)	
	Geometric mean	77.17		78.53	
	Median	79.00		78.00	
	IQR	7.00		6.00	
	Q1 / Q3	74.00 / 81.00		75.00 / 81.00	
	Min / Max	64.0 / 87.0		73.0 / 86.0	
	Abnormal <sup>1</sup> , n (%)	3 (5.8%)		2 (9.5%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	78.42 (4.623)	0.86 (3.744)	79.33 (5.474)	0.70 (3.627)
	Geometric mean	78.29		79.14	
	Median	78.70	0.00	80.00	1.00
	IQR	6.70	3.80	7.00	4.00
	Q1 / Q3	75.00 / 81.70	-1.70 / 2.10	76.00 / 83.00	-1.00 / 3.00
	Min / Max	65.0 / 89.0	-5.4 / 14.0	63.0 / 88.0	-10.0 / 6.0
	Abnormal <sup>1</sup> , n (%)	2 (3.9%)		1 (4.8%)	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	77.25 (5.601)	-0.03 (3.642)	79.99 (3.235)	1.30 (3.251)
	Geometric mean	77.05		79.93	
	Median	77.00	-0.65	80.20	2.00
	IQR	9.00	6.00	5.70	4.60
	Q1 / Q3	73.00 / 82.00	-3.00 / 3.00	77.30 / 83.00	-1.60 / 3.00
	Min / Max	64.0 / 89.0	-8.0 / 8.0	71.7 / 84.0	-4.0 / 9.0
	Abnormal <sup>1</sup> , n (%)	4 (8.7%)		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.3 Summary statistics, changes from baseline and frequencies of abnormal values of CD4 [ $10^9/L$ ] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	0.928 (0.2809)		1.035 (0.2964)	
	Geometric mean	0.887		0.995	
	Median	0.890		0.960	
	IQR	0.279		0.410	
	Q1 / Q3	0.762 / 1.040		0.780 / 1.190	
	Min / Max	0.37 / 1.80		0.59 / 1.61	
	Abnormal <sup>1</sup> , n (%)	4 (7.7%)		1 (4.8%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	0.977 (0.3214)	0.049 (0.3012)	1.221 (0.3581)	0.186 (0.3058)
	Geometric mean	0.927		1.178	
	Median	0.950	0.060	1.170	0.090
	IQR	0.510	0.310	0.390	0.362
	Q1 / Q3	0.690 / 1.200	-0.140 / 0.170	1.000 / 1.390	-0.012 / 0.350
	Min / Max	0.43 / 1.88	-0.76 / 0.83	0.65 / 2.29	-0.31 / 0.83
	Abnormal <sup>1</sup> , n (%)	4 (7.8%)		2 (9.5%)	
Visit 5	n / missing	45 / 2	45 / 2	19 / 0	19 / 0
	Mean (SD)	0.951 (0.3095)	0.002 (0.3162)	1.199 (0.3240)	0.136 (0.2790)
	Geometric mean	0.903		1.158	
	Median	0.920	0.010	1.200	0.120
	IQR	0.290	0.309	0.300	0.430
	Q1 / Q3	0.780 / 1.070	-0.180 / 0.129	1.040 / 1.340	-0.100 / 0.330
	Min / Max	0.35 / 1.85	-0.73 / 1.00	0.57 / 2.15	-0.28 / 0.63
	Abnormal <sup>1</sup> , n (%)	5 (11.1%)		1 (5.3%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.4 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD4 [%] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	48.17 (5.876)		51.14 (6.097)	
	Geometric mean	47.80		50.78	
	Median	48.20		51.00	
	IQR	10.00		8.00	
	Q1 / Q3	43.00 / 53.00		47.00 / 55.00	
	Min / Max	33.6 / 59.0		37.0 / 62.0	
	Abnormal <sup>1</sup> , n (%)	0		1 (4.8%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	49.89 (5.345)	1.59 (4.169)	52.34 (6.035)	1.20 (5.176)
	Geometric mean	49.61		51.99	
	Median	50.00	0.90	54.00	3.00
	IQR	8.00	4.90	9.00	5.50
	Q1 / Q3	46.00 / 54.00	-1.00 / 3.90	48.00 / 57.00	-1.00 / 4.50
	Min / Max	37.3 / 61.0	-5.0 / 16.0	42.0 / 62.0	-12.1 / 8.0
	Abnormal <sup>1</sup> , n (%)	0		1 (4.8%)	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	48.86 (6.512)	0.45 (4.782)	52.16 (5.962)	0.96 (4.702)
	Geometric mean	48.41		51.83	
	Median	50.00	0.00	52.90	2.00
	IQR	9.90	5.30	10.20	6.30
	Q1 / Q3	43.20 / 53.10	-2.30 / 3.00	47.00 / 57.20	-2.30 / 4.00
	Min / Max	33.0 / 60.0	-9.0 / 14.0	41.0 / 63.0	-9.4 / 10.0
	Abnormal <sup>1</sup> , n (%)	0		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.5 Summary statistics, changes from baseline and frequencies of abnormal values of CD8 [10<sup>9</sup>/L] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	0.500 (0.1887)		0.484 (0.1563)	
	Geometric mean	0.469		0.464	
	Median	0.445		0.450	
	IQR	0.260		0.150	
	Q1 / Q3	0.365 / 0.625		0.380 / 0.530	
	Min / Max	0.23 / 1.11		0.31 / 0.87	
	Abnormal <sup>1</sup> , n (%)	1 (1.9%)		0	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	0.501 (0.1801)	0.001 (0.1947)	0.551 (0.1745)	0.066 (0.1242)
	Geometric mean	0.470		0.531	
	Median	0.460	0.000	0.500	0.080
	IQR	0.270	0.164	0.150	0.172
	Q1 / Q3	0.370 / 0.640	-0.090 / 0.074	0.440 / 0.590	-0.020 / 0.152
	Min / Max	0.17 / 1.03	-0.47 / 0.52	0.39 / 1.15	-0.22 / 0.29
	Abnormal <sup>1</sup> , n (%)	1 (2.0%)		1 (4.8%)	
Visit 5	n / missing	45 / 2	45 / 2	19 / 0	19 / 0
	Mean (SD)	0.496 (0.2002)	-0.013 (0.2063)	0.577 (0.2437)	0.081 (0.1567)
	Geometric mean	0.463		0.537	
	Median	0.448	-0.010	0.500	0.020
	IQR	0.219	0.150	0.200	0.190
	Q1 / Q3	0.360 / 0.579	-0.090 / 0.060	0.430 / 0.630	-0.020 / 0.170
	Min / Max	0.24 / 1.20	-0.63 / 0.78	0.26 / 1.17	-0.11 / 0.50
	Abnormal <sup>1</sup> , n (%)	1 (2.2%)		2 (10.5%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.6 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD8 [%] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	25.92 (5.607)		24.23 (5.025)	
	Geometric mean	25.32		23.69	
	Median	25.45		25.00	
	IQR	7.00		3.00	
	Q1 / Q3	22.00 / 29.00		23.00 / 26.00	
	Min / Max	15.2 / 38.0		14.0 / 35.0	
	Abnormal <sup>1</sup> , n (%)	0		0	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	25.60 (4.886)	-0.40 (3.070)	23.93 (4.734)	-0.30 (2.034)
	Geometric mean	25.15		23.43	
	Median	25.00	0.00	24.00	-0.80
	IQR	8.10	2.90	4.00	1.00
	Q1 / Q3	21.60 / 29.70	-1.90 / 1.00	22.00 / 26.00	-1.00 / 0.00
	Min / Max	17.0 / 35.0	-12.0 / 11.9	13.5 / 33.2	-5.0 / 5.4
	Abnormal <sup>1</sup> , n (%)	0		0	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	25.34 (5.413)	-0.45 (2.825)	24.72 (5.781)	0.56 (2.498)
	Geometric mean	24.79		24.04	
	Median	24.35	-0.30	24.00	0.00
	IQR	6.50	2.20	8.00	3.00
	Q1 / Q3	21.50 / 28.00	-1.20 / 1.00	22.00 / 30.00	-1.00 / 2.00
	Min / Max	16.0 / 36.6	-10.0 / 7.7	14.0 / 34.0	-4.0 / 6.1
	Abnormal <sup>1</sup> , n (%)	0		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.7 Summary statistics, changes from baseline and frequencies of abnormal values of CD16 [10<sup>9</sup>/L] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	0.220 (0.1122)		0.202 (0.1030)	
	Geometric mean	0.193		0.175	
	Median	0.210		0.190	
	IQR	0.150		0.120	
	Q1 / Q3	0.130 / 0.280		0.130 / 0.250	
	Min / Max	0.05 / 0.60		0.04 / 0.40	
	Abnormal <sup>1</sup> , n (%)	3 (5.8%)		2 (9.5%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	0.214 (0.0973)	-0.002 (0.1102)	0.235 (0.1731)	0.033 (0.1186)
	Geometric mean	0.193		0.194	
	Median	0.190	-0.004	0.170	0.010
	IQR	0.124	0.120	0.110	0.110
	Q1 / Q3	0.150 / 0.274	-0.060 / 0.060	0.140 / 0.250	-0.050 / 0.060
	Min / Max	0.04 / 0.55	-0.31 / 0.29	0.07 / 0.77	-0.13 / 0.37
	Abnormal <sup>1</sup> , n (%)	2 (3.9%)		2 (9.5%)	
Visit 5	n / missing	45 / 2	45 / 2	19 / 0	19 / 0
	Mean (SD)	0.234 (0.1238)	0.010 (0.1187)	0.204 (0.0820)	0.007 (0.0891)
	Geometric mean	0.203		0.188	
	Median	0.220	0.000	0.180	0.010
	IQR	0.150	0.080	0.122	0.130
	Q1 / Q3	0.140 / 0.290	-0.030 / 0.050	0.158 / 0.280	-0.040 / 0.090
	Min / Max	0.05 / 0.65	-0.27 / 0.32	0.08 / 0.37	-0.23 / 0.12
	Abnormal <sup>1</sup> , n (%)	5 (11.1%)		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.8 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD16 [%] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	11.41 (5.071)		10.15 (4.899)	
	Geometric mean	10.43		8.91	
	Median	10.20		9.60	
	IQR	5.50		6.00	
	Q1 / Q3	8.00 / 13.50		7.00 / 13.00	
	Min / Max	3.0 / 27.0		3.0 / 19.0	
	Abnormal <sup>1</sup> , n (%)	1 (1.9%)		4 (19.0%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	11.02 (3.844)	-0.24 (3.606)	9.81 (5.515)	-0.33 (3.827)
	Geometric mean	10.31		8.59	
	Median	11.00	-0.20	8.40	-1.00
	IQR	6.00	5.00	6.00	3.60
	Q1 / Q3	8.00 / 14.00	-2.00 / 3.00	6.00 / 12.00	-3.00 / 0.60
	Min / Max	3.0 / 19.3	-10.0 / 8.0	3.0 / 27.0	-5.6 / 9.4
	Abnormal <sup>1</sup> , n (%)	1 (2.0%)		3 (14.3%)	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	11.90 (4.790)	0.56 (3.487)	8.86 (2.571)	-0.77 (3.394)
	Geometric mean	10.88		8.45	
	Median	12.00	1.00	9.00	0.00
	IQR	7.00	3.00	3.00	4.60
	Q1 / Q3	8.00 / 15.00	-1.00 / 2.00	7.00 / 10.00	-3.00 / 1.60
	Min / Max	3.0 / 24.0	-8.4 / 8.5	4.0 / 13.0	-9.0 / 5.3
	Abnormal <sup>1</sup> , n (%)	2 (4.3%)		2 (10.5%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1



**Table 14.2.3.1.9 Summary statistics, changes from baseline and frequencies of abnormal values of CD19 [10<sup>9</sup>/L] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	0.188 (0.0786)		0.215 (0.1178)	
	Geometric mean	0.171		0.188	
	Median	0.170		0.180	
	IQR	0.109		0.100	
	Q1 / Q3	0.132 / 0.240		0.130 / 0.230	
	Min / Max	0.05 / 0.37		0.07 / 0.47	
	Abnormal <sup>1</sup> , n (%)	3 (5.8%)		1 (4.8%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	0.187 (0.0913)	0.001 (0.0744)	0.237 (0.1512)	0.023 (0.0978)
	Geometric mean	0.165		0.202	
	Median	0.168	-0.010	0.210	0.010
	IQR	0.130	0.080	0.140	0.086
	Q1 / Q3	0.120 / 0.250	-0.040 / 0.040	0.130 / 0.270	-0.036 / 0.050
	Min / Max	0.03 / 0.41	-0.16 / 0.19	0.08 / 0.64	-0.18 / 0.22
	Abnormal <sup>1</sup> , n (%)	5 (9.8%)		3 (14.3%)	
Visit 5	n / missing	45 / 2	45 / 2	19 / 0	19 / 0
	Mean (SD)	0.183 (0.0852)	-0.009 (0.0710)	0.227 (0.0902)	-0.001 (0.0772)
	Geometric mean	0.164		0.208	
	Median	0.162	-0.010	0.230	0.000
	IQR	0.120	0.061	0.170	0.080
	Q1 / Q3	0.110 / 0.230	-0.040 / 0.021	0.140 / 0.310	-0.030 / 0.050
	Min / Max	0.07 / 0.42	-0.18 / 0.16	0.09 / 0.40	-0.17 / 0.12
	Abnormal <sup>1</sup> , n (%)	5 (11.1%)		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.10 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD19 [%] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	9.79 (3.424)		10.21 (3.842)	
	Geometric mean	9.21		9.56	
	Median	9.00		10.00	
	IQR	4.00		4.00	
	Q1 / Q3	8.00 / 12.00		7.00 / 11.00	
	Min / Max	3.5 / 20.0		5.0 / 18.0	
	Abnormal <sup>1</sup> , n (%)	4 (7.7%)		1 (4.8%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	9.32 (2.894)	-0.39 (2.142)	9.60 (3.774)	-0.61 (1.729)
	Geometric mean	8.86		8.96	
	Median	9.00	-0.10	8.90	-0.50
	IQR	4.10	2.40	4.00	1.90
	Q1 / Q3	7.00 / 11.10	-1.40 / 1.00	7.00 / 11.00	-1.00 / 0.90
	Min / Max	4.0 / 15.5	-8.0 / 3.2	5.0 / 18.0	-4.1 / 2.0
	Abnormal <sup>1</sup> , n (%)	4 (7.8%)		2 (9.5%)	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	9.30 (3.063)	-0.53 (2.333)	9.78 (3.273)	-0.93 (2.112)
	Geometric mean	8.79		9.31	
	Median	8.70	0.00	9.00	-1.00
	IQR	5.00	2.20	5.00	2.40
	Q1 / Q3	7.00 / 12.00	-1.20 / 1.00	7.00 / 12.00	-1.40 / 1.00
	Min / Max	4.0 / 16.0	-10.0 / 4.0	5.8 / 16.4	-6.0 / 2.0
	Abnormal <sup>1</sup> , n (%)	6 (13.0%)		1 (5.3%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.11 Summary statistics, changes from baseline and frequencies of abnormal values of CD3+/HLA-DR3+ [ $10^9/L$ ] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	50 / 2		21 / 0	
	Mean (SD)	0.084 (0.0656)		0.094 (0.0719)	
	Geometric mean <sup>2</sup>	0.067		0.075	
	Median	0.067		0.080	
	IQR	0.060		0.080	
	Q1 / Q3	0.040 / 0.100		0.040 / 0.120	
	Min / Max	0.00 / 0.34		0.01 / 0.31	
	Abnormal <sup>1</sup> , n (%)	9 (18.0%)		3 (14.3%)	
Visit 3	n / missing	48 / 3	48 / 3	21 / 0	21 / 0
	Mean (SD)	0.076 (0.0533)	-0.010 (0.0490)	0.092 (0.0484)	-0.002 (0.0556)
	Geometric mean <sup>2</sup>	0.064		0.079	
	Median	0.065	-0.010	0.090	0.010
	IQR	0.052	0.055	0.080	0.050
	Q1 / Q3	0.040 / 0.092	-0.035 / 0.020	0.060 / 0.140	-0.020 / 0.030
	Min / Max	0.01 / 0.26	-0.19 / 0.17	0.02 / 0.17	-0.16 / 0.08
	Abnormal <sup>1</sup> , n (%)	5 (10.4%)		3 (14.3%)	
Visit 5	n / missing	44 / 3	44 / 3	18 / 1	18 / 1
	Mean (SD)	0.081 (0.0555)	-0.006 (0.0589)	0.101 (0.0687)	0.007 (0.0335)
	Geometric mean <sup>2</sup>	0.066		0.085	
	Median	0.060	-0.009	0.080	0.000
	IQR	0.065	0.050	0.060	0.040
	Q1 / Q3	0.050 / 0.115	-0.030 / 0.020	0.060 / 0.120	-0.020 / 0.020
	Min / Max	0.00 / 0.26	-0.26 / 0.16	0.02 / 0.27	-0.04 / 0.10
	Abnormal <sup>1</sup> , n (%)	7 (15.9%)		3 (16.7%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

<sup>2</sup>Because the parameter could take a value of 0, an offset value of 0.01 was added to all values for the calculation of the geometric mean. After back-transformation, this offset value was subtracted again.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.12 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD3+/HLA-DR3+ [%] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	4.38 (3.217)		4.69 (2.920)	
	Geometric mean	3.38		3.72	
	Median	3.90		4.00	
	IQR	2.65		4.70	
	Q1 / Q3	2.35 / 5.00		2.30 / 7.00	
	Min / Max	0.1 / 18.0		0.6 / 11.0	
	Abnormal <sup>1</sup> , n (%)	15 (28.8%)		5 (23.8%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	4.14 (2.844)	-0.32 (2.124)	4.28 (2.507)	-0.41 (2.166)
	Geometric mean	3.34		3.32	
	Median	3.60	0.00	4.20	-0.40
	IQR	2.70	2.30	3.00	3.00
	Q1 / Q3	2.30 / 5.00	-1.30 / 1.00	3.00 / 6.00	-2.00 / 1.00
	Min / Max	0.5 / 13.9	-6.0 / 6.3	0.7 / 9.8	-6.0 / 3.0
	Abnormal <sup>1</sup> , n (%)	17 (33.3%)		5 (23.8%)	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	4.38 (2.739)	-0.15 (3.141)	4.58 (2.573)	-0.11 (2.213)
	Geometric mean	3.43		3.77	
	Median	4.00	-0.15	4.00	-0.10
	IQR	3.70	2.90	4.50	2.10
	Q1 / Q3	2.30 / 6.00	-1.50 / 1.40	2.20 / 6.70	-1.10 / 1.00
	Min / Max	0.1 / 13.0	-13.0 / 10.0	0.6 / 9.3	-4.0 / 4.8
	Abnormal <sup>1</sup> , n (%)	14 (30.4%)		4 (21.1%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.13 Summary statistics, changes from baseline and frequencies of abnormal values of total IgG [g/L] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	11.060 (2.2439)		10.679 (1.7209)	
	Geometric mean	10.834		10.545	
	Median	10.740		10.500	
	IQR	2.915		2.430	
	Q1 / Q3	9.485 / 12.400		9.400 / 11.830	
	Min / Max	6.60 / 15.92		8.23 / 13.93	
	Abnormal <sup>1</sup> , n (%)	1 (1.9%)		0	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	11.336 (2.1362)	0.203 (1.3182)	10.581 (1.7753)	-0.097 (1.0652)
	Geometric mean	11.137		10.443	
	Median	11.000	0.100	10.400	0.000
	IQR	2.910	1.390	2.370	1.010
	Q1 / Q3	9.890 / 12.800	-0.690 / 0.700	9.100 / 11.470	-0.610 / 0.400
	Min / Max	6.80 / 16.02	-2.29 / 3.83	8.00 / 14.75	-2.70 / 2.60
	Abnormal <sup>1</sup> , n (%)	2 (3.9%)		0	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	11.244 (2.2239)	0.142 (1.5836)	10.741 (2.0338)	0.224 (1.3731)
	Geometric mean	11.033		10.553	
	Median	11.050	-0.150	10.800	0.020
	IQR	2.900	1.900	4.480	2.440
	Q1 / Q3	9.800 / 12.700	-0.900 / 1.000	8.320 / 12.800	-1.140 / 1.300
	Min / Max	6.80 / 16.55	-2.60 / 4.39	7.78 / 14.00	-1.74 / 2.70
	Abnormal <sup>1</sup> , n (%)	2 (4.3%)		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.14 Summary statistics, changes from baseline and frequencies of abnormal values of total IgM [g/L] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	1.249 (0.4692)		1.339 (1.0216)	
	Geometric mean	1.169		1.126	
	Median	1.140		1.090	
	IQR	0.480		0.680	
	Q1 / Q3	0.990 / 1.470		0.860 / 1.540	
	Min / Max	0.46 / 2.79		0.35 / 5.35	
	Abnormal <sup>1</sup> , n (%)	2 (3.8%)		3 (14.3%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	1.210 (0.4691)	-0.030 (0.2090)	1.336 (1.1386)	-0.003 (0.1807)
	Geometric mean	1.123		1.090	
	Median	1.140	-0.030	1.110	-0.020
	IQR	0.450	0.160	0.870	0.210
	Q1 / Q3	0.960 / 1.410	-0.100 / 0.060	0.710 / 1.580	-0.130 / 0.080
	Min / Max	0.42 / 2.55	-0.97 / 0.43	0.31 / 5.88	-0.25 / 0.53
	Abnormal <sup>1</sup> , n (%)	4 (7.8%)		3 (14.3%)	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	1.218 (0.4569)	-0.043 (0.2333)	1.402 (1.1996)	0.048 (0.2442)
	Geometric mean	1.137		1.133	
	Median	1.150	-0.030	1.060	0.010
	IQR	0.580	0.230	0.940	0.320
	Q1 / Q3	0.900 / 1.480	-0.170 / 0.060	0.810 / 1.750	-0.150 / 0.170
	Min / Max	0.43 / 2.70	-0.84 / 0.85	0.33 / 5.89	-0.21 / 0.65
	Abnormal <sup>1</sup> , n (%)	2 (4.3%)		3 (15.8%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.15 Summary statistics, changes from baseline and frequencies of abnormal values of total IgA [g/L] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	2.084 (0.7128)		2.131 (0.7126)	
	Geometric mean	1.974		2.026	
	Median	1.985		1.880	
	IQR	0.805		1.190	
	Q1 / Q3	1.615 / 2.420		1.560 / 2.750	
	Min / Max	0.91 / 4.15		1.31 / 3.91	
	Abnormal <sup>1</sup> , n (%)	1 (1.9%)		1 (4.8%)	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	2.037 (0.6886)	-0.066 (0.1978)	2.081 (0.6800)	-0.050 (0.2505)
	Geometric mean	1.939		1.981	
	Median	1.830	-0.080	1.950	-0.050
	IQR	0.760	0.200	1.030	0.220
	Q1 / Q3	1.570 / 2.330	-0.190 / 0.010	1.520 / 2.550	-0.140 / 0.080
	Min / Max	1.12 / 4.20	-0.56 / 0.57	1.11 / 3.81	-0.78 / 0.46
	Abnormal <sup>1</sup> , n (%)	2 (3.9%)		1 (4.8%)	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	1.985 (0.5897)	-0.049 (0.2433)	2.125 (0.6267)	-0.004 (0.3094)
	Geometric mean	1.907		2.040	
	Median	1.820	-0.035	1.800	0.000
	IQR	0.650	0.300	1.180	0.420
	Q1 / Q3	1.600 / 2.250	-0.230 / 0.070	1.620 / 2.800	-0.160 / 0.260
	Min / Max	1.06 / 3.89	-0.49 / 0.68	1.22 / 3.17	-0.77 / 0.51
	Abnormal <sup>1</sup> , n (%)	1 (2.2%)		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.16 Summary statistics, changes from baseline and frequencies of abnormal values of Anti-D-Mannan IgM [U/mL] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	50 / 0		21 / 0	
	Mean (SD)	20.798 (18.9490)		20.624 (22.7586)	
	Geometric mean	14.205		13.131	
	Median	13.850		12.700	
	IQR	14.600		21.300	
	Q1 / Q3	9.100 / 23.700		7.500 / 28.800	
	Min / Max	0.10 / 98.50		1.10 / 100.00	
	Abnormal <sup>1</sup> , n (%)	22 (44.0%)		7 (33.3%)	
Visit 3	n / missing	49 / 0	49 / 0	21 / 0	21 / 0
	Mean (SD)	19.918 (15.1198)	-0.043 (9.9950)	19.638 (15.1029)	-0.986 (15.1021)
	Geometric mean	15.490		13.834	
	Median	17.400	0.700	15.500	1.300
	IQR	14.600	7.900	28.400	9.900
	Q1 / Q3	8.800 / 23.400	-3.500 / 4.400	8.400 / 36.800	-1.900 / 8.000
	Min / Max	1.90 / 68.60	-40.80 / 23.90	2.40 / 52.30	-47.70 / 19.80
	Abnormal <sup>1</sup> , n (%)	26 (53.1%)		11 (52.4%)	
Visit 5	n / missing	44 / 2	44 / 2	19 / 0	19 / 0
	Mean (SD)	17.268 (12.2924)	-2.216 (14.8520)	16.626 (14.7514)	-5.247 (20.3044)
	Geometric mean	13.849		11.446	
	Median	13.950	-0.050	12.700	-0.700
	IQR	10.950	5.900	14.000	7.000
	Q1 / Q3	9.600 / 20.550	-3.150 / 2.750	6.000 / 20.000	-4.700 / 2.300
	Min / Max	1.80 / 52.30	-72.60 / 26.30	1.60 / 52.50	-68.20 / 21.60
	Abnormal <sup>1</sup> , n (%)	19 (43.2%)		6 (31.6%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

Patients from study centers 004 and 005 were not included due to different measurement unit.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1



**Table 14.2.3.1.17 Summary statistics, changes from baseline and frequencies of abnormal values of Anti-D-Glucan IgG [U/mL] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	50 / 0		21 / 0	
	Mean (SD)	26.884 (22.9759)		31.862 (28.0225)	
	Geometric mean	18.679		21.674	
	Median	21.750		25.300	
	IQR	22.000		31.700	
	Q1 / Q3	10.800 / 32.800		11.200 / 42.900	
	Min / Max	2.50 / 100.00		4.20 / 100.00	
	Abnormal <sup>1</sup> , n (%)	30 (60.0%)		13 (61.9%)	
Visit 3	n / missing	49 / 0	49 / 0	21 / 0	21 / 0
	Mean (SD)	30.359 (27.6834)	3.202 (18.9801)	29.467 (30.3826)	-2.395 (15.9648)
	Geometric mean	18.778		19.532	
	Median	20.500	-0.800	19.100	0.000
	IQR	36.800	7.100	16.000	12.500
	Q1 / Q3	9.300 / 46.100	-4.300 / 2.800	11.500 / 27.500	-10.300 / 2.200
	Min / Max	0.70 / 100.00	-28.40 / 74.30	4.20 / 100.00	-38.50 / 44.70
	Abnormal <sup>1</sup> , n (%)	30 (61.2%)		12 (57.1%)	
Visit 5	n / missing	44 / 2	44 / 2	19 / 0	19 / 0
	Mean (SD)	21.677 (21.2119)	-4.945 (14.8008)	23.000 (25.2827)	-10.274 (19.5773)
	Geometric mean	13.180		15.229	
	Median	14.400	-4.000	11.500	-5.400
	IQR	22.400	10.950	23.200	16.500
	Q1 / Q3	6.400 / 28.800	-10.100 / 0.850	7.900 / 31.100	-16.300 / 0.200
	Min / Max	0.10 / 100.00	-39.70 / 51.80	4.40 / 96.60	-68.90 / 22.50
	Abnormal <sup>1</sup> , n (%)	21 (47.7%)		8 (42.1%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

Patients from study centers 004 and 005 were not included due to different measurement unit.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.1.18 Summary statistics, changes from baseline and frequencies of abnormal values of phagocyte activity [%] by scheduled visit and treatment (PPS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	52 / 0		21 / 0	
	Mean (SD)	97.546 (2.6073)		97.271 (4.5734)	
	Geometric mean	97.511		97.157	
	Median	98.100		99.000	
	IQR	2.000		1.400	
	Q1 / Q3	97.000 / 99.000		97.600 / 99.000	
	Min / Max	87.00 / 99.90		78.80 / 100.00	
	Abnormal <sup>1</sup> , n (%)	0		0	
Visit 3	n / missing	51 / 0	51 / 0	21 / 0	21 / 0
	Mean (SD)	96.723 (5.1851)	-0.777 (5.5778)	97.410 (2.1522)	0.138 (4.2756)
	Geometric mean	96.566		97.387	
	Median	98.000	0.000	98.000	0.000
	IQR	3.000	2.000	2.000	2.900
	Q1 / Q3	96.100 / 99.100	-1.000 / 1.000	97.000 / 99.000	-1.900 / 1.000
	Min / Max	69.60 / 100.00	-30.30 / 7.20	91.70 / 99.80	-5.90 / 16.80
	Abnormal <sup>1</sup> , n (%)	1 (2.0%)		0	
Visit 5	n / missing	46 / 1	46 / 1	19 / 0	19 / 0
	Mean (SD)	96.905 (3.1553)	-0.687 (3.4794)	95.721 (4.5385)	-1.384 (4.8891)
	Geometric mean	96.852		95.611	
	Median	97.660	0.000	97.000	-1.000
	IQR	2.600	3.000	2.400	1.500
	Q1 / Q3	96.400 / 99.000	-2.000 / 1.000	95.600 / 98.000	-1.500 / 0.000
	Min / Max	83.50 / 100.00	-15.30 / 8.00	80.80 / 99.00	-19.00 / 8.20
	Abnormal <sup>1</sup> , n (%)	0		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; PPS = Per-protocol set; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.1 Summary statistics, changes from baseline and frequencies of abnormal values of CD3 [ $10^9/L$ ] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	1.485 (0.4306)		1.539 (0.4382)	
	Geometric mean	1.425		1.477	
	Median	1.440		1.475	
	IQR	0.430		0.680	
	Q1 / Q3	1.220 / 1.650		1.230 / 1.910	
	Min / Max	0.63 / 2.81		0.69 / 2.71	
	Abnormal <sup>1</sup> , n (%)	6 (6.7%)		5 (10.9%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	1.527 (0.5101)	0.034 (0.4700)	1.662 (0.5000)	0.123 (0.4490)
	Geometric mean	1.443		1.596	
	Median	1.520	0.000	1.545	0.055
	IQR	0.730	0.430	0.608	0.590
	Q1 / Q3	1.120 / 1.850	-0.220 / 0.210	1.322 / 1.930	-0.200 / 0.390
	Min / Max	0.58 / 2.97	-1.52 / 1.50	0.87 / 3.10	-0.84 / 1.10
	Abnormal <sup>1</sup> , n (%)	9 (10.2%)		5 (10.9%)	
Visit 5	n / missing	77 / 2	77 / 2	42 / 0	42 / 0
	Mean (SD)	1.494 (0.5214)	-0.004 (0.4686)	1.668 (0.5214)	0.121 (0.4716)
	Geometric mean	1.411		1.595	
	Median	1.380	-0.059	1.595	0.090
	IQR	0.590	0.411	0.457	0.587
	Q1 / Q3	1.150 / 1.740	-0.251 / 0.160	1.383 / 1.840	-0.187 / 0.400
	Min / Max	0.67 / 3.03	-1.09 / 1.88	0.84 / 3.40	-0.88 / 1.71
	Abnormal <sup>1</sup> , n (%)	11 (14.3%)		3 (7.1%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.2 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD3 [%] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	76.54 (6.227)		76.82 (7.363)	
	Geometric mean	76.28		76.43	
	Median	77.50		77.00	
	IQR	8.00		8.30	
	Q1 / Q3	73.00 / 81.00		73.70 / 82.00	
	Min / Max	57.5 / 88.0		45.4 / 87.8	
	Abnormal <sup>1</sup> , n (%)	5 (5.6%)		6 (13.0%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	77.55 (5.232)	0.97 (3.812)	78.01 (6.164)	1.18 (5.758)
	Geometric mean	77.37		77.76	
	Median	78.00	0.40	78.85	1.00
	IQR	7.85	4.20	9.60	3.60
	Q1 / Q3	74.00 / 81.85	-1.20 / 3.00	73.40 / 83.00	-1.00 / 2.60
	Min / Max	62.1 / 89.0	-8.0 / 14.0	63.0 / 89.5	-10.0 / 33.3
	Abnormal <sup>1</sup> , n (%)	3 (3.4%)		4 (8.7%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	76.49 (5.970)	0.21 (3.878)	78.06 (5.840)	1.47 (5.149)
	Geometric mean	76.26		77.84	
	Median	76.10	0.00	79.50	0.85
	IQR	9.30	5.90	6.90	4.60
	Q1 / Q3	72.00 / 81.30	-2.90 / 3.00	75.10 / 82.00	-2.00 / 2.60
	Min / Max	64.0 / 89.0	-8.0 / 13.6	66.0 / 90.4	-4.2 / 26.9
	Abnormal <sup>1</sup> , n (%)	6 (7.7%)		2 (4.8%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.3 Summary statistics, changes from baseline and frequencies of abnormal values of CD4 [ $10^9/L$ ] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	0.922 (0.2998)		0.997 (0.3067)	
	Geometric mean	0.878		0.949	
	Median	0.870		0.960	
	IQR	0.386		0.420	
	Q1 / Q3	0.724 / 1.110		0.770 / 1.190	
	Min / Max	0.37 / 2.05		0.35 / 1.61	
	Abnormal <sup>1</sup> , n (%)	6 (6.7%)		2 (4.3%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	0.957 (0.3474)	0.032 (0.2901)	1.096 (0.3595)	0.099 (0.3030)
	Geometric mean	0.899		1.046	
	Median	0.909	0.010	1.035	0.041
	IQR	0.470	0.297	0.430	0.430
	Q1 / Q3	0.670 / 1.140	-0.127 / 0.170	0.860 / 1.290	-0.100 / 0.330
	Min / Max	0.35 / 2.23	-1.02 / 0.83	0.57 / 2.29	-0.55 / 0.83
	Abnormal <sup>1</sup> , n (%)	8 (9.1%)		3 (6.5%)	
Visit 5	n / missing	77 / 2	77 / 2	42 / 0	42 / 0
	Mean (SD)	0.940 (0.3466)	0.010 (0.3008)	1.096 (0.3384)	0.094 (0.3187)
	Geometric mean	0.883		1.047	
	Median	0.880	0.009	1.050	0.045
	IQR	0.436	0.270	0.406	0.390
	Q1 / Q3	0.694 / 1.130	-0.170 / 0.100	0.844 / 1.250	-0.101 / 0.289
	Min / Max	0.35 / 2.03	-0.73 / 1.00	0.57 / 2.15	-0.61 / 1.26
	Abnormal <sup>1</sup> , n (%)	8 (10.4%)		2 (4.8%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.4 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD4 [%] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	47.38 (5.901)		49.75 (7.442)	
	Geometric mean	47.01		49.11	
	Median	47.50		50.00	
	IQR	7.20		10.30	
	Q1 / Q3	43.80 / 51.00		44.70 / 55.00	
	Min / Max	33.1 / 61.3		23.1 / 62.0	
	Abnormal <sup>1</sup> , n (%)	0		2 (4.3%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	48.52 (5.512)	1.21 (3.902)	51.31 (5.989)	1.56 (4.626)
	Geometric mean	48.21		50.95	
	Median	48.00	0.90	52.00	2.00
	IQR	8.40	4.00	11.00	4.00
	Q1 / Q3	44.60 / 53.00	-1.00 / 3.00	46.00 / 57.00	0.00 / 4.00
	Min / Max	36.0 / 61.0	-9.0 / 16.0	38.3 / 62.0	-12.1 / 18.9
	Abnormal <sup>1</sup> , n (%)	0		1 (2.2%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	48.16 (6.585)	0.82 (4.633)	51.44 (5.822)	1.78 (4.983)
	Geometric mean	47.71		51.11	
	Median	48.80	0.20	51.65	1.00
	IQR	8.80	5.00	8.40	4.90
	Q1 / Q3	43.20 / 52.00	-2.00 / 3.00	46.90 / 55.30	-0.90 / 4.00
	Min / Max	33.0 / 62.4	-9.0 / 14.0	41.0 / 63.0	-9.4 / 18.6
	Abnormal <sup>1</sup> , n (%)	0		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.5 Summary statistics, changes from baseline and frequencies of abnormal values of CD8 [10<sup>9</sup>/L] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	0.498 (0.1884)		0.473 (0.1717)	
	Geometric mean	0.466		0.442	
	Median	0.460		0.450	
	IQR	0.230		0.170	
	Q1 / Q3	0.370 / 0.600		0.380 / 0.550	
	Min / Max	0.19 / 1.11		0.13 / 0.97	
	Abnormal <sup>1</sup> , n (%)	3 (3.4%)		1 (2.2%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	0.512 (0.1999)	0.005 (0.1843)	0.499 (0.1742)	0.026 (0.1330)
	Geometric mean	0.475		0.470	
	Median	0.483	-0.003	0.470	0.002
	IQR	0.247	0.150	0.150	0.170
	Q1 / Q3	0.380 / 0.627	-0.080 / 0.070	0.410 / 0.560	-0.040 / 0.130
	Min / Max	0.16 / 1.19	-0.47 / 0.67	0.14 / 1.15	-0.22 / 0.31
	Abnormal <sup>1</sup> , n (%)	3 (3.4%)		1 (2.2%)	
Visit 5	n / missing	77 / 2	77 / 2	42 / 0	42 / 0
	Mean (SD)	0.492 (0.2208)	-0.012 (0.1888)	0.513 (0.2168)	0.038 (0.1558)
	Geometric mean	0.453		0.472	
	Median	0.440	-0.020	0.454	0.010
	IQR	0.211	0.146	0.187	0.160
	Q1 / Q3	0.349 / 0.560	-0.100 / 0.046	0.393 / 0.580	-0.060 / 0.100
	Min / Max	0.22 / 1.33	-0.63 / 0.78	0.14 / 1.17	-0.30 / 0.50
	Abnormal <sup>1</sup> , n (%)	3 (3.9%)		2 (4.8%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.6 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD8 [%] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	25.81 (6.826)		23.72 (6.241)	
	Geometric mean	24.95		22.88	
	Median	25.00		24.25	
	IQR	9.00		6.10	
	Q1 / Q3	21.00 / 30.00		19.90 / 26.00	
	Min / Max	14.0 / 49.2		10.0 / 42.4	
	Abnormal <sup>1</sup> , n (%)	2 (2.2%)		2 (4.3%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	26.18 (6.248)	0.14 (3.219)	23.69 (5.883)	-0.02 (3.193)
	Geometric mean	25.46		22.94	
	Median	25.00	0.00	23.90	-0.40
	IQR	9.70	2.00	6.50	2.60
	Q1 / Q3	21.15 / 30.85	-1.00 / 1.00	20.00 / 26.50	-1.60 / 1.00
	Min / Max	15.0 / 45.6	-12.0 / 16.2	10.0 / 39.6	-5.0 / 17.0
	Abnormal <sup>1</sup> , n (%)	2 (2.3%)		1 (2.2%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	25.32 (6.574)	-0.43 (3.132)	23.89 (6.234)	0.35 (3.544)
	Geometric mean	24.50		23.06	
	Median	24.00	-0.30	23.15	0.00
	IQR	8.90	3.00	6.60	2.70
	Q1 / Q3	21.00 / 29.90	-2.00 / 1.00	20.30 / 26.90	-1.30 / 1.40
	Min / Max	13.6 / 44.5	-10.7 / 12.1	11.0 / 38.7	-6.0 / 14.6
	Abnormal <sup>1</sup> , n (%)	2 (2.6%)		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1



**Table 14.2.3.2.7 Summary statistics, changes from baseline and frequencies of abnormal values of CD16 [10<sup>9</sup>/L] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	0.228 (0.1158)		0.225 (0.1122)	
	Geometric mean	0.202		0.197	
	Median	0.210		0.213	
	IQR	0.150		0.160	
	Q1 / Q3	0.140 / 0.290		0.140 / 0.300	
	Min / Max	0.05 / 0.60		0.04 / 0.58	
	Abnormal <sup>1</sup> , n (%)	5 (5.6%)		4 (8.7%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	0.221 (0.1167)	-0.005 (0.1085)	0.234 (0.1467)	0.009 (0.1275)
	Geometric mean	0.194		0.199	
	Median	0.190	-0.010	0.190	0.000
	IQR	0.131	0.110	0.123	0.103
	Q1 / Q3	0.147 / 0.278	-0.060 / 0.050	0.147 / 0.270	-0.050 / 0.053
	Min / Max	0.04 / 0.61	-0.33 / 0.35	0.04 / 0.77	-0.43 / 0.37
	Abnormal <sup>1</sup> , n (%)	5 (5.7%)		5 (10.9%)	
Visit 5	n / missing	77 / 2	77 / 2	42 / 0	42 / 0
	Mean (SD)	0.226 (0.1151)	-0.007 (0.1122)	0.219 (0.0974)	-0.008 (0.0963)
	Geometric mean	0.198		0.196	
	Median	0.220	-0.010	0.204	0.000
	IQR	0.157	0.080	0.120	0.090
	Q1 / Q3	0.133 / 0.290	-0.050 / 0.030	0.160 / 0.280	-0.050 / 0.040
	Min / Max	0.05 / 0.65	-0.34 / 0.32	0.06 / 0.43	-0.39 / 0.17
	Abnormal <sup>1</sup> , n (%)	6 (7.8%)		4 (9.5%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.8 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD16 [%] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	12.17 (5.864)		11.70 (6.175)	
	Geometric mean	10.93		10.22	
	Median	11.40		10.95	
	IQR	7.00		8.00	
	Q1 / Q3	8.00 / 15.00		7.00 / 15.00	
	Min / Max	3.0 / 32.5		3.0 / 37.7	
	Abnormal <sup>1</sup> , n (%)	4 (4.5%)		7 (15.2%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	11.35 (4.807)	-0.69 (4.009)	10.94 (5.188)	-0.76 (5.324)
	Geometric mean	10.37		9.70	
	Median	10.50	-0.30	10.45	-0.45
	IQR	6.55	4.80	6.80	3.00
	Q1 / Q3	8.00 / 14.55	-2.80 / 2.00	7.00 / 13.80	-2.00 / 1.00
	Min / Max	3.0 / 30.3	-16.5 / 8.0	2.5 / 27.0	-29.0 / 9.4
	Abnormal <sup>1</sup> , n (%)	4 (4.5%)		4 (8.7%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	11.98 (5.374)	-0.13 (3.898)	10.44 (3.843)	-1.29 (5.082)
	Geometric mean	10.74		9.69	
	Median	12.00	0.30	10.00	0.00
	IQR	8.20	4.00	4.70	4.50
	Q1 / Q3	7.70 / 15.90	-2.00 / 2.00	8.30 / 13.00	-3.00 / 1.50
	Min / Max	3.0 / 26.3	-15.8 / 8.5	3.8 / 19.0	-25.9 / 5.3
	Abnormal <sup>1</sup> , n (%)	4 (5.1%)		4 (9.5%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.9 Summary statistics, changes from baseline and frequencies of abnormal values of CD19 [10<sup>9</sup>/L] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	0.197 (0.0900)		0.212 (0.1047)	
	Geometric mean	0.178		0.190	
	Median	0.180		0.185	
	IQR	0.107		0.119	
	Q1 / Q3	0.133 / 0.240		0.129 / 0.248	
	Min / Max	0.05 / 0.48		0.07 / 0.48	
	Abnormal <sup>1</sup> , n (%)	9 (10.1%)		1 (2.2%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	0.197 (0.1002)	-0.003 (0.0715)	0.220 (0.1228)	0.007 (0.0783)
	Geometric mean	0.174		0.194	
	Median	0.178	0.000	0.195	-0.009
	IQR	0.130	0.067	0.135	0.066
	Q1 / Q3	0.120 / 0.250	-0.040 / 0.027	0.130 / 0.265	-0.036 / 0.030
	Min / Max	0.03 / 0.56	-0.20 / 0.22	0.08 / 0.64	-0.18 / 0.22
	Abnormal <sup>1</sup> , n (%)	8 (9.1%)		3 (6.5%)	
Visit 5	n / missing	77 / 2	77 / 2	42 / 0	42 / 0
	Mean (SD)	0.199 (0.1009)	-0.002 (0.0723)	0.221 (0.0957)	0.003 (0.0785)
	Geometric mean	0.176		0.200	
	Median	0.170	-0.010	0.191	-0.006
	IQR	0.120	0.060	0.160	0.070
	Q1 / Q3	0.130 / 0.250	-0.040 / 0.020	0.140 / 0.300	-0.030 / 0.040
	Min / Max	0.07 / 0.48	-0.18 / 0.29	0.09 / 0.42	-0.17 / 0.28
	Abnormal <sup>1</sup> , n (%)	11 (14.3%)		2 (4.8%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.10 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD19 [%] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	10.18 (3.582)		10.48 (3.648)	
	Geometric mean	9.53		9.86	
	Median	10.00		10.35	
	IQR	4.10		5.10	
	Q1 / Q3	8.00 / 12.10		8.00 / 13.10	
	Min / Max	3.5 / 20.0		4.9 / 18.0	
	Abnormal <sup>1</sup> , n (%)	8 (9.0%)		3 (6.5%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	9.95 (3.338)	-0.30 (2.219)	10.02 (3.400)	-0.46 (1.717)
	Geometric mean	9.39		9.48	
	Median	9.10	0.00	9.00	0.00
	IQR	5.65	3.00	4.60	2.90
	Q1 / Q3	7.30 / 12.95	-2.00 / 1.00	7.40 / 12.00	-2.00 / 0.90
	Min / Max	4.0 / 21.0	-8.0 / 4.2	5.0 / 18.0	-4.1 / 4.0
	Abnormal <sup>1</sup> , n (%)	4 (4.5%)		3 (6.5%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	10.08 (3.393)	-0.15 (2.277)	10.19 (3.375)	-0.52 (2.046)
	Geometric mean	9.51		9.66	
	Median	10.00	0.00	9.70	-0.25
	IQR	4.70	2.00	5.30	2.10
	Q1 / Q3	7.30 / 12.00	-1.00 / 1.00	7.00 / 12.30	-1.20 / 0.90
	Min / Max	4.0 / 18.0	-10.0 / 5.0	5.2 / 16.4	-6.0 / 4.0
	Abnormal <sup>1</sup> , n (%)	6 (7.7%)		2 (4.8%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.11 Summary statistics, changes from baseline and frequencies of abnormal values of CD3+/HLA-DR3+ [10<sup>9</sup>/L] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	73 / 17		37 / 9	
	Mean (SD)	0.087 (0.0603)		0.081 (0.0606)	
	Geometric mean <sup>2</sup>	0.071		0.066	
	Median	0.070		0.070	
	IQR	0.079		0.060	
	Q1 / Q3	0.041 / 0.120		0.040 / 0.100	
	Min / Max	0.00 / 0.34		0.01 / 0.31	
	Abnormal <sup>1</sup> , n (%)	11 (15.1%)		5 (13.5%)	
Visit 3	n / missing	71 / 17	71 / 17	37 / 9	37 / 9
	Mean (SD)	0.078 (0.0511)	-0.011 (0.0472)	0.077 (0.0464)	-0.004 (0.0451)
	Geometric mean <sup>2</sup>	0.066		0.065	
	Median	0.070	-0.010	0.070	0.000
	IQR	0.060	0.060	0.074	0.040
	Q1 / Q3	0.040 / 0.100	-0.040 / 0.020	0.036 / 0.110	-0.020 / 0.020
	Min / Max	0.01 / 0.26	-0.19 / 0.17	0.01 / 0.17	-0.16 / 0.08
	Abnormal <sup>1</sup> , n (%)	9 (12.7%)		6 (16.2%)	
Visit 5	n / missing	64 / 15	64 / 15	33 / 9	33 / 9
	Mean (SD)	0.082 (0.0559)	-0.008 (0.0554)	0.086 (0.0701)	0.005 (0.0453)
	Geometric mean <sup>2</sup>	0.067		0.067	
	Median	0.069	-0.010	0.060	0.000
	IQR	0.065	0.050	0.078	0.030
	Q1 / Q3	0.050 / 0.115	-0.030 / 0.020	0.040 / 0.118	-0.020 / 0.010
	Min / Max	0.00 / 0.26	-0.26 / 0.16	0.02 / 0.27	-0.08 / 0.17
	Abnormal <sup>1</sup> , n (%)	9 (14.1%)		10 (30.3%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

<sup>2</sup>Because the parameter could take a value of 0, an offset value of 0.01 was added to all values for the calculation of the geometric mean. After back-transformation, this offset value was subtracted again.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.12 Summary statistics, changes from baseline and frequencies of abnormal values of differential count of CD3+/HLA-DR3+ [%] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	4.41 (2.879)		4.40 (2.972)	
	Geometric mean	3.54		3.51	
	Median	3.90		3.05	
	IQR	2.90		4.70	
	Q1 / Q3	2.40 / 5.30		2.10 / 6.80	
	Min / Max	0.1 / 18.0		0.6 / 12.1	
	Abnormal <sup>1</sup> , n (%)	24 (27.0%)		13 (28.3%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	4.24 (2.745)	-0.32 (2.020)	3.78 (2.408)	-0.62 (2.057)
	Geometric mean	3.43		2.96	
	Median	3.65	-0.20	3.50	-0.40
	IQR	3.05	2.50	3.70	2.80
	Q1 / Q3	2.40 / 5.45	-1.50 / 1.00	1.70 / 5.40	-2.00 / 0.80
	Min / Max	0.5 / 13.9	-6.0 / 6.3	0.7 / 9.8	-7.4 / 3.0
	Abnormal <sup>1</sup> , n (%)	27 (30.7%)		18 (39.1%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	4.17 (2.605)	-0.28 (2.901)	3.84 (2.537)	-0.65 (2.182)
	Geometric mean	3.30		3.05	
	Median	3.45	-0.20	3.25	-0.55
	IQR	3.70	2.60	4.40	1.90
	Q1 / Q3	2.30 / 6.00	-1.60 / 1.00	1.60 / 6.00	-1.50 / 0.40
	Min / Max	0.1 / 13.0	-13.0 / 10.0	0.6 / 10.2	-8.0 / 4.8
	Abnormal <sup>1</sup> , n (%)	23 (29.5%)		16 (38.1%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.13 Summary statistics, changes from baseline and frequencies of abnormal values of total IgG [g/L] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	10.842 (2.2169)		10.624 (1.5716)	
	Geometric mean	10.621		10.509	
	Median	10.660		10.700	
	IQR	2.840		2.300	
	Q1 / Q3	9.360 / 12.200		9.400 / 11.700	
	Min / Max	6.60 / 17.10		7.81 / 13.93	
	Abnormal <sup>1</sup> , n (%)	3 (3.4%)		0	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	11.062 (2.1714)	0.167 (1.1681)	10.488 (1.6173)	-0.136 (0.9285)
	Geometric mean	10.853		10.367	
	Median	10.785	0.100	10.470	-0.050
	IQR	3.015	1.280	2.300	1.200
	Q1 / Q3	9.525 / 12.540	-0.680 / 0.600	9.100 / 11.400	-0.800 / 0.400
	Min / Max	6.80 / 17.20	-2.29 / 3.83	7.54 / 14.75	-2.70 / 2.60
	Abnormal <sup>1</sup> , n (%)	3 (3.4%)		0	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	11.123 (2.1822)	0.185 (1.3968)	10.769 (2.3199)	0.232 (1.4908)
	Geometric mean	10.921		10.546	
	Median	11.060	0.120	10.565	0.055
	IQR	2.550	1.910	3.200	1.900
	Q1 / Q3	9.660 / 12.210	-0.770 / 1.140	8.700 / 11.900	-0.830 / 1.070
	Min / Max	6.80 / 18.50	-2.60 / 4.39	7.61 / 18.83	-1.74 / 5.29
	Abnormal <sup>1</sup> , n (%)	3 (3.8%)		1 (2.4%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.14 Summary statistics, changes from baseline and frequencies of abnormal values of total IgM [g/L] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	1.258 (0.4594)		1.464 (0.8126)	
	Geometric mean	1.176		1.301	
	Median	1.160		1.260	
	IQR	0.510		0.830	
	Q1 / Q3	0.990 / 1.500		1.000 / 1.830	
	Min / Max	0.46 / 2.79		0.35 / 5.35	
	Abnormal <sup>1</sup> , n (%)	2 (2.2%)		8 (17.4%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	1.226 (0.4503)	-0.024 (0.1944)	1.455 (0.8749)	-0.010 (0.1672)
	Geometric mean	1.141		1.269	
	Median	1.180	-0.020	1.255	-0.010
	IQR	0.505	0.180	0.800	0.160
	Q1 / Q3	0.965 / 1.470	-0.110 / 0.070	1.020 / 1.820	-0.120 / 0.040
	Min / Max	0.41 / 2.55	-0.97 / 0.43	0.31 / 5.88	-0.31 / 0.53
	Abnormal <sup>1</sup> , n (%)	4 (4.5%)		4 (8.7%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	1.219 (0.4630)	-0.014 (0.2458)	1.480 (0.9116)	-0.009 (0.2248)
	Geometric mean	1.138		1.286	
	Median	1.175	-0.030	1.315	-0.065
	IQR	0.590	0.230	0.980	0.260
	Q1 / Q3	0.890 / 1.480	-0.120 / 0.110	0.960 / 1.940	-0.150 / 0.110
	Min / Max	0.43 / 2.89	-0.84 / 1.10	0.33 / 5.89	-0.41 / 0.65
	Abnormal <sup>1</sup> , n (%)	3 (3.8%)		5 (11.9%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1



**Table 14.2.3.2.15 Summary statistics, changes from baseline and frequencies of abnormal values of total IgA [g/L] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	89 / 1		46 / 0	
	Mean (SD)	2.041 (0.8211)		2.236 (0.8391)	
	Geometric mean	1.877		2.095	
	Median	1.890		2.055	
	IQR	0.720		1.220	
	Q1 / Q3	1.590 / 2.310		1.560 / 2.780	
	Min / Max	0.10 / 5.72		1.15 / 4.37	
	Abnormal <sup>1</sup> , n (%)	4 (4.5%)		3 (6.5%)	
Visit 3	n / missing	88 / 0	87 / 1	46 / 0	46 / 0
	Mean (SD)	2.026 (0.8123)	-0.049 (0.1955)	2.217 (0.8290)	-0.018 (0.2512)
	Geometric mean	1.868		2.084	
	Median	1.830	-0.050	2.105	-0.005
	IQR	0.755	0.240	1.200	0.280
	Q1 / Q3	1.555 / 2.310	-0.190 / 0.050	1.520 / 2.720	-0.160 / 0.120
	Min / Max	0.10 / 5.71	-0.56 / 0.57	1.11 / 4.73	-0.78 / 0.91
	Abnormal <sup>1</sup> , n (%)	5 (5.7%)		3 (6.5%)	
Visit 5	n / missing	78 / 1	78 / 1	42 / 0	42 / 0
	Mean (SD)	1.982 (0.8195)	-0.044 (0.3381)	2.215 (0.7795)	-0.043 (0.3009)
	Geometric mean	1.805		2.091	
	Median	1.805	-0.020	2.145	0.005
	IQR	0.690	0.330	1.030	0.340
	Q1 / Q3	1.590 / 2.280	-0.230 / 0.100	1.570 / 2.600	-0.160 / 0.180
	Min / Max	0.10 / 6.20	-1.92 / 0.70	1.14 / 4.29	-0.77 / 0.51
	Abnormal <sup>1</sup> , n (%)	5 (6.4%)		3 (7.1%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.16.1 Summary statistics, changes from baseline and frequencies of abnormal values of Anti-D-Mannan IgM [U/mL] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	73 / 0		37 / 0	
	Mean (SD)	22.348 (19.7320)		27.427 (28.5503)	
	Geometric mean	15.755		16.566	
	Median	16.400		14.300	
	IQR	16.900		22.000	
	Q1 / Q3	9.800 / 26.700		8.900 / 30.900	
	Min / Max	0.10 / 100.00		1.10 / 100.00	
	Abnormal <sup>1</sup> , n (%)	39 (53.4%)		17 (45.9%)	
Visit 3	n / missing	72 / 0	72 / 0	37 / 0	37 / 0
	Mean (SD)	19.625 (15.4581)	-2.175 (11.1535)	21.773 (16.8968)	-5.654 (21.0164)
	Geometric mean	15.043		15.568	
	Median	14.750	0.000	17.100	0.700
	IQR	15.600	8.150	27.000	11.800
	Q1 / Q3	8.750 / 24.350	-5.250 / 2.900	9.500 / 36.500	-6.300 / 5.500
	Min / Max	1.90 / 77.70	-47.30 / 23.90	2.40 / 80.70	-63.70 / 19.80
	Abnormal <sup>1</sup> , n (%)	36 (50.0%)		22 (59.5%)	
Visit 5	n / missing	65 / 2	65 / 2	34 / 0	34 / 0
	Mean (SD)	18.874 (16.4249)	-2.445 (17.1771)	20.926 (18.0967)	-8.071 (21.3012)
	Geometric mean	14.355		13.912	
	Median	13.800	-0.500	13.200	-2.050
	IQR	12.400	7.700	25.500	10.500
	Q1 / Q3	9.600 / 22.000	-4.800 / 2.900	6.300 / 31.800	-9.400 / 1.100
	Min / Max	1.80 / 100.00	-72.60 / 65.80	1.60 / 70.80	-68.20 / 21.60
	Abnormal <sup>1</sup> , n (%)	29 (44.6%)		15 (44.1%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

Patients from study centers 004 and 005 were not included due to different measurement unit.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.16.2 Summary statistics, changes from baseline and frequencies of abnormal values of Anti-D-Mannan IgM [NTU] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	10 / 7		6 / 3	
	Mean (SD)	4.149 (3.2963)		7.250 (7.0121)	
	Geometric mean	3.059		5.043	
	Median	2.430		5.450	
	IQR	5.000		6.580	
	Q1 / Q3	1.870 / 6.870		1.840 / 8.420	
	Min / Max	0.95 / 10.70		1.74 / 20.60	
	Abnormal <sup>1</sup> , n (%)	1 (10.0%)		1 (16.7%)	
Visit 3	n / missing	14 / 2	9 / 7	7 / 2	5 / 4
	Mean (SD)	3.306 (2.1410)	-0.826 (1.4927)	4.507 (2.0865)	-0.160 (1.8721)
	Geometric mean	2.811		4.082	
	Median	2.395	-0.510	4.500	0.350
	IQR	2.520	2.050	3.940	1.550
	Q1 / Q3	1.700 / 4.220	-1.780 / 0.270	2.710 / 6.650	-0.680 / 0.870
	Min / Max	1.36 / 8.92	-3.16 / 1.22	2.09 / 7.49	-3.11 / 1.77
	Abnormal <sup>1</sup> , n (%)	1 (7.1%)		0	
Visit 5	n / missing	10 / 2	8 / 4	7 / 1	5 / 3
	Mean (SD)	4.639 (2.3915)	0.839 (3.7923)	6.453 (4.7492)	-0.819 (7.1851)
	Geometric mean	4.063		2.295	
	Median	4.555	0.545	5.910	-1.836
	IQR	3.520	5.827	7.250	3.100
	Q1 / Q3	2.280 / 5.800	-1.815 / 4.012	2.750 / 10.000	-2.090 / 1.010
	Min / Max	1.78 / 8.36	-4.90 / 6.13	0.00 / 14.60	-10.60 / 9.42
	Abnormal <sup>1</sup> , n (%)	0		2 (28.6%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

Only values for patients from study centers 004 and 005 were included due to different measurement unit.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.17.1 Summary statistics, changes from baseline and frequencies of abnormal values of Anti-D-Glucan IgG [U/mL] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	73 / 0		37 / 0	
	Mean (SD)	27.332 (22.9171)		27.089 (25.1012)	
	Geometric mean	19.709		18.200	
	Median	20.000		20.300	
	IQR	23.200		24.900	
	Q1 / Q3	12.300 / 35.500		10.400 / 35.300	
	Min / Max	2.50 / 100.00		3.70 / 100.00	
	Abnormal <sup>1</sup> , n (%)	44 (60.3%)		19 (51.4%)	
Visit 3	n / missing	72 / 0	72 / 0	37 / 0	37 / 0
	Mean (SD)	29.353 (26.9274)	1.829 (18.5656)	24.424 (25.4732)	-2.665 (14.2688)
	Geometric mean	19.145		16.203	
	Median	20.200	-0.750	14.500	-0.800
	IQR	30.950	8.000	20.300	12.200
	Q1 / Q3	10.400 / 41.350	-4.850 / 3.150	7.500 / 27.800	-10.000 / 2.200
	Min / Max	0.70 / 100.00	-30.30 / 74.30	4.20 / 100.00	-38.50 / 44.70
	Abnormal <sup>1</sup> , n (%)	45 (62.5%)		18 (48.6%)	
Visit 5	n / missing	65 / 2	65 / 2	34 / 0	34 / 0
	Mean (SD)	22.565 (23.4687)	-4.580 (17.2566)	19.809 (20.3833)	-8.391 (17.9039)
	Geometric mean	13.890		13.545	
	Median	14.200	-5.000	13.750	-3.050
	IQR	20.800	11.900	19.600	15.700
	Q1 / Q3	7.300 / 28.100	-11.200 / 0.700	6.800 / 26.400	-14.600 / 1.100
	Min / Max	0.10 / 100.00	-50.10 / 51.80	2.80 / 96.60	-68.90 / 22.50
	Abnormal <sup>1</sup> , n (%)	30 (46.2%)		15 (44.1%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

Patients from study centers 004 and 005 were not included due to different measurement unit.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.17.2 Summary statistics, changes from baseline and frequencies of abnormal values of Anti-D-Glucan IgG [NTU] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	10 / 7		6 / 3	
	Mean (SD)	33.140 (15.1836)		34.217 (10.1907)	
	Geometric mean	30.580		33.037	
	Median	28.850		29.800	
	IQR	11.300		18.600	
	Q1 / Q3	24.900 / 36.200		27.800 / 46.400	
	Min / Max	14.40 / 70.00		23.80 / 47.70	
	Abnormal <sup>1</sup> , n (%)	6 (60.0%)		5 (83.3%)	
Visit 3	n / missing	15 / 1	9 / 7	8 / 1	5 / 4
	Mean (SD)	32.333 (7.8324)	-2.911 (7.5976)	28.238 (13.6414)	1.380 (3.3700)
	Geometric mean	31.501		25.308	
	Median	31.900	-1.300	29.250	1.700
	IQR	11.300	1.100	16.600	2.100
	Q1 / Q3	24.800 / 36.100	-1.500 / -0.400	17.050 / 33.650	-0.400 / 1.700
	Min / Max	23.40 / 47.30	-22.70 / 3.50	11.80 / 54.20	-2.60 / 6.50
	Abnormal <sup>1</sup> , n (%)	11 (73.3%)		5 (62.5%)	
Visit 5	n / missing	10 / 2	8 / 4	6 / 2	4 / 4
	Mean (SD)	27.630 (7.7873)	-7.950 (9.1748)	26.117 (7.6006)	-9.250 (7.4933)
	Geometric mean	26.766		25.025	
	Median	25.800	-5.600	29.800	-9.750
	IQR	5.700	4.950	13.300	12.400
	Q1 / Q3	21.800 / 27.500	-8.400 / -3.450	18.400 / 31.700	-15.450 / -3.050
	Min / Max	19.90 / 42.30	-29.40 / 0.70	14.70 / 32.30	-16.80 / -0.70
	Abnormal <sup>1</sup> , n (%)	7 (70.0%)		4 (66.7%)	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

Only values for patients from study centers 004 and 005 were included due to different measurement unit.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1

**Table 14.2.3.2.18 Summary statistics, changes from baseline and frequencies of abnormal values of phagocyte activity [%] by scheduled visit and treatment (FAS)**

Visit	Statistics	IMUNOR		PLACEBO	
		Value	Change from BL	Value	Change from BL
Visit 1	n / missing	83 / 7		42 / 4	
	Mean (SD)	97.286 (3.5053)		97.568 (3.3785)	
	Geometric mean	97.215		97.504	
	Median	98.000		98.850	
	IQR	2.000		2.200	
	Q1 / Q3	97.000 / 99.000		96.800 / 99.000	
	Min / Max	73.40 / 100.00		78.80 / 100.00	
	Abnormal <sup>1</sup> , n (%)	1 (1.2%)		0	
Visit 3	n / missing	83 / 5	82 / 6	41 / 5	40 / 6
	Mean (SD)	96.760 (4.5385)	-0.437 (5.5507)	96.951 (3.0612)	-0.517 (4.1638)
	Geometric mean	96.640		96.901	
	Median	98.000	0.000	98.000	-0.300
	IQR	3.000	2.000	3.240	2.495
	Q1 / Q3	96.000 / 99.000	-1.000 / 1.000	95.760 / 99.000	-1.495 / 1.000
	Min / Max	69.60 / 100.00	-30.30 / 23.50	84.00 / 100.00	-13.23 / 16.80
	Abnormal <sup>1</sup> , n (%)	1 (1.2%)		0	
Visit 5	n / missing	78 / 1	73 / 6	42 / 0	39 / 3
	Mean (SD)	95.721 (7.3766)	-0.431 (4.4820)	95.892 (4.0825)	-1.464 (4.4745)
	Geometric mean	95.311		95.802	
	Median	97.475	0.000	96.950	-0.640
	IQR	4.000	3.000	3.410	2.000
	Q1 / Q3	95.000 / 99.000	-2.000 / 1.000	94.990 / 98.400	-1.500 / 0.500
	Min / Max	42.77 / 100.00	-15.64 / 22.40	80.80 / 100.00	-19.00 / 8.20
	Abnormal <sup>1</sup> , n (%)	2 (2.6%)		0	

% = Percentage of subjects (calculated out of non-missing values in specified group); BL = Baseline; FAS = Full analysis set; IQR = Interquartile range; Max = Maximum; Min = Minimum; n = Number of subjects with available value; missing = number of subjects with missing value; Q1 = lower quartile (25th percentile); Q3 = upper quartile (75th percentile); SD = Standard deviation.

<sup>1</sup>Abnormal value classification is based on laboratory normal ranges. The normal ranges for individual measurements can be found in Patients' data listings. Normal ranges may vary over time and between different centers.

Produced on 17-JAN-2025

Page 1 of 1