

**Clinical trial results:****RANDOMIZED, DOUBLE-BLIND, PHASE 3 STUDY EVALUATING TAS-102 PLUS BEST SUPPORTIVE CARE (BSC) VERSUS PLACEBO PLUS BSC IN PATIENTS WITH METASTATIC GASTRIC CANCER REFRACTORY TO STANDARD TREATMENTS****Summary**

EudraCT number	2015-002683-16
Trial protocol	GB DE BE ES IE PT CZ PL IT
Global end of trial date	19 December 2019

Results information

Result version number	v1 (current)
This version publication date	12 April 2020
First version publication date	12 April 2020
Summary attachment (see zip file)	TAS-120-302 CSR Synopsis (2015-002683-16_CSR Synopsis_04Dec2018.pdf)

Trial information**Trial identification**

Sponsor protocol code	TO-TAS-102-302
-----------------------	----------------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02500043
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	Taiho Oncology, Inc
Sponsor organisation address	101 Carnegie Center, Suite 101, Princeton, New Jersey, United States, NJ 08540
Public contact	Simon Ruini, Associate Director, Regulatory Affairs, Taiho Pharma Europe, Ltd, +01 609 750 5300, SRuini@taiho.eu
Scientific contact	Simon Ruini, Associate Director, Regulatory Affairs, Taiho Pharma Europe, Ltd, +01 609 750 5300, SRuini@taiho.eu

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	28 September 2018
Is this the analysis of the primary completion data?	Yes
Primary completion date	30 April 2018
Global end of trial reached?	Yes
Global end of trial date	19 December 2019
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

Overall survival (OS)

Protection of trial subjects:

This study was designed and conducted in accordance with the Sponsor procedures, which comply with the ethical principles of Good Clinical Practice (GCP) as required by the major regulatory authorities and in accordance with the Declaration of Helsinki, ICH E6 Guideline for GCP, and local regulations.

The protocol and amendments, Informed Consent Form (ICF), and the Investigator's Brochure (IB) provided to the Investigators, and any other documents that pertained to patient information (eg, patient diaries), recruitment methods, and advertisements received Institutional Review Board (IRB)/Independent Ethics Committee (IEC) approval before the first patient was enrolled at the investigational site. When necessary, all protocol amendments and changes to the ICF were submitted by the Investigator to the IRB/IEC for approval. The Investigators notified the IRB/IEC of deviations from the protocol or serious adverse events occurring at the site according to local policies and IRB/IEC requirements, as well as other adverse event reports, in accordance with local procedures.

The Investigator or a designee under the Investigator's responsibility (according to applicable regulatory requirements) fully informed patients of all pertinent aspects of the clinical study. All participants were informed to the fullest extent possible about the study in a language and in terms they were able to understand.

Prior to participation in the trial, the written ICF was signed and personally dated by the patient or by the patient's legal representative and by the person who conducted the informed consent discussion. A copy of the signed and dated ICF was provided to the patient.

Background therapy:

Best Supportive Care according to local clinical practice i.e.:

- regimens including fluoropyrimidines, platinum derivatives, and either a taxane- and/or irinotecan-containing regimen;
- a anti-HER2+ therapy for patients whose tumors are HER2-neupositive (HER2+);
- pre- or post-operative adjuvant chemotherapy or chemoradiotherapy,

Evidence for comparator:

This present study was designed as a randomized, double-blind, Phase 3 study comparing TAS-102 plus best supportive care (BSC) to placebo plus BSC in patients with metastatic gastric cancer who have received at least 2 prior regimens for advanced disease and were refractory or unable to tolerate their last prior therapy.

A placebo-controlled design was considered appropriate as there are currently no standard therapies for patients with metastatic gastric cancer who have failed first- and second-line therapies.

Actual start date of recruitment	11 February 2016
Long term follow-up planned	Yes
Long term follow-up rationale	Safety, Efficacy
Long term follow-up duration	2 Years
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Italy: 66
Country: Number of subjects enrolled	United States: 26
Country: Number of subjects enrolled	Turkey: 43
Country: Number of subjects enrolled	France: 24
Country: Number of subjects enrolled	Japan: 73
Country: Number of subjects enrolled	Israel: 12
Country: Number of subjects enrolled	Russian Federation: 46
Country: Number of subjects enrolled	Belarus: 30
Country: Number of subjects enrolled	Poland: 14
Country: Number of subjects enrolled	Portugal: 39
Country: Number of subjects enrolled	Romania: 8
Country: Number of subjects enrolled	Spain: 40
Country: Number of subjects enrolled	United Kingdom: 45
Country: Number of subjects enrolled	Belgium: 13
Country: Number of subjects enrolled	Czech Republic: 9
Country: Number of subjects enrolled	Germany: 11
Country: Number of subjects enrolled	Ireland: 8
Worldwide total number of subjects	507
EEA total number of subjects	277

Notes:

Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	279
From 65 to 84 years	159
85 years and over	69

Subject disposition

Recruitment

Recruitment details:

Patients were randomized at 110 centers in 17 countries.

A total of 625 patients signed informed consent for participation in the study.

Pre-assignment

Screening details:

All patients had to complete the following study procedures prior to a confirmation of eligibility: -Medical History, -Histologic Confirmation, -Human Epidermal Growth Factor Receptor 2 (HER2) Status, -Physical Examination, -Baseline Signs and Symptoms, -Height, Vital Signs, Weight, -ECOG performance status and - Clinical Laboratory Evaluations

Pre-assignment period milestones

Number of subjects started	625 ^[1]
Intermediate milestone: Number of subjects	Screening - Confirmation of Eligibility: 507
Number of subjects completed	507

Pre-assignment subject non-completion reasons

Reason: Number of subjects	Screen failure: 111
Reason: Number of subjects	Consent withdrawn by subject: 1
Reason: Number of subjects	Other: 4
Reason: Number of subjects	Death: 1
Reason: Number of subjects	Protocol deviation: 1

Notes:

[1] - The number of subjects reported to have started the pre-assignment period are not the same as the worldwide number enrolled in the trial. It is expected that these numbers will be the same.

Justification: A total of 625 patients provided signed informed consent for participation in the study. Of these, 118 (18.9%) were screened but not randomized primarily due to screen failures.

Period 1

Period 1 title	Baseline Period
Is this the baseline period?	Yes
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Arm title	Baseline
------------------	----------

Arm description:

Baseline period; screening for participation in the study

Arm type	Screening
Investigational medicinal product name	TAS-102
Investigational medicinal product code	FTD, F3TdR, F3dThd
Other name	5-TRIFLUOROTHYIMIDINE
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

TAS-102 is formulated as an immediate-release film-coated tablet, which is supplied in 2 strengths:

- The 15 mg white, round tablet
- The 20 mg pale-red, round tablet

The total dose is 70 mg/m² milligram(s)/square meter, per day within 1 hour after completion of

morning and evening meals, for 5 days a week with 2 days rest for 2 weeks, followed by a 14-day rest. This treatment cycle will be repeated every 4 weeks.

Number of subjects in period 1	Baseline
Started	507
Sign ICF	507
Enrollment	507
Medical History	507
Histological Confirmation	507
HER2 status (if available)	507
Physical Examination	507
Baseline Signs & Symptoms	507
Height	507
Vital Signs & Weight	507
ECOG Performance Status	507
Hematology	507
Serum Chemistry	507
Urinalysis	507
Pregnancy Test	507
Tumor Measurements	507
Quality of Life Assessment	507
Concomitant Medications	507
Completed	507

Period 2

Period 2 title	Treatment Period
Is this the baseline period?	No
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Investigator, Subject

Blinding implementation details:

Eligible patients who met all of the inclusion and none of the exclusion criteria were centrally randomized (2:1) to TAS-102 plus BSC (experimental arm) or placebo plus BSC (control arm) using a

dynamic allocation method (biased coin) via an Interactive Voice/Web Response System (IXRS). Randomization was stratified by: region (Rest of World (ROW) vs Japan); ECOG performance status (0 vs 1); and prior treatment with ramucirumab (yes vs no).

Arms

Are arms mutually exclusive?	Yes
Arm title	TAS-102 plus BSC

Arm description:

Experimental arm, evaluating the efficacy and safety of TAS-102 (Investigational Medicinal Product) plus Best Supportive Care (BSC)

Arm type	Experimental
Investigational medicinal product name	TAS-102
Investigational medicinal product code	FTD, F3TdR, F3dThd
Other name	5-TRIFLUOROTHYMIDINE
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

TAS-102 is formulated as an immediate-release film-coated tablet, which is supplied in 2 strengths:

- The 15 mg white, round tablet
- The 20 mg pale-red, round tablet

The total dose is 70 mg/m² milligram(s)/square meter, per day within 1 hour after completion of morning and evening meals, for 5 days a week with 2 days rest for 2 weeks, followed by a 14-day rest. This treatment cycle will be repeated every 4 weeks.

Arm title	Placebo plus BSC
------------------	------------------

Arm description:

Experimental arm, evaluating the efficacy and safety of placebo plus Best Supportive Care (BSC)

Arm type	Placebo
Investigational medicinal product name	TAS-102
Investigational medicinal product code	FTD, F3TdR, F3dThd
Other name	5-TRIFLUOROTHYMIDINE
Pharmaceutical forms	Film-coated tablet
Routes of administration	Oral use

Dosage and administration details:

TAS-102 is formulated as an immediate-release film-coated tablet, which is supplied in 2 strengths:

- The 15 mg white, round tablet
- The 20 mg pale-red, round tablet

The total dose is 70 mg/m² milligram(s)/square meter, per day within 1 hour after completion of morning and evening meals, for 5 days a week with 2 days rest for 2 weeks, followed by a 14-day rest. This treatment cycle will be repeated every 4 weeks.

Number of subjects in period 2	TAS-102 plus BSC	Placebo plus BSC
Started	337	170
Randomization	337	170
Physical Examination	337	170
Vital Signs & Weight	337	170
ECOG Performance Status	337	170

Hematology	337	170
Serum Chemistry	337	170
Tumor Measurements	337	170
Quality of Life Assessment	337	170
Concomitant Medications	337	170
AE/SAE Assessment	337	170
TAS-102 or Placebo treatment	337	170
Survival Status	337	170
Completed	335	168
Not completed	2	2
Adverse event, serious fatal	1	-
Consent withdrawn by subject	-	2
Protocol deviation	1	-

Period 3

Period 3 title	End of Treatment Period
Is this the baseline period?	No
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Arm title	End of Study
Arm description:	
End of Treatment	
Arm type	No intervention
No investigational medicinal product assigned in this arm	

Number of subjects in period 3	End of Study
Started	503
Physical Examination	22
Vital Signs & Weight	22
ECOG Performance Status	22
Hematology	22
Serum Chemistry	22
Pregnancy Test	22
Tumor Measurements	22

Quality of Life Assessment	22
Concomitant Medications	22
AE/SAE Assessment	22
Survival Status	22
Completed	22
Not completed	481
Clinical progression	89
Adverse event, serious fatal	2
Consent withdrawn by subject	18
Physician decision	14
Radiological progression	302
Adverse event, non-fatal	44
Death	11
Protocol deviation	1

Baseline characteristics

Reporting groups

Reporting group title	Baseline Period
-----------------------	-----------------

Reporting group description: -

Reporting group values	Baseline Period	Total	
Number of subjects	507	507	
Age categorical			
Units: Subjects			
Adults (18-64 years)	279	279	
From 65-84 years	159	159	
85 years and over	69	69	
Age continuous			
Units: years			
arithmetic mean	62.5		
full range (min-max)	24 to 89	-	
Gender categorical			
Units: Subjects			
Female	138	138	
Male	369	369	
Race			
Units: Subjects			
White	357	357	
Black/African American	3	3	
Asian	80	80	
Not collectable	62	62	
Other	5	5	
Region			
Units: Subjects			
Japan	73	73	
United States	26	26	
European Union	408	408	
Baseline renal function			
Units: Subjects			
Normal (CrCl \geq 90 mL/min)	202	202	
Mild impairment (CrCl 60-89 mL/min)	212	212	
Moderate impairment (CrCl 30-59 mL/min)	86	86	
Severe impairment (CrCl < 30 mL/min)	3	3	
Not recorded	4	4	
Body Surface Area			
Units: m ²			
arithmetic mean	1.749		
full range (min-max)	1.20 to 2.52	-	

Subject analysis sets

Subject analysis set title	Intention to treat (ITT)
Subject analysis set type	Intention-to-treat
Subject analysis set description: ITT population - patients who were randomized	
Subject analysis set title	As Treated (AT) - TAS-102 + BSC
Subject analysis set type	Full analysis
Subject analysis set description: As Treated (AT) population for the TAS-102 + BSC group	
Subject analysis set title	As Treated (AT) - Placebo + BSC
Subject analysis set type	Full analysis
Subject analysis set description: As Treated population for the Placebo + BSC group	

Reporting group values	Intention to treat (ITT)	As Treated (AT) - TAS-102 + BSC	As Treated (AT) - Placebo + BSC
Number of subjects	507	337	170
Age categorical Units: Subjects			
Adults (18-64 years)	279	183	96
From 65-84 years	159	103	56
85 years and over	69	51	18
Age continuous Units: years			
arithmetic mean	62.5	62.8	62.0
full range (min-max)	24 to 89	24 to 89	32 to 82
Gender categorical Units: Subjects			
Female	138	85	53
Male	369	252	117
Race Units: Subjects			
White	357	244	113
Black/African American	3	1	2
Asian	80	51	29
Not collectable	62	38	24
Other	5	3	2
Region Units: Subjects			
Japan	73	46	27
United States	26	21	5
European Union	408	270	138
Baseline renal function Units: Subjects			
Normal (CrCl \geq 90 mL/min)	202	134	68
Mild impairment (CrCl 60-89 mL/min)	212	141	71
Moderate impairment (CrCl 30-59 mL/min)	86	58	28
Severe impairment (CrCl < 30 mL/min)	3	2	1
Not recorded	4	2	2

Body Surface Area			
Units: m2			
arithmetic mean	1.749	1.747	1.754
full range (min-max)	1.20 to 2.52	1.20 to 2.37	1.29 to 2.52

End points

End points reporting groups

Reporting group title	Baseline
-----------------------	----------

Reporting group description:

Baseline period; screening for participation in the study

Reporting group title	TAS-102 plus BSC
-----------------------	------------------

Reporting group description:

Experimental arm, evaluating the efficacy and safety of TAS-102 (Investigational Medicinal Product) plus Best Supportive Care (BSC)

Reporting group title	Placebo plus BSC
-----------------------	------------------

Reporting group description:

Experimental arm, evaluating the efficacy and safety of placebo plus Best Supportive Care (BSC)

Reporting group title	End of Study
-----------------------	--------------

Reporting group description:

End of Treatment

Subject analysis set title	Intention to treat (ITT)
----------------------------	--------------------------

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

ITT population - patients who were randomized

Subject analysis set title	As Treated (AT) - TAS-102 + BSC
----------------------------	---------------------------------

Subject analysis set type	Full analysis
---------------------------	---------------

Subject analysis set description:

As Treated (AT) population for the TAS-102 + BSC group

Subject analysis set title	As Treated (AT) - Placebo + BSC
----------------------------	---------------------------------

Subject analysis set type	Full analysis
---------------------------	---------------

Subject analysis set description:

As Treated population for the Placebo + BSC group

Primary: Overall Survival (OS)

End point title	Overall Survival (OS)
-----------------	-----------------------

End point description:

Overall Survival (OS) was the primary endpoint of this study and was defined as the time from the date of randomization to the death date. In the absence of death confirmation or for patients alive on the survival cut-off date, survival was censored at the date of last study follow-up or the cut-off date, whichever was earlier.

End point type	Primary
----------------	---------

End point timeframe:

The overall survival (OS) cut-off date used for the primary analysis was based on the survival data obtained through the date of the 384th death observed in the study (27 Mar 2018).

End point values	TAS-102 plus BSC	Placebo plus BSC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	337	170		
Units: Subjects				
Total patients	337	170		
Not censored (dead)	244	140		
Censored	93	30		
Overall Survival by Region: Japan	46	27		
Overall Survival by Region: Rest of World	291	143		
Overall Survival by ECOG status at baseline: 0	123	68		
Overall Survival by ECOG status at baseline: 1	214	102		
OS by Prior treatment with ramucirumab: Yes	114	55		
OS by Prior treatment with ramucirumab: No	223	115		

Statistical analyses

Statistical analysis title	Overall Survival (months)
Statistical analysis description:	
Overall Survival (OS) in the Intent to Treat (ITT) population will be compared between the 2 treatment groups using the stratified log-rank test.	
Comparison groups	TAS-102 plus BSC v Placebo plus BSC
Number of subjects included in analysis	507
Analysis specification	Pre-specified
Analysis type	equivalence
P-value	= 0.0003
Method	t-test, 2-sided
Parameter estimate	Hazard ratio (HR)
Point estimate	0.6917
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.5597
upper limit	0.8548
Variability estimate	Standard deviation

Secondary: Progression-Free Survival (PFS)

End point title	Progression-Free Survival (PFS)
------------------------	---------------------------------

End point description:

Progression free survival was defined as the time from the date of randomization until the date of the investigator-assessed radiological disease progression or death due to any cause. Patients who were alive with no disease progression as of the analysis cut-off date were censored at the date of the last tumor assessment. Patients who received non-study cancer treatment before disease progression, or patients with clinical but not radiological evidence of progression was censored at the date of the last evaluable tumor assessment before the non-study cancer treatment was initiated.

End point type	Secondary
----------------	-----------

End point timeframe:

PFS was defined as the time from the date of randomization until the date of the investigator-assessed radiological disease progression or death due to any cause as of the pre-specified cut-off date of 31 Mar 2018 for non-survival data.

End point values	TAS-102 plus BSC	Placebo plus BSC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	337	170		
Units: Subjects				
Total patients	337	170		
PFS Events	287	156		
Progressed	209	113		
Death	78	43		
Censored	50	14		
Discontinued follow-up	12	1		
Initiated anti-tumor therapy	8	6		
Missed visit (> 91 days since last response)	10	3		
Follow-up ongoing at the time of analysis	20	4		
PFS by Region: Japan	46	27		
PFS by Region: Rest of World	291	143		
PFS by ECOG Performance Status at Baseline: 0	123	68		
PFS by ECOG Performance Status at Baseline: 1	214	102		
PFS by Prior Treatment with Ramucirumab: Yes	114	55		
PFS by Prior Treatment with Ramucirumab: No	223	115		

Statistical analyses

Statistical analysis title	Progression-Free Survival
-----------------------------------	---------------------------

Statistical analysis description:

Progression-free survival (PFS) was defined as the time from the date of randomization until the date of the investigator-assessed radiological disease progression or death due to any cause as of the pre-specified cut-off date of 31 Mar 2018 for non-survival data.

Comparison groups	TAS-102 plus BSC v Placebo plus BSC
Number of subjects included in analysis	507
Analysis specification	Pre-specified
Analysis type	equivalence
P-value	= 0.0001
Method	t-test, 2-sided
Parameter estimate	Hazard ratio (HR)
Point estimate	0.5723

Confidence interval	
level	95 %
sides	2-sided
lower limit	0.4674
upper limit	0.7008
Variability estimate	Standard deviation

Adverse events

Adverse events information

Timeframe for reporting adverse events:

Adverse Events (AE) were reported from the first dose of study medication through the period of patient follow-up (30 days after the last dose of study medication or until the start of new anti-tumor therapy, whichever was earlier).

Adverse event reporting additional description:

All adverse event reporting were performed using the as-treated (AT) analysis population.

The Common Terminology Criteria for Adverse Events (CTCAE Version 4.03) terms was used to assess severity/provide the grade for each adverse event (AE) that was reported.

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	CTCAE
Dictionary version	4.03

Reporting groups

Reporting group title	TAS-120 Population
-----------------------	--------------------

Reporting group description:

Adverse Events occurring in subjects within the treatment (TAS-120) group

Reporting group title	Placebo Population
-----------------------	--------------------

Reporting group description:

Adverse Events occurring in subjects within the placebo group

Serious adverse events	TAS-120 Population	Placebo Population	
Total subjects affected by serious adverse events			
subjects affected / exposed	143 / 335 (42.69%)	70 / 168 (41.67%)	
number of deaths (all causes)	253	142	
number of deaths resulting from adverse events	46	20	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Metastases to central nervous system			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Tumour haemorrhage			
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	2 / 2	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Brain neoplasm			

subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lymphangiosis carcinomatosa			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Malignant ascites			
subjects affected / exposed	1 / 335 (0.30%)	2 / 168 (1.19%)	
occurrences causally related to treatment / all	1 / 1	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Neoplasm malignant			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Cancer pain			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ureteric cancer metastatic			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vascular disorders			
Shock haemorrhagic			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	2 / 2	0 / 0	
Lymphoedema			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Venous thrombosis			

subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypotension			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
General disorders and administration site conditions			
General physical health deterioration			
subjects affected / exposed	21 / 335 (6.27%)	15 / 168 (8.93%)	
occurrences causally related to treatment / all	21 / 21	15 / 15	
deaths causally related to treatment / all	17 / 17	11 / 11	
Fatigue			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyrexia			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Asthenia			
subjects affected / exposed	1 / 335 (0.30%)	3 / 168 (1.79%)	
occurrences causally related to treatment / all	1 / 1	3 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Disease progression			
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
Pain			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Malaise			

subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Performance status decreased			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Reproductive system and breast disorders			
Breast pain			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Endometrial hyperplasia			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Respiratory, thoracic and mediastinal disorders			
Pleural effusion			
subjects affected / exposed	5 / 335 (1.49%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	5 / 5	1 / 1	
deaths causally related to treatment / all	2 / 2	1 / 1	
Pulmonary embolism			
subjects affected / exposed	5 / 335 (1.49%)	2 / 168 (1.19%)	
occurrences causally related to treatment / all	5 / 5	2 / 2	
deaths causally related to treatment / all	3 / 3	0 / 0	
Dyspnoea			
subjects affected / exposed	4 / 335 (1.19%)	2 / 168 (1.19%)	
occurrences causally related to treatment / all	4 / 4	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia aspiration			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Respiratory failure subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Investigations			
Neutrophil count decreased subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood bilirubin increased subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
White blood cell count decreased subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bilirubin conjugated increased subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injury, poisoning and procedural complications			
Accidental overdose subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Clavicle fracture subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Thoracic vertebral fracture subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Cardiac disorders			
Acute coronary syndrome			
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	2 / 2	1 / 1	
deaths causally related to treatment / all	2 / 2	0 / 0	
Atrial fibrillation			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
Altered state of consciousness			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Cerebral haemorrhage			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Hemiparesis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Presyncope			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Transient ischaemic attack			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cerebral infarction			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Cerebrovascular accident			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Headache			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Depressed level of consciousness			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ischaemic stroke			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	13 / 335 (3.88%)	4 / 168 (2.38%)	
occurrences causally related to treatment / all	13 / 13	4 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pancytopenia			
subjects affected / exposed	7 / 335 (2.09%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	7 / 7	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Febrile neutropenia			
subjects affected / exposed	4 / 335 (1.19%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	4 / 4	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Neutropenia			
subjects affected / exposed	4 / 335 (1.19%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	4 / 4	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Disseminated intravascular			

coagulation			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Thrombocytopenia			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Myocardial infarction			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardio-respiratory arrest			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Ear and labyrinth disorders			
Vertigo			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			
Vomiting			
subjects affected / exposed	9 / 335 (2.69%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	9 / 9	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Abdominal pain			
subjects affected / exposed	8 / 335 (2.39%)	6 / 168 (3.57%)	
occurrences causally related to treatment / all	8 / 8	6 / 6	
deaths causally related to treatment / all	0 / 0	0 / 0	
Diarrhoea			
subjects affected / exposed	6 / 335 (1.79%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	6 / 6	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Dysphagia			
subjects affected / exposed	6 / 335 (1.79%)	2 / 168 (1.19%)	
occurrences causally related to treatment / all	6 / 6	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal haemorrhage			
subjects affected / exposed	4 / 335 (1.19%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	4 / 4	1 / 1	
deaths causally related to treatment / all	1 / 1	0 / 0	
Intestinal obstruction			
subjects affected / exposed	4 / 335 (1.19%)	3 / 168 (1.79%)	
occurrences causally related to treatment / all	4 / 4	3 / 3	
deaths causally related to treatment / all	0 / 0	1 / 1	
Ascites			
subjects affected / exposed	3 / 335 (0.90%)	7 / 168 (4.17%)	
occurrences causally related to treatment / all	3 / 3	7 / 7	
deaths causally related to treatment / all	0 / 0	1 / 1	
Gastric haemorrhage			
subjects affected / exposed	3 / 335 (0.90%)	3 / 168 (1.79%)	
occurrences causally related to treatment / all	3 / 3	3 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haematemesis			
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	3 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Small intestinal obstruction			
subjects affected / exposed	3 / 335 (0.90%)	2 / 168 (1.19%)	
occurrences causally related to treatment / all	3 / 3	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ileus			
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	2 / 2	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Obstruction gastric			

subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	2 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Upper gastrointestinal haemorrhage		
subjects affected / exposed	2 / 335 (0.60%)	2 / 168 (1.19%)
occurrences causally related to treatment / all	2 / 2	2 / 2
deaths causally related to treatment / all	1 / 1	0 / 0
Nausea		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	2 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Constipation		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Gastric ulcer haemorrhage		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Large intestinal obstruction		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Lower gastrointestinal haemorrhage		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Oesophageal obstruction		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Abdominal distension		

subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastric stenosis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Abdominal pain lower			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Abdominal pain upper			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal obstruction			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
Melaena			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Oesophageal pain			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Oesophagitis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pancreatitis			

subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Stomatitis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Subileus			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ulcerative gastritis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatobiliary disorders			
Hepatic failure			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	2 / 2	0 / 0	
Jaundice			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cholangitis			
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cholestasis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Jaundice cholestatic			

subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatitis toxic			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
Renal and urinary disorders			
Acute kidney injury			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal failure			
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hydronephrosis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Musculoskeletal and connective tissue disorders			
Joint swelling			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Back pain			
subjects affected / exposed	0 / 335 (0.00%)	3 / 168 (1.79%)	
occurrences causally related to treatment / all	0 / 0	3 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
Neutropenic sepsis			
subjects affected / exposed	4 / 335 (1.19%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	4 / 4	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Pneumonia			
subjects affected / exposed	4 / 335 (1.19%)	2 / 168 (1.19%)	
occurrences causally related to treatment / all	4 / 4	2 / 2	
deaths causally related to treatment / all	1 / 1	0 / 0	
Septic shock			
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	3 / 3	0 / 0	
deaths causally related to treatment / all	3 / 3	0 / 0	
Infection			
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	2 / 2	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Escherichia sepsis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
Influenza			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Salmonellosis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Sepsis			
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urosepsis			
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Clostridium difficile colitis			

subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Clostridium difficile infection			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Typhoid fever			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Upper respiratory tract infection			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Biliary sepsis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cellulitis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Peritonitis bacterial			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
Urinary tract infection			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Metabolism and nutrition disorders			
Decreased appetite			

subjects affected / exposed	11 / 335 (3.28%)	4 / 168 (2.38%)
occurrences causally related to treatment / all	11 / 11	4 / 4
deaths causally related to treatment / all	0 / 0	0 / 0
Failure to thrive		
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences causally related to treatment / all	2 / 2	1 / 1
deaths causally related to treatment / all	2 / 2	1 / 1
Dehydration		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences causally related to treatment / all	1 / 1	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0
Hypoalbuminaemia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Hypoglycaemia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Alkalosis hypochloraemic		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Cachexia		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences causally related to treatment / all	1 / 1	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0
Hyponatraemia		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences causally related to treatment / all	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0

Frequency threshold for reporting non-serious adverse events: 1 %

Non-serious adverse events	TAS-120 Population	Placebo Population	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	326 / 335 (97.31%)	157 / 168 (93.45%)	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Cancer pain			
subjects affected / exposed	3 / 335 (0.90%)	2 / 168 (1.19%)	
occurrences (all)	3	2	
Malignant ascites			
subjects affected / exposed	2 / 335 (0.60%)	2 / 168 (1.19%)	
occurrences (all)	2	2	
Metastases to central nervous system			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences (all)	2	0	
Tumour pain			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences (all)	2	0	
Brain neoplasm			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Lymphangiosis carcinomatosa			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Neoplasm malignant			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Lipoma			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Tumour associated fever			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Skin papilloma			

subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Ureteric cancer metastatic			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Vascular disorders			
Hypotension			
subjects affected / exposed	7 / 335 (2.09%)	2 / 168 (1.19%)	
occurrences (all)	7	2	
Hypertension			
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)	
occurrences (all)	3	0	
Shock haemorrhagic			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences (all)	2	0	
Deep vein thrombosis			
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)	
occurrences (all)	2	1	
Embolism			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences (all)	2	0	
Pallor			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences (all)	2	0	
Lymphoedema			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Phlebitis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Raynaud's phenomenon			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Thrombophlebitis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	

Venous thrombosis subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Hot flush subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	2 / 168 (1.19%) 2	
General disorders and administration site conditions			
Fatigue subjects affected / exposed occurrences (all)	89 / 335 (26.57%) 89	35 / 168 (20.83%) 35	
Asthenia subjects affected / exposed occurrences (all)	65 / 335 (19.40%) 65	40 / 168 (23.81%) 40	
Pyrexia subjects affected / exposed occurrences (all)	25 / 335 (7.46%) 25	8 / 168 (4.76%) 8	
General physical health deterioration subjects affected / exposed occurrences (all)	23 / 335 (6.87%) 23	17 / 168 (10.12%) 17	
Oedema peripheral subjects affected / exposed occurrences (all)	17 / 335 (5.07%) 17	12 / 168 (7.14%) 12	
Malaise subjects affected / exposed occurrences (all)	9 / 335 (2.69%) 9	9 / 168 (5.36%) 9	
Mucosal inflammation subjects affected / exposed occurrences (all)	8 / 335 (2.39%) 8	3 / 168 (1.79%) 3	
Oedema subjects affected / exposed occurrences (all)	8 / 335 (2.39%) 8	2 / 168 (1.19%) 2	
Pain subjects affected / exposed occurrences (all)	6 / 335 (1.79%) 6	8 / 168 (4.76%) 8	
Chills			

subjects affected / exposed	4 / 335 (1.19%)	0 / 168 (0.00%)
occurrences (all)	4	0
Chest pain		
subjects affected / exposed	3 / 335 (0.90%)	1 / 168 (0.60%)
occurrences (all)	3	1
Chest discomfort		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Disease progression		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Catheter site pain		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Drug intolerance		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Facial pain		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Feeling abnormal		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Gait disturbance		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Hypothermia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Localised oedema		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Non-cardiac chest pain		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Performance status decreased		

subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1	
Peripheral swelling subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Early satiety subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Face oedema subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Feeling cold subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Feeling hot subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Immune system disorders Perfume sensitivity subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Autoimmune disorder subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Reproductive system and breast disorders Breast pain subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Pelvic pain subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	2 / 168 (1.19%) 2	
Endometrial hyperplasia subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Pelvic discomfort			

subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Vaginal haemorrhage subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	3 / 168 (1.79%) 3	
Respiratory, thoracic and mediastinal disorders			
Dyspnoea subjects affected / exposed occurrences (all)	24 / 335 (7.16%) 24	17 / 168 (10.12%) 17	
Pleural effusion subjects affected / exposed occurrences (all)	13 / 335 (3.88%) 13	5 / 168 (2.98%) 5	
Cough subjects affected / exposed occurrences (all)	11 / 335 (3.28%) 11	6 / 168 (3.57%) 6	
Pulmonary embolism subjects affected / exposed occurrences (all)	10 / 335 (2.99%) 10	3 / 168 (1.79%) 3	
Epistaxis subjects affected / exposed occurrences (all)	4 / 335 (1.19%) 4	1 / 168 (0.60%) 1	
Hiccups subjects affected / exposed occurrences (all)	4 / 335 (1.19%) 4	5 / 168 (2.98%) 5	
Productive cough subjects affected / exposed occurrences (all)	4 / 335 (1.19%) 4	1 / 168 (0.60%) 1	
Dysphonia subjects affected / exposed occurrences (all)	3 / 335 (0.90%) 3	0 / 168 (0.00%) 0	
Pneumonia aspiration subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0	
Dyspnoea exertional			

subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Oropharyngeal pain		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Respiratory failure		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Nasal congestion		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Pleurisy		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Pleuritic pain		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Rhinitis allergic		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Rhinorrhoea		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Sputum discoloured		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Upper respiratory tract inflammation		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Wheezing		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Respiratory disorder		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Tachypnoea		

subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	2 / 168 (1.19%) 2	
Psychiatric disorders			
Insomnia			
subjects affected / exposed occurrences (all)	11 / 335 (3.28%) 11	10 / 168 (5.95%) 10	
Anxiety			
subjects affected / exposed occurrences (all)	9 / 335 (2.69%) 9	4 / 168 (2.38%) 4	
Depression			
subjects affected / exposed occurrences (all)	3 / 335 (0.90%) 3	2 / 168 (1.19%) 2	
Agitation			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	2 / 168 (1.19%) 2	
Confusional state			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	3 / 168 (1.79%) 3	
Delirium			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0	
Disturbance in social behaviour			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Drug abuse			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Hallucination			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Nervousness			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Product issues			
Device dislocation			

subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Investigations			
Neutrophil count decreased subjects affected / exposed occurrences (all)	51 / 335 (15.22%) 51	1 / 168 (0.60%) 1	
Blood alkaline phosphatase increased subjects affected / exposed occurrences (all)	30 / 335 (8.96%) 30	14 / 168 (8.33%) 14	
Platelet count decreased subjects affected / exposed occurrences (all)	28 / 335 (8.36%) 28	6 / 168 (3.57%) 6	
White blood cell count decreased subjects affected / exposed occurrences (all)	23 / 335 (6.87%) 23	0 / 168 (0.00%) 0	
Aspartate aminotransferase increased subjects affected / exposed occurrences (all)	21 / 335 (6.27%) 21	13 / 168 (7.74%) 13	
Weight decreased subjects affected / exposed occurrences (all)	20 / 335 (5.97%) 20	12 / 168 (7.14%) 12	
Blood bilirubin increased subjects affected / exposed occurrences (all)	17 / 335 (5.07%) 17	7 / 168 (4.17%) 7	
Alanine aminotransferase increased subjects affected / exposed occurrences (all)	16 / 335 (4.78%) 16	8 / 168 (4.76%) 8	
Blood creatinine increased subjects affected / exposed occurrences (all)	10 / 335 (2.99%) 10	0 / 168 (0.00%) 0	
Blood urea increased subjects affected / exposed occurrences (all)	7 / 335 (2.09%) 7	7 / 168 (4.17%) 7	
Gamma-glutamyltransferase increased			

subjects affected / exposed occurrences (all)	4 / 335 (1.19%) 4	5 / 168 (2.98%) 5
Hepatic enzyme increased subjects affected / exposed occurrences (all)	3 / 335 (0.90%) 3	0 / 168 (0.00%) 0
Haemoglobin decreased subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0
Lymphocyte count decreased subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0
Enzyme level increased subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0
Protein total decreased subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0
Red blood cell count decreased subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0
Bilirubin conjugated increased subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	2 / 168 (1.19%) 2
Blood bilirubin subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Vital capacity abnormal subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Alanine aminotransferase subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1
Aspartate aminotransferase subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Blood bilirubin unconjugated		

increased		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Blood creatinine decreased		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Blood potassium increased		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Creatinine renal clearance decreased		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Haematocrit decreased		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
International normalised ratio increased		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Neutrophil count		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Red cell distribution width increased		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Blood iron decreased		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Liver function test increased		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Platelet count increased		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Weight increased		

subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Injury, poisoning and procedural complications			
Fall			
subjects affected / exposed occurrences (all)	3 / 335 (0.90%) 3	3 / 168 (1.79%) 3	
Accidental overdose			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0	
Foreign body in gastrointestinal tract			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Procedural pain			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1	
Radiation injury			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Transfusion reaction			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Clavicle fracture			
subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Thoracic vertebral fracture			
subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Congenital, familial and genetic disorders			
Hydrocele			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Cardiac disorders			
Palpitations			
subjects affected / exposed occurrences (all)	6 / 335 (1.79%) 6	2 / 168 (1.19%) 2	
Acute coronary syndrome			

subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences (all)	2	1
Myocardial infarction		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Tachycardia		
subjects affected / exposed	2 / 335 (0.60%)	2 / 168 (1.19%)
occurrences (all)	2	2
Cardio-respiratory arrest		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Angina pectoris		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Atrial fibrillation		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Atrial flutter		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Bradycardia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Cardiovascular insufficiency		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Mitral valve incompetence		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Pericardial effusion		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Atrial tachycardia		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Cardiac disorder		

subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Sinus tachycardia			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Nervous system disorders			
Dysgeusia			
subjects affected / exposed	11 / 335 (3.28%)	1 / 168 (0.60%)	
occurrences (all)	11	1	
Dizziness			
subjects affected / exposed	8 / 335 (2.39%)	4 / 168 (2.38%)	
occurrences (all)	8	4	
Paraesthesia			
subjects affected / exposed	8 / 335 (2.39%)	0 / 168 (0.00%)	
occurrences (all)	8	0	
Headache			
subjects affected / exposed	7 / 335 (2.09%)	4 / 168 (2.38%)	
occurrences (all)	7	4	
Somnolence			
subjects affected / exposed	5 / 335 (1.49%)	1 / 168 (0.60%)	
occurrences (all)	5	1	
Neuropathy peripheral			
subjects affected / exposed	4 / 335 (1.19%)	1 / 168 (0.60%)	
occurrences (all)	4	1	
Lethargy			
subjects affected / exposed	2 / 335 (0.60%)	3 / 168 (1.79%)	
occurrences (all)	2	3	
Peripheral sensory neuropathy			
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)	
occurrences (all)	2	0	
Altered state of consciousness			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Cerebral haemorrhage			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	

Hemiparesis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Presyncope		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Transient ischaemic attack		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Agnosia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Amnesia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Burning sensation		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Cerebral infarction		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Cerebrovascular accident		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Myoclonus		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Neuralgia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Peroneal nerve palsy		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Sciatica		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0

Depressed level of consciousness subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Encephalopathy subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Ischaemic stroke subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Monoplegia subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Blood and lymphatic system disorders			
Anaemia subjects affected / exposed occurrences (all)	149 / 335 (44.48%) 149	32 / 168 (19.05%) 32	
Neutropenia subjects affected / exposed occurrences (all)	129 / 335 (38.51%) 129	6 / 168 (3.57%) 6	
Leukopenia subjects affected / exposed occurrences (all)	57 / 335 (17.01%) 57	3 / 168 (1.79%) 3	
Thrombocytopenia subjects affected / exposed occurrences (all)	33 / 335 (9.85%) 33	2 / 168 (1.19%) 2	
Lymphopenia subjects affected / exposed occurrences (all)	20 / 335 (5.97%) 20	8 / 168 (4.76%) 8	
Pancytopenia subjects affected / exposed occurrences (all)	7 / 335 (2.09%) 7	0 / 168 (0.00%) 0	
Febrile neutropenia subjects affected / exposed occurrences (all)	6 / 335 (1.79%) 6	0 / 168 (0.00%) 0	
Disseminated intravascular coagulation			

subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0	
Bone marrow failure subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Lymph node pain subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Lymphadenopathy subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Neutrophilia subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Ear and labyrinth disorders			
Vertigo subjects affected / exposed occurrences (all)	4 / 335 (1.19%) 4	1 / 168 (0.60%) 1	
Ear discomfort subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Ear pain subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Eye disorders			
Dry eye subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Eye irritation subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Eye pain subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Eye pruritus			

subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Lacrimation increased subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Visual impairment subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Gastrointestinal disorders			
Nausea subjects affected / exposed occurrences (all)	124 / 335 (37.01%) 124	53 / 168 (31.55%) 53	
Vomiting subjects affected / exposed occurrences (all)	83 / 335 (24.78%) 83	34 / 168 (20.24%) 34	
Diarrhoea subjects affected / exposed occurrences (all)	76 / 335 (22.69%) 76	24 / 168 (14.29%) 24	
Abdominal pain subjects affected / exposed occurrences (all)	54 / 335 (16.12%) 54	31 / 168 (18.45%) 31	
Constipation subjects affected / exposed occurrences (all)	45 / 335 (13.43%) 45	25 / 168 (14.88%) 25	
Abdominal pain upper subjects affected / exposed occurrences (all)	22 / 335 (6.57%) 22	15 / 168 (8.93%) 15	
Dysphagia subjects affected / exposed occurrences (all)	20 / 335 (5.97%) 20	8 / 168 (4.76%) 8	
Ascites subjects affected / exposed occurrences (all)	19 / 335 (5.67%) 19	16 / 168 (9.52%) 16	
Stomatitis subjects affected / exposed occurrences (all)	15 / 335 (4.48%) 15	4 / 168 (2.38%) 4	

Abdominal distension		
subjects affected / exposed	13 / 335 (3.88%)	9 / 168 (5.36%)
occurrences (all)	13	9
Gastrointestinal haemorrhage		
subjects affected / exposed	5 / 335 (1.49%)	1 / 168 (0.60%)
occurrences (all)	5	1
Intestinal obstruction		
subjects affected / exposed	5 / 335 (1.49%)	4 / 168 (2.38%)
occurrences (all)	5	4
Haematemesis		
subjects affected / exposed	5 / 335 (1.49%)	0 / 168 (0.00%)
occurrences (all)	5	0
Dyspepsia		
subjects affected / exposed	5 / 335 (1.49%)	3 / 168 (1.79%)
occurrences (all)	5	3
Gastroesophageal reflux disease		
subjects affected / exposed	4 / 335 (1.19%)	2 / 168 (1.19%)
occurrences (all)	4	2
Gastric haemorrhage		
subjects affected / exposed	3 / 335 (0.90%)	4 / 168 (2.38%)
occurrences (all)	3	4
Ileus		
subjects affected / exposed	3 / 335 (0.90%)	1 / 168 (0.60%)
occurrences (all)	3	1
Small intestinal obstruction		
subjects affected / exposed	3 / 335 (0.90%)	2 / 168 (1.19%)
occurrences (all)	3	2
Melaena		
subjects affected / exposed	3 / 335 (0.90%)	1 / 168 (0.60%)
occurrences (all)	3	1
Abdominal discomfort		
subjects affected / exposed	3 / 335 (0.90%)	1 / 168 (0.60%)
occurrences (all)	3	1
Abdominal pain lower		
subjects affected / exposed	3 / 335 (0.90%)	3 / 168 (1.79%)
occurrences (all)	3	3

Dry mouth		
subjects affected / exposed	3 / 335 (0.90%)	4 / 168 (2.38%)
occurrences (all)	3	4
Obstruction gastric		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Upper gastrointestinal haemorrhage		
subjects affected / exposed	2 / 335 (0.60%)	2 / 168 (1.19%)
occurrences (all)	2	2
Impaired gastric emptying		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Lower gastrointestinal haemorrhage		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Flatulence		
subjects affected / exposed	2 / 335 (0.60%)	2 / 168 (1.19%)
occurrences (all)	2	2
Haemorrhoidal haemorrhage		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Odynophagia		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Oesophageal pain		
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences (all)	2	1
Rectal haemorrhage		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Toothache		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Gastric ulcer haemorrhage		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0

Large intestinal obstruction subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Oesophageal obstruction subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Anal inflammation subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Anal pruritus subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Aphthous ulcer subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Cheilitis subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1
Dyschezia subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Epigastric discomfort subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1
Eructation subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Faeces discoloured subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	2 / 168 (1.19%) 2
Gastric stenosis subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1
Haematochezia subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0

Haemorrhoids			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Intra-abdominal fluid collection			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Mouth haemorrhage			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Proctalgia			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Gastrointestinal obstruction			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Oesophagitis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Pancreatitis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Subileus			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Ulcerative gastritis			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Hepatobiliary disorders			
Hyperbilirubinaemia			
subjects affected / exposed	10 / 335 (2.99%)	3 / 168 (1.79%)	
occurrences (all)	10	3	
Jaundice			
subjects affected / exposed	4 / 335 (1.19%)	2 / 168 (1.19%)	
occurrences (all)	4	2	
Liver disorder			

subjects affected / exposed	4 / 335 (1.19%)	0 / 168 (0.00%)
occurrences (all)	4	0
Hepatic failure		
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences (all)	2	1
Cholangitis		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Cholestasis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Hypertransaminasaemia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Jaundice cholestatic		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Biliary colic		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Hepatic function abnormal		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Hepatic pain		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Portal vein thrombosis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Cholecystitis acute		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Hepatitis toxic		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Hepatomegaly		

subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Skin and subcutaneous tissue disorders			
Alopecia			
subjects affected / exposed occurrences (all)	12 / 335 (3.58%) 12	1 / 168 (0.60%) 1	
Pruritus			
subjects affected / exposed occurrences (all)	8 / 335 (2.39%) 8	2 / 168 (1.19%) 2	
Dry skin			
subjects affected / exposed occurrences (all)	5 / 335 (1.49%) 5	1 / 168 (0.60%) 1	
Rash			
subjects affected / exposed occurrences (all)	4 / 335 (1.19%) 4	1 / 168 (0.60%) 1	
Night sweats			
subjects affected / exposed occurrences (all)	3 / 335 (0.90%) 3	0 / 168 (0.00%) 0	
Decubitus ulcer			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	1 / 168 (0.60%) 1	
Dermatitis			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0	
Nail disorder			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0	
Palmar-plantar erythrodysesthesia syndrome			
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	0 / 168 (0.00%) 0	
Skin lesion			
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Acne			

subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Blister subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Dermatitis allergic subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Hand dermatitis subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Hyperhidrosis subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Onychoclasia subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Petechiae subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Urticaria subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Xeroderma subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Nail discolouration subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Rash maculo-papular subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Renal and urinary disorders Dysuria subjects affected / exposed occurrences (all)	5 / 335 (1.49%) 5	3 / 168 (1.79%) 3	

Acute kidney injury		
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences (all)	2	1
Choluria		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Proteinuria		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Urinary retention		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Albuminuria		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Leukocyturia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Micturition disorder		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Nocturia		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Pollakiuria		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Renal failure		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Urinary incontinence		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Haematuria		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1

Hydronephrosis			
subjects affected / exposed	0 / 335 (0.00%)	2 / 168 (1.19%)	
occurrences (all)	0	2	
Microalbuminuria			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Renal colic			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Renal pain			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	25 / 335 (7.46%)	11 / 168 (6.55%)	
occurrences (all)	25	11	
Arthralgia			
subjects affected / exposed	6 / 335 (1.79%)	2 / 168 (1.19%)	
occurrences (all)	6	2	
Musculoskeletal pain			
subjects affected / exposed	5 / 335 (1.49%)	2 / 168 (1.19%)	
occurrences (all)	5	2	
Myalgia			
subjects affected / exposed	4 / 335 (1.19%)	0 / 168 (0.00%)	
occurrences (all)	4	0	
Groin pain			
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)	
occurrences (all)	3	0	
Bone pain			
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)	
occurrences (all)	3	0	
Pain in extremity			
subjects affected / exposed	3 / 335 (0.90%)	1 / 168 (0.60%)	
occurrences (all)	3	1	
Muscle atrophy			

subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Spinal pain		
subjects affected / exposed	2 / 335 (0.60%)	2 / 168 (1.19%)
occurrences (all)	2	2
Soft tissue disorder		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Bone swelling		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Flank pain		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Intervertebral disc compression		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Intervertebral disc disorder		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Joint swelling		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Limb discomfort		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Muscular weakness		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Musculoskeletal chest pain		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Musculoskeletal stiffness		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Neck pain		

subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Osteoarthritis			
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)	
occurrences (all)	1	1	
Osteoporosis			
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)	
occurrences (all)	1	0	
Muscle spasms			
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)	
occurrences (all)	0	1	
Tumour haemorrhage			
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)	
occurrences (all)	2	1	
Infections and infestations			
Upper respiratory tract infection			
subjects affected / exposed	9 / 335 (2.69%)	0 / 168 (0.00%)	
occurrences (all)	9	0	
Urinary tract infection			
subjects affected / exposed	9 / 335 (2.69%)	5 / 168 (2.98%)	
occurrences (all)	9	5	
Pneumonia			
subjects affected / exposed	8 / 335 (2.39%)	3 / 168 (1.79%)	
occurrences (all)	8	3	
Oral candidiasis			
subjects affected / exposed	6 / 335 (1.79%)	2 / 168 (1.19%)	
occurrences (all)	6	2	
Nasopharyngitis			
subjects affected / exposed	5 / 335 (1.49%)	0 / 168 (0.00%)	
occurrences (all)	5	0	
Neutropenic sepsis			
subjects affected / exposed	4 / 335 (1.19%)	0 / 168 (0.00%)	
occurrences (all)	4	0	
Septic shock			
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)	
occurrences (all)	3	0	

Conjunctivitis		
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)
occurrences (all)	3	0
Cystitis		
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)
occurrences (all)	3	0
Lower respiratory tract infection		
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)
occurrences (all)	3	0
Respiratory tract infection		
subjects affected / exposed	3 / 335 (0.90%)	2 / 168 (1.19%)
occurrences (all)	3	2
Rhinitis		
subjects affected / exposed	3 / 335 (0.90%)	0 / 168 (0.00%)
occurrences (all)	3	0
Infection		
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences (all)	2	1
Influenza		
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences (all)	2	1
Bronchitis		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Herpes zoster		
subjects affected / exposed	2 / 335 (0.60%)	1 / 168 (0.60%)
occurrences (all)	2	1
Tonsillitis		
subjects affected / exposed	2 / 335 (0.60%)	0 / 168 (0.00%)
occurrences (all)	2	0
Escherichia sepsis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Salmonellosis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0

Sepsis		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Urosepsis		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Angular cheilitis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Cellulitis		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Clostridium difficile colitis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Clostridium difficile infection		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Erysipelas		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Gastrointestinal infection		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Gingival abscess		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Herpes virus infection		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Laryngitis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Localised infection		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0

Lung infection		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Pyuria		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Sinusitis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Skin infection		
subjects affected / exposed	1 / 335 (0.30%)	1 / 168 (0.60%)
occurrences (all)	1	1
Staphylococcal infection		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Typhoid fever		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Vulvovaginal candidiasis		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Bacterial infection		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Biliary sepsis		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Candida infection		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Gingivitis		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Oesophageal candidiasis		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1

Peritonitis bacterial subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Tracheobronchitis subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Urinary tract infection bacterial subjects affected / exposed occurrences (all)	0 / 335 (0.00%) 0	1 / 168 (0.60%) 1	
Blood alkaline phosphatase subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0	
Metabolism and nutrition disorders			
Decreased appetite subjects affected / exposed occurrences (all)	115 / 335 (34.33%) 115	52 / 168 (30.95%) 52	
Hypoalbuminaemia subjects affected / exposed occurrences (all)	22 / 335 (6.57%) 22	10 / 168 (5.95%) 10	
Hypokalaemia subjects affected / exposed occurrences (all)	9 / 335 (2.69%) 9	3 / 168 (1.79%) 3	
Hyperglycaemia subjects affected / exposed occurrences (all)	9 / 335 (2.69%) 9	5 / 168 (2.98%) 5	
Hypocalcaemia subjects affected / exposed occurrences (all)	9 / 335 (2.69%) 9	1 / 168 (0.60%) 1	
Hypomagnesaemia subjects affected / exposed occurrences (all)	7 / 335 (2.09%) 7	3 / 168 (1.79%) 3	
Hyponatraemia subjects affected / exposed occurrences (all)	5 / 335 (1.49%) 5	8 / 168 (4.76%) 8	
Dehydration			

subjects affected / exposed occurrences (all)	5 / 335 (1.49%) 5	3 / 168 (1.79%) 3
Cachexia		
subjects affected / exposed occurrences (all)	4 / 335 (1.19%) 4	1 / 168 (0.60%) 1
Failure to thrive		
subjects affected / exposed occurrences (all)	2 / 335 (0.60%) 2	1 / 168 (0.60%) 1
Hyperkalaemia		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	3 / 168 (1.79%) 3
Hypoglycaemia		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Hypophagia		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Alkalosis hypochloraemic		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Electrolyte imbalance		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Feeding intolerance		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	0 / 168 (0.00%) 0
Hyperuricaemia		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1
Hypophosphataemia		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1
Hypoproteinaemia		
subjects affected / exposed occurrences (all)	1 / 335 (0.30%) 1	1 / 168 (0.60%) 1
Iron deficiency		

subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Malnutrition		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Vitamin D deficiency		
subjects affected / exposed	1 / 335 (0.30%)	0 / 168 (0.00%)
occurrences (all)	1	0
Hypercalcaemia		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1
Hypercreatininaemia		
subjects affected / exposed	0 / 335 (0.00%)	1 / 168 (0.60%)
occurrences (all)	0	1

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
19 February 2016	Various sections of the protocol was amended (namely Inclusion/Exclusion criteria etc) in response to Health Authority requests and in order to make changes for consistency and/or clarification.
05 May 2016	The following sections of the study protocol were updated in response to Health Authority requests, and in order to make changes for consistency and/or clarification: <ul style="list-style-type: none">- Section 8.8.1, Prohibited Medications and Therapies- Section 9.2.4.3, Dose Modification in Response to Hematologic Toxicities- Section 9.5, Study Drug Accountability- Section 12.1.1, Adverse Events

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported