



## Clinical trial results:

### A Phase III, Open-Label, Randomized Study of Atezolizumab (MPDL3280A, Anti-Pd-L1 Antibody) in Combination With Carboplatin or Cisplatin + Pemetrexed Compared With Carboplatin or Cisplatin + Pemetrexed in Patients Who Are Chemotherapy-Naive and Have Stage IV Non-Squamous Non-Small Cell Lung Cancer

#### Summary

EudraCT number	2015-003605-42
Trial protocol	ES RO BE LT SK PT HU AT LV FR NL HR IT
Global end of trial date	

#### Results information

Result version number	v1
This version publication date	23 July 2020
First version publication date	23 July 2020

#### Trial information

##### Trial identification

Sponsor protocol code	GO29438
-----------------------	---------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02657434
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	Hoffmann-La Roche
Sponsor organisation address	Grenzacherstrasse 124, Basel, Switzerland,
Public contact	F. Hoffmann-La Roche AG, F. Hoffmann-La Roche AG, 41 616878333, global.trial_information@roche.com
Scientific contact	F. Hoffmann-La Roche AG, F. Hoffmann-La Roche AG, 41 616878333, global.trial_information@roche.com

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Interim
Date of interim/final analysis	18 July 2019
Is this the analysis of the primary completion data?	Yes
Primary completion date	18 July 2019
Global end of trial reached?	No

Notes:

## General information about the trial

Main objective of the trial:

The main objective of this study is to evaluate the efficacy and safety of atezolizumab in combination with carboplatin or cisplatin + pemetrexed compared with carboplatin or cisplatin+ pemetrexed in subjects who are chemotherapy-naïve and have Stage IV non-squamous non-small cell lung cancer (NSCLC).

Protection of trial subjects:

All study subjects were required to read and sign an Informed Consent Form.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	07 April 2016
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Argentina: 3
Country: Number of subjects enrolled	Australia: 25
Country: Number of subjects enrolled	Austria: 1
Country: Number of subjects enrolled	Belgium: 6
Country: Number of subjects enrolled	Bulgaria: 1
Country: Number of subjects enrolled	Chile: 12
Country: Number of subjects enrolled	Spain: 118
Country: Number of subjects enrolled	France: 47
Country: Number of subjects enrolled	United Kingdom: 40
Country: Number of subjects enrolled	Hungary: 17
Country: Number of subjects enrolled	Ireland: 2
Country: Number of subjects enrolled	Israel: 9
Country: Number of subjects enrolled	Italy: 30
Country: Number of subjects enrolled	Japan: 101
Country: Number of subjects enrolled	Korea, Republic of: 14
Country: Number of subjects enrolled	Lithuania: 3
Country: Number of subjects enrolled	Latvia: 5
Country: Number of subjects enrolled	Malaysia: 7
Country: Number of subjects enrolled	Netherlands: 12
Country: Number of subjects enrolled	Peru: 1
Country: Number of subjects enrolled	Portugal: 8
Country: Number of subjects enrolled	Romania: 5

Country: Number of subjects enrolled	Russian Federation: 9
Country: Number of subjects enrolled	Taiwan: 10
Country: Number of subjects enrolled	Ukraine: 33
Country: Number of subjects enrolled	United States: 59
Worldwide total number of subjects	578
EEA total number of subjects	295

Notes:

---

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	321
From 65 to 84 years	256
85 years and over	1

## Subject disposition

### Recruitment

Recruitment details: -

### Pre-assignment

Screening details:

This study included chemotherapy-naïve subjects with histologically or cytologically confirmed Stage IV non-squamous non-small cell lung cancer (NSCLC).

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Not blinded

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Arm B (Carboplatin or Cisplatin + Pemetrexed)

Arm description:

Participants received IV infusion of 500 mg/m<sup>2</sup> pemetrexed on Day 1 q3w, and as per investigator's choice of either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain AUC =6 mg/mL/min or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period for 4 or 6 cycles (Cycle length=21 days). Participants who did not experience disease progression during the induction phase began maintenance therapy. Participants will receive IV infusion of 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

Arm type	Active comparator
Investigational medicinal product name	Carboplatin
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

Carboplatin - induction treatment: AUC of 6 mg/mL/min by IV infusion q3w for 4 or 6 cycles

Investigational medicinal product name	Cisplatin
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

Cisplatin - induction treatment: 75 mg/m<sup>2</sup> by IV infusion q3w for 4 or 6 cycles

Investigational medicinal product name	Pemetrexed
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

Pemetrexed - induction treatment: 500 mg/m<sup>2</sup> q3w by IV infusion for 4 or 6 cycles.

Pemetrexed - maintenance treatment: 500 mg/m<sup>2</sup> q3w by IV infusion until progressive disease.

<b>Arm title</b>	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
------------------	--

#### Arm description:

Participants received intravenous (IV) infusion of 1200 milligrams (mg) of atezolizumab on Day 1 every 3 weeks (q3w), IV infusion of 500 milligrams per meter square (mg/m<sup>2</sup>) pemetrexed on Day 1 q3w, and as per investigator's choice either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain area under concentration versus time (AUC) = 6 milligrams per milliliter per minute (mg/mL/min) or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period of 4 or 6 cycles (Cycle length=21 days). Participants who experienced clinical benefit during the induction phase began maintenance therapy. Participants will receive IV infusion of 1200 mg of atezolizumab and 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

Arm type	Experimental
Investigational medicinal product name	Atezolizumab
Investigational medicinal product code	
Other name	Tecentriq
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

#### Dosage and administration details:

Atezolizumab - induction treatment: 1200 mg by IV infusion for 4 – 6 cycles of 21 days (q3w). Atezolizumab - maintenance treatment: 1200 mg by IV infusion q3w until progressive disease or loss of clinical benefit.

Investigational medicinal product name	Carboplatin
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

#### Dosage and administration details:

Carboplatin - induction treatment: AUC of 6 mg/mL/min by IV infusion q3w for 4 or 6 cycles

Investigational medicinal product name	Cisplatin
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

#### Dosage and administration details:

Cisplatin - induction treatment: 75 mg/m<sup>2</sup> by IV infusion q3w for 4 or 6 cycles

Investigational medicinal product name	Pemetrexed
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

#### Dosage and administration details:

Pemetrexed - induction treatment: 500 mg/m<sup>2</sup> q3w by IV infusion for 4 or 6 cycles.  
Pemetrexed - maintenance treatment: 500 mg/m<sup>2</sup> q3w by IV infusion until progressive disease.

Number of subjects in period 1	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
Started	286	292
Completed	0	0
Not completed	286	292

Adverse event, serious fatal	177	183
Consent withdrawn by subject	27	16
On-going in study	77	91
Lost to follow-up	2	2
Randomization by error	2	-
Protocol deviation	1	-

## Baseline characteristics

### Reporting groups

Reporting group title	Arm B (Carboplatin or Cisplatin + Pemetrexed)
-----------------------	---

Reporting group description:

Participants received IV infusion of 500 mg/m<sup>2</sup> pemetrexed on Day 1 q3w, and as per investigator's choice of either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain AUC = 6 mg/mL/min or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period for 4 or 6 cycles (Cycle length=21 days). Participants who did not experience disease progression during the induction phase began maintenance therapy. Participants will receive IV infusion of 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

Reporting group title	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
-----------------------	--

Reporting group description:

Participants received intravenous (IV) infusion of 1200 milligrams (mg) of atezolizumab on Day 1 every 3 weeks (q3w), IV infusion of 500 milligrams per meter square (mg/m<sup>2</sup>) pemetrexed on Day 1 q3w, and as per investigator's choice either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain area under concentration versus time (AUC) = 6 milligrams per milliliter per minute (mg/mL/min) or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period of 4 or 6 cycles (Cycle length=21 days). Participants who experienced clinical benefit during the induction phase began maintenance therapy. Participants will receive IV infusion of 1200 mg of atezolizumab and 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

Reporting group values	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)	Total
Number of subjects	286	292	578
Age categorical Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	0	0	0
Infants and toddlers (28 days-23 months)	0	0	0
Children (2-11 years)	0	0	0
Adolescents (12-17 years)	0	0	0
Adults (18-64 years)	168	153	321
From 65-84 years	118	138	256
85 years and over	0	1	1
Age Continuous Units: Years			
arithmetic mean	61.8	63.3	-
standard deviation	± 9.3	± 9.4	-
Sex: Female, Male Units: Participants			
Female	94	100	194
Male	192	192	384
Ethnicity (NIH/OMB) Units: Subjects			
Hispanic or Latino	21	17	38
Not Hispanic or Latino	241	243	484
Unknown or Not Reported	24	32	56

Race (NIH/OMB)			
Units: Subjects			
American Indian or Alaska Native	1	1	2
Asian	65	71	136
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	4	2	6
White	203	193	396
More than one race	0	0	0
Unknown or Not Reported	13	25	38



## End points

### End points reporting groups

Reporting group title	Arm B (Carboplatin or Cisplatin + Pemetrexed)
-----------------------	---

Reporting group description:

Participants received IV infusion of 500 mg/m<sup>2</sup> pemetrexed on Day 1 q3w, and as per investigator's choice of either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain AUC =6 mg/mL/min or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period for 4 or 6 cycles (Cycle length=21 days). Participants who did not experience disease progression during the induction phase began maintenance therapy. Participants will receive IV infusion of 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

Reporting group title	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
-----------------------	--

Reporting group description:

Participants received intravenous (IV) infusion of 1200 milligrams (mg) of atezolizumab on Day 1 every 3 weeks (q3w), IV infusion of 500 milligrams per meter square (mg/m<sup>2</sup>) pemetrexed on Day 1 q3w, and as per investigator's choice either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain area under concentration versus time (AUC) = 6 milligrams per milliliter per minute (mg/mL/min) or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period of 4 or 6 cycles (Cycle length=21 days). Participants who experienced clinical benefit during the induction phase began maintenance therapy. Participants will receive IV infusion of 1200 mg of atezolizumab and 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

### Primary: Progression Free Survival (PFS) as assessed by the Investigator Using Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST v1.1)

End point title	Progression Free Survival (PFS) as assessed by the Investigator Using Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST v1.1)
-----------------	--

End point description:

PFS is defined as the time from randomization to the first occurrence of disease progression as determined by the investigator using RECIST v1.1 or death from any cause, whichever occurred first.

End point type	Primary
----------------	---------

End point timeframe:

Randomization up to approximately 39 months

End point values	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	286	292		
Units: Months				
median (confidence interval 95%)	5.2 (4.3 to 5.6)	7.7 (6.7 to 8.5)		

### Statistical analyses

Statistical analysis title	PFS Statistical Analysis
----------------------------	--------------------------

**Statistical analysis description:****Unstratified Analysis**

Comparison groups	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed v Arm B (Carboplatin or Cisplatin + Pemetrexed)
Number of subjects included in analysis	578
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.0001
Method	Logrank
Parameter estimate	Log hazard ratio
Point estimate	0.562
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.471
upper limit	0.671

**Primary: Overall Survival (OS)**

End point title	Overall Survival (OS)
End point description:	OS is defined as time from randomization to death from any cause.
End point type	Primary
End point timeframe:	Randomization up to approximately 39 months

<b>End point values</b>	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	286	292		
Units: Months				
median (confidence interval 95%)	13.6 (11.0 to 15.7)	17.5 (13.2 to 19.6)		

**Statistical analyses**

<b>Statistical analysis title</b>	OS Statistical Analysis (Unstratified Analysis)
Comparison groups	Arm B (Carboplatin or Cisplatin + Pemetrexed) v Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)

Number of subjects included in analysis	578
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.1559
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.866
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.709
upper limit	1.056

<b>Statistical analysis title</b>	OS Statistical Analysis (Stratified Analysis)
Comparison groups	Arm B (Carboplatin or Cisplatin + Pemetrexed) v Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
Number of subjects included in analysis	578
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.1546
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.864
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.707
upper limit	1.056

## Secondary: Overall Survival Rate at Year 1

End point title	Overall Survival Rate at Year 1
End point description:	
The Overall Survival Rate at the 1-year landmark time point is defined as the probabilities that participants are alive 1-year after randomization.	
End point type	Secondary
End point timeframe:	
Year 1	

End point values	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	286	292		
Units: Percentage				
number (confidence interval 95%)	55.04 (49.21 to 60.87)	59.72 (54.02 to 65.41)		

## Statistical analyses

Statistical analysis title	OS Rate at 1 Year Statistical Analysis
Comparison groups	Arm B (Carboplatin or Cisplatin + Pemetrexed) v Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
Number of subjects included in analysis	578
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.2606
Method	z test
Parameter estimate	Difference in event free rate
Point estimate	4.68
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.47
upper limit	12.83

## Secondary: Overall Survival Rate Year 2

End point title	Overall Survival Rate Year 2
End point description:	The Overall Survival Rate at the 2-year landmark time point is defined as the probabilities that participants are alive 2-years after randomization.
End point type	Secondary
End point timeframe:	Year 2

End point values	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	91	109		
Units: Percentage				
number (confidence interval 95%)	34.01 (28.40	39.13 (33.44		

**Statistical analyses**

<b>Statistical analysis title</b>	OS Rate at 2 Year Statistical Analysis
Comparison groups	Arm B (Carboplatin or Cisplatin + Pemetrexed) v Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
Number of subjects included in analysis	200
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.209
Method	Z-test
Parameter estimate	Difference in Event Free Rate
Point estimate	5.12
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.87
upper limit	13.11

**Secondary: Percentage of Participants With an Objective Response (Complete Response [CR] or Partial Response [PR]) Assessed by the Investigator Using RECIST V1.1**

End point title	Percentage of Participants With an Objective Response (Complete Response [CR] or Partial Response [PR]) Assessed by the Investigator Using RECIST V1.1
End point description:	
An objective response is defined as either an unconfirmed CR or a PR, as determined by the investigator using RECIST v1.1. Objective Response Rate is defined as the proportion of patients who had an objective response.	
End point type	Secondary
End point timeframe:	
Randomization up to approximately 25 months	

<b>End point values</b>	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	286	292		
Units: Percentage of Participants				
number (not applicable)				
Responders	37.4	51.7		
Non-Responders	62.6	48.3		

## Statistical analyses

<b>Statistical analysis title</b>	Objective Response Statistical Analysis
Comparison groups	Arm B (Carboplatin or Cisplatin + Pemetrexed) v Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
Number of subjects included in analysis	578
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.0005
Method	Cochran-Mantel-Haenszel
Parameter estimate	Difference in response rate
Point estimate	14.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	5.9
upper limit	22.7

## Secondary: Duration of Response (DOR) as Determined by the Investigator Using RECIST v1.1

End point title	Duration of Response (DOR) as Determined by the Investigator Using RECIST v1.1
End point description: DOR is defined as the time interval from the date of the first occurrence of a CR or PR (whichever status is recorded first) until the first date that progressive disease or death is documented, whichever occurs first.	
End point type	Secondary
End point timeframe: Randomization up to approximately 25 months	

End point values	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	286	292		
Units: Months				
number (confidence interval 95%)	6.4 (4.4 to 7.6)	9.5 (6.9 to 12.2)		

## Statistical analyses

<b>Statistical analysis title</b>	DOR Statistical Analysis
Comparison groups	Arm B (Carboplatin or Cisplatin + Pemetrexed) v Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
Number of subjects included in analysis	578
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.0024
Method	Logrank
Parameter estimate	Hazard ratio (HR)
Point estimate	0.62
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.45
upper limit	0.85

## Secondary: Change From Baseline in Patient-Reported Lung Cancer Symptoms as Assessed by European Organization for the Research and Treatment of Cancer (EORTC) Quality-of-Life Questionnaire-Core 30 (QLQ-C30) Symptom Score

End point title	Change From Baseline in Patient-Reported Lung Cancer Symptoms as Assessed by European Organization for the Research and Treatment of Cancer (EORTC) Quality-of-Life Questionnaire-Core 30 (QLQ-C30) Symptom Score
End point description:	
Note: 999999=not available. FU=Follow-Up.	
End point type	Secondary
End point timeframe:	
Baseline up to 3 and 6 months after disease progression or loss of clinical benefit (up to approximately 25 months)	

<b>End point values</b>	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	232	242		
Units: Units on a scale				
arithmetic mean (standard deviation)				
Dyspnoea: Week 3 (Arm B n=222)(Arm A n=216)	-1.95 (± 23.97)	-1.39 (± 25.17)		

Dyspnoea: Week 6 (Arm B n=189)(Arm A n=198)	-1.23 (± 23.16)	-4.71 (± 26.66)		
Dyspnoea: Week 9 (Arm B n=166)(Arm A n=179)	0.20 (± 26.34)	-6.33 (± 26.62)		
Dyspnoea: Week 12 (Arm B n=147)(Arm A n=177)	-0.91 (± 27.57)	-3.39 (± 29.10)		
Dyspnoea: Week 15 (Arm B n=140)(Arm A n=164)	-1.67 (± 28.92)	-2.44 (± 28.73)		
Dyspnoea: Week 18 (Arm B n=126)(Arm A n=155)	-2.12 (± 24.85)	-3.44 (± 30.43)		
Dyspnoea: Week 21 (Arm B n=100)(Arm A n=150)	-5.67 (± 29.61)	-2.89 (± 31.83)		
Dyspnoea: Week 24 (Arm B n=97)(Arm A n=136)	-2.75 (± 28.74)	-2.45 (± 31.85)		
Dyspnoea: Week 27 (Arm B n=79)(Arm A n=126)	-2.53 (± 31.47)	-1.32 (± 27.13)		
Dyspnoea: Week 30 (Arm B n=80)(Arm A n=123)	-1.25 (± 32.45)	-2.71 (± 28.50)		
Dyspnoea: Week 33 (Arm B n=73)(Arm A n=107)	-1.83 (± 27.15)	-3.74 (± 30.14)		
Dyspnoea: Week 36 (Arm B n=61)(Arm A n=105)	-3.83 (± 30.49)	-6.67 (± 31.49)		
Dyspnoea: Week 39 (Arm B n=60)(Arm A n=95)	-1.67 (± 31.55)	-8.42 (± 29.16)		
Dyspnoea: Week 42 (Arm B n=51)(Arm A n=95)	-3.27 (± 28.48)	-9.12 (± 28.12)		
Dyspnoea: Week 45 (Arm B n=48)(Arm A n=79)	-6.25 (± 32.00)	-2.11 (± 30.82)		
Dyspnoea: Week 48 (Arm B n=39)(Arm A n=81)	-5.13 (± 32.03)	-7.41 (± 26.87)		
Dyspnoea: Week 51 (Arm B n=36)(Arm A n=80)	-14.81 (± 30.28)	-2.92 (± 26.09)		
Dyspnoea: Week 54 (Arm B n=31)(Arm A n=72)	-9.68 (± 35.69)	-2.78 (± 27.26)		
Dyspnoea: Week 57 (Arm B n=26)(Arm A n=71)	-7.69 (± 28.76)	-5.63 (± 28.72)		
Dyspnoea: Week 60 (Arm B n=18)(Arm A n=56)	-11.11 (± 25.57)	0.00 (± 25.43)		
Dyspnoea: Week 63 (Arm B n=13)(Arm A n=39)	-7.69 (± 14.62)	-5.13 (± 23.62)		
Dyspnoea: Week 66 (Arm B n=7)(Arm A n=37)	-9.52 (± 16.27)	-1.80 (± 24.78)		
Dyspnoea: Week 69 (Arm B n=6)(Arm A n=28)	-16.67 (± 18.26)	-3.57 (± 24.58)		
Dyspnoea: Week 72 (Arm B n=8)(Arm A n=22)	-4.17 (± 27.82)	0.00 (± 27.22)		
Dyspnoea: Week 75 (Arm B n=5)(Arm A n=23)	-6.67 (± 14.91)	-5.80 (± 19.21)		
Dyspnoea: Week 78 (Arm B n=2)(Arm A n=14)	0.00 (± 0.00)	-4.76 (± 25.68)		
Dyspnoea: Week 81 (Arm B n=2)(Arm A n=9)	0.00 (± 0.00)	-11.11 (± 23.57)		
Dyspnoea: Week 84 (Arm B n=2)(Arm A n=8)	0.00 (± 0.00)	-4.17 (± 21.36)		
Dyspnoea: Week 87 (Arm B n=1)(Arm A n=4)	0.00 (± 999999)	8.33 (± 16.67)		
Dyspnoea: Week 90 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	0.00 (± 0.00)		
Dyspnoea: Week 93 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	0.00 (± 999999)		
Dyspnoea: Time of First Pd (Arm B n=0)(Arm A n=0)	999999 (± 999999)	999999 (± 999999)		



Dyspnoea:Time of Last Tx Dose Arm B n=0;Arm A n=0	999999 (± 999999)	999999 (± 999999)		
Dyspnoea:Survival FU Wk 12 Arm B n=29;Arm A n=25	9.20 (± 35.52)	8.00 (± 42.25)		
Dyspnoea:Survival FU Wk 24 Arm B n=19;Arm A n=7	5.26 (± 25.49)	-14.29 (± 17.82)		

## Statistical analyses

No statistical analyses for this end point

## Secondary: Change from Baseline in Patient-Reported Lung Cancer Symptoms as Assessed by EORTC Quality-of-Life Lung Cancer Module (QLQ-LC13) Symptom Score

End point title	Change from Baseline in Patient-Reported Lung Cancer Symptoms as Assessed by EORTC Quality-of-Life Lung Cancer Module (QLQ-LC13) Symptom Score
-----------------	--

End point description:

Note: 999999=not available. ToL=Time of Last. ToF=Time of First. Sur=survival.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to 3 and 6 months after disease progression or loss of clinical benefit (up to approximately 25 months)

End point values	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	230	234		
Units: Units on a scale				
arithmetic mean (standard deviation)				
Coughing: Week 3 (Arm B n=213)(Arm A n=205)	-2.50 (± 25.37)	-3.41 (± 26.90)		
Coughing: Week 6 (Arm B n=187)(Arm A n=190)	-3.57 (± 27.20)	-9.82 (± 26.50)		
Coughing: Week 9 (Arm B n=162)(Arm A n=173)	-1.85 (± 24.43)	-8.29 (± 31.37)		
Coughing: Week 12 (Arm B n=145)(Arm A n=168)	-3.91 (± 28.46)	-10.71 (± 31.66)		
Coughing: Week 15 (Arm B n=137)(Arm A n=159)	-6.33 (± 30.39)	-11.53 (± 29.05)		
Coughing: Week 18 (Arm B n=125)(Arm A n=149)	-4.00 (± 26.97)	-11.63 (± 27.92)		
Coughing: Week 21 (Arm B n=100)(Arm A n=146)	-6.00 (± 27.78)	-10.50 (± 30.76)		
Coughing: Week 24 (Arm B n=97)(Arm A n=133)	-4.81 (± 28.05)	-10.03 (± 33.08)		
Coughing: Week 27 (Arm B n=78)(Arm A n=123)	-3.85 (± 26.85)	-11.65 (± 34.40)		
Coughing: Week 30 (Arm B n=80)(Arm A n=120)	0.42 (± 29.76)	-11.39 (± 31.31)		

Coughing: Week 33 (Arm B n=72)(Arm A n=103)	-0.46 (± 26.53)	-11.97 (± 30.91)		
Coughing: Week 36 (Arm B n=60)(Arm A n=103)	-6.11 (± 31.59)	-12.94 (± 30.33)		
Coughing: Week 39 (Arm B n=60)(Arm A n=92)	-5.00 (± 30.58)	-13.41 (± 32.43)		
Coughing: Week 42 (Arm B n=52)(Arm A n=93)	-8.97 (± 27.31)	-14.70 (± 29.68)		
Coughing: Week 45 (Arm B n=48)(Arm A n=77)	-13.19 (± 28.13)	-14.72 (± 27.83)		
Coughing: Week 48 (Arm B n=39)(Arm A n=79)	-7.69 (± 31.96)	-13.92 (± 32.29)		
Coughing: Week 51 (Arm B n=36)(Arm A n=79)	-17.59 (± 25.80)	-10.97 (± 34.48)		
Coughing: Week 54 (Arm B n=31)(Arm A n=70)	-17.20 (± 27.04)	-13.81 (± 33.81)		
Coughing: Week 57 (Arm B n=26)(Arm A n=69)	-11.54 (± 22.98)	-13.53 (± 29.33)		
Coughing: Week 60 (Arm B n=19)(Arm A n=56)	-14.04 (± 27.92)	-10.71 (± 31.21)		
Coughing: Week 63 (Arm B n=13)(Arm A n=39)	-10.26 (± 21.01)	-12.82 (± 27.16)		
Coughing: Week 66 (Arm B n=7)(Arm A n=37)	-9.52 (± 25.20)	-12.61 (± 28.71)		
Coughing: Week 69 (Arm B n=6)(Arm A n=28)	-11.11 (± 27.22)	-11.90 (± 30.38)		
Coughing: Week 72 (Arm B n=8)(Arm A n=22)	-8.33 (± 23.57)	-16.67 (± 30.43)		
Coughing: Week 75 (Arm B n=5)(Arm A n=23)	-13.33 (± 18.26)	-18.84 (± 28.12)		
Coughing: Week 78 (Arm B n=2)(Arm A n=14)	-16.67 (± 23.57)	-23.81 (± 27.51)		
Coughing: Week 81 (Arm B n=2)(Arm A n=9)	-16.67 (± 23.57)	-18.52 (± 33.79)		
Coughing: Week 84 (Arm B n=2)(Arm A n=8)	-33.33 (± 0.00)	-29.17 (± 27.82)		
Coughing: Week 87 (Arm B n=1)(Arm A n=4)	0.00 (± 999999)	-25.00 (± 31.91)		
Coughing: Week 90 (Arm B n=1)(Arm A n=2)	-33.33 (± 999999)	-16.67 (± 23.57)		
Coughing: Week 93 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	0.00 (± 999999)		
Coughing:Time of First Pd Arm B n=61;Arm A n=70	-6.01 (± 30.13)	-10.48 (± 28.68)		
Coughing:ToL Tx Dose Arm B n=199;Arm A n=201	-5.19 (± 28.05)	-8.13 (± 28.78)		
Coughing:Survival FU Wk 12 Arm B n=28,Arm A n=24	3.57 (± 29.17)	-6.94 (± 34.02)		
Coughing: Survival FU Wk 24 Arm B n=19,Arm A n=7	-8.77 (± 24.45)	-14.29 (± 32.53)		
Dyspnoea: Week 3 (Arm B n=213)(Arm A n=205)	0.21 (± 18.69)	-1.41 (± 17.68)		
Dyspnoea: Week 6 (Arm B n=187)(Arm A n=190)	0.12 (± 17.13)	-1.17 (± 18.85)		
Dyspnoea: Week 9 (Arm B n=162)(Arm A n=173)	0.89 (± 20.68)	-3.34 (± 18.65)		
Dyspnoea: Week 12 (Arm B n=145)(Arm A n=168)	1.15 (± 21.82)	-0.46 (± 21.76)		
Dyspnoea: Week 15 (Arm B n=137)(Arm A n=159)	0.00 (± 21.30)	-1.12 (± 20.78)		
Dyspnoea: Week 18 (Arm B n=125)(Arm A n=149)	-1.07 (± 18.90)	-3.88 (± 21.92)		

Dyspnoea: Week 21 (Arm B n=100)(Arm A n=146)	-3.56 (± 22.77)	-1.45 (± 21.85)		
Dyspnoea: Week 24 (Arm B n=97)(Arm A n=133)	-2.29 (± 21.75)	-0.42 (± 23.82)		
Dyspnoea: Week 27 (Arm B n=78)(Arm A n=123)	-0.71 (± 22.32)	-2.08 (± 19.83)		
Dyspnoea: Week 30 (Arm B n=80)(Arm A n=120)	1.25 (± 22.57)	-2.78 (± 20.33)		
Dyspnoea: Week 33 (Arm B n=72)(Arm A n=103)	0.77 (± 21.45)	-3.02 (± 22.82)		
Dyspnoea: Week 36 (Arm B n=60)(Arm A n=103)	-1.11 (± 20.93)	-2.91 (± 23.70)		
Dyspnoea: Week 39 (Arm B n=60)(Arm A n=92)	1.30 (± 26.91)	-4.95 (± 22.31)		
Dyspnoea: Week 42 (Arm B n=52)(Arm A n=93)	-0.43 (± 22.65)	-6.33 (± 18.64)		
Dyspnoea: Week 45 (Arm B n=48)(Arm A n=77)	-2.08 (± 26.60)	-3.61 (± 21.36)		
Dyspnoea: Week 48 (Arm B n=39)(Arm A n=79)	-4.56 (± 24.94)	-5.34 (± 21.49)		
Dyspnoea: Week 51 (Arm B n=36)(Arm A n=79)	-9.88 (± 21.87)	-1.69 (± 18.41)		
Dyspnoea: Week 54 (Arm B n=31)(Arm A n=70)	-6.45 (± 24.47)	-1.90 (± 19.10)		
Dyspnoea: Week 57 (Arm B n=26)(Arm A n=69)	-2.99 (± 23.42)	-4.99 (± 22.99)		
Dyspnoea: Week 60 (Arm B n=19)(Arm A n=56)	-11.11 (± 23.42)	-0.40 (± 22.32)		
Dyspnoea: Week 63 (Arm B n=13)(Arm A n=39)	-6.84 (± 15.41)	-6.55 (± 23.04)		
Dyspnoea: Week 66 (Arm B n=7)(Arm A n=37)	-9.52 (± 16.27)	-4.50 (± 21.98)		
Dyspnoea: Week 69 (Arm B n=6)(Arm A n=28)	-16.67 (± 21.94)	-4.76 (± 23.12)		
Dyspnoea: Week 72 (Arm B n=8)(Arm A n=22)	-5.56 (± 19.70)	-1.52 (± 24.56)		
Dyspnoea: Week 75 (Arm B n=5)(Arm A n=23)	-6.67 (± 14.91)	-6.76 (± 17.64)		
Dyspnoea: Week 78 (Arm B n=2)(Arm A n=14)	0.00 (± 0.00)	-3.17 (± 21.09)		
Dyspnoea: Week 81 (Arm B n=2)(Arm A n=9)	11.11 (± 15.71)	-9.88 (± 22.53)		
Dyspnoea: Week 84 (Arm B n=2)(Arm A n=8)	0.00 (± 0.00)	-6.94 (± 22.17)		
Dyspnoea: Week 87 (Arm B n=1)(Arm A n=4)	0.00 (± 999999)	-2.78 (± 18.98)		
Dyspnoea: Week 90 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	5.56 (± 23.57)		
Dyspnoea: Week 93 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	22.22 (± 999999)		
Dyspnoea:Time of First PD Arm B n=61,Arm A n=70	2.37 (± 22.05)	-0.16 (± 24.48)		
Dyspnoea:ToL Tx Dose Arm B n=199, Arm A n=201	3.96 (± 22.74)	0.44 (± 21.83)		
Dyspnoea:Survival FU Wk 12 Arm B n=28,Arm A n=24	9.92 (± 25.90)	13.43 (± 33.73)		
Dyspnoea:Survival FU Wk 24 Arm B n=19,Arm A n=7	4.68 (± 17.50)	4.76 (± 23.00)		
Pain In Chest: Week 3 (Arm B n=213)(Arm A n=205)	-1.25 (± 21.19)	0.81 (± 23.67)		
Pain In Chest: Week 6 (Arm B n=187)(Arm A n=190)	0.71 (± 25.15)	-6.32 (± 24.39)		

Pain In Chest: Week 9 (Arm B n=162)(Arm A n=173)	0.82 (± 24.63)	-4.24 (± 25.57)		
Pain In Chest: Week 12 (Arm B n=145)(Arm A n=168)	-2.07 (± 24.60)	-0.60 (± 27.65)		
Pain In Chest: Week 15 (Arm B n=137)(Arm A n=159)	-1.46 (± 24.88)	-3.56 (± 25.60)		
Pain In Chest: Week 18 (Arm B n=125)(Arm A n=149)	-3.47 (± 26.38)	-3.36 (± 26.77)		
Pain In Chest: Week 21 (Arm B n=100)(Arm A n=146)	-3.33 (± 26.17)	-2.74 (± 25.22)		
Pain In Chest: Week 24 (Arm B n=97)(Arm A n=133)	-2.41 (± 26.89)	-4.76 (± 29.05)		
Pain In Chest: Week 27 (Arm B n=78)(Arm A n=123)	-5.13 (± 29.95)	-2.71 (± 22.82)		
Pain In Chest: Week 30 (Arm B n=80)(Arm A n=120)	-4.17 (± 29.71)	-3.06 (± 27.33)		
Pain In Chest: Week 33 (Arm B n=72)(Arm A n=103)	-0.46 (± 29.86)	-3.88 (± 28.12)		
Pain In Chest: Week 36 (Arm B n=60)(Arm A n=103)	-1.67 (± 31.55)	-2.91 (± 27.26)		
Pain In Chest: Week 39 (Arm B n=60)(Arm A n=92)	-3.89 (± 30.12)	-3.99 (± 27.44)		
Pain In Chest: Week 42 (Arm B n=52)(Arm A n=93)	-7.69 (± 29.24)	-7.17 (± 26.40)		
Pain In Chest: Week 45 (Arm B n=48)(Arm A n=77)	-6.25 (± 29.70)	-5.63 (± 25.02)		
Pain In Chest: Week 48 (Arm B n=39)(Arm A n=79)	-7.69 (± 29.08)	-4.64 (± 24.88)		
Pain In Chest: Week 51 (Arm B n=36)(Arm A n=79)	-11.11 (± 28.73)	-5.49 (± 24.13)		
Pain In Chest: Week 54 (Arm B n=31)(Arm A n=70)	-7.53 (± 29.45)	-4.29 (± 25.33)		
Pain In Chest: Week 57 (Arm B n=26)(Arm A n=69)	-1.28 (± 31.95)	-5.80 (± 26.17)		
Pain In Chest: Week 60 (Arm B n=19)(Arm A n=56)	0.00 (± 22.22)	-3.57 (± 27.47)		
Pain In Chest: Week 63 (Arm B n=13)(Arm A n=39)	-5.13 (± 18.49)	-6.84 (± 25.57)		
Pain In Chest: Week 66 (Arm B n=7)(Arm A n=37)	-4.76 (± 23.00)	-4.50 (± 27.40)		
Pain In Chest: Week 69 (Arm B n=6)(Arm A n=28)	-16.67 (± 18.26)	-5.95 (± 31.50)		
Pain In Chest: Week 72 (Arm B n=8)(Arm A n=22)	-8.33 (± 23.57)	-4.55 (± 23.67)		
Pain In Chest: Week 75 (Arm B n=5)(Arm A n=23)	-6.67 (± 14.91)	0.00 (± 20.10)		
Pain In Chest: Week 78 (Arm B n=2)(Arm A n=14)	0.00 (± 0.00)	0.00 (± 26.15)		
Pain In Chest: Week 81 (Arm B n=2)(Arm A n=9)	0.00 (± 47.14)	7.41 (± 14.70)		
Pain In Chest: Week 84 (Arm B n=2)(Arm A n=8)	0.00 (± 0.00)	4.17 (± 21.36)		
Pain In Chest: Week 87 (Arm B n=1)(Arm A n=4)	-33.33 (± 999999)	8.33 (± 16.67)		
Pain In Chest: Week 90 (Arm B n=1)(Arm A n=2)	-33.33 (± 999999)	0.00 (± 0.00)		
Pain In Chest: Week 93 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	0.00 (± 999999)		
Pain In Chest: ToF Pd (Arm B n=61)(Arm A n=70)	6.56 (± 22.62)	-3.81 (± 29.24)		
Pain In Chest: ToL Tx Dose Arm B n=199 Arm A n=201	1.34 (± 25.26)	-0.83 (± 26.96)		

Pain In Chest: Sur FU Wk12 Arm B n=28 Arm A n=24	5.95 (± 36.35)	-2.78 (± 32.48)		
Pain In Chest: Sur FU Wk 24 Arm B n=19 Arm A n=7	1.75 (± 17.48)	-4.76 (± 29.99)		
Pain In Arm/Shoulder: Wk 3 Arm B n=213 Arm A n=205	-4.69 (± 25.26)	-2.60 (± 24.78)		
Pain In Arm/Shoulder: Wk 6 Arm B n=187 Arm A n=190	-4.46 (± 23.14)	-5.44 (± 25.19)		
Pain In Arm/Shoulder: Wk 9 Arm B n=162 Arm A n=173	-2.88 (± 25.85)	-8.09 (± 28.73)		
Pain In Arm/Shoulder: Wk12 Arm B n=145 Arm A n=168	-6.90 (± 23.54)	-4.37 (± 27.19)		
Pain In Arm/Shoulder: Wk 15 Arm B n=137 Arm A n=159	-3.89 (± 23.24)	-7.34 (± 30.15)		
Pain In Arm/Shoulder: Wk18 Arm B n=125 Arm A n=149	-5.07 (± 25.77)	-6.26 (± 29.09)		
Pain In Arm/Shoulder: Wk21 Arm B n=100 Arm A n=146	-6.00 (± 21.91)	-1.83 (± 28.71)		
Pain In Arm/Shoulder: Wk24 Arm B n=97 Arm A n=133	-2.41 (± 28.16)	-5.26 (± 30.11)		
Pain In Arm/Shoulder: Wk27 Arm B n=78 Arm A n=123	-5.56 (± 28.13)	-3.79 (± 26.72)		
Pain In Arm/Shoulder: Wk30 Arm B n=80 Arm A n=120	1.25 (± 27.27)	-3.33 (± 29.12)		
Pain In Arm/Shoulder: Wk33 Arm B n=72 Arm A n=103	-5.56 (± 25.02)	-1.94 (± 27.54)		
Pain In Arm/Shoulder: Wk36 Arm B n=60 Arm A n=103	-1.67 (± 30.33)	-3.24 (± 31.49)		
Pain In Arm/Shoulder: Wk39 (Arm B n=60)(Arm A n=92)	-5.00 (± 30.58)	-3.99 (± 30.80)		
Pain In Arm/Shoulder: Wk42 (Arm B n=52)(Arm A n=93)	0.00 (± 28.77)	-8.60 (± 25.49)		
Pain In Arm/Shoulder: Wk45 (Arm B n=48)(Arm A n=77)	-0.69 (± 27.06)	-6.06 (± 28.47)		
Pain In Arm/Shoulder: Wk48 (Arm B n=39)(Arm A n=79)	-1.71 (± 27.52)	-4.22 (± 29.89)		
Pain In Arm/Shoulder: Wk51 Arm B n=36 Arm A n=79	-2.78 (± 25.67)	-3.38 (± 27.53)		
Pain In Arm/Shoulder: Wk54 Arm B n=31 Arm A n=70	-1.08 (± 21.92)	-6.19 (± 25.56)		
Pain In Arm/Shoulder: Wk57 Arm B n=26 Arm A n=69	-1.28 (± 22.07)	-2.90 (± 26.65)		
Pain In Arm/Shoulder: Wk 60 Arm B n=19 Arm A n=56	-5.26 (± 20.07)	-4.17 (± 27.75)		
Pain In Arm/Shoulder: Wk 63 Arm B n=13 Arm A n=39	-5.13 (± 18.49)	-3.42 (± 27.35)		
Pain In Arm/Shoulder: Wk 66 Arm B n=7 Arm A n=37	4.76 (± 12.60)	-1.80 (± 31.37)		
Pain In Arm/Shoulder: Wk 69 Arm B n=6 Arm A n=28	0.00 (± 0.00)	-7.14 (± 22.87)		
Pain In Arm/Shoulder: Wk 72 Arm B n=8 Arm A n=22	-8.33 (± 15.43)	-7.58 (± 27.08)		
Pain In Arm/Shoulder: Wk 75 Arm B n=5 Arm A n=23	-6.67 (± 14.91)	-8.70 (± 25.06)		
Pain In Arm/Shoulder: Wk78 Arm B n=2 Arm A n=14	0.00 (± 0.00)	-2.38 (± 24.33)		
Pain In Arm/Shoulder: Wk 81 Arm B n=2 Arm A n=9	0.00 (± 0.00)	3.70 (± 11.11)		
Pain In Arm/Shoulder: Wk84 Arm B n=2 Arm A n=8	0.00 (± 0.00)	4.17 (± 21.36)		
Pain In Arm/Shoulder: Wk 87 Arm B n=1 Arm A n=4	0.00 (± 999999)	-16.67 (± 33.33)		

Pain In Arm/Shoulder:Wk 90 Arm B n=1 Arm A n=2	0.00 (± 999999)	0.00 (± 0.00)		
Pain In Arm/Shoulder:Wk 93 Arm B n=0 Arm A n=1	999999 (± 999999)	0.00 (± 999999)		
Pain In Arm/Shoulder: ToF Pd Arm B n=61 Arm A n=70	2.19 (± 28.46)	1.90 (± 25.94)		
Pain In Arm/Shoulder:ToLTxDose ArmBn=199 ArmAn=201	-0.50 (± 27.11)	0.33 (± 30.55)		
Pain In Arm/Shoulder:Sur FU Wk12 ArmBn=28 ArmAn=24	13.10 (± 30.55)	5.56 (± 40.13)		
Pain In Arm/Shoulder:Sur FU Wk24 ArmBn=19 ArmAn=7	1.75 (± 28.27)	0.00 (± 0.00)		

## Statistical analyses

No statistical analyses for this end point

## Secondary: Change from Baseline in Patient-Reported Lung Cancer Symptoms as Reported Using the Symptoms in Lung Cancer (SILC) Scale Score

End point title	Change from Baseline in Patient-Reported Lung Cancer Symptoms as Reported Using the Symptoms in Lung Cancer (SILC) Scale Score
End point description:	Change from baseline per SILC scale will be analyzed for each of the lung cancer symptom scores (chest pain, cough, dyspnea). Note: 999999=not available. ToF=Time of First. ToL=Time of Last.
End point type	Secondary
End point timeframe:	Baseline up to 3 and 6 months after disease progression or loss of clinical benefit (up to approximately 25 months)

End point values	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	176	200		
Units: Units on a scale				
arithmetic mean (standard deviation)				
Chest Pain: Week 1 (Arm B n=160)(Arm A n=186)	0.20 (± 0.83)	0.30 (± 0.88)		
Chest Pain: Week 2 (Arm B n=152)(Arm A n=176)	-0.01 (± 0.88)	0.21 (± 0.85)		
Chest Pain: Week 3 (Arm B n=151)(Arm A n=165)	-0.05 (± 0.92)	0.06 (± 0.95)		
Chest Pain: Week 4 (Arm B n=144)(Arm A n=160)	0.03 (± 0.91)	0.03 (± 0.93)		
Chest Pain: Week 5 (Arm B n=143)(Arm A n=166)	-0.02 (± 0.93)	-0.01 (± 0.88)		
Chest Pain: Week 6 (Arm B n=132)(Arm A n=166)	-0.05 (± 0.88)	-0.02 (± 0.92)		
Chest Pain: Week 7 (Arm B n=132)(Arm A n=156)	0.07 (± 1.00)	-0.04 (± 1.00)		

Chest Pain: Week 8 (Arm B n=124)(Arm A n=157)	0.03 (± 0.94)	-0.07 (± 1.02)		
Chest Pain: Week 9 (Arm B n=125)(Arm A n=159)	0.06 (± 1.00)	-0.03 (± 1.03)		
Chest Pain: Week 10 (Arm B n=120)(Arm A n=152)	0.03 (± 1.02)	0.02 (± 1.05)		
Chest Pain: Week 11 (Arm B n=117)(Arm A n=156)	0.02 (± 0.96)	0.04 (± 1.09)		
Chest Pain: Week 12 (Arm B n=116)(Arm A n=142)	-0.02 (± 0.96)	-0.01 (± 1.02)		
Chest Pain: Week 13 (Arm B n=108)(Arm A n=142)	-0.05 (± 1.02)	0.05 (± 0.97)		
Chest Pain: Week 14 (Arm B n=103)(Arm A n=132)	-0.07 (± 1.12)	0.06 (± 1.06)		
Chest Pain: Week 15 (Arm B n=101)(Arm A n=133)	0.13 (± 1.16)	0.08 (± 0.94)		
Chest Pain: Week 16 (Arm B n=100)(Arm A n=133)	0.02 (± 1.08)	0.08 (± 1.04)		
Chest Pain: Week 17 (Arm B n=97)(Arm A n=131)	0.02 (± 1.08)	0.00 (± 1.08)		
Chest Pain: Week 18 (Arm B n=88)(Arm A n=121)	-0.06 (± 1.11)	0.12 (± 1.07)		
Chest Pain: Week 19 (Arm B n=81)(Arm A n=122)	-0.15 (± 1.12)	0.02 (± 1.01)		
Chest Pain: Week 20 (Arm B n=71)(Arm A n=112)	-0.02 (± 1.22)	-0.01 (± 1.03)		
Chest Pain: Week 21 (Arm B n=74)(Arm A n=114)	0.03 (± 1.28)	0.01 (± 1.03)		
Chest Pain: Week 22 (Arm B n=65)(Arm A n=106)	0.04 (± 1.21)	0.02 (± 1.02)		
Chest Pain: Week 23 (Arm B n=65)(Arm A n=106)	-0.08 (± 1.14)	0.04 (± 0.96)		
Chest Pain: Week 24 (Arm B n=64)(Arm A n=106)	-0.02 (± 1.16)	0.09 (± 0.99)		
Chest Pain: Week 25 (Arm B n=60)(Arm A n=100)	0.05 (± 1.24)	0.14 (± 1.03)		
Chest Pain: Week 26 (Arm B n=55)(Arm A n=100)	-0.05 (± 1.14)	0.09 (± 1.00)		
Chest Pain: Week 27 (Arm B n=55)(Arm A n=99)	0.04 (± 1.17)	0.04 (± 1.01)		
Chest Pain: Week 28 (Arm B n=55)(Arm A n=96)	0.24 (± 1.18)	0.03 (± 0.93)		
Chest Pain: Week 29 (Arm B n=54)(Arm A n=99)	0.06 (± 1.29)	0.04 (± 0.95)		
Chest Pain: Week 30 (Arm B n=52)(Arm A n=95)	0.01 (± 1.37)	0.04 (± 0.97)		
Chest Pain: Week 31 (Arm B n=44)(Arm A n=93)	0.02 (± 1.09)	0.05 (± 0.88)		
Chest Pain: Week 32 (Arm B n=39)(Arm A n=94)	0.21 (± 1.21)	-0.01 (± 0.98)		
Chest Pain: Week 33 (Arm B n=41)(Arm A n=85)	0.04 (± 1.17)	0.03 (± 0.92)		
Chest Pain: Week 34 (Arm B n=40)(Arm A n=84)	0.21 (± 1.27)	0.11 (± 0.99)		
Chest Pain: Week 35 (Arm B n=39)(Arm A n=87)	0.33 (± 1.30)	0.11 (± 0.98)		
Chest Pain: Week 36 (Arm B n=34)(Arm A n=86)	0.15 (± 1.07)	0.17 (± 1.04)		
Chest Pain: Week 37 (Arm B n=33)(Arm A n=86)	0.15 (± 1.21)	0.06 (± 0.94)		
Chest Pain: Week 38 (Arm B n=30)(Arm A n=80)	0.12 (± 1.16)	0.04 (± 1.03)		

Chest Pain: Week 39 (Arm B n=32)(Arm A n=72)	0.20 (± 1.23)	0.13 (± 1.01)		
Chest Pain: Week 40 (Arm B n=31)(Arm A n=77)	0.15 (± 1.31)	0.04 (± 0.96)		
Chest Pain: Week 41 (Arm B n=30)(Arm A n=74)	-0.05 (± 1.06)	0.09 (± 1.03)		
Chest Pain: Week 42 (Arm B n=27)(Arm A n=73)	-0.06 (± 1.07)	0.03 (± 1.02)		
Chest Pain: Week 43 (Arm B n=23)(Arm A n=76)	-0.22 (± 0.82)	0.13 (± 1.02)		
Chest Pain: Week 44 (Arm B n=25)(Arm A n=72)	-0.10 (± 1.29)	0.10 (± 1.04)		
Chest Pain: Week 45 (Arm B n=25)(Arm A n=69)	-0.20 (± 1.24)	0.03 (± 0.95)		
Chest Pain: Week 46 (Arm B n=24)(Arm A n=67)	-0.06 (± 1.18)	0.08 (± 0.99)		
Chest Pain: Week 47 (Arm B n=23)(Arm A n=63)	0.02 (± 1.09)	0.09 (± 0.91)		
Chest Pain: Week 48 (Arm B n=25)(Arm A n=64)	0.02 (± 1.27)	0.05 (± 1.00)		
Chest Pain: Week 49 (Arm B n=23)(Arm A n=62)	-0.09 (± 1.07)	0.03 (± 0.90)		
Chest Pain: Week 50 (Arm B n=23)(Arm A n=58)	-0.13 (± 0.99)	0.00 (± 1.03)		
Chest Pain: Week 51 (Arm B n=21)(Arm A n=62)	-0.02 (± 0.77)	0.12 (± 0.94)		
Chest Pain: Week 52 (Arm B n=21)(Arm A n=55)	0.14 (± 1.06)	0.00 (± 1.11)		
Chest Pain: Week 53 (Arm B n=23)(Arm A n=54)	0.13 (± 1.35)	-0.01 (± 1.06)		
Chest Pain: Week 54 (Arm B n=22)(Arm A n=54)	0.11 (± 1.16)	-0.08 (± 1.10)		
Chest Pain: Week 55 (Arm B n=21)(Arm A n=49)	0.12 (± 1.38)	0.03 (± 1.14)		
Chest Pain: Week 56 (Arm B n=20)(Arm A n=53)	0.13 (± 1.46)	0.04 (± 1.18)		
Chest Pain: Week 57 (Arm B n=17)(Arm A n=46)	0.12 (± 1.21)	-0.01 (± 1.15)		
Chest Pain: Week 58 (Arm B n=15)(Arm A n=43)	0.27 (± 0.98)	-0.12 (± 1.10)		
Chest Pain: Week 59 (Arm B n=13)(Arm A n=40)	0.23 (± 0.73)	-0.23 (± 1.01)		
Chest Pain: Week 60 (Arm B n=11)(Arm A n=41)	0.14 (± 0.67)	-0.15 (± 1.17)		
Chest Pain: Week 61 (Arm B n=13)(Arm A n=43)	0.23 (± 0.83)	0.00 (± 1.06)		
Chest Pain: Week 62 (Arm B n=13)(Arm A n=37)	0.46 (± 0.72)	-0.04 (± 1.08)		
Chest Pain: Week 63 (Arm B n=12)(Arm A n=36)	0.25 (± 0.75)	-0.06 (± 1.09)		
Chest Pain: Week 64 (Arm B n=10)(Arm A n=35)	0.15 (± 0.78)	0.09 (± 0.95)		
Chest Pain: Week 65 (Arm B n=9)(Arm A n=33)	0.11 (± 0.82)	-0.03 (± 0.99)		
Chest Pain: Week 66 (Arm B n=6)(Arm A n=34)	0.42 (± 0.92)	0.13 (± 1.05)		
Chest Pain: Week 67 (Arm B n=7)(Arm A n=30)	0.36 (± 0.85)	0.02 (± 1.00)		
Chest Pain: Week 68 (Arm B n=6)(Arm A n=30)	0.33 (± 0.88)	0.12 (± 1.18)		
Chest Pain: Week 69 (Arm B n=7)(Arm A n=29)	0.07 (± 0.93)	-0.05 (± 1.10)		



Chest Pain: Week 70 (Arm B n=6)(Arm A n=30)	0.25 (± 0.88)	0.03 (± 1.11)		
Chest Pain: Week 71 (Arm B n=5)(Arm A n=26)	0.30 (± 0.97)	0.04 (± 1.09)		
Chest Pain: Week 72 (Arm B n=4)(Arm A n=25)	-0.13 (± 0.25)	-0.22 (± 0.94)		
Chest Pain: Week 73 (Arm B n=4)(Arm A n=24)	0.13 (± 1.31)	-0.10 (± 1.01)		
Chest Pain: Week 74 (Arm B n=4)(Arm A n=23)	0.38 (± 1.11)	-0.02 (± 1.14)		
Chest Pain: Week 75 (Arm B n=4)(Arm A n=21)	0.38 (± 1.11)	-0.07 (± 1.14)		
Chest Pain: Week 76 (Arm B n=3)(Arm A n=17)	0.00 (± 0.50)	-0.06 (± 1.20)		
Chest Pain: Week 77 (Arm B n=2)(Arm A n=17)	-0.25 (± 0.35)	-0.21 (± 1.03)		
Chest Pain: Week 78 (Arm B n=2)(Arm A n=14)	-0.75 (± 0.35)	-0.25 (± 1.14)		
Chest Pain: Week 79 (Arm B n=2)(Arm A n=14)	0.00 (± 0.71)	-0.18 (± 1.20)		
Chest Pain: Week 80 (Arm B n=2)(Arm A n=13)	-0.25 (± 0.35)	-0.12 (± 0.96)		
Chest Pain: Week 81 (Arm B n=2)(Arm A n=12)	-0.25 (± 0.35)	-0.25 (± 0.89)		
Chest Pain: Week 82 (Arm B n=2)(Arm A n=9)	-0.75 (± 0.35)	0.00 (± 1.25)		
Chest Pain: Week 83 (Arm B n=2)(Arm A n=4)	-0.25 (± 0.35)	-0.50 (± 1.73)		
Chest Pain: Week 84 (Arm B n=2)(Arm A n=5)	0.00 (± 0.71)	-0.30 (± 1.57)		
Chest Pain: Week 85 (Arm B n=1)(Arm A n=5)	-1.00 (± 999999)	-0.20 (± 1.68)		
Chest Pain: Week 86 (Arm B n=1)(Arm A n=4)	0.00 (± 999999)	-0.50 (± 1.73)		
Chest Pain: Week 87 (Arm B n=1)(Arm A n=3)	-1.00 (± 999999)	0.50 (± 0.87)		
Chest Pain: Week 88 (Arm B n=1)(Arm A n=4)	-1.00 (± 999999)	0.00 (± 0.00)		
Chest Pain: Week 89 (Arm B n=1)(Arm A n=2)	-1.00 (± 999999)	0.00 (± 0.00)		
Chest Pain: Week 90 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	0.00 (± 0.00)		
Chest Pain: Week 91 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	0.00 (± 0.00)		
Chest Pain: Week 92 (Arm B n=0)(Arm A n=2)	999999 (± 999999)	0.00 (± 0.00)		
Chest Pain: Week 93 (Arm B n=0)(Arm A n=2)	999999 (± 999999)	0.00 (± 0.00)		
Chest Pain: Week 94 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	1.00 (± 999999)		
Chest Pain: Week 95 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	0.50 (± 999999)		
Chest Pain: ToF Pd (Arm B n=110)(Arm A n=96)	0.28 (± 1.06)	0.20 (± 0.91)		
Chest Pain: ToLTx Dose (Arm B n=141)(Arm A n=150)	0.17 (± 1.17)	0.13 (± 0.98)		
Cough: Week 1 (Arm B n=160)(Arm A n=186)	-0.08 (± 0.74)	-0.06 (± 0.89)		
Cough: Week 2 (Arm B n=152)(Arm A n=176)	-0.13 (± 0.73)	-0.08 (± 0.90)		
Cough: Week 3 (Arm B n=151)(Arm A n=165)	-0.05 (± 0.77)	-0.07 (± 0.81)		

Cough: Week 4 (Arm B n=144)(Arm A n=160)	-0.11 (± 0.89)	-0.22 (± 0.85)		
Cough: Week 5 (Arm B n=143)(Arm A n=166)	-0.14 (± 0.83)	-0.33 (± 0.89)		
Cough: Week 6 (Arm B n=132)(Arm A n=166)	-0.16 (± 0.85)	-0.33 (± 0.91)		
Cough: Week 7 (Arm B n=132)(Arm A n=156)	-0.22 (± 0.89)	-0.22 (± 0.97)		
Cough: Week 8 (Arm B n=124)(Arm A n=157)	-0.20 (± 0.89)	-0.28 (± 0.98)		
Cough: Week 9 (Arm B n=125)(Arm A n=159)	-0.17 (± 0.91)	-0.29 (± 0.97)		
Cough: Week 10 (Arm B n=120)(Arm A n=152)	-0.22 (± 0.95)	-0.35 (± 1.06)		
Cough: Week 11 (Arm B n=117)(Arm A n=156)	-0.21 (± 0.99)	-0.33 (± 1.02)		
Cough: Week 12 (Arm B n=116)(Arm A n=142)	-0.13 (± 1.05)	-0.40 (± 0.96)		
Cough: Week 13 (Arm B n=108)(Arm A n=142)	-0.24 (± 1.05)	-0.37 (± 1.04)		
Cough: Week 14 (Arm B n=103)(Arm A n=132)	-0.23 (± 0.98)	-0.34 (± 1.03)		
Cough: Week 15 (Arm B n=101)(Arm A n=133)	-0.11 (± 1.03)	-0.33 (± 1.03)		
Cough: Week 16 (Arm B n=100)(Arm A n=133)	-0.20 (± 1.05)	-0.33 (± 1.04)		
Cough: Week 17 (Arm B n=97)(Arm A n=131)	-0.20 (± 1.11)	-0.31 (± 0.99)		
Cough: Week 18 (Arm B n=88)(Arm A n=121)	-0.19 (± 1.12)	-0.31 (± 1.00)		
Cough: Week 19 (Arm B n=81)(Arm A n=122)	-0.25 (± 1.07)	-0.36 (± 1.06)		
Cough: Week 20 (Arm B n=71)(Arm A n=112)	-0.25 (± 1.16)	-0.39 (± 0.99)		
Cough: Week 21 (Arm B n=74)(Arm A n=114)	-0.19 (± 1.09)	-0.36 (± 1.07)		
Cough: Week 22 (Arm B n=65)(Arm A n=106)	-0.29 (± 1.11)	-0.41 (± 1.08)		
Cough: Week 23 (Arm B n=65)(Arm A n=106)	-0.34 (± 1.07)	-0.44 (± 1.04)		
Cough: Week 24 (Arm B n=64)(Arm A n=106)	-0.27 (± 0.96)	-0.29 (± 1.13)		
Cough: Week 25 (Arm B n=60)(Arm A n=100)	-0.24 (± 1.11)	-0.27 (± 1.10)		
Cough: Week 26 (Arm B n=55)(Arm A n=100)	-0.25 (± 1.16)	-0.22 (± 1.22)		
Cough: Week 27 (Arm B n=55)(Arm A n=99)	-0.19 (± 0.99)	-0.32 (± 1.11)		
Cough: Week 28 (Arm B n=55)(Arm A n=96)	-0.19 (± 1.04)	-0.32 (± 1.12)		
Cough: Week 29 (Arm B n=54)(Arm A n=99)	-0.25 (± 1.13)	-0.34 (± 1.11)		
Cough: Week 30 (Arm B n=52)(Arm A n=95)	-0.08 (± 1.12)	-0.34 (± 1.00)		
Cough: Week 31 (Arm B n=44)(Arm A n=93)	-0.13 (± 1.13)	-0.37 (± 0.99)		
Cough: Week 32 (Arm B n=39)(Arm A n=94)	0.01 (± 1.09)	-0.45 (± 0.94)		
Cough: Week 33 (Arm B n=41)(Arm A n=85)	0.05 (± 1.17)	-0.44 (± 0.99)		
Cough: Week 34 (Arm B n=40)(Arm A n=84)	-0.10 (± 1.09)	-0.29 (± 1.02)		

Cough: Week 35 (Arm B n=39)(Arm A n=87)	-0.27 (± 1.05)	-0.32 (± 0.99)		
Cough: Week 36 (Arm B n=34)(Arm A n=86)	0.01 (± 1.10)	-0.31 (± 0.99)		
Cough: Week 37 (Arm B n=33)(Arm A n=86)	-0.11 (± 1.16)	-0.37 (± 1.01)		
Cough: Week 38 (Arm B n=30)(Arm A n=80)	0.02 (± 1.03)	-0.31 (± 1.03)		
Cough: Week 39 (Arm B n=32)(Arm A n=72)	0.02 (± 1.19)	-0.33 (± 0.98)		
Cough: Week 40 (Arm B n=31)(Arm A n=77)	-0.10 (± 1.15)	-0.35 (± 1.07)		
Cough: Week 41 (Arm B n=30)(Arm A n=74)	-0.27 (± 1.10)	-0.30 (± 1.09)		
Cough: Week 42 (Arm B n=27)(Arm A n=73)	-0.31 (± 1.04)	-0.31 (± 1.04)		
Cough: Week 43 (Arm B n=23)(Arm A n=76)	-0.20 (± 1.04)	-0.37 (± 0.98)		
Cough: Week 44 (Arm B n=25)(Arm A n=72)	0.00 (± 1.04)	-0.36 (± 1.03)		
Cough: Week 45 (Arm B n=25)(Arm A n=69)	-0.18 (± 1.14)	-0.41 (± 1.04)		
Cough: Week 46 (Arm B n=24)(Arm A n=67)	-0.25 (± 1.04)	-0.34 (± 1.17)		
Cough: Week 47 (Arm B n=23)(Arm A n=63)	-0.43 (± 1.09)	-0.29 (± 1.17)		
Cough: Week 48 (Arm B n=25)(Arm A n=64)	-0.26 (± 1.16)	-0.38 (± 1.08)		
Cough: Week 49 (Arm B n=23)(Arm A n=62)	-0.50 (± 1.07)	-0.40 (± 1.10)		
Cough: Week 50 (Arm B n=23)(Arm A n=58)	-0.37 (± 0.84)	-0.40 (± 1.06)		
Cough: Week 51 (Arm B n=21)(Arm A n=62)	-0.45 (± 1.16)	-0.15 (± 1.10)		
Cough: Week 52 (Arm B n=21)(Arm A n=55)	-0.29 (± 1.07)	-0.10 (± 1.17)		
Cough: Week 53 (Arm B n=23)(Arm A n=54)	-0.24 (± 1.21)	-0.13 (± 1.17)		
Cough: Week 54 (Arm B n=22)(Arm A n=54)	-0.27 (± 1.10)	-0.16 (± 1.15)		
Cough: Week 55 (Arm B n=21)(Arm A n=49)	-0.10 (± 1.15)	-0.35 (± 1.13)		
Cough: Week 56 (Arm B n=20)(Arm A n=53)	-0.30 (± 1.14)	-0.26 (± 1.17)		
Cough: Week 57 (Arm B n=17)(Arm A n=46)	-0.41 (± 0.92)	-0.25 (± 1.10)		
Cough: Week 58 (Arm B n=15)(Arm A n=43)	-0.17 (± 1.05)	-0.45 (± 0.96)		
Cough: Week 59 (Arm B n=13)(Arm A n=40)	-0.31 (± 0.93)	-0.50 (± 1.02)		
Cough: Week 60 (Arm B n=11)(Arm A n=41)	-0.55 (± 1.04)	-0.52 (± 1.03)		
Cough: Week 61 (Arm B n=13)(Arm A n=43)	-0.23 (± 1.15)	-0.38 (± 1.07)		
Cough: Week 62 (Arm B n=13)(Arm A n=37)	0.04 (± 1.03)	-0.53 (± 1.15)		
Cough: Week 63 (Arm B n=12)(Arm A n=36)	-0.29 (± 1.18)	-0.57 (± 1.10)		
Cough: Week 64 (Arm B n=10)(Arm A n=35)	-0.45 (± 1.04)	-0.49 (± 1.01)		
Cough: Week 65 (Arm B n=9)(Arm A n=33)	-0.61 (± 0.96)	-0.39 (± 0.94)		

Cough: Week 66 (Arm B n=6)(Arm A n=34)	-0.17 (± 0.82)	-0.43 (± 1.18)		
Cough: Week 67 (Arm B n=7)(Arm A n=30)	-0.64 (± 1.07)	-0.47 (± 0.84)		
Cough: Week 68 (Arm B n=6)(Arm A n=30)	-0.75 (± 0.88)	-0.35 (± 1.04)		
Cough: Week 69 (Arm B n=7)(Arm A n=29)	-0.57 (± 1.10)	-0.53 (± 0.90)		
Cough: Week 70 (Arm B n=6)(Arm A n=30)	-0.83 (± 0.93)	-0.43 (± 0.98)		
Cough: Week 71 (Arm B n=5)(Arm A n=26)	-0.90 (± 0.89)	-0.56 (± 1.04)		
Cough: Week 72 (Arm B n=4)(Arm A n=25)	-1.25 (± 0.50)	-0.72 (± 1.15)		
Cough: Week 73 (Arm B n=4)(Arm A n=24)	-0.63 (± 1.11)	-0.77 (± 1.00)		
Cough: Week 74 (Arm B n=4)(Arm A n=23)	-0.63 (± 1.11)	-0.54 (± 0.94)		
Cough: Week 75 (Arm B n=4)(Arm A n=21)	-0.63 (± 1.11)	-0.67 (± 1.00)		
Cough: Week 76 (Arm B n=3)(Arm A n=17)	-1.00 (± 1.00)	-0.65 (± 1.22)		
Cough: Week 77 (Arm B n=2)(Arm A n=17)	-0.50 (± 0.71)	-0.82 (± 1.21)		
Cough: Week 78 (Arm B n=2)(Arm A n=14)	-1.00 (± 0.00)	-0.57 (± 1.04)		
Cough: Week 79 (Arm B n=2)(Arm A n=14)	-0.25 (± 0.35)	-0.64 (± 1.23)		
Cough: Week 80 (Arm B n=2)(Arm A n=13)	-0.50 (± 0.71)	-1.04 (± 1.11)		
Cough: Week 81 (Arm B n=2)(Arm A n=12)	-0.50 (± 0.71)	-0.79 (± 1.05)		
Cough: Week 82 (Arm B n=2)(Arm A n=9)	-0.50 (± 0.71)	-0.89 (± 1.27)		
Cough: Week 83 (Arm B n=2)(Arm A n=4)	-1.00 (± 0.00)	0.13 (± 1.03)		
Cough: Week 84 (Arm B n=2)(Arm A n=5)	-0.50 (± 0.71)	-0.20 (± 1.04)		
Cough: Week 85 (Arm B n=1)(Arm A n=5)	0.00 (± 999999)	0.00 (± 0.87)		
Cough: Week 86 (Arm B n=1)(Arm A n=4)	0.00 (± 999999)	-0.38 (± 0.48)		
Cough: Week 87 (Arm B n=1)(Arm A n=3)	0.00 (± 999999)	-0.83 (± 0.58)		
Cough: Week 88 (Arm B n=1)(Arm A n=4)	0.00 (± 999999)	-0.88 (± 0.48)		
Cough: Week 89 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	0.00 (± 0.71)		
Cough: Week 90 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	0.25 (± 1.06)		
Cough: Week 91 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	-0.50 (± 0.00)		
Cough: Week 92 (Arm B n=0)(Arm A n=2)	999999 (± 999999)	0.50 (± 0.71)		
Cough: Week 93 (Arm B n=0)(Arm A n=2)	999999 (± 999999)	0.50 (± 0.71)		
Cough: Week 94 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	0.50 (± 999999)		
Cough: Week 95 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	0.00 (± 999999)		
Cough: ToF Pd (Arm B n=110)(Arm A n=96)	-0.08 (± 1.12)	-0.45 (± 0.93)		

Cough: ToL Tx Dose (Arm B n=141)(Arm A n=150)	-0.15 (± 0.95)	-0.29 (± 0.91)		
Dyspnoea: Week 1 (Arm B n=160)(Arm A n=186)	0.20 (± 0.82)	0.16 (± 0.81)		
Dyspnoea: Week 2 (Arm B n=152)(Arm A n=176)	0.11 (± 0.76)	0.15 (± 0.82)		
Dyspnoea: Week 3 (Arm B n=151)(Arm A n=165)	0.10 (± 0.71)	0.12 (± 0.70)		
Dyspnoea: Week 4 (Arm B n=144)(Arm A n=160)	0.21 (± 0.84)	0.17 (± 0.95)		
Dyspnoea: Week 5 (Arm B n=143)(Arm A n=166)	0.22 (± 0.82)	0.17 (± 0.91)		
Dyspnoea: Week 6 (Arm B n=132)(Arm A n=166)	0.23 (± 0.82)	0.16 (± 0.95)		
Dyspnoea: Week 7 (Arm B n=132)(Arm A n=156)	0.30 (± 0.93)	0.25 (± 0.92)		
Dyspnoea: Week 8 (Arm B n=124)(Arm A n=157)	0.35 (± 0.87)	0.24 (± 0.96)		
Dyspnoea: Week 9 (Arm B n=125)(Arm A n=159)	0.34 (± 0.89)	0.17 (± 0.91)		
Dyspnoea: Week 10 (Arm B n=120)(Arm A n=152)	0.46 (± 0.93)	0.16 (± 0.92)		
Dyspnoea: Week 11 (Arm B n=117)(Arm A n=156)	0.39 (± 1.01)	0.26 (± 0.96)		
Dyspnoea: Week 12 (Arm B n=116)(Arm A n=142)	0.38 (± 0.96)	0.22 (± 0.90)		
Dyspnoea: Week 13 (Arm B n=108)(Arm A n=142)	0.34 (± 0.96)	0.30 (± 1.02)		
Dyspnoea: Week 14 (Arm B n=103)(Arm A n=132)	0.44 (± 0.95)	0.22 (± 0.99)		
Dyspnoea: Week 15 (Arm B n=101)(Arm A n=133)	0.45 (± 0.94)	0.19 (± 0.95)		
Dyspnoea: Week 16 (Arm B n=100)(Arm A n=133)	0.37 (± 1.06)	0.26 (± 1.02)		
Dyspnoea: Week 17 (Arm B n=97)(Arm A n=131)	0.35 (± 1.04)	0.15 (± 1.00)		
Dyspnoea: Week 18 (Arm B n=88)(Arm A n=121)	0.35 (± 0.99)	0.22 (± 0.99)		
Dyspnoea: Week 19 (Arm B n=81)(Arm A n=122)	0.30 (± 1.02)	0.22 (± 0.97)		
Dyspnoea: Week 20 (Arm B n=71)(Arm A n=112)	0.30 (± 1.09)	0.27 (± 1.06)		
Dyspnoea: Week 21 (Arm B n=74)(Arm A n=114)	0.22 (± 1.03)	0.22 (± 1.05)		
Dyspnoea: Week 22 (Arm B n=65)(Arm A n=106)	0.16 (± 1.10)	0.28 (± 1.03)		
Dyspnoea: Week 23 (Arm B n=65)(Arm A n=106)	0.24 (± 1.05)	0.22 (± 1.03)		
Dyspnoea: Week 24 (Arm B n=64)(Arm A n=106)	0.30 (± 0.98)	0.24 (± 0.99)		
Dyspnoea: Week 25 (Arm B n=60)(Arm A n=100)	0.50 (± 1.05)	0.28 (± 0.99)		
Dyspnoea: Week 26 (Arm B n=55)(Arm A n=100)	0.30 (± 1.05)	0.24 (± 1.04)		
Dyspnoea: Week 27 (Arm B n=55)(Arm A n=99)	0.43 (± 1.17)	0.24 (± 1.06)		
Dyspnoea: Week 28 (Arm B n=55)(Arm A n=96)	0.46 (± 1.02)	0.25 (± 0.97)		
Dyspnoea: Week 29 (Arm B n=54)(Arm A n=99)	0.43 (± 1.06)	0.20 (± 1.01)		
Dyspnoea: Week 30 (Arm B n=52)(Arm A n=95)	0.45 (± 1.08)	0.21 (± 0.99)		

Dyspnoea: Week 31 (Arm B n=44)(Arm A n=93)	0.35 (± 1.03)	0.15 (± 0.99)		
Dyspnoea: Week 32 (Arm B n=39)(Arm A n=94)	0.48 (± 1.03)	0.16 (± 1.01)		
Dyspnoea: Week 33 (Arm B n=41)(Arm A n=85)	0.36 (± 0.96)	0.17 (± 0.95)		
Dyspnoea: Week 34 (Arm B n=40)(Arm A n=84)	0.47 (± 1.15)	0.24 (± 1.04)		
Dyspnoea: Week 35 (Arm B n=39)(Arm A n=87)	0.42 (± 1.00)	0.23 (± 1.03)		
Dyspnoea: Week 36 (Arm B n=34)(Arm A n=86)	0.59 (± 1.06)	0.22 (± 1.02)		
Dyspnoea: Week 37 (Arm B n=33)(Arm A n=86)	0.53 (± 1.17)	0.24 (± 1.03)		
Dyspnoea: Week 38 (Arm B n=30)(Arm A n=80)	0.59 (± 1.06)	0.22 (± 1.07)		
Dyspnoea: Week 39 (Arm B n=32)(Arm A n=72)	0.65 (± 1.04)	0.19 (± 1.01)		
Dyspnoea: Week 40 (Arm B n=31)(Arm A n=77)	0.62 (± 1.02)	0.19 (± 0.95)		
Dyspnoea: Week 41 (Arm B n=30)(Arm A n=74)	0.63 (± 1.01)	0.16 (± 0.95)		
Dyspnoea: Week 42 (Arm B n=27)(Arm A n=73)	0.50 (± 1.01)	0.17 (± 0.92)		
Dyspnoea: Week 43 (Arm B n=23)(Arm A n=76)	0.32 (± 0.89)	0.22 (± 0.90)		
Dyspnoea: Week 44 (Arm B n=25)(Arm A n=72)	0.58 (± 0.83)	0.23 (± 0.92)		
Dyspnoea: Week 45 (Arm B n=25)(Arm A n=69)	0.49 (± 0.96)	0.21 (± 0.90)		
Dyspnoea: Week 46 (Arm B n=24)(Arm A n=67)	0.55 (± 1.02)	0.23 (± 0.89)		
Dyspnoea: Week 47 (Arm B n=23)(Arm A n=63)	0.53 (± 1.09)	0.14 (± 0.90)		
Dyspnoea: Week 48 (Arm B n=25)(Arm A n=64)	0.55 (± 0.96)	0.14 (± 0.85)		
Dyspnoea: Week 49 (Arm B n=23)(Arm A n=62)	0.63 (± 1.04)	0.16 (± 0.90)		
Dyspnoea: Week 50 (Arm B n=23)(Arm A n=58)	0.52 (± 0.98)	0.24 (± 0.94)		
Dyspnoea: Week 51 (Arm B n=21)(Arm A n=62)	0.67 (± 1.04)	0.30 (± 0.97)		
Dyspnoea: Week 52 (Arm B n=21)(Arm A n=55)	0.72 (± 1.02)	0.27 (± 1.02)		
Dyspnoea: Week 53 (Arm B n=23)(Arm A n=54)	0.81 (± 1.06)	0.22 (± 0.95)		
Dyspnoea: Week 54 (Arm B n=22)(Arm A n=54)	0.82 (± 1.06)	0.23 (± 0.95)		
Dyspnoea: Week 55 (Arm B n=21)(Arm A n=49)	0.77 (± 1.17)	0.16 (± 0.99)		
Dyspnoea: Week 56 (Arm B n=20)(Arm A n=53)	0.84 (± 1.15)	0.18 (± 0.95)		
Dyspnoea: Week 57 (Arm B n=17)(Arm A n=46)	0.68 (± 1.16)	0.11 (± 0.91)		
Dyspnoea: Week 58 (Arm B n=15)(Arm A n=43)	0.77 (± 1.06)	0.16 (± 0.86)		
Dyspnoea: Week 59 (Arm B n=13)(Arm A n=40)	0.66 (± 0.90)	0.07 (± 0.85)		
Dyspnoea: Week 60 (Arm B n=11)(Arm A n=41)	0.73 (± 0.96)	0.27 (± 0.94)		
Dyspnoea: Week 61 (Arm B n=13)(Arm A n=43)	0.85 (± 1.04)	0.23 (± 0.90)		

Dyspnoea: Week 62 (Arm B n=13)(Arm A n=37)	0.95 (± 1.05)	0.16 (± 0.92)		
Dyspnoea: Week 63 (Arm B n=12)(Arm A n=36)	0.67 (± 1.01)	0.27 (± 0.95)		
Dyspnoea: Week 64 (Arm B n=10)(Arm A n=35)	0.46 (± 0.78)	0.14 (± 0.93)		
Dyspnoea: Week 65 (Arm B n=9)(Arm A n=33)	0.40 (± 0.66)	0.12 (± 0.95)		
Dyspnoea: Week 66 (Arm B n=6)(Arm A n=34)	0.67 (± 0.74)	0.16 (± 0.94)		
Dyspnoea: Week 67 (Arm B n=7)(Arm A n=30)	0.37 (± 0.76)	0.26 (± 1.01)		
Dyspnoea: Week 68 (Arm B n=6)(Arm A n=30)	0.47 (± 0.80)	0.25 (± 1.03)		
Dyspnoea: Week 69 (Arm B n=7)(Arm A n=29)	0.43 (± 0.80)	0.16 (± 1.11)		
Dyspnoea: Week 70 (Arm B n=6)(Arm A n=30)	0.33 (± 0.83)	0.07 (± 0.89)		
Dyspnoea: Week 71 (Arm B n=5)(Arm A n=26)	0.40 (± 0.91)	0.13 (± 0.89)		
Dyspnoea: Week 72 (Arm B n=4)(Arm A n=25)	-0.05 (± 0.25)	0.06 (± 0.91)		
Dyspnoea: Week 73 (Arm B n=4)(Arm A n=24)	0.45 (± 1.04)	0.09 (± 0.99)		
Dyspnoea: Week 74 (Arm B n=4)(Arm A n=23)	0.50 (± 1.01)	0.27 (± 1.19)		
Dyspnoea: Week 75 (Arm B n=4)(Arm A n=21)	0.55 (± 0.97)	0.45 (± 1.31)		
Dyspnoea: Week 76 (Arm B n=3)(Arm A n=17)	-0.07 (± 0.12)	0.02 (± 1.10)		
Dyspnoea: Week 77 (Arm B n=2)(Arm A n=17)	0.00 (± 0.00)	0.09 (± 1.17)		
Dyspnoea: Week 78 (Arm B n=2)(Arm A n=14)	0.00 (± 0.00)	0.03 (± 1.19)		
Dyspnoea: Week 79 (Arm B n=2)(Arm A n=14)	0.00 (± 0.00)	0.19 (± 1.18)		
Dyspnoea: Week 80 (Arm B n=2)(Arm A n=13)	0.10 (± 0.14)	-0.26 (± 0.80)		
Dyspnoea: Week 81 (Arm B n=2)(Arm A n=12)	0.10 (± 0.14)	-0.17 (± 1.15)		
Dyspnoea: Week 82 (Arm B n=2)(Arm A n=9)	0.00 (± 0.00)	0.18 (± 1.54)		
Dyspnoea: Week 83 (Arm B n=2)(Arm A n=4)	0.00 (± 0.00)	0.45 (± 1.22)		
Dyspnoea: Week 84 (Arm B n=2)(Arm A n=5)	0.00 (± 0.00)	0.60 (± 1.10)		
Dyspnoea: Week 85 (Arm B n=1)(Arm A n=5)	0.20 (± 999999)	0.32 (± 1.53)		
Dyspnoea: Week 86 (Arm B n=1)(Arm A n=4)	0.00 (± 999999)	-0.15 (± 0.34)		
Dyspnoea: Week 87 (Arm B n=1)(Arm A n=3)	0.20 (± 999999)	-0.47 (± 0.23)		
Dyspnoea: Week 88 (Arm B n=1)(Arm A n=4)	0.20 (± 999999)	-0.70 (± 0.20)		
Dyspnoea: Week 89 (Arm B n=1)(Arm A n=2)	0.20 (± 999999)	-0.20 (± 0.57)		
Dyspnoea: Week 90 (Arm B n=1)(Arm A n=2)	0.20 (± 999999)	-0.50 (± 0.14)		
Dyspnoea: Week 91 (Arm B n=1)(Arm A n=2)	0.00 (± 999999)	-0.50 (± 0.14)		
Dyspnoea: Week 92 (Arm B n=0)(Arm A n=2)	999999 (± 999999)	-0.70 (± 0.14)		

Dyspnoea: Week 93 (Arm B n=0)(Arm A n=2)	999999 (± 999999)	-0.40 (± 0.28)		
Dyspnoea: Week 94 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	-0.80 (± 999999)		
Dyspnoea: Week 95 (Arm B n=0)(Arm A n=1)	999999 (± 999999)	-1.00 (± 999999)		
Dyspnoea: ToF Pd (Arm B n=110)(Arm A n=96)	0.58 (± 0.99)	0.40 (± 1.06)		
Dyspnoea: ToL Tx Dose (Arm B n=141)(Arm A n=150)	0.47 (± 0.95)	0.18 (± 0.94)		

## Statistical analyses

No statistical analyses for this end point

## Secondary: Minimum Observed Serum Atezolizumab Concentration (Cmin) Prior to Infusion

End point title	Minimum Observed Serum Atezolizumab Concentration (Cmin) Prior to Infusion <sup>[1]</sup>
-----------------	---

End point description:

Minimum observed serum atezolizumab concentration (Cmin) prior to infusion at selected cycles (Arm A). Note: 999999=not available.

End point type	Secondary
----------------	-----------

End point timeframe:

Predose (Prd; 0 hour [h]) on D1 of Cy1,2,3,4,8,16 (Cy length=21 days) and thereafter on D1 of every 8th cycle (up to approximately 25 months)

Notes:

[1] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: No statistical analysis for this end point.

End point values	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)			
Subject group type	Reporting group			
Number of subjects analysed	281			
Units: µg/mL				
number (not applicable)				
Cy1D1 Nominal Time From First Dose Day 0	999999			
Cy1D1 Nominal Time From First Dose Day 0.0625	0.0300			
Cy2D1 Nominal Time From First Dose Day 21	0.0300			
Cy3D1 Nominal Time From First Dose Day 42	0.0300			
Cy4D1 Nominal Time From First Dose Day 63	0.0300			
Cy8D1 Nominal Time From First Dose Day is Day 147	49.0			
Cy16D1 Nominal Time From First Dose Day is Day 315	29.3			
Cy24D1 Nominal Time From First Dose Day is Day 483	110			



Treatment Discontinuation Visit	0.0300			
Day 120 Post Last Dose	0.0300			

## Statistical analyses

No statistical analyses for this end point

## Secondary: Maximum Observed Serum Atezolizumab Concentration (Cmax) Prior to Infusion

End point title	Maximum Observed Serum Atezolizumab Concentration (Cmax) Prior to Infusion <sup>[2]</sup>
-----------------	---

End point description:

Maximum observed serum atezolizumab concentration (Cmax) after infusion (Arm A)

End point type	Secondary
----------------	-----------

End point timeframe:

Prd(0h) on D1 of Cy1,2,3,4,8,16(Cy length=21 days)&thereafter on D1 of every 8th cy;0.5h post infusion (infusion duration=1h) on Cy1D1;at treatment discontinuation &then every 30 days (up to 120 days) after atezolizumab last dose (up to approx 25 months)

Notes:

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: No statistical analysis for this end point.

End point values	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)			
Subject group type	Reporting group			
Number of subjects analysed	281			
Units: µg/mL				
number (not applicable)				
Cy1D1 Nominal Time From First Dose Day 0	0.00			
Cy1D1 Nominal Time From First Dose Day 0.0625	691			
Cy2D1 Nominal Time From First Dose Day 21	196			
Cy3D1 Nominal Time From First Dose Day 42	269			
Cy4D1 Nominal Time From First Dose Day 63	362			
Cy8D1 Nominal Time From First Dose Day is Day 147	556			
Cy16D1 Nominal Time From First Dose Day is Day 315	480			
Cy24D1 Nominal Time From First Dose Day is Day 483	386			
Treatment Discontinuation Visit	361			
Day 120 Post Last Dose	96.8			

## Statistical analyses

No statistical analyses for this end point

### Secondary: Plasma Concentrations for Carboplatin in Arm A(Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)

End point title	Plasma Concentrations for Carboplatin in Arm A(Atezolizumab + Carboplatin or Cisplatin + Pemetrexed) <sup>[3]</sup>
-----------------	---

End point description:

Note: 999999 = not available.

End point type	Secondary
----------------	-----------

End point timeframe:

Prd (0 h), 5-10 minutes (mins) before end of carboplatin infusion (infusion duration=1-2 h), 1 h post-infusion on D1 of Cy1,3 (Cy length=21 days)

Notes:

[3] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: No statistical analysis for this end point.

End point values	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)			
Subject group type	Reporting group			
Number of subjects analysed	27			
Units: ng/mL				
arithmetic mean (standard deviation)				
Cy1D1 Nominal Time From First Dose Day 0	999999 (± 999999)			
Cy1D1 Nominal time from first dose Day 0.01389	14900 (± 4260)			
Cy1D1 Nominal Time From First Dose Day 0.0625	12800 (± 4470)			
Cy3D1 Nominal Time From First Dose Day 42	220 (± 83.8)			
Cy3D1 Nominal Time From First Dose Day 42.01389	17900 (± 4390)			
Cy3D1 Nominal Time From First Dose Day 42.0625	13900 (± 4080)			

## Statistical analyses

No statistical analyses for this end point

### Secondary: Plasma Concentrations for Cisplatin in Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)

End point title	Plasma Concentrations for Cisplatin in Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed) <sup>[4]</sup>
-----------------	--

End point description:

Note: 999999=not available.

End point type	Secondary
----------------	-----------

End point timeframe:

Prd (0 h), 5-10 mins before end of cisplatin infusion (infusion duration=30-60 mins), 1 h post-infusion

## Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: No statistical analysis for this end point.

End point values	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)			
Subject group type	Reporting group			
Number of subjects analysed	10			
Units: ng/mL				
arithmetic mean (standard deviation)				
Cy1D1 Nominal Time From First Dose Day 0	999999 (± 999999)			
Cy1D1 Nominal Time From First Dose Day 0.05556	3630 (± 589)			
Cy1D1 Nominal Time From First Dose Day 0.10417	2400 (± 360)			
Cy3D1 Nominal Time From First Dose Day 42	290 (± 86.1)			
Cy3D1 Nominal Time From First Dose Day 42.05556	3020 (± 968)			
Cy3D1 Nominal Time From First Dose Day 42.10417	2740 (± 543)			

## Statistical analyses

No statistical analyses for this end point

## Secondary: Plasma Concentrations for Pemetrexed in Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)

End point title	Plasma Concentrations for Pemetrexed in Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed) <sup>[5]</sup>
-----------------	---

End point description:

Note: 999999=not available.

End point type	Secondary
----------------	-----------

End point timeframe:

Prd (0 h), 5-10 mins before end of pemetrexed infusion (infusion duration=10 mins), 1 h post-infusion on D1 of Cy1,3 (Cy length=21 days)

## Notes:

[5] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: No statistical analysis for this end point.

End point values	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)			
Subject group type	Reporting group			
Number of subjects analysed	36			
Units: ng/mL				

arithmetic mean (standard deviation)				
Cy1D1 Nominal Time From First Dose Day 0	999999 (± 999999)			
Cy1D1 Nominal Time From First Dose Day 0.00347	86500 (± 41600)			
Cy1D1 Nominal Time From First Dose Day 0.04861	43600 (± 15800)			
Cy3D1 Nominal Time From First Dose Day 42	1.83 (± 0.681)			
Cy3D1 Nominal Time From First Dose Day 42.00347	79400 (± 44400)			
Cy3D1 Nominal Time From First Dose Day 42.04861	50100 (± 26100)			

## Statistical analyses

No statistical analyses for this end point

## Secondary: Percentage of Participants With Anti-Therapeutic Antibodies (ATAs) of Atezolizumab

End point title	Percentage of Participants With Anti-Therapeutic Antibodies (ATAs) of Atezolizumab <sup>[6]</sup>
-----------------	---

End point description:

Baseline prevalence and post-baseline incidence of anti-drug antibodies (ADA) to Atezolizumab in the Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)

End point type	Secondary
----------------	-----------

End point timeframe:

Prd (0 h) on D1 of Cy1,2,3,4,8,16 (Cy length=21 days) and thereafter on D1 of every 8th cycle, at treatment discontinuation & then every 30 days (up to 120 days) after last dose of atezolizumab (up to app 25 months)

Notes:

[6] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: No statistical analysis for this end point.

<b>End point values</b>	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)			
Subject group type	Reporting group			
Number of subjects analysed	291			
Units: Percentage of Participants				
number (not applicable)				
Baseline Evaluable Participants	1.8			
Post-Baseline Evaluable Participants	35.4			

## Statistical analyses

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

From the first study drug to the data cutoff date: 18 July 2019 (up to approximately 39 months).

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	22.0
--------------------	------

### Reporting groups

Reporting group title	Arm B (Carboplatin or Cisplatin + Pemetrexed)
-----------------------	---

Reporting group description:

Participants received IV infusion of 500 mg/m<sup>2</sup> pemetrexed on Day 1 q3w, and as per investigator's choice of either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain AUC =6 mg/mL/min or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period for 4 or 6 cycles (Cycle length=21 days). Participants who do not experience disease progression during the induction phase will begin maintenance therapy. Participants will receive IV infusion of 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

Reporting group title	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)
-----------------------	--

Reporting group description:

Participants received intravenous (IV) infusion of 1200 milligrams (mg) of atezolizumab on Day 1 every 3 weeks (q3w), IV infusion of 500 milligrams per meter square (mg/m<sup>2</sup>) pemetrexed on Day 1 q3w, and as per investigator's choice either IV infusion of carboplatin on Day 1 q3w with a dose calculated using 'Calvert formula' to obtain area under concentration versus time (AUC) = 6 milligrams per milliliter per minute (mg/mL/min) or IV infusion of 75 mg/m<sup>2</sup> cisplatin q3w on Day 1 q3w, during induction dosing period of 4 or 6 cycles (Cycle length=21 days). Participants who experienced clinical benefit during the induction phase began maintenance therapy. Participants will receive IV infusion of 1200 mg of atezolizumab and 500 mg/m<sup>2</sup> of pemetrexed on Day 1 q3w until disease progression in the maintenance period.

Serious adverse events	Arm B (Carboplatin or Cisplatin + Pemetrexed)	Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)	
Total subjects affected by serious adverse events			
subjects affected / exposed	89 / 274 (32.48%)	146 / 291 (50.17%)	
number of deaths (all causes)	187	192	
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
ADENOCARCINOMA GASTRIC			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
CANCER PAIN			

subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HISTIOCYTIC NECROTISING LYMPHADENITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
METASTASES TO CENTRAL NERVOUS SYSTEM			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
PROSTATE CANCER RECURRENT			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
TUMOUR EMBOLISM			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 1	0 / 1	
Vascular disorders			
AORTIC EMBOLUS			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
DEEP VEIN THROMBOSIS			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
EMBOLISM			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HYPERTENSION			

subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPOTENSION</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>PERIPHERAL ARTERY THROMBOSIS</b>			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>PERIPHERAL ISCHAEMIA</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>THROMBOPHLEBITIS</b>			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>VASCULITIS</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>General disorders and administration site conditions</b>			
<b>ASTHENIA</b>			
subjects affected / exposed	2 / 274 (0.73%)	5 / 291 (1.72%)	
occurrences causally related to treatment / all	0 / 2	3 / 5	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>CHEST PAIN</b>			
subjects affected / exposed	2 / 274 (0.73%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	1 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>DEATH</b>			

subjects affected / exposed	1 / 274 (0.36%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	1 / 1	1 / 2	
deaths causally related to treatment / all	1 / 1	1 / 2	
<b>DISEASE SUSCEPTIBILITY</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>FATIGUE</b>			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>GAIT DISTURBANCE</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>GENERAL PHYSICAL HEALTH DETERIORATION</b>			
subjects affected / exposed	1 / 274 (0.36%)	3 / 291 (1.03%)	
occurrences causally related to treatment / all	1 / 1	2 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>INFLAMMATION</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>MALAISE</b>			
subjects affected / exposed	2 / 274 (0.73%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 2	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>MUCOSAL INFLAMMATION</b>			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>NON-CARDIAC CHEST PAIN</b>			



subjects affected / exposed	2 / 274 (0.73%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>PAIN</b>			
subjects affected / exposed	2 / 274 (0.73%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>PYREXIA</b>			
subjects affected / exposed	1 / 274 (0.36%)	12 / 291 (4.12%)	
occurrences causally related to treatment / all	0 / 2	6 / 12	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Immune system disorders</b>			
<b>CYTOKINE RELEASE SYNDROME</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPERSENSITIVITY</b>			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Reproductive system and breast disorders</b>			
<b>BENIGN PROSTATIC HYPERPLASIA</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Respiratory, thoracic and mediastinal disorders</b>			
<b>ATELECTASIS</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>BRONCHIAL HYPERREACTIVITY</b>			

subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
CHRONIC OBSTRUCTIVE PULMONARY DISEASE			
subjects affected / exposed	2 / 274 (0.73%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 2	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
DYSPNOEA			
subjects affected / exposed	3 / 274 (1.09%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 3	0 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
EPISTAXIS			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HAEMOPTYSIS			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HYPERVENTILATION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
INTERSTITIAL LUNG DISEASE			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
ORGANISING PNEUMONIA			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
PHARYNGEAL INFLAMMATION			

subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
PLEURAL EFFUSION			
subjects affected / exposed	3 / 274 (1.09%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 4	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
PNEUMONIA ASPIRATION			
subjects affected / exposed	1 / 274 (0.36%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 1	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
PNEUMONITIS			
subjects affected / exposed	4 / 274 (1.46%)	9 / 291 (3.09%)	
occurrences causally related to treatment / all	4 / 4	9 / 9	
deaths causally related to treatment / all	1 / 1	1 / 1	
PNEUMOTHORAX			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
PULMONARY EMBOLISM			
subjects affected / exposed	4 / 274 (1.46%)	4 / 291 (1.37%)	
occurrences causally related to treatment / all	0 / 4	0 / 4	
deaths causally related to treatment / all	0 / 0	0 / 2	
PULMONARY HAEMORRHAGE			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
RESPIRATORY FAILURE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
Psychiatric disorders			
ANXIETY			

subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
CATATONIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
COMPLETED SUICIDE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
Product issues			
DEVICE MALFUNCTION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Investigations			
ALANINE AMINOTRANSFERASE INCREASED			
subjects affected / exposed	1 / 274 (0.36%)	3 / 291 (1.03%)	
occurrences causally related to treatment / all	1 / 1	2 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
ASPARTATE AMINOTRANSFERASE INCREASED			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
BLOOD CREATININE INCREASED			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
CREATININE RENAL CLEARANCE DECREASED			

subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
NEUTROPHIL COUNT DECREASED			
subjects affected / exposed	3 / 274 (1.09%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	5 / 5	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
PLATELET COUNT DECREASED			
subjects affected / exposed	3 / 274 (1.09%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	3 / 3	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
WEIGHT DECREASED			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
WHITE BLOOD CELL COUNT DECREASED			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injury, poisoning and procedural complications			
FEMUR FRACTURE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
INFUSION RELATED REACTION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
INTENTIONAL OVERDOSE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
LUMBAR VERTEBRAL FRACTURE			

subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
SKIN INJURY			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
STAB WOUND			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
SUBDURAL HAEMATOMA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
TOXICITY TO VARIOUS AGENTS			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac disorders			
ACUTE MYOCARDIAL INFARCTION			
subjects affected / exposed	2 / 274 (0.73%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 3	0 / 1	
deaths causally related to treatment / all	0 / 1	0 / 1	
ATRIAL FIBRILLATION			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
CARDIAC FAILURE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
CARDIAC FAILURE ACUTE			

subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	1 / 1	0 / 0	
MYOCARDITIS			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
STRESS CARDIOMYOPATHY			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
SUPRAVENTRICULAR TACHYCARDIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
TACHYCARDIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
APHASIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
ATAXIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
BRAIN OEDEMA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
CAROTID ARTERY OCCLUSION			

subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
CAROTID ARTERY STENOSIS			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
CEREBRAL HAEMORRHAGE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
CEREBRAL INFARCTION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
CEREBROVASCULAR ACCIDENT			
subjects affected / exposed	1 / 274 (0.36%)	4 / 291 (1.37%)	
occurrences causally related to treatment / all	1 / 1	0 / 4	
deaths causally related to treatment / all	1 / 1	0 / 1	
HEADACHE			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
LETHARGY			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
SEIZURE			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
SYNCOPE			



subjects affected / exposed	2 / 274 (0.73%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
TRANSIENT ISCHAEMIC ATTACK			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood and lymphatic system disorders			
ANAEMIA			
subjects affected / exposed	7 / 274 (2.55%)	10 / 291 (3.44%)	
occurrences causally related to treatment / all	9 / 10	13 / 13	
deaths causally related to treatment / all	0 / 0	0 / 0	
BONE MARROW FAILURE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
FEBRILE BONE MARROW APLASIA			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
FEBRILE NEUTROPENIA			
subjects affected / exposed	5 / 274 (1.82%)	12 / 291 (4.12%)	
occurrences causally related to treatment / all	3 / 5	12 / 12	
deaths causally related to treatment / all	0 / 0	0 / 0	
LEUKOPENIA			
subjects affected / exposed	1 / 274 (0.36%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	1 / 1	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
NEUTROPENIA			
subjects affected / exposed	3 / 274 (1.09%)	3 / 291 (1.03%)	
occurrences causally related to treatment / all	3 / 3	3 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
PANCYTOPENIA			

subjects affected / exposed	4 / 274 (1.46%)	4 / 291 (1.37%)	
occurrences causally related to treatment / all	6 / 6	4 / 4	
deaths causally related to treatment / all	1 / 1	0 / 0	
THROMBOCYTOPENIA			
subjects affected / exposed	4 / 274 (1.46%)	11 / 291 (3.78%)	
occurrences causally related to treatment / all	4 / 4	14 / 15	
deaths causally related to treatment / all	0 / 0	0 / 0	
Eye disorders			
CATARACT			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	0 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			
ABDOMINAL PAIN			
subjects affected / exposed	2 / 274 (0.73%)	4 / 291 (1.37%)	
occurrences causally related to treatment / all	0 / 2	3 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
AUTOIMMUNE COLITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
COLITIS			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
CONSTIPATION			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
DIARRHOEA			
subjects affected / exposed	3 / 274 (1.09%)	9 / 291 (3.09%)	
occurrences causally related to treatment / all	2 / 3	8 / 9	
deaths causally related to treatment / all	0 / 0	0 / 0	
DIVERTICULUM INTESTINAL			

HAEMORRHAGIC			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
ENTEROCOLITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
GASTRIC PERFORATION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
GASTROINTESTINAL DISORDER			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
INGUINAL HERNIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
INTESTINAL OBSTRUCTION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
NAUSEA			
subjects affected / exposed	2 / 274 (0.73%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	2 / 2	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
OESOPHAGITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	3 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
PANCREATITIS			

subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
PROCTITIS ULCERATIVE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
SMALL INTESTINAL PERFORATION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
STOMATITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
UPPER GASTROINTESTINAL HAEMORRHAGE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
VOMITING			
subjects affected / exposed	0 / 274 (0.00%)	6 / 291 (2.06%)	
occurrences causally related to treatment / all	0 / 0	4 / 6	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatobiliary disorders			
AUTOIMMUNE HEPATITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
BILE DUCT OBSTRUCTION			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
CHOLANGITIS			

subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
CHOLELITHIASIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
DRUG-INDUCED LIVER INJURY			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
HEPATIC CONGESTION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HEPATITIS			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HEPATITIS ACUTE			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
HEPATOTOXICITY			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	1 / 1	
Skin and subcutaneous tissue disorders			
DERMATITIS			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
RASH			

subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
RASH MACULO-PAPULAR			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal and urinary disorders			
ACUTE KIDNEY INJURY			
subjects affected / exposed	1 / 274 (0.36%)	3 / 291 (1.03%)	
occurrences causally related to treatment / all	1 / 1	2 / 3	
deaths causally related to treatment / all	0 / 0	0 / 1	
CHRONIC KIDNEY DISEASE			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	2 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
GLYCOSURIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
NEPHRITIS ALLERGIC			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
RENAL FAILURE			
subjects affected / exposed	2 / 274 (0.73%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 2	1 / 1	
deaths causally related to treatment / all	0 / 1	0 / 0	
RENAL IMPAIRMENT			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
TUBULOINTERSTITIAL NEPHRITIS			

subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
Endocrine disorders			
HYPOTHYROIDISM			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Musculoskeletal and connective tissue disorders			
BACK PAIN			
subjects affected / exposed	2 / 274 (0.73%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
BONE PAIN			
subjects affected / exposed	2 / 274 (0.73%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
FRACTURE PAIN			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
MUSCULOSKELETAL CHEST PAIN			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
MYALGIA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
NECK PAIN			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Infections and infestations BACTERAEMIA subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 274 (0.36%) 0 / 1 0 / 0	0 / 291 (0.00%) 0 / 0 0 / 0	
BRONCHITIS subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 274 (0.00%) 0 / 0 0 / 0	2 / 291 (0.69%) 0 / 2 0 / 0	
BRONCHOPULMONARY ASPERGILLOSIS subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 274 (0.00%) 0 / 0 0 / 0	1 / 291 (0.34%) 0 / 1 0 / 0	
CAMPYLOBACTER GASTROENTERITIS subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 274 (0.00%) 0 / 0 0 / 0	1 / 291 (0.34%) 0 / 1 0 / 0	
CATHETER SITE CELLULITIS subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 274 (0.00%) 0 / 0 0 / 0	1 / 291 (0.34%) 0 / 1 0 / 0	
CELLULITIS subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	2 / 274 (0.73%) 1 / 2 0 / 0	3 / 291 (1.03%) 2 / 3 0 / 0	
DIVERTICULITIS subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 274 (0.36%) 0 / 1 0 / 0	1 / 291 (0.34%) 1 / 1 0 / 0	
ENCEPHALITIS subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 274 (0.00%) 0 / 0 0 / 0	1 / 291 (0.34%) 1 / 1 0 / 0	



ERYSIPELAS			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
ESCHERICHIA URINARY TRACT INFECTION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
H3N2 INFLUENZA			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 1	
INFECTIOUS PLEURAL EFFUSION			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
INFLUENZA			
subjects affected / exposed	1 / 274 (0.36%)	3 / 291 (1.03%)	
occurrences causally related to treatment / all	0 / 1	1 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
LOWER RESPIRATORY TRACT INFECTION			
subjects affected / exposed	2 / 274 (0.73%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
LUNG INFECTION			
subjects affected / exposed	3 / 274 (1.09%)	5 / 291 (1.72%)	
occurrences causally related to treatment / all	0 / 3	2 / 5	
deaths causally related to treatment / all	0 / 0	0 / 0	
MENINGITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
NEUTROPENIC SEPSIS			

subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
OESOPHAGEAL CANDIDIASIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
PERIPHERAL NERVE INFECTION			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
PHARYNGITIS			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
PNEUMONIA			
subjects affected / exposed	13 / 274 (4.74%)	13 / 291 (4.47%)	
occurrences causally related to treatment / all	5 / 14	4 / 13	
deaths causally related to treatment / all	3 / 4	1 / 2	
PULMONARY SEPSIS			
subjects affected / exposed	0 / 274 (0.00%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 0	1 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
PYELONEPHRITIS			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
RESPIRATORY TRACT INFECTION			
subjects affected / exposed	2 / 274 (0.73%)	5 / 291 (1.72%)	
occurrences causally related to treatment / all	0 / 3	1 / 8	
deaths causally related to treatment / all	0 / 0	0 / 1	
SEPSIS			

subjects affected / exposed	2 / 274 (0.73%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 2	2 / 2	
deaths causally related to treatment / all	0 / 1	0 / 0	
SOFT TISSUE INFECTION			
subjects affected / exposed	1 / 274 (0.36%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 1	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
TRACHEITIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
URINARY TRACT INFECTION			
subjects affected / exposed	2 / 274 (0.73%)	6 / 291 (2.06%)	
occurrences causally related to treatment / all	1 / 2	2 / 6	
deaths causally related to treatment / all	0 / 0	0 / 0	
UROSEPSIS			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Metabolism and nutrition disorders			
DECREASED APPETITE			
subjects affected / exposed	2 / 274 (0.73%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	1 / 2	1 / 1	
deaths causally related to treatment / all	0 / 0	1 / 1	
DEHYDRATION			
subjects affected / exposed	4 / 274 (1.46%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	0 / 4	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	
DIABETES MELLITUS INADEQUATE CONTROL			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
GOUT			

subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPERCALCAEMIA</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPERGLYCAEMIA</b>			
subjects affected / exposed	0 / 274 (0.00%)	3 / 291 (1.03%)	
occurrences causally related to treatment / all	0 / 0	0 / 4	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPERKALAEMIA</b>			
subjects affected / exposed	1 / 274 (0.36%)	0 / 291 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPOKALAEMIA</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPOMAGNEAEMIA</b>			
subjects affected / exposed	0 / 274 (0.00%)	1 / 291 (0.34%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>HYPONATRAEMIA</b>			
subjects affected / exposed	1 / 274 (0.36%)	2 / 291 (0.69%)	
occurrences causally related to treatment / all	1 / 1	2 / 2	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	<b>Arm B (Carboplatin or Cisplatin + Pemetrexed)</b>	<b>Arm A (Atezolizumab + Carboplatin or Cisplatin + Pemetrexed)</b>	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	254 / 274 (92.70%)	275 / 291 (94.50%)	
Vascular disorders			
HYPERTENSION			
subjects affected / exposed	6 / 274 (2.19%)	17 / 291 (5.84%)	
occurrences (all)	8	20	
General disorders and administration site conditions			
ASTHENIA			
subjects affected / exposed	54 / 274 (19.71%)	79 / 291 (27.15%)	
occurrences (all)	81	116	
CHEST PAIN			
subjects affected / exposed	18 / 274 (6.57%)	21 / 291 (7.22%)	
occurrences (all)	18	24	
FATIGUE			
subjects affected / exposed	68 / 274 (24.82%)	69 / 291 (23.71%)	
occurrences (all)	98	110	
MALAISE			
subjects affected / exposed	17 / 274 (6.20%)	14 / 291 (4.81%)	
occurrences (all)	23	23	
MUCOSAL INFLAMMATION			
subjects affected / exposed	19 / 274 (6.93%)	23 / 291 (7.90%)	
occurrences (all)	28	24	
OEDEMA PERIPHERAL			
subjects affected / exposed	33 / 274 (12.04%)	45 / 291 (15.46%)	
occurrences (all)	38	63	
PYREXIA			
subjects affected / exposed	37 / 274 (13.50%)	56 / 291 (19.24%)	
occurrences (all)	50	84	
Respiratory, thoracic and mediastinal disorders			
COUGH			
subjects affected / exposed	30 / 274 (10.95%)	43 / 291 (14.78%)	
occurrences (all)	36	54	
DYSPNOEA			

subjects affected / exposed occurrences (all)	40 / 274 (14.60%) 42	42 / 291 (14.43%) 48	
EPISTAXIS subjects affected / exposed occurrences (all)	19 / 274 (6.93%) 22	13 / 291 (4.47%) 16	
HICCUPS subjects affected / exposed occurrences (all)	17 / 274 (6.20%) 24	16 / 291 (5.50%) 35	
RHINORRHOEA subjects affected / exposed occurrences (all)	14 / 274 (5.11%) 15	8 / 291 (2.75%) 10	
Psychiatric disorders INSOMNIA subjects affected / exposed occurrences (all)	18 / 274 (6.57%) 18	28 / 291 (9.62%) 35	
Investigations ALANINE AMINOTRANSFERASE INCREASED subjects affected / exposed occurrences (all)	23 / 274 (8.39%) 29	52 / 291 (17.87%) 73	
ASPARTATE AMINOTRANSFERASE INCREASED subjects affected / exposed occurrences (all)	27 / 274 (9.85%) 34	51 / 291 (17.53%) 78	
BLOOD CREATININE INCREASED subjects affected / exposed occurrences (all)	20 / 274 (7.30%) 26	34 / 291 (11.68%) 42	
NEUTROPHIL COUNT DECREASED subjects affected / exposed occurrences (all)	48 / 274 (17.52%) 108	45 / 291 (15.46%) 115	
PLATELET COUNT DECREASED subjects affected / exposed occurrences (all)	38 / 274 (13.87%) 69	38 / 291 (13.06%) 63	
WEIGHT DECREASED subjects affected / exposed occurrences (all)	16 / 274 (5.84%) 17	26 / 291 (8.93%) 26	
WHITE BLOOD CELL COUNT DECREASED			

subjects affected / exposed occurrences (all)	29 / 274 (10.58%) 67	20 / 291 (6.87%) 55	
Nervous system disorders			
DIZZINESS			
subjects affected / exposed	24 / 274 (8.76%)	19 / 291 (6.53%)	
occurrences (all)	27	22	
DYSGEUSIA			
subjects affected / exposed	19 / 274 (6.93%)	28 / 291 (9.62%)	
occurrences (all)	19	33	
HEADACHE			
subjects affected / exposed	23 / 274 (8.39%)	35 / 291 (12.03%)	
occurrences (all)	25	38	
PARAESTHESIA			
subjects affected / exposed	14 / 274 (5.11%)	10 / 291 (3.44%)	
occurrences (all)	16	10	
Blood and lymphatic system disorders			
ANAEMIA			
subjects affected / exposed	113 / 274 (41.24%)	128 / 291 (43.99%)	
occurrences (all)	140	182	
NEUTROPENIA			
subjects affected / exposed	38 / 274 (13.87%)	47 / 291 (16.15%)	
occurrences (all)	65	94	
THROMBOCYTOPENIA			
subjects affected / exposed	23 / 274 (8.39%)	39 / 291 (13.40%)	
occurrences (all)	37	60	
Eye disorders			
LACRIMATION INCREASED			
subjects affected / exposed	18 / 274 (6.57%)	16 / 291 (5.50%)	
occurrences (all)	21	16	
Gastrointestinal disorders			
ABDOMINAL PAIN UPPER			
subjects affected / exposed	5 / 274 (1.82%)	15 / 291 (5.15%)	
occurrences (all)	7	20	
CONSTIPATION			
subjects affected / exposed	79 / 274 (28.83%)	92 / 291 (31.62%)	
occurrences (all)	96	104	
DIARRHOEA			

subjects affected / exposed occurrences (all)	47 / 274 (17.15%) 56	58 / 291 (19.93%) 81	
DYSPEPSIA subjects affected / exposed occurrences (all)	9 / 274 (3.28%) 10	16 / 291 (5.50%) 18	
NAUSEA subjects affected / exposed occurrences (all)	114 / 274 (41.61%) 206	109 / 291 (37.46%) 262	
STOMATITIS subjects affected / exposed occurrences (all)	23 / 274 (8.39%) 26	35 / 291 (12.03%) 44	
VOMITING subjects affected / exposed occurrences (all)	49 / 274 (17.88%) 64	58 / 291 (19.93%) 80	
Skin and subcutaneous tissue disorders DRY SKIN subjects affected / exposed occurrences (all)	7 / 274 (2.55%) 7	19 / 291 (6.53%) 22	
PRURITUS subjects affected / exposed occurrences (all)	16 / 274 (5.84%) 17	27 / 291 (9.28%) 34	
RASH subjects affected / exposed occurrences (all)	21 / 274 (7.66%) 23	41 / 291 (14.09%) 53	
Endocrine disorders HYPOTHYROIDISM subjects affected / exposed occurrences (all)	2 / 274 (0.73%) 2	16 / 291 (5.50%) 17	
Musculoskeletal and connective tissue disorders ARTHRALGIA subjects affected / exposed occurrences (all)	17 / 274 (6.20%) 19	33 / 291 (11.34%) 38	
BACK PAIN subjects affected / exposed occurrences (all)	25 / 274 (9.12%) 28	38 / 291 (13.06%) 45	
PAIN IN EXTREMITY			



subjects affected / exposed occurrences (all)	13 / 274 (4.74%) 13	22 / 291 (7.56%) 22	
Infections and infestations			
CONJUNCTIVITIS			
subjects affected / exposed	15 / 274 (5.47%)	17 / 291 (5.84%)	
occurrences (all)	18	22	
UPPER RESPIRATORY TRACT INFECTION			
subjects affected / exposed	10 / 274 (3.65%)	16 / 291 (5.50%)	
occurrences (all)	13	22	
Metabolism and nutrition disorders			
DECREASED APPETITE			
subjects affected / exposed	64 / 274 (23.36%)	77 / 291 (26.46%)	
occurrences (all)	84	100	
HYPERGLYCAEMIA			
subjects affected / exposed	17 / 274 (6.20%)	13 / 291 (4.47%)	
occurrences (all)	20	15	
HYPOKALAEMIA			
subjects affected / exposed	4 / 274 (1.46%)	21 / 291 (7.22%)	
occurrences (all)	4	29	
HYPOMAGNESAEMIA			
subjects affected / exposed	14 / 274 (5.11%)	19 / 291 (6.53%)	
occurrences (all)	19	31	

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
24 November 2015	Protocol was amended to change OS rate at 3 years from Secondary Efficacy Objective to Exploratory Outcome Measure. Exclusion Criteria Treatment with systemic immunostimulatory agents within 4 weeks prior to randomization updated to include or 5 half-lives of the drug, whichever is longer.
09 June 2016	Protocol was amended to change the assessment of the patients for the objectives to chemotherapy-naïve and have stage IV non-squamous NSCLC (the ITT population) treated with atezolizumab + carboplatin or cisplatin + pemetrexed (Arm A) in comparison to carboplatin or cisplatin + pemetrexed (Arm B). Previous primary efficacy objectives were removed with the exception of investigator-assessed PFS and OS was added. Secondary Efficacy Objectives were updated with the change in patient assessment. Secondary Efficacy Objective investigator-assessed PFS replaced with IRF-assessed PFS. Time to response added to Secondary Efficacy Objective. "Patients with a sensitizing mutation in the EGFR gene or an ALK fusion oncogene" added to cancer specific exclusion criteria. "Or cerebrovascular accident" added to "Significant cardiovascular disease, such as New York Heart Association cardiac disease (Class II or greater), myocardial infarction within 3 months prior to randomization, unstable arrhythmias, or unstable angina." General Medical Exclusions. "Illness or condition that may interfere with a patient's capacity to understand, follow, and/or comply with study procedures" added to General Medical Exclusions. "Prior treatment with EGFR inhibitors or ALK inhibitors" added to Exclusion Criteria Related to Medications. Exclusion Criteria Related to Medications updated with "Any approved anti-cancer therapy, including hormonal therapy within 21 days prior to initiation of study treatment".
09 October 2017	Protocol was amended to remove all objectives and outcome measures based on review by IRF. Secondary efficacy objective and outcome measure for SILC scale symptom severity scores has been updated so that it will be measured from baseline instead of time to deterioration (TTD).

Notes:

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported