



## Clinical trial results:

**A Phase III, randomised, double-blind, multicentre, parallel-group, non-inferiority study evaluating the efficacy, safety, and tolerability of dolutegravir plus lamivudine compared to dolutegravir plus tenofovir/emtricitabine in HIV-1-infected treatment-naïve adults**

### Summary

EudraCT number	2015-004418-95
Trial protocol	DE ES BE NL PT FR IT
Global end of trial date	15 August 2022

### Results information

Result version number	v2 (current)
This version publication date	17 August 2023
First version publication date	11 April 2019
Version creation reason	

### Trial information

#### Trial identification

Sponsor protocol code	204861
-----------------------	--------

#### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	-
WHO universal trial number (UTN)	-

Notes:

### Sponsors

Sponsor organisation name	ViiV Healthcare
Sponsor organisation address	980 Great West Road, Brentford, Middlesex, United Kingdom, TW8 9GS
Public contact	GSK Response Center, ViiV Healthcare, 1 8664357343, GSKClinicalSupportHD@gsk.com
Scientific contact	GSK Response Center, ViiV Healthcare, 1 8664357343, GSKClinicalSupportHD@gsk.com

Notes:

### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	22 September 2022
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	15 August 2022
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

To demonstrate non-inferior antiviral activity of dolutegravir plus lamivudine ( DTG + 3TC) versus dolutegravir plus tenofovir/emtricitabine (DTG + TDF/FTC) at 48 weeks in HIV-1-infected, ART-naïve participants

Protection of trial subjects:

Not applicable

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	21 July 2016
Long term follow-up planned	Yes
Long term follow-up rationale	Ethical reason
Long term follow-up duration	4 Years
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Russian Federation: 75
Country: Number of subjects enrolled	South Africa: 7
Country: Number of subjects enrolled	Spain: 68
Country: Number of subjects enrolled	Taiwan: 62
Country: Number of subjects enrolled	United States: 113
Country: Number of subjects enrolled	United Kingdom: 9
Country: Number of subjects enrolled	Argentina: 105
Country: Number of subjects enrolled	Australia: 6
Country: Number of subjects enrolled	Belgium: 22
Country: Number of subjects enrolled	Canada: 33
Country: Number of subjects enrolled	France: 27
Country: Number of subjects enrolled	Germany: 19
Country: Number of subjects enrolled	Italy: 78
Country: Number of subjects enrolled	Korea, Republic of: 7
Country: Number of subjects enrolled	Mexico: 60
Country: Number of subjects enrolled	Netherlands: 4
Country: Number of subjects enrolled	Portugal: 16
Country: Number of subjects enrolled	Romania: 8

Worldwide total number of subjects	719
EEA total number of subjects	242

Notes:

<b>Subjects enrolled per age group</b>	
In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	715
From 65 to 84 years	4
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

This study is a randomized, double-blind, parallel-group, non-inferiority study. The study consisted of double blind, open label, and continuation phase.

### Pre-assignment

Screening details:

Total of 719 participants were enrolled and randomized. Five participants were randomized but not treated. A total of 714 participants received at least one dose of study treatment, following randomization creating the intent to treat exposed (ITT-E) population.

### Period 1

Period 1 title	Overall study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	DTG + 3TC-Double blind phase

Arm description:

Participants received a two-drug regimen of DTG + 3TC administered orally, once daily for 96 weeks in a double-blind phase; from Week 96 to Week 148 in an open-label phase; and from Week 148 to Week 280 in a continuation phase.

Arm type	Experimental
Investigational medicinal product name	Lamivudine
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

3TC 300 mg capsule, oral administration, once daily.

Investigational medicinal product name	Dolutegravir
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

DTG 50 mg tablet, oral administration, once daily.

<b>Arm title</b>	DTG + TDF/FTC-Double blind phase
------------------	----------------------------------

Arm description:

Participants received a three-drug regimen of DTG + TDF/FTC fixed dose combination (FDC) administered orally, once daily for 96 weeks in a double-blind phase and from Week 96 to Week 148 in an open-label phase.

Arm type	Active comparator
Investigational medicinal product name	Dolutegravir
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

DTG 50 mg tablet, oral administration, once daily.

Investigational medicinal product name	Tenofovir disoproxil fumarate/emtricitabine fixed-dose combination
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

300 mg TDF/ 200 mg FTC capsule, oral administration, once daily.

<b>Number of subjects in period 1<sup>[1]</sup></b>	<b>DTG + 3TC-Double blind phase</b>	<b>DTG + TDF/FTC-Double blind phase</b>
Started	356	358
Completed	257	293
Not completed	99	65
Physician decision	12	8
Consent withdrawn by subject	24	11
Randomized, but did not receive treatment	3	2
Adverse event, non-fatal	14	13
Protocol Deviation	7	7
Protocol Withdrawal Criterion Met	4	7
Lost to follow-up	30	13
Lack of efficacy	5	4

Notes:

[1] - The number of subjects reported to be in the baseline period are not the same as the worldwide number enrolled in the trial. It is expected that these numbers will be the same.

Justification: Total of 719 participants were enrolled and randomized. Five participants were randomized but not treated. A total of 714 participants received at least one dose of study treatment, following randomization creating the intent to treat exposed (ITT-E) population

## Baseline characteristics

### Reporting groups

Reporting group title	Overall study
Reporting group description:	
Participants received a two-drug regimen of DTG + 3TC administered orally, once daily for 96 weeks in a double-blind phase; from Week 96 to Week 148 in an open-label phase; and from Week 148 to Week 280 in a continuation phase.	

Reporting group values	Overall study	Total	
Number of subjects	714	714	
Age categorical			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			
Age Continuous			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: years			
arithmetic mean	34.5		
standard deviation	± 10.31	-	
Sex: Female, Male			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			
Female	111	111	
Male	603	603	
Race/Ethnicity, Customized			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Subjects			
American (Am) Indian or Alaska (Al.) native	59	59	
Asian-Central/South Asian heritage (H.)	4	4	
Asian - East Asian H.	69	69	
Asian - South East Asian H.	6	6	
Black or African Am	75	75	
Native Hawaiian or other Pacific Islander	2	2	
White (Wt)-Arabic/North African H.	10	10	
Wt-Wt/Caucasian (Ca.)/European (Eu.) H.	481	481	
Multiple	8	8	

### Subject analysis sets

Subject analysis set title	DTG + 3TC - Double-blind Phase + Open-label Phase
Subject analysis set type	Sub-group analysis

Subject analysis set description:

Participants received a two-drug regimen of DTG + 3TC administered orally, once daily until Week 96 in

double-blind phase and participants continued to receive DTG + 3TC from Week 96 to Week 148 in an open-label phase.

Subject analysis set title	DTG + TDF/FTC - Double-blind Phase + Open-label Phase
----------------------------	---

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a three-drug regimen of DTG + TDF/FTC FDC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + TDF/FTC FDC from Week 96 to Week 148 in an open-label phase.

Subject analysis set title	DTG + 3TC Double blind Phase
----------------------------	------------------------------

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a two-drug regimen of dolutegravir plus lamivudine (DTG + 3TC) once daily for 96 weeks in double blind phase

Subject analysis set title	DTG + 3TC Double blind Phase
----------------------------	------------------------------

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a two-drug regimen of dolutegravir plus lamivudine (DTG + 3TC) once daily for 96 weeks in the double blind phase

Subject analysis set title	DTG + 3TC - Double-blind Phase
----------------------------	--------------------------------

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a two-drug regimen of dolutegravir plus lamivudine (DTG + 3TC) once daily for 96 weeks in the double blind phase

Subject analysis set title	DTG + 3TC - Double-blind Phase + Open-label Phase
----------------------------	---

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a two-drug regimen of DTG + 3TC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + 3TC from Week 96 to Week 148 in an open-label phase.

Subject analysis set title	DTG + TDF/FTC - Double-blind Phase + Open-label Phase
----------------------------	---

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a three-drug regimen of DTG + TDF/FTC FDC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + TDF/FTC FDC from Week 96 to Week 148 in an open-label phase.

Subject analysis set title	DTG + 3TC - Double-blind Phase + Open-label Phase
----------------------------	---

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a two-drug regimen of DTG + 3TC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + 3TC from Week 96 to Week 148 in an open-label phase.

Subject analysis set title	DTG + TDF/FTC - Double-blind Phase + Open-label Phase
----------------------------	---

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

Participants received a three-drug regimen of DTG + TDF/FTC FDC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + TDF/FTC FDC from Week 96 to Week 148 in an open-label phase.

Reporting group values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase	DTG + 3TC Double blind Phase
Number of subjects	356	358	356
Age categorical			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			

Age Continuous			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: years			
arithmetic mean			
standard deviation	±	±	±
Sex: Female, Male			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			
Female	59	52	
Male	297	306	
Race/Ethnicity, Customized			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Subjects			
American (Am) Indian or Alaska (Al.) native	31	28	
Asian-Central/South Asian heritage (H.)	0	4	
Asian - East Asian H.	33	36	
Asian - South East Asian H.	4	2	
Black or African Am	39	36	
Native Hawaiian or other Pacific Islander	2	0	
White (Wt)-Arabic/North African H.	5	5	
Wt-Wt/Caucasian (Ca.)/European (Eu.) H.	239	242	
Multiple	3	5	

Reporting group values	DTG + 3TC Double blind Phase	DTG + 3TC - Double-blind Phase	DTG + 3TC - Double-blind Phase + Open-label Phase
Number of subjects	301	301	8
Age categorical			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			

Age Continuous			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: years			
arithmetic mean			
standard deviation	±	±	±
Sex: Female, Male			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			
Female			
Male			
Race/Ethnicity, Customized			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Subjects			



American (Am) Indian or Alaska (Al.) native Asian-Central/South Asian heritage (H.) Asian - East Asian H. Asian - South East Asian H. Black or African Am Native Hawaiian or other Pacific Islander White (Wt)-Arabic/North African H. Wt-Wt/Caucasian (Ca.)/European (Eu.) H. Multiple			
---	--	--	--

<b>Reporting group values</b>	DTG + TDF/FTC - Double-blind Phase + Open-label Phase	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects	8	5	6
Age categorical			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			

Age Continuous			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: years			
arithmetic mean			
standard deviation	±	±	±
Sex: Female, Male			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Participants			
Female			
Male			
Race/Ethnicity, Customized			
Baseline Characteristic data are reported for the Intent-to-Treat Exposed (ITT-E) Population which consisted of all randomized participants who received at least one dose of study treatment.			
Units: Subjects			
American (Am) Indian or Alaska (Al.) native Asian-Central/South Asian heritage (H.) Asian - East Asian H. Asian - South East Asian H. Black or African Am Native Hawaiian or other Pacific Islander White (Wt)-Arabic/North African H. Wt-Wt/Caucasian (Ca.)/European (Eu.) H. Multiple			

## End points

### End points reporting groups

Reporting group title	DTG + 3TC-Double blind phase
Reporting group description: Participants received a two-drug regimen of DTG + 3TC administered orally, once daily for 96 weeks in a double-blind phase; from Week 96 to Week 148 in an open-label phase; and from Week 148 to Week 280 in a continuation phase.	
Reporting group title	DTG + TDF/FTC-Double blind phase
Reporting group description: Participants received a three-drug regimen of DTG + TDF/FTC fixed dose combination (FDC) administered orally, once daily for 96 weeks in a double-blind phase and from Week 96 to Week 148 in an open-label phase.	
Subject analysis set title	DTG + 3TC - Double-blind Phase + Open-label Phase
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received a two-drug regimen of DTG + 3TC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + 3TC from Week 96 to Week 148 in an open-label phase.	
Subject analysis set title	DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received a three-drug regimen of DTG + TDF/FTC FDC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + TDF/FTC FDC from Week 96 to Week 148 in an open-label phase.	
Subject analysis set title	DTG + 3TC Double blind Phase
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received a two-drug regimen of dolutegravir plus lamivudine (DTG + 3TC) once daily for 96 weeks in double blind phase	
Subject analysis set title	DTG + 3TC Double blind Phase
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received a two-drug regimen of dolutegravir plus lamivudine (DTG + 3TC) once daily for 96 weeks in the double blind phase	
Subject analysis set title	DTG + 3TC - Double-blind Phase
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received a two-drug regimen of dolutegravir plus lamivudine (DTG + 3TC) once daily for 96 weeks in the double blind phase	
Subject analysis set title	DTG + 3TC - Double-blind Phase + Open-label Phase
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received a two-drug regimen of DTG + 3TC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + 3TC from Week 96 to Week 148 in an open-label phase.	
Subject analysis set title	DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received a three-drug regimen of DTG + TDF/FTC FDC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + TDF/FTC FDC from Week 96 to Week 148 in an open-label phase.	
Subject analysis set title	DTG + 3TC - Double-blind Phase + Open-label Phase
Subject analysis set type	Sub-group analysis

Subject analysis set description:

Participants received a two-drug regimen of DTG + 3TC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + 3TC from Week 96 to Week 148 in an open-label phase.

Subject analysis set title	DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Subject analysis set type	Sub-group analysis

Subject analysis set description:

Participants received a three-drug regimen of DTG + TDF/FTC FDC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + TDF/FTC FDC from Week 96 to Week 148 in an open-label phase.

### **Primary: Percentage of Participants With Plasma Human Immunodeficiency Virus Type 1 (HIV-1) Ribonucleic Acid (RNA) <50 Copies/mL (c/mL) at Week 48**

End point title	Percentage of Participants With Plasma Human Immunodeficiency Virus Type 1 (HIV-1) Ribonucleic Acid (RNA) <50 Copies/mL (c/mL) at Week 48
-----------------	---

End point description:

Percentage of participants with HIV-1 RNA<50 c/mL was obtained using Food and Drug Administration (FDA) Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant antiretroviral therapy (ART) prior to the visit of interest. This endpoint was analyzed using a stratified analysis with Cochran-Mantel-Haenszel (CMH) weights. Intent-To-Treat Exposed (ITT-E) Population was used which comprised of all randomized participants who received at least one dose of study treatment. Percentage values are rounded to the nearest whole digit.

End point type	Primary
----------------	---------

End point timeframe:

Week 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[1]</sup>	358 <sup>[2]</sup>		
Units: Percentage of participants				
number (confidence interval 95%)	90 (86.8 to 93.0)	93 (90.0 to 95.4)		

Notes:

[1] - ITT-E Population

[2] - ITT-E Population

### **Statistical analyses**

Statistical analysis title	Statistical Analysis
----------------------------	----------------------

Statistical analysis description:

Difference in proportion was based on CMH stratified analysis adjusting for Baseline stratification factors: Plasma HIV-1 RNA (<=versus [vs.]>100,000 c/mL) and cluster of differentiation 4+ (CD4+) cell count (<= vs. >200 cells per cubic millimeter).

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
-------------------	---

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
Parameter estimate	Adjusted difference in proportion
Point estimate	-2.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-6.7
upper limit	1.5

## Secondary: Percentage of participants with plasma HIV-1 RNA <50 c/mL at Week 24

End point title	Percentage of participants with plasma HIV-1 RNA <50 c/mL at Week 24
End point description: Percentage of participants with HIV-1 RNA<50 c/mL was obtained using FDA Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant ART prior to the visit of interest. This endpoint was analyzed using a stratified analysis with Cochran-Mantel-Haenszel weights. Percentage values are rounded to the nearest whole digit.	
End point type	Secondary
End point timeframe: Week 24	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[3]</sup>	358 <sup>[4]</sup>		
Units: Percentage of participants				
number (confidence interval 95%)	92 (89.7 to 95.2)	93 (90.4 to 95.7)		

Notes:

[3] - ITT-E Population

[4] - ITT-E Population

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description: Difference in proportion was based on CMH stratified analysis adjusting for Baseline stratification factors: Plasma HIV-1 RNA (<=vs.>100,000 c/mL) and CD4+ cell count (<= vs. >200 cells per cubic millimeter	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Adjusted difference in proportion
Point estimate	-0.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.2
upper limit	3.4

## Secondary: Percentage of Participants With Plasma HIV-1 RNA <50 c/mL at Week 96

End point title	Percentage of Participants With Plasma HIV-1 RNA <50 c/mL at Week 96
End point description:	
Percentage of participants with HIV-1 RNA<50 c/mL was obtained using FDA Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant ART prior to the visit of interest. This endpoint was analyzed using a stratified analysis with Cochran-Mantel-Haenszel weights. Percentage values are rounded to the nearest whole digit.	
End point type	Secondary
End point timeframe:	
Week 96	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[5]</sup>	358 <sup>[6]</sup>		
Units: Percentage of participants				
number (confidence interval 95%)	84 (80.5 to 88.1)	89 (86.2 to 92.6)		

Notes:

[5] - ITT-E Population

[6] - ITT-E Population

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	
Week 96. Difference in proportion was based on CMH stratified analysis adjusting for Baseline stratification factors: Plasma HIV-1 RNA (<= vs. >100,000 c/mL) and CD4+ cell count (<= vs. >200 cells/mm <sup>3</sup> ).	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Adjusted difference in proportion
Point estimate	-4.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9.8
upper limit	0

## Secondary: Percentage of Participants With Plasma HIV-1 RNA <50 c/mL at Week 144

End point title	Percentage of Participants With Plasma HIV-1 RNA <50 c/mL at Week 144
End point description:	Percentage of participants with HIV-1 RNA<50 c/mL was obtained using FDA Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant ART prior to the visit of interest. This endpoint was analyzed using a stratified analysis with CMH weights. Percentage values are rounded to the nearest whole digit.
End point type	Secondary
End point timeframe:	Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[7]</sup>	358 <sup>[8]</sup>		
Units: Percentage of participants				
number (confidence interval 95%)	79 (74.7 to 83.2)	83 (78.8 to 86.6)		

Notes:

[7] - ITT-E Population

[8] - ITT-E Population

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	Week 144. Difference in proportion was based on CMH stratified analysis adjusting for Baseline stratification factors: Plasma HIV-1 RNA (<= vs. >100,000 c/mL) and CD4+ cell count (<= vs. >200 cells/mm <sup>3</sup> ).
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Adjusted difference in proportion
Point estimate	-3.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9.4
upper limit	2.1

### Secondary: Time to Viral Suppression (HIV-1 RNA <50 c/mL) up to Week 144

End point title	Time to Viral Suppression (HIV-1 RNA <50 c/mL) up to Week 144
End point description:	Time of viral suppression is defined as the first viral load value <50 c/mL. Nonparametric Kaplan-Meier method was performed. Participants who withdrew for any reason without being suppressed were censored at date of withdrawal. Participants who have not been withdrawn and have not had viral suppression at time of the analysis were censored at last viral load date. Confidence Interval (CI) was estimated using the Brookmeyer-Crowley method.
End point type	Secondary
End point timeframe:	Up to Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[9]</sup>	358 <sup>[10]</sup>		
Units: Days				
median (inter-quartile range (Q1-Q3))	29.0 (29.0 to 53.0)	29.0 (29.0 to 56.0)		

Notes:

[9] - ITT-E Population

[10] - ITT-E Population

### Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	Hazard ratios were estimated using the Cox proportional hazard regression model.
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.658 <sup>[11]</sup>
Method	Generalized Wilcoxon procedure
Parameter estimate	Hazard ratio (HR)
Point estimate	1
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.86
upper limit	1.16

Notes:

[11] - The generalized Wilcoxon procedure was used to estimate a p-value for detecting a difference in cumulative incidence curves between treatment groups.

### Secondary: CD4+ cell counts at Weeks 24 and 48

End point title	CD4+ cell counts at Weeks 24 and 48
End point description:	
CD4+ cells are type of white blood cells that fight infection and as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+. It was evaluated by flow cytometry. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).	
End point type	Secondary
End point timeframe:	
Weeks 24 and 48	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[12]</sup>	358 <sup>[13]</sup>		
Units: Cells/mm <sup>3</sup>				
arithmetic mean (standard deviation)				
Week 24, n=340,341	655.3 (± 288.32)	632.8 (± 262.61)		
Week 48, n=324,334	687.7 (± 275.47)	675.3 (± 274.46)		

Notes:

[12] - ITT-E Population.

[13] - ITT-E Population.

### Statistical analyses

No statistical analyses for this end point

### Secondary: CD4+ Cell Counts at Week 96

End point title	CD4+ Cell Counts at Week 96
End point description:	
CD4+ cells are type of white blood cells that fight infection and as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+ cells.	



Analysis was performed by flow cytometry. Only those participants available at the specified time points were analyzed.

End point type	Secondary
End point timeframe:	
Week 96	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	301 <sup>[14]</sup>	320 <sup>[15]</sup>		
Units: Cells per cubic millimeter (cells/mm <sup>3</sup> )				
arithmetic mean (standard deviation)	732.8 (± 298.05)	711.5 (± 284.15)		

Notes:

[14] - ITT-E Population.

[15] - ITT-E Population.

### Statistical analyses

No statistical analyses for this end point

### Secondary: CD4+ Cell Counts at Week 144

End point title	CD4+ Cell Counts at Week 144
End point description:	
CD4+ cells are type of white blood cells that fight infection and as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+ cells. Analysis was performed by flow cytometry. Only those participants available at the specified time points were analyzed.	
End point type	Secondary
End point timeframe:	
Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	270 <sup>[16]</sup>	287 <sup>[17]</sup>		
Units: Cells per cubic millimeter (cells/mm <sup>3</sup> )				
arithmetic mean (standard deviation)	767.8 (± 274.00)	758.2 (± 285.35)		

Notes:

[16] - ITT-E Population.

[17] - ITT-E Population.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Changes from Baseline in CD4+ cell counts at Week 24 and 48

End point title	Changes from Baseline in CD4+ cell counts at Week 24 and 48
-----------------	---

End point description:

CD4+ cells are type of white blood cells that fight infection and as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+. It was evaluated by flow cytometry. Baseline value is defined as the the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Adjusted mean and standard error has been presented. Adjusted mean is the estimated mean change from Baseline at each visit in each arm calculated from a repeated measures model adjusting for the following covariates/factors: treatment, visit, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, treatment and visit interaction, and Baseline CD4+ cell count and visit interaction, with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and Weeks 24, 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[18]</sup>	358 <sup>[19]</sup>		
Units: Cells per cubic millimeter				
arithmetic mean (standard error)				
Week 24, n=340,341	192.2 (± 9.67)	175.1 (± 9.41)		
Week 48, n=324,334	222.2 (± 9.87)	217.7 (± 10.64)		

Notes:

[18] - ITT-E Population.

[19] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 2
----------------------------	------------------------

Statistical analysis description:

Week 48. Following covariates/factors were adjusted: treatment, visit, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, treatment and visit interaction and Baseline CD4+ cell count and visit interaction with visit as the repeated factor

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.754
Method	Mixed Model Repeated Measures (MMRM)
Parameter estimate	Mean difference (net)
Point estimate	4.6

Confidence interval	
level	95 %
sides	2-sided
lower limit	-23.9
upper limit	33

<b>Statistical analysis title</b>	Statistical Analysis 1
-----------------------------------	------------------------

Statistical analysis description:

Week 24. Following covariates/factors were adjusted: treatment, visit, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, treatment and visit interaction and Baseline CD4+ cell count and visit interaction with visit as the repeated factor.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.206
Method	Mixed Model Repeated Measures (MMRM)
Parameter estimate	Mean difference (net)
Point estimate	17.1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9.4
upper limit	43.6

## Secondary: Changes From Baseline in CD4+ Cell Counts at Week 144

End point title	Changes From Baseline in CD4+ Cell Counts at Week 144
-----------------	---

End point description:

CD4+ cells are type of white blood cells that fight infection and as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+ cells. Analysis was performed by flow cytometry. Baseline value is defined as the the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Adjusted mean and standard error has been presented. Adjusted mean is the estimated mean change from Baseline at each visit in each arm calculated from a repeated measures model adjusting for the following covariates/factors: treatment, visit, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, treatment and visit interaction, and Baseline CD4+ cell count and visit interaction, with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	270 <sup>[20]</sup>	287 <sup>[21]</sup>		
Units: Cells per cubic millimeter				
arithmetic mean (standard error)	301.8 (± 11.51)	303.2 (± 12.13)		

Notes:

[20] - ITT-E Population.

[21] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
----------------------------	----------------------

Statistical analysis description:

Week 144. Following covariates/factors were adjusted: treatment, visit, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, treatment and visit interaction and Baseline CD4+ cell count and visit interaction with visit as the repeated factor.

Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	557
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.934
Method	Mixed Model Repeated Measures (MMRM)
Parameter estimate	Mean difference (net)
Point estimate	-1.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-34.2
upper limit	31.5

## Secondary: Changes From Baseline in CD4+ Cell Counts at Week 96

End point title	Changes From Baseline in CD4+ Cell Counts at Week 96
-----------------	--

End point description:

CD4+ cells are type of white blood cells that fight infection and as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+ cells. Analysis was performed by flow cytometry. Baseline value is defined as the the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Adjusted mean and standard error has been presented. Adjusted mean is the estimated mean change from Baseline at each visit in each arm calculated from a repeated measures model adjusting for the following covariates/factors: treatment, visit, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, treatment and visit interaction, and Baseline CD4+ cell count and visit interaction, with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and Week 96

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	301 <sup>[22]</sup>	320 <sup>[23]</sup>		
Units: Cells/mm <sup>3</sup>				
arithmetic mean (standard error)	264.7 (± 11.32)	253.8 (± 11.40)		

Notes:

[22] - ITT-E Population.

[23] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	
Week 96. Following covariates/factors were adjusted: treatment, visit, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, treatment and visit interaction and Baseline CD4+ cell count and visit interaction with visit as the repeated factor.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	621
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.5
Method	Mixed Model Repeated Measures (MMRM)
Parameter estimate	Mean difference (net)
Point estimate	10.8
Confidence interval	
level	95 %
sides	2-sided
lower limit	-20.7
upper limit	42.4

## Secondary: Number of Participants With HIV-1 Disease Progression up to Week 144

End point title	Number of Participants With HIV-1 Disease Progression up to Week 144
End point description:	
HIV-associated conditions were recorded during the study and was assessed according to the 2014 Centers for Disease Control and Prevention (CDC) Classification System for HIV Infection in Adults. Disease progressions summarize participants who had HIV infection stage 3 associated conditions or death. Indicators of clinical disease progression were defined as: CDC Category Stage 1 at enrollment to Stage 3 event; CDC Category Stage 2 at enrolment to Stage 3 event; CDC Category Stage 3 at enrollment to New Stage 3 Event; CDC Category Stage 1, 2 or 3 at enrolment to Death.	
End point type	Secondary
End point timeframe:	
Up to Week 144	

<b>End point values</b>	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[24]</sup>	358 <sup>[25]</sup>		
Units: Participants				
No HIV-1 disease progression	352	356		
From CDC Stage 1 to CDC Stage 3 Event	0	0		
From CDC Stage 2 to CDC Stage 3 Event	2	2		
From CDC Stage 3 to New CDC Stage 3 Event	1	0		
From CDC Stage 1, 2 or 3 to Death	1	0		

Notes:

[24] - ITT-E Population

[25] - ITT-E Population

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of Participants With Treatment-emergent Genotypic Resistance up to Week 144

End point title	Number of Participants With Treatment-emergent Genotypic Resistance up to Week 144
-----------------	--

End point description:

Number of participants, who met confirmed virologic withdrawal (CVW) criteria, with treatment emergent genotypic resistance to Integrase strand transfer inhibitor (INSTI) and/or Nucleoside reverse transcriptase inhibitor (NRTI) was summarized. The Viral Genotypic Population comprised of all participants in the ITT-E population who have available on-treatment genotypic resistance data. Only those participants available at the specified time points were analyzed

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Week 144

<b>End point values</b>	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	8 <sup>[26]</sup>	8 <sup>[27]</sup>		
Units: Participants				
INSTI Mutations	0	0		
Major mutations of NRTI	0	0		

Notes:

[26] - Viral Genotypic Population.

[27] - Viral Genotypic Population.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Number of Participants With Treatment-emergent Phenotypic Resistance up to Week 144

End point title	Number of Participants With Treatment-emergent Phenotypic Resistance up to Week 144
-----------------	---

End point description:

Number of participants, who meet CVW criteria, with treatment emergent phenotypic resistance to INSTI and/or NRTI were summarized. Assessment of antiviral activity of anti-retroviral therapy (ART) using phenotypic test results was interpreted through a proprietary algorithm (from Monogram Biosciences) and provides the overall susceptibility of the drug. Partially sensitive and resistant calls were considered resistant in this analysis. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Week 144

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	5 <sup>[28]</sup>	6 <sup>[29]</sup>		
Units: Participants				
INSTI, DTG, Sensitive, n=,5,4	5	4		
INSTI, DTG, Resistant, n=5,4	0	0		
INSTI, EGV, Sensitive, n=5,4	5	4		
INSTI, EGV, Resistant, n=5,4	0	0		
INSTI, RAL, Sensitive, n=5,4	5	4		
INSTI, RAL, Resistant, n=5,4	0	0		
NRTI, 3TC, Sensitive, n=5,5	5	5		
NRTI, 3TC, Resistant, n=5,5	0	0		
NRTI, ABC, Sensitive, n=5,5	5	5		
NRTI, ABC, Resistant, n=5,5	0	0		
NRTI, AZT, Sensitive, n=5,5	5	5		
NRTI, AZT, Resistant, n=5,5	0	0		
NRTI, D4T, Sensitive, n=5,5	5	5		
NRTI, D4T, Resistant, n=5,5	0	0		
NRTI, DDI, Sensitive, n=5,5	5	5		
NRTI, DDI, Resistant, n=5,5	0	0		
NRTI, FTC, Sensitive, n=5,5	5	5		
NRTI, FTC, Resistant, n=5,5	0	0		

NRTI, TDF, Sensitive, n=5,5	5	5		
NRTI, TDF, Resistant, n=5,5	0	0		

Notes:

[28] - Viral Phenotypic Population.

[29] - Viral Phenotypic Population.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Number of Participants With Any AE and SAE up to Week 148

End point title	Number of Participants With Any AE and SAE up to Week 148
-----------------	---

End point description:

An AE is any untoward medical occurrence in a clinical investigation participant, temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. Any untoward event resulting in death, life threatening, requires hospitalization or prolongation of existing hospitalization, results in disability/incapacity, congenital anomaly/birth defect, any other situation according to medical or scientific judgment that may not be immediately life-threatening or result in death or hospitalization but may jeopardize the participant or may require medical or surgical intervention or protocol defined event associated with liver injury and impaired liver function were categorized as SAE. Safety Population was used which comprised of all participants who received at least one dose of study treatment.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Week 148

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[30]</sup>	358 <sup>[31]</sup>		
Units: Participants				
Any AE	307	316		
Any SAE	37	38		

Notes:

[30] - Safety Population

[31] - Safety Population

## Statistical analyses

No statistical analyses for this end point

## Secondary: Number of Participants With AEs by Maximum Severity Grades up to Week 144

End point title	Number of Participants With AEs by Maximum Severity Grades up to Week 144
-----------------	---

End point description:

An AE is any untoward medical occurrence in a clinical investigation participant, temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. AEs were evaluated by the investigator and graded according to the DAIDS toxicity scales from Grade 1 to 5 (1=Mild, 2=Moderate, 3=Severe, 4=Potentially life threatening, 5=Death). The higher the grade, the



more severe the symptoms. Number of participants with adverse events by maximum grade have been presented.

End point type	Secondary
End point timeframe:	
Up to Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[32]</sup>	358 <sup>[33]</sup>		
Units: Participants				
Grade 1 AEs	33	35		
Grade 2 AEs	229	242		
Grade 3 AEs	37	34		
Grade 4 AEs	7	5		
Grade 5 AEs	1	0		

Notes:

[32] - Safety Population

[33] - Safety Population

## Statistical analyses

No statistical analyses for this end point

## Secondary: Number of Participants With Any Drug Related AEs and Drug Related AEs by Maximum Grade up to Week 144

End point title	Number of Participants With Any Drug Related AEs and Drug Related AEs by Maximum Grade up to Week 144
-----------------	---

End point description:

An AE is any untoward medical occurrence in a clinical investigation participant, temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. AEs were evaluated by the investigator and graded according to the DAIDS toxicity scales from Grade 1 to 5. (1=Mild, 2=Moderate, 3=Severe, 4=Potentially life threatening, 5=Death). The higher the grade, the more severe the symptoms. Number of participants with drug related AEs and drug related AEs by maximum grade have been presented.

End point type	Secondary
End point timeframe:	
Up to Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[34]</sup>	358 <sup>[35]</sup>		
Units: Participants				
Any drug related AE	77	101		

Drug related AEs with maximum toxicity Grade 1	51	70		
Drug related AEs with maximum toxicity Grade 2	19	28		
Drug related AEs with maximum toxicity Grade 3	7	3		
Drug related AEs with maximum toxicity Grade 4	0	0		
Drug related AEs with maximum toxicity Grade 5	0	0		

Notes:

[34] - Safety Population

[35] - Safety Population

## Statistical analyses

No statistical analyses for this end point

## Secondary: Number of Participants With Maximum Post-Baseline Emergent Hematology Toxicities up to Week 144

End point title	Number of Participants With Maximum Post-Baseline Emergent Hematology Toxicities up to Week 144
-----------------	---

End point description:

Blood samples were collected up to Week 144 for assessment of hemoglobin, leukocytes, neutrophils and platelets. Any abnormality was graded according to DAIDS toxicity scales from Grade 1 to 4 (1=Mild, 2=Moderate, 3=Severe, 4=Potentially life threatening). The higher the grade, the more severe the symptoms. Only those participants with maximum post-Baseline emergent hematology toxicities in any of the listed hematology parameters have been presented.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[36]</sup>	358 <sup>[37]</sup>		
Units: Participants				
Hemoglobin, Grades 1 to 4	9	7		
Hemoglobin, Grades 2 to 4	1	1		
Hemoglobin, Grades 3 to 4	0	0		
Hemoglobin, Grade 1	8	6		
Hemoglobin, Grade 2	1	1		
Hemoglobin, Grade 3	0	0		
Hemoglobin, Grade 4	0	0		
Leukocytes, Grades 1 to 4	16	3		
Leukocytes, Grades 2 to 4	4	2		
Leukocytes, Grades 3 to 4	0	0		
Leukocytes, Grade 1	12	1		
Leukocytes, Grade 2	4	2		
Leukocytes, Grade 3	0	0		

Leukocytes, Grade 4	0	0		
Neutrophils, Grades 1 to 4	25	18		
Neutrophils, Grades 2 to 4	19	11		
Neutrophils, Grades 3 to 4	7	6		
Neutrophils, Grade 1	6	7		
Neutrophils, Grade 2	12	5		
Neutrophils, Grade 3	5	5		
Neutrophils, Grade 4	2	1		
Platelets, Grades 1 to 4	14	11		
Platelets, Grades 2 to 4	8	4		
Platelets, Grades 3 to 4	1	1		
Platelets, Grade 1	6	7		
Platelets, Grade 2	7	3		
Platelets, Grade 3	0	1		
Platelets, Grade 4	1	0		

Notes:

[36] - Safety Population

[37] - Safety Population

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of Participants With Maximum Post-Baseline Emergent Chemistry Toxicities up to Week 144

End point title	Number of Participants With Maximum Post-Baseline Emergent Chemistry Toxicities up to Week 144
-----------------	--

End point description:

Blood samples were collected up to Week 144 for assessment of Alanine Aminotransferase (ALT), Albumin, Alkaline Phosphatase (ALP), Aspartate aminotransferase (AST), Bilirubin, Carbon dioxide (CO<sub>2</sub>), Cholesterol, Creatine kinase (CK), Creatinine, Direct Bilirubin, Glomerular filtration rate (GFR) from creatinine adjusted for body surface area (BSA), Hypercalcemia, Hyperglycemia, Hyperkalemia, Hyponatremia, Hypocalcemia, Hypoglycemia, Hypokalemia, Hyponatremia, Low density lipid (LDL) Cholesterol, Lactate Dehydrogenase, Lipase, Phosphate, and Triglycerides. Any abnormality was graded according to DAIDS toxicity scales from Grade 1 to 4 (1=Mild, 2=Moderate, 3=Severe, 4=Potentially life threatening). The higher the grade, the more severe the symptoms. Only those participants with maximum post-Baseline emergent chemistry toxicities in any of the chemistry parameters have been presented.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[38]</sup>	358 <sup>[39]</sup>		
Units: Participants				
ALT, Grades 1 to 4	55	81		
ALT, Grades 2 to 4	23	25		
ALT, Grades 3 to 4	14	9		

ALT, Grade 1	32	56		
ALT, Grade 2	9	16		
ALT, Grade 3	7	4		
ALT, Grade 4	7	5		
Albumin, Grades 1 to 4	1	1		
Albumin, Grades 2 to 4	1	0		
Albumin, Grades 3 to 4	0	0		
Albumin, Grade 1	0	1		
Albumin, Grade 2	1	0		
Albumin, Grade 3	0	0		
Albumin, Grade 4	0	0		
ALP, Grades 1 to 4	8	10		
ALP, Grades 2 to 4	3	1		
ALP, Grades 3 to 4	0	0		
ALP, Grade 1	5	9		
ALP, Grade 2	3	1		
ALP, Grade 3	0	0		
ALP, Grade 4	0	0		
AST, Grades 1 to 4	52	79		
AST, Grades 2 to 4	27	31		
AST, Grades 3 to 4	8	13		
AST, Grade 1	25	48		
AST, Grade 2	19	18		
AST, Grade 3	6	9		
AST, Grade 4	2	4		
Bilirubin, Grades 1 to 4	42	51		
Bilirubin, Grades 2 to 4	14	17		
Bilirubin, Grades 3 to 4	4	4		
Bilirubin, Grade 1	28	34		
Bilirubin, Grade 2	10	13		
Bilirubin, Grade 3	2	4		
Bilirubin, Grade 4	2	0		
CO2, Grades 1 to 4	126	116		
CO2, Grades 2 to 4	8	9		
CO2, Grades 3 to 4	0	0		
CO2, Grade 1	118	107		
CO2, Grade 2	8	9		
CO2, Grade 3	0	0		
CO2, Grade 4	0	0		
Cholesterol, Grades 1 to 4	77	40		
Cholesterol, Grades 2 to 4	24	13		
Cholesterol, Grades 3 to 4	0	1		
Cholesterol, Grade 1	53	27		
Cholesterol, Grade 2	24	12		
Cholesterol, Grade 3	0	1		
Cholesterol, Grade 4	0	0		
CK, Grades 1 to 4	76	75		
CK, Grades 2 to 4	43	52		
CK, Grades 3 to 4	26	36		
CK, Grade 1	33	23		
CK, Grade 2	17	16		
CK, Grade 3	15	21		

CK, Grade 4	11	15		
Creatinine, Grades 1 to 4	21	31		
Creatinine, Grades 2 to 4	1	3		
Creatinine, Grades 3 to 4	0	2		
Creatinine, Grade 1	20	28		
Creatinine, Grade 2	1	1		
Creatinine, Grade 3	0	2		
Creatinine, Grade 4	0	0		
Direct Bilirubin, Grades 1 to 4	14	13		
Direct Bilirubin, Grades 2 to 4	14	13		
Direct Bilirubin, Grades 3 to 4	14	13		
Direct Bilirubin, Grade 1	0	0		
Direct Bilirubin, Grade 2	0	0		
Direct Bilirubin, Grade 3	14	13		
Direct Bilirubin, Grade 4	0	0		
GFR frm creatinine adjusted for BSA Grades 1 to 4	185	226		
GFR from creatinine adjusted for BSA Grades 2 to 4	185	226		
GFR from creatinine adjusted for BSA Grades 3 to 4	13	27		
GFR from creatinine adjusted for BSA, Grade 1	0	0		
GFR from creatinine adjusted for BSA, Grade 2	172	199		
GFR from creatinine adjusted for BSA Grades 3	13	25		
GFR from creatinine adjusted for BSA, Grade 4	0	2		
Hypercalcaemia, Grades 1 to 4	7	4		
Hypercalcaemia, Grades 2 to 4	0	0		
Hypercalcaemia, Grades 3 to 4	0	0		
Hypercalcaemia, Grade 1	7	4		
Hypercalcaemia, Grade 2	0	0		
Hypercalcaemia, Grade 3	0	0		
Hypercalcaemia, Grade 4	0	0		
Hyperglycemia, Grades 1 to 4	91	81		
Hyperglycemia, Grades 2 to 4	38	25		
Hyperglycemia, Grades 3 to 4	3	2		
Hyperglycemia, Grade 1	53	56		
Hyperglycemia, Grade 2	35	23		
Hyperglycemia, Grade 3	3	2		
Hyperglycemia, Grade 4	0	0		
Hyperkalemia, Grades 1 to 4	1	4		
Hyperkalemia, Grades 2 to 4	0	1		
Hyperkalemia, Grades 3 to 4	0	1		
Hyperkalemia, Grade 1	1	3		
Hyperkalemia, Grade 2	0	0		
Hyperkalemia, Grade 3	0	0		
Hyperkalemia, Grade 4	0	1		
Hypernatremia, Grades 1 to 4	6	3		
Hypernatremia, Grades 2 to 4	0	0		
Hypernatremia, Grades 3 to 4	0	0		
Hypernatremia, Grade 1	6	3		

Hypernatremia, Grade 2	0	0		
Hypernatremia, Grade 3	0	0		
Hypernatremia, Grade 4	0	0		
Hypocalcaemia, Grades 1 to 4	14	13		
Hypocalcaemia, Grades 2 to 4	4	3		
Hypocalcaemia, Grades 3 to 4	0	1		
Hypocalcaemia, Grade 1	10	10		
Hypocalcaemia, Grade 2	4	2		
Hypocalcaemia, Grade 3	0	1		
Hypocalcaemia, Grade 4	0	0		
Hypoglycemia, Grades 1 to 4	22	17		
Hypoglycemia, Grades 2 to 4	8	3		
Hypoglycemia, Grades 3 to 4	2	1		
Hypoglycemia, Grade 1	14	14		
Hypoglycemia, Grade 2	6	2		
Hypoglycemia, Grade 3	1	0		
Hypoglycemia, Grade 4	1	1		
Hypokalemia, Grades 1 to 4	6	7		
Hypokalemia, Grades 2 to 4	0	1		
Hypokalemia, Grades 3 to 4	0	0		
Hypokalemia, Grade 1	6	6		
Hypokalemia, Grade 2	0	1		
Hypokalemia, Grade 3	0	0		
Hypokalemia, Grade 4	0	0		
Hyponatremia, Grades 1 to 4	25	28		
Hyponatremia, Grades 2 to 4	2	0		
Hyponatremia, Grades 3 to 4	0	0		
Hyponatremia, Grade 1	23	28		
Hyponatremia, Grade 2	2	0		
Hyponatremia, Grade 3	0	0		
Hyponatremia, Grade 4	0	0		
LDL Cholesterol, Grades 1 to 4	57	35		
LDL Cholesterol, Grades 2 to 4	17	14		
LDL Cholesterol, Grades 3 to 4	5	4		
LDL Cholesterol, Grade 1	40	21		
LDL Cholesterol, Grade 2	12	10		
LDL Cholesterol, Grade 3	5	4		
LDL Cholesterol, Grade 4	0	0		
Lactate Dehydrogenase, Grades 1 to 4	3	5		
Lactate Dehydrogenase, Grades 2 to 4	0	1		
Lactate Dehydrogenase, Grades 3 to 4	0	0		
Lactate Dehydrogenase, Grade 1	3	4		
Lactate Dehydrogenase, Grade 2	0	1		
Lactate Dehydrogenase, Grade 3	0	0		
Lactate Dehydrogenase, Grade 4	0	0		
Lipase, Grades 1 to 4	65	80		
Lipase, Grades 2 to 4	35	49		
Lipase, Grades 3 to 4	10	18		
Lipase, Grade 1	30	31		
Lipase, Grade 2	25	31		
Lipase, Grade 3	8	10		
Lipase, Grade 4	2	8		

Phosphate, Grades 1 to 4	70	68		
Phosphate, Grades 2 to 4	35	47		
Phosphate, Grades 3 to 4	2	6		
Phosphate, Grade 1	35	21		
Phosphate, Grade 2	33	41		
Phosphate, Grade 3	2	6		
Phosphate, Grade 4	0	0		
Triglycerides, Grades 1 to 4	80	62		
Triglycerides, Grades 2 to 4	12	14		
Triglycerides, Grades 3 to 4	7	3		
Triglycerides, Grade 1	68	48		
Triglycerides, Grade 2	5	11		
Triglycerides, Grade 3	6	2		
Triglycerides, Grade 4	1	1		

Notes:

[38] - Safety Population

[39] - Safety Population

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of Participants Who Discontinue Treatment Due to AEs Over Week 144

End point title	Number of Participants Who Discontinue Treatment Due to AEs Over Week 144
-----------------	---

End point description:

An AE is any untoward medical occurrence in a clinical investigation participant, temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. Number of participants who discontinued treatment due to AEs have been reported.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[40]</sup>	358 <sup>[41]</sup>		
Units: Participants	18	17		

Notes:

[40] - Safety Population

[41] - Safety Population

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of Participants Who Discontinue Treatment Due to AEs Over Weeks 24, 48, 96

End point title	Number of Participants Who Discontinue Treatment Due to AEs Over Weeks 24, 48, 96
End point description: An AE is any untoward medical occurrence in a clinical investigation participant, temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product. Number of participants who discontinued treatment due to AEs have been reported.	
End point type	Secondary
End point timeframe: Up to Week 24, Week 48 and Week 96	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[42]</sup>	358 <sup>[43]</sup>		
Units: Participants				
Up to Week 24	6	4		
Up to Week 48	7	8		
Up to Week 96	14	11		

Notes:

[42] - Safety Population

[43] - Safety Population

## Statistical analyses

No statistical analyses for this end point

## Secondary: Change from Baseline in renal biomarkers-Serum Cystatin C and Serum Retinol Binding Protein (RBP) at Weeks 24, 48

End point title	Change from Baseline in renal biomarkers-Serum Cystatin C and Serum Retinol Binding Protein (RBP) at Weeks 24, 48
End point description: Blood and/or urine were collected to perform evaluation of renal inflammation biomarkers which included Serum Cystatin C and Serum Retinol Binding Protein (RBP). Baseline value is the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), presence of diabetes mellitus (factor), presence of hypertension (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).	
End point type	Secondary
End point timeframe: Baseline (Day 1) and at Weeks 24, 48	



End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[44]</sup>	358 <sup>[45]</sup>		
Units: Milligrams per Liter (mg/L)				
arithmetic mean (standard error)				
Serum Cystatin C, Week 24, n=338, 336	-0.05 (± 0.007)	-0.03 (± 0.007)		
Serum Cystatin C, Week 48, n=324, 332	-0.07 (± 0.007)	-0.04 (± 0.006)		
Serum RBP, Week 24, n=332, 334	1.6 (± 0.41)	1.9 (± 0.51)		
Serum RBP, Week 48, n=322, 332	0.5 (± 0.47)	0.6 (± 0.46)		

Notes:

[44] - Safety Population.

[45] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description: Serum Cystatin C, Week 24	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.025
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.02
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.04
upper limit	0

Statistical analysis title	Statistical Analysis 3
Statistical analysis description: Serum RBP, Week 24	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.683
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.3

Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.6
upper limit	1

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Serum RBP, Week 48	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.93
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.4
upper limit	1.2

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: Serum Cystatin C, Week 48	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.03
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.05
upper limit	-0.01

## Secondary: Change From Baseline in Renal Biomarker-Serum Cystatin C at Week 144

End point title	Change From Baseline in Renal Biomarker-Serum Cystatin C at
-----------------	---

## End point description:

Blood samples were collected to perform evaluation of renal biomarkers which included Serum Cystatin C. Baseline value is the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Biomarkers were adjusted for treatment, visit, Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, age, sex, race, presence of diabetes mellitus, presence of hypertension, Baseline biomarker value, treatment and visit interaction, and Baseline biomarker value and visit interaction; with visit as the repeated factor. Adjusted mean and standard error is presented. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

## End point timeframe:

Baseline (Day 1) and at Week 144
----------------------------------

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	283 <sup>[46]</sup>	298 <sup>[47]</sup>		
Units: Milligrams per Liter (mg/L)				
arithmetic mean (standard error)	-0.12 (± 0.006)	-0.11 (± 0.007)		

## Notes:

[46] - Safety Population.

[47] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description: Week 144. Serum Cystatin C.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	581
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.108
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.01
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.03
upper limit	0

## Secondary: Change From Baseline in Renal Biomarker-Serum Cystatin C at Week 96

End point title	Change From Baseline in Renal Biomarker-Serum Cystatin C at Week 96
-----------------	---

**End point description:**

Blood samples were collected to perform evaluation of renal inflammation biomarkers which included Serum Cystatin C. Baseline value is the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), presence of diabetes mellitus (factor), presence of hypertension (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

**End point timeframe:**

Baseline (Day 1) and at Week 96

End point values	DTG + 3TC-Double blind phase	DTG + TDF/FTC-Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	300 <sup>[48]</sup>	319 <sup>[49]</sup>		
Units: Milligrams per Liter (mg/L)				
arithmetic mean (standard error)	-0.11 (± 0.006)	-0.09 (± 0.006)		

**Notes:**

[48] - Safety Population.

[49] - Safety Population.

**Statistical analyses**

<b>Statistical analysis title</b>	Statistical Analysis
-----------------------------------	----------------------

**Statistical analysis description:**

Week 96. Serum Cystatin C.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	619
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.009
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.02
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.04
upper limit	-0.01

**Secondary: Change From Baseline in Renal Biomarker-Serum RBP at Week 96**

End point title	Change From Baseline in Renal Biomarker-Serum RBP at Week 96
-----------------	--

**End point description:**

Blood and/or urine samples were collected to perform evaluation of renal biomarkers which included Serum RBP. Baseline value is the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Biomarkers were adjusted for treatment, visit, Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, age, sex, race, presence of diabetes mellitus, presence of hypertension, Baseline biomarker value, treatment and visit interaction, and Baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

**End point timeframe:**

Baseline (Day 1) and at Week 96

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	288 <sup>[50]</sup>	310 <sup>[51]</sup>		
Units: Microgram per millimoles (ug/mmol)				
arithmetic mean (standard deviation)	1.535 (± 8.5872)	7.704 (± 41.9650)		

**Notes:**

[50] - Safety Population.

[51] - Safety Population.

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Change From Baseline in Renal Biomarker-Serum RBP at Week 144**

End point title	Change From Baseline in Renal Biomarker-Serum RBP at Week 144
-----------------	---

**End point description:**

Blood and/or urine samples were collected to perform evaluation of renal biomarkers which included Serum RBP. Baseline value is the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Biomarkers were adjusted for treatment, visit, Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, age, sex, race, presence of diabetes mellitus, presence of hypertension, Baseline biomarker value, treatment and visit interaction, and Baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

**End point timeframe:**

Baseline (Day 1) and at Week 144

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	276 <sup>[52]</sup>	292 <sup>[53]</sup>		

Units: Microgram per millimoles (ug/mmol)				
arithmetic mean (standard deviation)	1.760 (± 5.7909)	8.855 (± 35.5147)		

Notes:

[52] - Safety Population.

[53] - Safety Population.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in renal biomarkers-Serum GFR from cystatin C Adjusted using Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) and Serum or Plasma GFR from creatinine adjusted using CKD-EPI at Weeks 24, 48

End point title	Change from Baseline in renal biomarkers-Serum GFR from cystatin C Adjusted using Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) and Serum or Plasma GFR from creatinine adjusted using CKD-EPI at Weeks 24, 48
-----------------	--

End point description:

Blood &/or urine collected for evaluation of renal inflammation biomarkers: Serum GFR from cystatin C adjusted using CKD-EPI (GFR-cystatin C adjusted) and Serum or Plasma GFR from creatinine adjusted using CKD-EPI. Baseline value is the latest pre-dose assessment (Day 1). Change from Baseline was post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), presence of diabetes mellitus (factor), presence of hypertension (factor), baseline biomarker value, treatment and visit interaction, baseline biomarker value & visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Weeks 24, 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[54]</sup>	358 <sup>[55]</sup>		
Units: Milliliter/minute/1.73* $\text{meter}^2$				
arithmetic mean (standard error)				
GFR-cystatin C adjusted, Week 24, n=338, 336	4.4 (± 0.63)	2.2 (± 0.60)		
GFR-cystatin C adjusted, Week 48, n=324, 332	7.0 (± 0.60)	4.1 (± 0.59)		
GFR-creatinine adjusted, Week 24, n=340, 341	-13.5 (± 0.59)	-16.7 (± 0.56)		
GFR-creatinine adjusted, Week 48, n=326, 335	-12.1 (± 0.56)	-15.6 (± 0.55)		

Notes:

[54] - Safety Population.

[55] - Safety Population.

## Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1
Statistical analysis description: GFR-cystatin C adjusted, Week 24	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.011
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	2.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.5
upper limit	4

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: GFR- creatinine adjusted, Week 48	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	3.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	2
upper limit	5.1

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: GFR-creatinine adjusted, Week 24	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	3.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.6
upper limit	4.8

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: GFR-cystatin C adjusted, Week 48	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	2.8
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.2
upper limit	4.5

**Secondary: Change From Baseline in Renal Biomarkers-Serum GFR From Cystatin C Adjusted Using CKD-EPI and Serum or Plasma GFR From Creatinine Adjusted for BSA Using CKD-EPI method at Week 96**

End point title	Change From Baseline in Renal Biomarkers-Serum GFR From Cystatin C Adjusted Using CKD-EPI and Serum or Plasma GFR From Creatinine Adjusted for BSA Using CKD-EPI method at Week 96
-----------------	--

End point description:

Blood samples were collected to perform evaluation of renal biomarkers which included Serum GFR from cystatin C adjusted using CKD-EPI and Serum or Plasma GFR from creatinine adjusted for BSA using CKD-EPI. Baseline value is the latest pre-dose Assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Biomarkers were adjusted for treatment, visit, Baseline plasma HIV-1 RNA, baseline CD4+ cell count, age, sex, race, presence of diabetes mellitus, presence of hypertension, Baseline biomarker value, treatment and visit interaction, and Baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
End point timeframe: Baseline (Day 1) and at Week 96	



End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	300 <sup>[56]</sup>	319 <sup>[57]</sup>		
Units: Milliliter/minute/1.73* $\text{meter}^2$				
arithmetic mean (standard deviation)				
GFR Cystatin C adjusted, Week 96	11.3 ( $\pm$ 14.54)	9.3 ( $\pm$ 13.78)		
GFR creatinine adjusted, Week 96	-15.3 ( $\pm$ 11.50)	-19.0 ( $\pm$ 11.25)		

Notes:

[56] - Safety Population.

[57] - Safety Population.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in renal biomarker-Serum or Plasma Creatinine at Weeks 24, 48

End point title	Change from Baseline in renal biomarker-Serum or Plasma Creatinine at Weeks 24, 48
-----------------	--

End point description:

Blood &/or urine were for evaluation of renal inflammation biomarker which included Serum or Plasma Creatinine. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean & standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), presence of diabetes mellitus (factor), presence of hypertension (factor), baseline biomarker value, treatment and visit interaction, baseline biomarker value & visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=x in the category

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Weeks 24, 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[58]</sup>	358 <sup>[59]</sup>		
Units: Micromoles per Liter ( $\mu\text{mol/L}$ )				
arithmetic mean (standard error)				
Serum or Plasma Creatinine, Week 24, n=340, 343	11.88 ( $\pm$ 0.510)	15.07 ( $\pm$ 0.520)		
Serum or Plasma Creatinine, Week 48, n=326, 335	10.39 ( $\pm$ 0.466)	13.61 ( $\pm$ 0.480)		

Notes:

[58] - Safety Population.

[59] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description: Week 24	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-3.19
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.62
upper limit	-1.75

Statistical analysis title	Statistical Analysis 2
Statistical analysis description: Week 48	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-3.22
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.54
upper limit	-1.91

**Secondary: Change From Baseline in Renal Biomarkers-Serum GFR From Cystatin C Adjusted Using CKD-EPI and Serum or Plasma GFR From Creatinine Adjusted for BSA Using CKD-EPI method at Week 144**

End point title	Change From Baseline in Renal Biomarkers-Serum GFR From Cystatin C Adjusted Using CKD-EPI and Serum or Plasma GFR From Creatinine Adjusted for BSA Using CKD-EPI method at Week 144
End point description: Blood samples were collected to perform evaluation of renal biomarkers which included Serum GFR from cystatin C adjusted using CKD-EPI and Serum or Plasma GFR from creatinine adjusted for BSA using CKD-EPI. Baseline value is the latest pre-dose Assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Biomarkers were adjusted for treatment, visit, Baseline plasma HIV-1 RNA, baseline CD4+ cell count, age, sex, race, presence of diabetes mellitus, presence of hypertension, Baseline biomarker value, treatment and visit interaction, and Baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).	
End point type	Secondary
End point timeframe: Baseline (Day 1) and at Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[60]</sup>	358 <sup>[61]</sup>		
Units: Milliliter/minute/1.73*meter^2				
arithmetic mean (standard deviation)				
GFR Cystatin C adjusted, Week 144, n=283,298	13.0 (± 13.64)	12.1 (± 15.24)		
GFR creatinine adjusted, Week 144, n=271, 289	-16.7 (± 11.65)	-19.3 (± 11.07)		

Notes:

[60] - Safety Population.

[61] - Safety Population.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Change From Baseline in Renal Biomarker-Serum or Plasma Creatinine at Week 96

End point title	Change From Baseline in Renal Biomarker-Serum or Plasma Creatinine at Week 96
End point description: Blood and/or urine were collected to perform evaluation of renal inflammation biomarker which included Serum or Plasma Creatinine. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), presence of diabetes mellitus (factor), presence of hypertension (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.	
End point type	Secondary
End point timeframe: Baseline (Day 1) and at Week 96	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	300 <sup>[62]</sup>	319 <sup>[63]</sup>		
Units: Micromoles per Liter (umol/L)				
arithmetic mean (standard error)	12.75 (± 0.623)	16.10 (± 0.539)		

Notes:

[62] - Safety Population.

[63] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description: Week 96. Serum or Plasma creatinine	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	619
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-3.34
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.96
upper limit	-1.72

## Secondary: Change From Baseline in Renal Biomarker-Serum or Plasma Creatinine at Week 144

End point title	Change From Baseline in Renal Biomarker-Serum or Plasma Creatinine at Week 144
End point description: Blood and/or urine were collected to perform evaluation of renal inflammation biomarker which included Serum or Plasma Creatinine. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), presence of diabetes mellitus (factor), presence of hypertension (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.	
End point type	Secondary
End point timeframe: Baseline (Day 1) and at Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	271 <sup>[64]</sup>	289 <sup>[65]</sup>		
Units: Micromoles per Liter (umol/L)				
arithmetic mean (standard error)	12.89 (± 0.583)	15.87 (± 0.560)		

Notes:

[64] - Safety Population.

[65] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description: Week 144. Serum or Plasma creatinine	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	560
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-2.98
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4.57
upper limit	-1.4

## Secondary: Ratio to Baseline in renal biomarkers-Urine and Serum Beta-2 Microglobulin (B2M), Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine, Urine RBP 4 and Urine RBP 4/Urine Creatinine at Weeks 24, 48

End point title	Ratio to Baseline in renal biomarkers-Urine and Serum Beta-2 Microglobulin (B2M), Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine, Urine RBP 4 and Urine RBP 4/Urine Creatinine at Weeks 24, 48
-----------------	---

End point description:

Blood and/or urine were collected to perform evaluation of renal inflammation biomarkers which included Urine and Serum B2M, Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine, Urine RBP 4 and Urine RBP 4/Urine Creatinine. Baseline value is defined as the latest pre-dose assessment (Day 1). Ratio to Baseline was calculated as ratio of post-dose visit value over Baseline value. Statistical analysis of changes from baseline were performed on log-transformed data. Results were transformed back via exponential transformation such that treatment comparisons are assessed via odds ratios. Estimated ratio of geometric means (each visit over Baseline) and 95%

confidence interval (CI) have been presented. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles)

End point type	Secondary
End point timeframe:	
Baseline (Day 1) and at Weeks 24, 48	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[66]</sup>	358 <sup>[67]</sup>		
Units: Ratio				
geometric mean (confidence interval 95%)				
Serum B2M, Week 24, n=338, 335	0.798 (0.779 to 0.817)	0.872 (0.856 to 0.890)		
Serum B2M, Week 48, n=324, 332	0.806 (0.790 to 0.823)	0.892 (0.876 to 0.908)		
Urine B2M, Week 24, n=121, 95	0.887 (0.756 to 1.039)	1.351 (1.060 to 1.722)		
Urine B2M, Week 48, n=119, 103	0.900 (0.792 to 1.022)	1.338 (1.148 to 1.560)		
Urine Albumin/Creatinine, Week 24, n=254, 252	1.014 (0.927 to 1.109)	1.050 (0.964 to 1.144)		
Urine Albumin/Creatinine , Week 48, n=237, 244	0.934 (0.857 to 1.017)	1.048 (0.968 to 1.134)		
Urine B2M/Urine Creatinine, Week 24, n=121, 95	0.852 (0.737 to 0.985)	1.331 (1.071 to 1.655)		
Urine B2M/Urine Creatinine, Week 48, n=114, 100	0.888 (0.777 to 1.015)	1.278 (1.119 to 1.458)		
Urine Phosphate, Week 24, n=330, 332	1.115 (1.025 to 1.212)	1.012 (0.934 to 1.095)		
Urine Phosphate, Week 48, n=316, 330	1.061 (0.983 to 1.145)	1.075 (0.996 to 1.159)		
Urine Protein/Creatinine , Week 24, n=269, 265	0.850 (0.806 to 0.895)	1.016 (0.960 to 1.075)		
Urine Protein/Creatinine , Week 48, n=252, 269	0.879 (0.838 to 0.922)	1.061 (1.009 to 1.115)		
Urine RBP 4, Week 24, n=332, 330	0.934 (0.842 to 1.036)	1.073 (0.951 to 1.209)		
Urine RBP 4, Week 48, n=318, 328	1.115 (1.009 to 1.233)	1.490 (1.332 to 1.667)		
Urine RBP 4/Urine Creatinine , Week 24, n=329, 330	0.919 (0.846 to 0.998)	1.110 (1.003 to 1.228)		
Urine RBP 4/Urine Creatinine , Week 48, n=304, 318	1.147 (1.060 to 1.241)	1.500 (1.367 to 1.646)		

Notes:

[66] - Safety Population.

[67] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description:	
Week 24. Serum B2M	

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.915
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.887
upper limit	0.943

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: Week 48. Serum B2M	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.904
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.88
upper limit	0.929

<b>Statistical analysis title</b>	Statistical Analysis 9
Statistical analysis description: Week 24. Urine Phosphate	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.099
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	1.102

Confidence interval	
level	95 %
sides	2-sided
lower limit	0.982
upper limit	1.237

<b>Statistical analysis title</b>	Statistical Analysis 8
Statistical analysis description: Week 48. Urine B2M/Urine Creatinine	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.695
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.576
upper limit	0.839

<b>Statistical analysis title</b>	Statistical Analysis 7
Statistical analysis description: Week 24. Urine B2M/Urine Creatinine	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.64
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.493
upper limit	0.831

<b>Statistical analysis title</b>	Statistical Analysis 6
-----------------------------------	------------------------



Statistical analysis description: Week 48. Urine Albumin/Creatinine	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.051
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.891
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.793
upper limit	1.001

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description: Week 24. Urine Albumin/Creatinine	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.575
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.965
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.853
upper limit	1.092

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Week 48. Urine B2M	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.672

Confidence interval	
level	95 %
sides	2-sided
lower limit	0.551
upper limit	0.821

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Week 24. Urine B2M	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.005
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.656
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.491
upper limit	0.877

<b>Statistical analysis title</b>	Statistical Analysis 14
Statistical analysis description: Week 48. Urine RBP 4	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.748
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.644
upper limit	0.87

<b>Statistical analysis title</b>	Statistical Analysis 13
-----------------------------------	-------------------------

Statistical analysis description:

Week 24. Urine RBP 4

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.087
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.871
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.743
upper limit	1.02

**Statistical analysis title**

Statistical Analysis 12

Statistical analysis description:

Week 48. Urine Protein/Creatinine

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.829
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.773
upper limit	0.888

**Statistical analysis title**

Statistical Analysis 11

Statistical analysis description:

Week 24. Urine Protein/Creatinine

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.836

Confidence interval	
level	95 %
sides	2-sided
lower limit	0.774
upper limit	0.904

<b>Statistical analysis title</b>	Statistical Analysis 10
Statistical analysis description: Week 48. Urine Phosphate	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.816
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.987
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.886
upper limit	1.1

<b>Statistical analysis title</b>	Statistical Analysis 15
Statistical analysis description: Week 24. Urine RBP 4/Urine Creatinine	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.005
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.828
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.727
upper limit	0.944

<b>Statistical analysis title</b>	Statistical Analysis 16
-----------------------------------	-------------------------

## Statistical analysis description:

Week 48. Urine RBP 4/Urine Creatinine

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.765
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.677
upper limit	0.864

**Secondary: Ratio to Baseline in Renal Biomarkers- Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine and Urine RBP 4/Urine Creatinine at Week 96**

End point title	Ratio to Baseline in Renal Biomarkers- Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine and Urine RBP 4/Urine Creatinine at Week 96
-----------------	--

## End point description:

Blood and/or urine were collected to perform evaluation of renal inflammation biomarkers which included Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine, Urine RBP 4 and Urine RBP 4/Urine Creatinine. Baseline value is defined as the latest pre-dose assessment (Day 1). Ratio to Baseline was calculated as ratio of post-dose visit value over Baseline value. Statistical analysis of changes from baseline were performed on log-transformed data. Results were transformed back via exponential transformation such that treatment comparisons are assessed via odds ratios. Estimated ratio of geometric means (each visit over Baseline) and 95% confidence interval (CI) have been presented. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).

End point type	Secondary
----------------	-----------

## End point timeframe:

Baseline (Day 1) and Week 96

End point values	DTG + 3TC-Double blind phase	DTG + TDF/FTC-Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[68]</sup>	358 <sup>[69]</sup>		
Units: Ratio				
geometric mean (confidence interval 95%)				
Urine Albumin/Creatinine, Week 96, n=222, 243	0.924 (0.847 to 1.008)	1.101 (1.008 to 1.202)		
Urine B2M/Urine Creatinine, Week 96, n=107, 104	0.794 (0.720 to 0.877)	1.441 (1.193 to 1.740)		

Urine Phosphate, Week 96, n=292, 316	1.113 (1.021 to 1.214)	1.066 (0.984 to 1.155)		
Urine Protein/Creatinine, Week 96, n=238, 258	0.868 (0.818 to 0.920)	1.053 (1.004 to 1.105)		
Urine RBP 4/Urine Creatinine, Week 96, n=289, 311	1.310 (1.207 to 1.420)	1.771 (1.604 to 1.955)		

Notes:

[68] - Safety Population.

[69] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description: Week 96. Urine Albumin/Creatinine.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.006
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.839
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.742
upper limit	0.95

Statistical analysis title	Statistical Analysis 2
Statistical analysis description: Week 96. Urine B2M/Urine Creatinine.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.551
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.445
upper limit	0.682

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description: Week 96. Urine RBP 4/Urine Creatinine	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.74
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.651
upper limit	0.84

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Week 96. Urine Protein/Creatinine.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.824
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.764
upper limit	0.889

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Week 96. Urine Phosphate.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.467
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	1.045
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.928
upper limit	1.175

### Secondary: Ratio to Baseline in Renal Biomarkers- Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine and Urine RBP 4/Urine Creatinine at Week 144

End point title	Ratio to Baseline in Renal Biomarkers- Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine and Urine RBP 4/Urine Creatinine at Week 144
-----------------	---

#### End point description:

Blood and/or urine were collected to perform evaluation of renal inflammation biomarkers which included Urine and Serum B2M, Urine Albumin/Creatinine, Urine B2M/Urine Creatinine, Urine Phosphate, Urine Protein/Creatinine, Urine RBP 4 and Urine RBP 4/Urine Creatinine. Baseline value is defined as the latest pre-dose assessment (Day 1). Ratio to Baseline was calculated as ratio of post-dose visit value over Baseline value. Statistical analysis of changes from baseline were performed on log-transformed data. Results were transformed back via exponential transformation such that treatment comparisons are assessed via odds ratios. Estimated ratio of geometric means (each visit over Baseline) and 95% confidence interval (CI) have been presented. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).

End point type	Secondary
----------------	-----------

#### End point timeframe:

Baseline (Day 1) and Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[70]</sup>	358 <sup>[71]</sup>		
Units: Ratio				
geometric mean (confidence interval 95%)				
Urine Albumin/Creatinine , Week 144, n=207, 212	1.050 (0.954 to 1.155)	1.146 (1.039 to 1.264)		
Urine B2M/Urine Creatinine , Week 144, n=100, 102	0.751 (0.679 to 0.830)	1.518 (1.281 to 1.799)		
Urine Phosphate, Week 144, n=274, 294	1.040 (0.954 to 1.133)	0.955 (0.879 to 1.037)		
Urine Protein/Creatinine , Week 144, n=225,232	0.988 (0.932 to 1.047)	1.210 (1.144 to 1.280)		



Urine RBP 4/Urine Creatinine, Week 144, n=276, 292	1.648 (1.550 to 1.752)	2.425 (2.208 to 2.663)		
--	------------------------	------------------------	--	--

Notes:

[70] - Safety Population.

[71] - Safety Population.

## Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1
Statistical analysis description: Week 144. Urine Albumin/Creatinine.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.205
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.916
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.799
upper limit	1.05

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: Week 144. Urine B2M/Urine Creatinine.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.495
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.406
upper limit	0.603

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Week 144. Urine Phosphate.	

Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.16
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	1.089
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.967
upper limit	1.226

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Week 144. Urine Protein/Creatinine.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.817
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.753
upper limit	0.885

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description: Week 144. Urine RBP 4/Urine Creatinine	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Ratio of geometric means
Point estimate	0.679

Confidence interval	
level	95 %
sides	2-sided
lower limit	0.607
upper limit	0.76

**Secondary: Change from Baseline in bone biomarkers-Serum Bone Specific Alkaline Phosphatase (bone-ALP), Serum Osteocalcin, Serum Procollagen 1 N-Terminal Propeptide (PINP) and Serum Type I Collagen C-Telopeptides (CTX-1) at Weeks 24, 48**

End point title	Change from Baseline in bone biomarkers-Serum Bone Specific Alkaline Phosphatase (bone-ALP), Serum Osteocalcin, Serum Procollagen 1 N-Terminal Propeptide (PINP) and Serum Type I Collagen C-Telopeptides (CTX-1) at Weeks 24, 48
-----------------	---

End point description:

Blood samples were collected to perform evaluation of bone biomarkers which included bone-ALP, Serum Osteocalcin, PINP and CTX-1. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), BMI (factor), smoking status (factor), current Vitamin D use (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Weeks 24, 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[72]</sup>	358 <sup>[73]</sup>		
Units: Micrograms per Liter (ug/L)				
arithmetic mean (standard error)				
Bone-ALP, Week 24, n=334, 332	0.91 (± 0.179)	3.13 (± 0.199)		
Bone-ALP, Week 48, n=321, 331	1.21 (± 0.193)	3.79 (± 0.239)		
Serum Osteocalcin, Week 24, n=335, 334	2.56 (± 0.341)	6.74 (± 0.347)		
Serum Osteocalcin, Week 48, n=322, 330	0.78 (± 0.311)	6.01 (± 0.400)		
PINP, Week 24, n=337, 336	4.5 (± 0.91)	18.3 (± 1.06)		
PINP, Week 48, n=321, 334	0.5 (± 0.83)	13.1 (± 0.84)		
CTX-1, Week 24, n=337, 334	0.1192 (± 0.01304)	0.2820 (± 0.01472)		
CTX-1, Week 48, n=323, 331	0.1338 (± 0.01258)	0.3352 (± 0.01885)		

Notes:

[72] - Safety Population.

**Statistical analyses**

<b>Statistical analysis title</b>	Statistical Analysis 1
Statistical analysis description: Week 24. Bone ALP	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-2.23
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.75
upper limit	-1.7

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: Week 48. Bone ALP	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-2.58
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.19
upper limit	-1.98

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Week 24. Serum Osteocalcin	

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-4.19
Confidence interval	
level	95 %
sides	2-sided
lower limit	-5.15
upper limit	-3.23

<b>Statistical analysis title</b>	Statistical Analysis 8
Statistical analysis description: Week 48. CTX-1	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.2015
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.246
upper limit	-0.1569

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description: Week 24. Serum PINP	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-13.8

Confidence interval	
level	95 %
sides	2-sided
lower limit	-16.5
upper limit	-11.1

<b>Statistical analysis title</b>	Statistical Analysis 6
Statistical analysis description: Week 48. Serum PINP	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-12.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-15
upper limit	-10.3

<b>Statistical analysis title</b>	Statistical Analysis 7
Statistical analysis description: Week 24. CTX-1	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.1628
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.2015
upper limit	-0.1241

<b>Statistical analysis title</b>	Statistical Analysis 4
-----------------------------------	------------------------

## Statistical analysis description:

Week 48. Serum Osteocalcin

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-5.23
Confidence interval	
level	95 %
sides	2-sided
lower limit	-6.22
upper limit	-4.23

**Secondary: Change From Baseline in Bone Biomarkers-Serum Bone-ALP, Serum Osteocalcin, Serum PINP and Serum Type I CTX-1 at Week 96**

End point title	Change From Baseline in Bone Biomarkers-Serum Bone-ALP, Serum Osteocalcin, Serum PINP and Serum Type I CTX-1 at Week 96
-----------------	---

## End point description:

Blood samples were collected to perform evaluation of bone biomarkers which included bone-ALP, Serum Osteocalcin, PINP and CTX-1. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), BMI (factor), smoking status (factor), current Vitamin D use (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).

End point type	Secondary
----------------	-----------

## End point timeframe:

Baseline (Day 1) and at Week 96

End point values	DTG + 3TC-Double blind phase	DTG + TDF/FTC-Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[74]</sup>	358 <sup>[75]</sup>		
Units: Micrograms per Liter (ug/L)				
arithmetic mean (standard error)				
Bone-ALP, Week 96, n=296, 317	0.30 (± 0.191)	2.37 (± 0.216)		
Serum Osteocalcin, Week 96, n=297, 320	0.40 (± 0.345)	4.57 (± 0.391)		
PINP, Week 96, n=297, 319	15.0 (± 1.63)	28.3 (± 1.50)		
CTX-1, Week 96, n=297, 315	0.1351 (± 0.01580)	0.2943 (± 0.01916)		

Notes:

[74] - Safety Population.

[75] - Safety Population.

## Statistical analyses

<b>Statistical analysis title</b>	Statistical Analysis 1
Statistical analysis description: Week 96, Bone ALP	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-2.06
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.63
upper limit	-1.5

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Week 96, CTX-1	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.1592
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.208
upper limit	-0.1104

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Week 96, Serum PINP	



Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-13.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-17.6
upper limit	-8.9

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: Week 96, Serum Osteocalcin	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-4.17
Confidence interval	
level	95 %
sides	2-sided
lower limit	-5.2
upper limit	-3.14

### **Secondary: Change From Baseline in Bone Biomarkers-Serum Bone-ALP, Serum Osteocalcin, Serum PINP and Serum Type I CTX-1 at Week 144**

End point title	Change From Baseline in Bone Biomarkers-Serum Bone-ALP, Serum Osteocalcin, Serum PINP and Serum Type I CTX-1 at Week 144
End point description: Blood samples were collected to perform evaluation of bone biomarkers which included bone-ALP, Serum Osteocalcin, PINP and CTX-1. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), BMI (factor), smoking status (factor), current Vitamin D use (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).	
End point type	Secondary

End point timeframe:

Baseline (Day 1) and at Week 144

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[76]</sup>	358 <sup>[77]</sup>		
Units: Micrograms per Liter (ug/L)				
arithmetic mean (standard error)				
Bone-ALP, Week 144, n=281, 295	-0.25 (± 0.172)	1.43 (± 0.217)		
Serum Osteocalcin, Week 144, n=281, 299	0.29 (± 0.374)	3.21 (± 0.403)		
PINP, Week 144, n=281,299	4.6 (± 1.04)	13.8 (± 1.14)		
CTX-1, Week 144, n=281, 296	0.0750 (± 0.01150)	0.2164 (± 0.01407)		

Notes:

[76] - Safety Population.

[77] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description: Week 144, Bone ALP	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-1.68
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.23
upper limit	-1.14

Statistical analysis title	Statistical Analysis 2
Statistical analysis description: Week 144, Serum Osteocalcin	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-2.91
Confidence interval	
level	95 %
sides	2-sided
lower limit	-4
upper limit	-1.83

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Week 144, Serum PINP	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-9.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-12.3
upper limit	-6.2

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Week 144, CTX-1	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-0.1414

Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.1771
upper limit	-0.1056

## Secondary: Change From Baseline in Bone Biomarker-Serum Vitamin D at Week 96

End point title	Change From Baseline in Bone Biomarker-Serum Vitamin D at Week 96
-----------------	---

End point description:

Blood samples were collected to perform evaluation of bone biomarker serum vitamin D. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), BMI (factor), smoking status (factor), current Vitamin D use (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Week 96

End point values	DTG + 3TC-Double blind phase	DTG + TDF/FTC-Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	298 <sup>[78]</sup>	320 <sup>[79]</sup>		
Units: Nanomoles per Liter (nmol/L)				
arithmetic mean (standard error)	-2.2 (± 1.05)	0.7 (± 1.04)		

Notes:

[78] - Safety Population.

[79] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
----------------------------	----------------------

Statistical analysis description:

Week 96

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	618
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.048
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-2.9

Confidence interval	
level	95 %
sides	2-sided
lower limit	-5.8
upper limit	0

## Secondary: Change from Baseline in bone biomarker-Serum Vitamin D at Weeks 24, 48

End point title	Change from Baseline in bone biomarker-Serum Vitamin D at Weeks 24, 48
-----------------	--

### End point description:

Blood samples were collected to perform evaluation of bone biomarker serum vitamin D. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), BMI (factor), smoking status (factor), current Vitamin D use (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).

End point type	Secondary
----------------	-----------

### End point timeframe:

Baseline (Day 1) and at Weeks 24, 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[80]</sup>	358 <sup>[81]</sup>		
Units: Nanomoles per Liter (nmol/L)				
arithmetic mean (standard error)				
Serum Vitamin D, Week 24, n=337, 337	5.9 (± 1.15)	12.4 (± 1.33)		
Serum Vitamin D, Week 48, n=322, 333	-3.1 (± 0.89)	3.1 (± 1.10)		

### Notes:

[80] - Safety Population.

[81] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 2
----------------------------	------------------------

### Statistical analysis description:

Week 48

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
-------------------	---

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-6.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9
upper limit	-3.4

<b>Statistical analysis title</b>	Statistical Analysis 1
Statistical analysis description: Week 24	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	< 0.001
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-6.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9.9
upper limit	-3

**Secondary: Percentage change from Baseline in fasting lipids-Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides at Weeks 24, 48**

End point title	Percentage change from Baseline in fasting lipids-Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides at Weeks 24, 48
-----------------	---

End point description:

Blood samples were collected to perform evaluation of fasting lipids which included Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides. Baseline value is defined as the latest pre-dose assessment (Day 1). Percentage change from Baseline was calculated as 100 multiplied by ([post-dose visit value minus Baseline value] divided by Baseline value). Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Weeks 24, 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[82]</sup>	358 <sup>[83]</sup>		
Units: Percentage change				
arithmetic mean (standard deviation)				
Serum or Plasma Cholesterol, Week 24, n=294, 297	9.4 (± 17.44)	-4.7 (± 16.12)		
Serum or Plasma Cholesterol, Week 48, n=280, 289	10.5 (± 18.89)	-2.4 (± 17.14)		
HDL Cholesterol, Direct, Week 24, n=294, 297	16.4 (± 22.58)	3.4 (± 21.55)		
HDL Cholesterol, Direct, Week 48, n=280, 289	15.0 (± 25.07)	5.0 (± 33.04)		
LDL Cholesterol, Week 24, n=294, 297	12.4 (± 45.05)	-8.1 (± 23.70)		
LDL Cholesterol, Week 48, n=280, 289	14.8 (± 48.74)	-4.0 (± 24.06)		
Triglycerides ,Week 24, n=294, 297	8.5 (± 46.57)	4.3 (± 72.35)		
Triglycerides , Week 48, n=280, 289	12.8 (± 68.99)	4.4 (± 70.43)		

Notes:

[82] - Safety Population.

[83] - Safety Population.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Change From Baseline in Bone Biomarker-Serum Vitamin D at Week 144

End point title	Change From Baseline in Bone Biomarker-Serum Vitamin D at Week 144
-----------------	--

End point description:

Blood samples were collected to perform evaluation of bone biomarker serum vitamin D. Baseline value is defined as the latest pre-dose assessment (Day 1). Change from Baseline was calculated as post-dose visit value minus Baseline value. Adjusted mean and standard error is presented. Adjusted mean is the estimated mean change from baseline at each visit in each arm calculated from a repeated measures model adjusting for: treatment, visit, baseline plasma HIV-1 RNA (factor), baseline CD4+ cell count (factor), age, sex (factor), race (factor), BMI (factor), smoking status (factor), current Vitamin D use (factor), baseline biomarker value, treatment and visit interaction, and baseline biomarker value and visit interaction; with visit as the repeated factor. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	281 <sup>[84]</sup>	298 <sup>[85]</sup>		
Units: Nanomoles per Liter (nmol/L)				
arithmetic mean (standard error)	-2.0 (± 1.16)	2.9 (± 1.28)		

Notes:

[84] - Safety Population.

[85] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description: Week 144	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	579
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.004
Method	Mixed Model Repeated Measures
Parameter estimate	Mean difference (net)
Point estimate	-4.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	-8.3
upper limit	-1.6

## Secondary: Change From Baseline in Fasting Lipids-Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides at Week 96

End point title	Change From Baseline in Fasting Lipids-Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides at Week 96
End point description: Blood samples were collected to perform evaluation of fasting lipids which included Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides. Baseline value was defined as the latest pre-dose assessment (Day 1). Change from Baseline was defined as value at the indicated time point minus Baseline value. Adjusted mean and standard error is presented. Only those participants available at the specified time points were analyzed.	
End point type	Secondary
End point timeframe: Baseline (Day 1) and at Week 96	



End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	253 <sup>[86]</sup>	277 <sup>[87]</sup>		
Units: Millimoles per liter				
arithmetic mean (standard error)				
Serum or Plasma Cholesterol, Week 96	0.379 (± 0.0376)	-0.104 (± 0.0378)		
HDL Cholesterol, Direct, Week 96	0.199 (± 0.0156)	0.090 (± 0.0149)		
LDL Cholesterol, Week 96,	0.147 (± 0.0338)	-0.154 (± 0.0303)		
Triglycerides, Week 96,	0.129 (± 0.0835)	-0.112 (± 0.0358)		

Notes:

[86] - Safety Population.

[87] - Safety Population.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage change from Baseline in fasting lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio at Weeks 24, 48

End point title	Percentage change from Baseline in fasting lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio at Weeks 24, 48
-----------------	--

End point description:

Blood samples were collected to perform evaluation of fasting lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio. Baseline value is the latest pre-dose assessment (Day 1). Percentage change from Baseline was calculated as 100 multiplied by ([post-dose visit value minus Baseline value] divided by Baseline value). Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Weeks 24, 48

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[88]</sup>	358 <sup>[89]</sup>		
Units: Percentage change				
arithmetic mean (standard deviation)				
Total/HDL Cholesterol Ratio, Week 24, n=294, 297	-4.0 (± 19.08)	-4.6 (± 27.52)		
Total/HDL Cholesterol Ratio, Week 48, n=280, 289	-0.2 (± 31.10)	-4.4 (± 16.96)		

Notes:

[88] - Safety Population.

[89] - Safety Population.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change From Baseline in Fasting Lipids-Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides at Week 144

End point title	Change From Baseline in Fasting Lipids-Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides at Week 144
-----------------	--

End point description:

Blood samples were collected to perform evaluation of fasting lipids which included Serum or Plasma Cholesterol, Serum or Plasma HDL Cholesterol (Direct), Serum or Plasma LDL Cholesterol (Calculated or Direct) and Serum or Plasma Triglycerides. Baseline value was defined as the latest pre-dose assessment (Day 1). Change from Baseline was defined as value at the indicated time point minus Baseline value. Adjusted mean and standard error is presented. Only those participants available at the specified time points were analyzed..

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	245 <sup>[90]</sup>	256 <sup>[91]</sup>		
Units: Millimoles per liter				
arithmetic mean (standard error)				
Serum or Plasma Cholesterol, Week 144,	0.367 (± 0.0408)	-0.037 (± 0.0406)		
HDL Cholesterol, Direct, Week 144	0.181 (± 0.0150)	0.098 (± 0.0147)		
LDL Cholesterol, Week 144,	0.170 (± 0.0336)	-0.105 (± 0.0348)		
Triglycerides, Week 144	0.117 (± 0.0872)	-0.104 (± 0.0385)		

Notes:

[90] - Safety Population.

[91] - Safety Population.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change From Baseline in Fasting Lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio at Week 144

End point title	Change From Baseline in Fasting Lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio at Week 144
-----------------	---

End point description:

Blood samples were collected to perform evaluation of fasting lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio. Baseline value was defined as the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Adjusted mean and standard error has been presented. Only those participants available at the specified time points were

analyzed.

End point type	Secondary
End point timeframe:	
Baseline (Day 1) and at Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	245 <sup>[92]</sup>	256 <sup>[93]</sup>		
Units: Ratio				
arithmetic mean (standard error)	-0.229 (± 0.0559)	-0.386 (± 0.0463)		

Notes:

[92] - Safety Population.

[93] - Safety Population.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change From Baseline in Fasting Lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio at Week 96

End point title	Change From Baseline in Fasting Lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio at Week 96
-----------------	--

End point description:

Blood samples were collected to perform evaluation of fasting lipid-Serum or Plasma Total Cholesterol/HDL Cholesterol Ratio. Baseline value was defined as the latest pre-dose assessment (Day 1). Change from Baseline was defined as post-dose visit value minus Baseline value. Adjusted mean and standard error has been presented. Only those participants available at the specified time points were analyzed.

End point type	Secondary
End point timeframe:	
Baseline (Day 1) and at Week 96	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	253 <sup>[94]</sup>	277 <sup>[95]</sup>		
Units: Ratio				
arithmetic mean (standard error)	-0.213 (± 0.0566)	-0.402 (± 0.0479)		

Notes:

[94] - Safety Population.

[95] - Safety Population.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of participants with Grade 2 or greater laboratory abnormalities in fasting LDL cholesterol by Weeks 24, 48

End point title	Percentage of participants with Grade 2 or greater laboratory abnormalities in fasting LDL cholesterol by Weeks 24, 48
-----------------	--

End point description:

Blood samples evaluation of fasting LDL cholesterol. Any abnormalities were evaluated by the investigator and graded according to DAIDS toxicity scales from Grade 1 to 4 (1=Mild,2=Moderate,3=Severe,4=Potentially life threatening). The higher the grade, the more severe the symptoms. Percentage of participants with Grade 2 or greater laboratory abnormalities in fasting LDL cholesterol by Weeks 24 and 48 have been presented. Participants without any post-Baseline fasting LDL cholesterol value prior to Week 48 or those who had Baseline lipids-lowering agents are not included. Lipid Last Observation Carried Forward (LOCF) data was used such that the last available fasted, on-treatment lipid value prior to the initiation of a lipid-lowering agent was used in place of future observed values. Percentage values are rounded to the nearest whole digit. Only those participants available at the specified time points were analyzed (represented by n=x in the category

End point type	Secondary
----------------	-----------

End point timeframe:

Weeks 24 and Week 48

End point values	DTG + 3TC-Double blind phase	DTG + TDF/FTC-Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[96]</sup>	358 <sup>[97]</sup>		
Units: Percentage of participants				
Week 24, n=309, 316	4	2		
Week 48, n=318, 320	4	3		

Notes:

[96] - Safety Population.

[97] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
----------------------------	------------------------

Statistical analysis description:

Week 24

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.157 <sup>[98]</sup>
Method	Fisher exact
Parameter estimate	Difference in percentage
Point estimate	2

Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.6
upper limit	4.6

Notes:

[98] - Fisher's exact p-value.

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description:	
Week 48	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.414 <sup>[99]</sup>
Method	Fisher exact
Parameter estimate	Difference in percentage
Point estimate	1.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.7
upper limit	4.2

Notes:

[99] - Fisher's exact p-value.

### **Secondary: Percentage of Participants With Grade 2 or Greater Laboratory Abnormalities in Fasting LDL Cholesterol by Week 96**

End point title	Percentage of Participants With Grade 2 or Greater Laboratory Abnormalities in Fasting LDL Cholesterol by Week 96
End point description:	
<p>Blood samples were collected for fasting LDL cholesterol. Any abnormalities were evaluated by investigator&amp; graded according to DAIDS toxicity scales from Grade 1 to 4 (1=Mild, 2=Moderate, 3=Severe, 4=Potentially life threatening). The higher the grade, more severe the symptoms. Percentage of participants with Grade 2 or greater laboratory abnormalities in fasting LDL cholesterol by Week 96 have been presented. Participants without any post-Baseline fasting LDL cholesterol value prior to Week 96 or those who had Baseline lipids-lowering agents were not included. Lipid Last Observation Carried Forward (LOCF) data was used such that the last available fasted on-treatment lipid value prior to the initiation of a lipid-lowering agent was used in place of future observed values. Percentage values are rounded to the nearest whole digit. Only those participants available at the specified time points were analyzed.</p>	
End point type	Secondary
End point timeframe:	
Week 96	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	320 <sup>[100]</sup>	323 <sup>[101]</sup>		
Units: Percentage of participants	5	4		

Notes:

[100] - Safety Population.

[101] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	
Week 96	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	643
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.562 <sup>[102]</sup>
Method	Fisher exact
Parameter estimate	Difference in percentage
Point estimate	1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.1
upper limit	4.1

Notes:

[102] - Fisher's exact p-value.

## Secondary: Percentage of Participants With Grade 2 or Greater Laboratory Abnormalities in Fasting LDL Cholesterol by Week 144

End point title	Percentage of Participants With Grade 2 or Greater Laboratory Abnormalities in Fasting LDL Cholesterol by Week 144
End point description:	
<p>Blood samples were collected to perform evaluation of fasting LDL cholesterol. Any abnormalities were evaluated by the investigator and graded according to DAIDS toxicity scales from Grade 1 to 4 (1=Mild, 2=Moderate, 3=Severe, 4=Potentially life threatening). Percentage of participants with Grade 2 or greater laboratory abnormalities in fasting LDL cholesterol by Week 144 have been presented. Participants without any post-Baseline fasting LDL cholesterol value prior to Week 144 or those who had Baseline lipids-lowering agents were not included. Lipid Last Observation Carried Forward (LOCF) data was used such that the last available fasted, on-treatment lipid value prior to the initiation of a lipid-lowering agent was used in place of future observed values. Percentage values are rounded to the nearest whole digit. Only those participants available at the specified time points were analyzed.</p>	
End point type	Secondary
End point timeframe:	
Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	321 <sup>[103]</sup>	322 <sup>[104]</sup>		
Units: Percentage of participants	5	4		

Notes:

[103] - Safety Population.

[104] - Safety Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	
Week 144	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	643
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.587 <sup>[105]</sup>
Method	Fisher exact
Parameter estimate	Difference in percentage
Point estimate	0.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.4
upper limit	4.3

Notes:

[105] - Fisher's exact p-value.

## Secondary: Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count, Baseline HIV-1 RNA, race) with plasma HIV-1 RNA <50 c/mL at Week 24

End point title	Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count, Baseline HIV-1 RNA, race) with plasma HIV-1 RNA <50 c/mL at Week 24
End point description:	
Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count, Baseline HIV-1 RNA, race) with HIV-1 RNA<50 c/mL was obtained using FDA Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant ART prior to the visit of interest. Data was presented by subgroups: age (<35, 35 to <50, >=50 years); gender (males and females), Baseline CD4+ cell count (<=200, >200), Baseline HIV-1 RNA (<=100000, >100000) and Race (White, African American/African H., Asian, Other). Percentage values are rounded to the nearest whole digit. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).	
End point type	Secondary
End point timeframe:	
Week 24	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[106]</sup>	358 <sup>[107]</sup>		
Units: Percentage of participants				
Baseline CD4+ cell count, <=200,n=31,29	90	86		
Baseline CD4+ cell count, >200,n=325,329	93	94		
Female, n=59, 52	93	96		
Male, n=297, 306	92	92		
Age, <35,n= 211, 205	93	95		
Age, 35 to <50,n=116, 107	91	93		
Age, >=50, n=29, 46	93	85		
Baseline plasma HIV-1 RNA, <=100000,n=282,282	93	95		
Baseline plasma HIV-1 RNA, >100000,n=74, 76	92	87		
Race, White, n=244,247	93	95		
Race, African American/African H., n=39, 36	92	81		
Race, Asian, n=37, 42	89	93		
Race, Other, n=36, 33	94	94		

Notes:

[106] - ITT-E Population.

[107] - ITT-E Population.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count Baseline HIV-1 RNA, race) with plasma HIV-1 RNA <50 c/mL at Week 48

End point title	Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count Baseline HIV-1 RNA, race) with plasma HIV-1 RNA <50 c/mL at Week 48
-----------------	---

End point description:

Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count, Baseline HIV-1 RNA, race) with HIV-1 RNA<50 c/mL was obtained using FDA Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant ART prior to the visit of interest. Data was presented by subgroups: age (<35, 35 to <50, >=50 years); gender (males and females), Baseline CD4+ cell count (<=200, >200), Baseline HIV-1 RNA (<=100000, >100000) and Race (White, African American/African H., Asian, Other). Percentage values are rounded to the nearest whole digit. ITT-E Population. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).

End point type	Secondary
End point timeframe:	
Week 48	



End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[108]</sup>	358 <sup>[109]</sup>		
Units: Percentage of participants				
Baseline CD4+ cell count, <=200,n=31,29	81	90		
Baseline CD4+ cell count, >200,n=325,329	91	93		
Female, n=59, 52	88	94		
Male, n=297, 306	90	92		
Age, <35,n= 211, 205	92	93		
Age, 35 to <50,n=116, 107	86	94		
Age, >=50, n=29, 46	90	87		
Baseline plasma HIV-1 RNA, <=100000,n=282,282	90	93		
Baseline plasma HIV-1 RNA, >100000,n=74, 76	88	91		
Race, White, n=244,247	90	94		
Race, African American/African H., n=39, 36	87	81		
Race, Asian, n=37, 42	92	98		
Race, Other, n=36, 33	89	94		

Notes:

[108] - ITT-E Population.

[109] - ITT-E Population.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Percentage of Participants by Subgroups (by Age, Gender, Baseline CD4+ Cell Count Baseline HIV-1 RNA, Race) With Plasma HIV-1 RNA <50 c/mL at Week 96

End point title	Percentage of Participants by Subgroups (by Age, Gender, Baseline CD4+ Cell Count Baseline HIV-1 RNA, Race) With Plasma HIV-1 RNA <50 c/mL at Week 96
-----------------	---

End point description:

Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count, Baseline HIV-1 RNA, race) with HIV-1 RNA<50 c/mL was obtained using FDA Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant ART prior to the visit of interest. Data was presented by subgroups: age (<35, 35 to <50, >=50 years); gender (males and females), Baseline CD4+ cell count (<=200 cells/mm<sup>3</sup>, >200 cells/mm<sup>3</sup>), Baseline HIV-1 RNA (<=100000, >100000) and Race (White, African American/African H., Asian and other). Percentage values are rounded to the nearest whole digit. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Week 96

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[110]</sup>	358 <sup>[111]</sup>		
Units: Percentage of participants				
Baseline CD4+ cell count, <=200,n=31, 29	65	90		
Baseline CD4+ cell count, >200,n=325,329	86	89		
Female, n=59, 52	83	88		
Male, n=297,306,	85	90		
Age, <35,n= 211, 205	84	90		
Age, 35 to <50,n=116, 107	84	89		
Age, >=50, n=29,46	86	87		
Baseline plasma HIV-1 RNA, <=100000,n=282,282	85	90		
Baseline plasma HIV-1 RNA, >100000,n=74, 76	81	88		
Race, White, n=244,247	86	90		
Race, African American/African H., n=39,36	79	81		
Race, Asian, n=37, 42	78	90		
Race, Other, n=36, 33	86	91		

Notes:

[110] - ITT-E Population.

[111] - ITT-E Population.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Percentage of Participants by Subgroups (by Age, Gender, Baseline CD4+ Cell Count Baseline HIV-1 RNA, Race) With Plasma HIV-1 RNA <50 c/mL at Week 144

End point title	Percentage of Participants by Subgroups (by Age, Gender, Baseline CD4+ Cell Count Baseline HIV-1 RNA, Race) With Plasma HIV-1 RNA <50 c/mL at Week 144
-----------------	--

End point description:

Percentage of participants by subgroups (by age, gender, Baseline CD4+ cell count, Baseline HIV-1 RNA, race) with HIV-1 RNA<50 c/mL was obtained using FDA Snapshot algorithm. The Snapshot algorithm treated all participants without HIV-1 RNA data at the visit of interest (due to missing data or discontinuation of investigational product prior to the visit window) as non-responders, as well as participants who switch their concomitant ART prior to the visit of interest. Data was presented by subgroups: age (<35, 35 to <50, >=50 years); gender (males and females), Baseline CD4+ cell count (<=200 cells/mm<sup>3</sup>, >200 cells/mm<sup>3</sup>), Baseline HIV-1 RNA (<=100000, >100000) and Race (White, African American/African H., Asian and other). Percentage values are rounded to the nearest whole digit. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Week 144

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[112]</sup>	358 <sup>[113]</sup>		
Units: Percentage of participants				
Baseline CD4+ cell count, ≤200, n=31, 29	58	83		
Baseline CD4+ cell count, >200, n=325, 329	81	83		
Female, n=59, 52	71	85		
Male, n=297, 306,	80	82		
Age, <35, n= 211, 205	77	83		
Age, 35 to <50, n=116, 107	81	83		
Age, ≥50, n=29, 46	83	78		
Baseline plasma HIV-1 RNA, ≤100000, n=282, 282	79	82		
Baseline plasma HIV-1 RNA, >100000, n=74, 76	78	87		
Race, White, n=244, 247	82	85		
Race, African American/African H., n=39, 36	69	72		
Race, Asian, n=37, 42	73	81		
Race, Other, n=36, 33	78	79		

Notes:

[112] - ITT-E Population

[113] - ITT-E Population.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Changes from Baseline in CD4+ cell counts at Week 24 by subgroups

End point title	Changes from Baseline in CD4+ cell counts at Week 24 by subgroups
-----------------	---

End point description:

CD4+ cells are type of white blood cells that fight infection& as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+. It was evaluated by flow cytometry. Baseline value is latest pre-dose assessment (Day 1). Change from Baseline as post-dose visit value minus Baseline value. Adjusted mean & standard error is presented for subgroups (Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, Age, Gender, and race). For each subgroup, adjusted mean is estimated mean change from Baseline in each arm calculated from ANCOVA model adjusting for following covariates/factors: treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, subgroup, treatment & relevant subgroup interaction. For CD4+ cell count subgroup, Baseline CD4+ cell count group is included as a factor only. Only those participants available at the specified time points were analyzed (represented by n=X in the category titles).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and Week 24

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[114]</sup>	358 <sup>[115]</sup>		
Units: Cells per cubic millimeter				
arithmetic mean (standard error)				
Baseline plasma HIV-1 RNA, <=100000, n=268,268	187.72 (± 10.860)	167.93 (± 10.842)		
Baseline plasma HIV-1 RNA, >100000, n=72,73	206.63 (± 21.107)	205.96 (± 20.990)		
Baseline CD4+ cell count, <=200, n=29,27	157.01 (± 33.113)	120.17 (± 34.151)		
Baseline CD4+ cell count, >200, n=311,314	195.11 (± 10.026)	180.73 (± 9.972)		
Age, <35, n= 203,199	202.76 (± 12.456)	177.62 (± 12.563)		
Age, 35 to <50, n=109, 100	172.05 (± 16.983)	179.87 (± 17.733)		
Age, >=50, n=28, 42	188.79 (± 33.534)	159.34 (± 27.344)		
Female, n=57,50	199.45 (± 23.498)	181.78 (± 25.263)		
Male, n=283,291	190.21 (± 10.538)	175.05 (± 10.400)		
Race, White, n=236,235	204.78 (± 11.531)	182.27 (± 11.582)		
Race, African Am/African H., n=36,33	143.84 (± 29.541)	170.51 (± 30.870)		
Race, Asian, n=34, 41	169.80 (± 30.491)	165.36 (± 27.782)		
Race, Other, n=34,32	174.30 (± 30.375)	149.34 (± 31.404)		

Notes:

[114] - ITT-E Population.

[115] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	
Baseline plasma HIV-1 RNA, <=100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count and treatment and HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	19.8
Confidence interval	
level	95 %
sides	2-sided
lower limit	-10.23
upper limit	49.83

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: Baseline plasma HIV-1 RNA,>100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count and treatment and HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	0.67
Confidence interval	
level	95 %
sides	2-sided
lower limit	-57.07
upper limit	58.4

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Baseline CD4+ cell count,<=200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	36.84
Confidence interval	
level	95 %
sides	2-sided
lower limit	-55.94
upper limit	129.63

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Baseline CD4+ cell count,>200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	14.37
Confidence interval	
level	95 %
sides	2-sided
lower limit	-13.38
upper limit	42.12

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description: Age<35. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	25.14
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9.56
upper limit	59.85

<b>Statistical analysis title</b>	Statistical Analysis 6
Statistical analysis description: Age 35 to <50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-7.82
Confidence interval	
level	95 %
sides	2-sided
lower limit	-55.98
upper limit	40.34

<b>Statistical analysis title</b>	Statistical Analysis 7
Statistical analysis description:	
Age ≥ 50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	29.45
Confidence interval	
level	95 %
sides	2-sided
lower limit	-55.47
upper limit	114.38

<b>Statistical analysis title</b>	Statistical Analysis 10
Statistical analysis description:	
Race group white. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	22.51
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9.52
upper limit	54.54

<b>Statistical analysis title</b>	Statistical Analysis 9
Statistical analysis description:	
Male. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	15.16

Confidence interval	
level	95 %
sides	2-sided
lower limit	-13.9
upper limit	44.21

<b>Statistical analysis title</b>	Statistical Analysis 8
-----------------------------------	------------------------

Statistical analysis description:

Female. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	17.67
Confidence interval	
level	95 %
sides	2-sided
lower limit	-49.89
upper limit	85.23

<b>Statistical analysis title</b>	Statistical Analysis 11
-----------------------------------	-------------------------

Statistical analysis description:

Race group African Am/African H.. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-26.67
Confidence interval	
level	95 %
sides	2-sided
lower limit	-110.64
upper limit	57.3

<b>Statistical analysis title</b>	Statistical Analysis 12
-----------------------------------	-------------------------

Statistical analysis description:

Race group Asian. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind
-------------------	---



	phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	4.44
Confidence interval	
level	95 %
sides	2-sided
lower limit	-76.18
upper limit	85.06

<b>Statistical analysis title</b>	Statistical Analysis 13
Statistical analysis description:	
Race group Other. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	24.96
Confidence interval	
level	95 %
sides	2-sided
lower limit	-60.68
upper limit	110.59

<b>Secondary: Changes from Baseline in CD4+ cell counts at Week 48 by subgroups</b>	
End point title	Changes from Baseline in CD4+ cell counts at Week 48 by subgroups
End point description:	
CD4+ cells are type of white blood cells that fight infection & as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+. It was evaluated by flow cytometry. Baseline value is latest pre-dose assessment (Day 1). Change from Baseline was post-dose visit value minus Baseline value. Adjusted mean & standard error is presented for subgroups (Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, Age group, Gender and race). For each subgroup, adjusted mean is estimated mean change from Baseline in each arm calculated from ANCOVA model adjusting for following covariates/factors: treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, subgroup, treatment & relevant subgroup interaction. For CD4+ cell count subgroup, Baseline CD4+ cell count group is included as factor only. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles)	
End point type	Secondary
End point timeframe:	
Baseline (Day 1) and Week 48	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[116]</sup>	358 <sup>[117]</sup>		
Units: Cells per cubic millimeter				
arithmetic mean (standard error)				
Baseline plasma HIV-1 RNA, <=100000, n=257,264	220.0 (± 11.72)	212.4 (± 11.56)		
Baseline plasma HIV-1 RNA, >100000, n=67,70	238.5 (± 23.09)	235.5 (± 22.70)		
Baseline CD4+ cell count, <=200, n=26, 27	200.5 (± 36.97)	177.9 (± 36.18)		
Baseline CD4+ cell count, >200, n=298, 307	225.9 (± 10.84)	220.7 (± 10.68)		
Age group-1, <35, n= 194, 192	233.6 (± 13.49)	225.2 (± 13.53)		
Age group-1, 35 to <50, n=104, 101	208.7 (± 18.40)	211.2 (± 18.67)		
Age group-1, >=50, n=26, 41	212.6 (± 36.84)	194.8 (± 29.27)		
Female, n=54, 49	237.1 (± 25.53)	226.8 (± 26.98)		
Male, n=270, 285	221.2 (± 11.41)	215.6 (± 11.11)		
Race, White, n=224, 231	226.0 (± 12.55)	219.7 (± 12.39)		
Race, African Am/African H., n=33, 31	209.8 (± 32.75)	239.9 (± 33.79)		
Race, Asian, n=34, 41	246.4 (± 32.35)	197.2 (± 29.47)		
Race, Other, n=33, 31	200.2 (± 32.69)	202.7 (± 33.83)		

Notes:

[116] - ITT-E Population.

[117] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 2
Statistical analysis description:	
Baseline plasma HIV-1 RNA, >100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count and treatment and HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-59.8
upper limit	65.9

<b>Statistical analysis title</b>	Statistical Analysis 1
Statistical analysis description: Baseline plasma HIV-1 RNA, <=100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count and treatment and HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	7.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-24.6
upper limit	39.8

<b>Statistical analysis title</b>	Statistical Analysis 6
Statistical analysis description: Age 35 to <50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-2.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-53.9
upper limit	48.9

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description: Age<35. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	8.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-29.1
upper limit	45.9

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description:	
Baseline CD4+ cell count,>200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction..	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	5.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-24.7
upper limit	35.1

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description:	
Baseline CD4+ cell count,<=200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	22.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-78.3
upper limit	123.5

<b>Statistical analysis title</b>	Statistical Analysis 12
Statistical analysis description:	
Race group Asian. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	49.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-36.3
upper limit	134.7

<b>Statistical analysis title</b>	Statistical Analysis 11
Statistical analysis description:	
Race group African Am/African H.. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-30.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-122.7
upper limit	62.4

<b>Statistical analysis title</b>	Statistical Analysis 10
Statistical analysis description:	
Race group white. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	6.3

Confidence interval	
level	95 %
sides	2-sided
lower limit	-28.2
upper limit	40.9

<b>Statistical analysis title</b>	Statistical Analysis 9
-----------------------------------	------------------------

Statistical analysis description:

Male. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	5.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-25.6
upper limit	36.9

<b>Statistical analysis title</b>	Statistical Analysis 8
-----------------------------------	------------------------

Statistical analysis description:

Female. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
Parameter estimate	Mean difference (net)
Point estimate	10.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-62.3
upper limit	83.1

<b>Statistical analysis title</b>	Statistical Analysis 13
-----------------------------------	-------------------------

Statistical analysis description:

Race group Other. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race group, and treatment and race group interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
-------------------	---

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
Parameter estimate	Mean difference (net)
Point estimate	-2.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-94.7
upper limit	89.7

<b>Statistical analysis title</b>	Statistical Analysis 7
Statistical analysis description: Age ≥ 50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	17.7
Confidence interval	
level	95 %
sides	2-sided
lower limit	-74.6
upper limit	110.1

## Secondary: Changes From Baseline in CD4+ Cell Counts at Week 96 by Subgroups

End point title	Changes From Baseline in CD4+ Cell Counts at Week 96 by Subgroups
End point description: CD4+ cells are type of white blood cells that fight infection & as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+. It was evaluated by flow cytometry. Baseline value is latest pre-dose assessment (Day 1). Change from Baseline was post-dose visit value minus Baseline value. Adjusted mean & standard error is presented for subgroups (Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, Age group, Gender and race). For each subgroup, adjusted mean is estimated mean change from Baseline in each arm calculated from ANCOVA model adjusting for following covariates/factors: treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, subgroup, treatment & relevant subgroup interaction. For CD4+ cell count subgroup, Baseline CD4+ cell count group is included as factor only. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles)	
End point type	Secondary
End point timeframe: Baseline (Day 1) and Week 96	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[118]</sup>	358 <sup>[119]</sup>		
Units: Cells per cubic millimeter				
arithmetic mean (standard error)				
Baseline plasma HIV-1 RNA, <=100000, n=240,253	254.8 (± 13.30)	252.9 (± 12.93)		
Baseline plasma HIV-1 RNA, >100000, n=61,67	300.2 (± 26.50)	260.1 (± 25.42)		
Baseline CD4+ cell count, <=200, n=21,26	240.5 (± 45.16)	244.4 (± 40.44)		
Baseline CD4+ cell count, >200, n=280,294	265.9 (± 12.28)	255.1 (± 11.98)		
Age group-1, <35, n= 179,185	270.2 (± 15.38)	263.0 (± 15.09)		
Age group-1, 35 to <50, n=97,95	259.5 (± 20.86)	262.0 (± 21.09)		
Age group-1, >=50, n=25, 40	237.6 (± 41.11)	195.9 (± 32.45)		
Female, n=49,46	277.9 (± 29.40)	259.1 (± 30.54)		
Male, n=252, 274	261.4 (± 12.95)	253.5 (± 12.43)		
Race group, White, n=210,223	275.2 (± 14.21)	260.0 (± 13.80)		
Race group, African Am/African H., n=31,29	228.5 (± 37.07)	230.2 (± 38.23)		
Race group, Asian, n=29,38	212.1 (± 38.34)	244.8 (± 33.53)		
Race group, Other, n=31,30	273.4 (± 36.97)	247.3 (± 37.67)		

Notes:

[118] - ITT-E Population.

[119] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description:	
Baseline plasma HIV-1 RNA, <=100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, Baseline plasma HIV-1 RNA, and treatment and relevant Baseline plasma HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	1.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	-34.5
upper limit	38.2



<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description: Baseline plasma HIV-1 RNA,>100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, Baseline plasma HIV-1 RNA, and treatment and Baseline plasma HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	40.1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-31.2
upper limit	111.5

<b>Statistical analysis title</b>	Statistical Analysis 3
Statistical analysis description: Baseline CD4+ cell count,<=200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-3.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	-122.3
upper limit	114.5

<b>Statistical analysis title</b>	Statistical Analysis 4
Statistical analysis description: Baseline CD4+ cell count,>200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	10.8
Confidence interval	
level	95 %
sides	2-sided
lower limit	-22.9
upper limit	44.5

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description:	
Age Group,<35. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	7.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-35
upper limit	49.6

<b>Statistical analysis title</b>	Statistical Analysis 6
Statistical analysis description:	
Age Group,35 to <50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-2.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-60.7
upper limit	55.7

<b>Statistical analysis title</b>	Statistical Analysis 7
Statistical analysis description: Age Group,>=50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	41.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-61.2
upper limit	144.5

<b>Statistical analysis title</b>	Statistical Analysis 10
Statistical analysis description: Race group-white. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	15.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-23.6
upper limit	54

<b>Statistical analysis title</b>	Statistical Analysis 9
Statistical analysis description: Male. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	7.8

Confidence interval	
level	95 %
sides	2-sided
lower limit	-27.4
upper limit	43.1

<b>Statistical analysis title</b>	Statistical Analysis 8
-----------------------------------	------------------------

Statistical analysis description:

Female. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	18.8
Confidence interval	
level	95 %
sides	2-sided
lower limit	-64.2
upper limit	101.8

<b>Statistical analysis title</b>	Statistical Analysis 11
-----------------------------------	-------------------------

Statistical analysis description:

Race group-African Am/African H. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-1.7
Confidence interval	
level	95 %
sides	2-sided
lower limit	-106.3
upper limit	103

<b>Statistical analysis title</b>	Statistical Analysis 12
-----------------------------------	-------------------------

Statistical analysis description:

Race group-Asian. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
-------------------	---

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Median difference (net)
Point estimate	-32.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-132.2
upper limit	66.9

<b>Statistical analysis title</b>	Statistical Analysis 13
-----------------------------------	-------------------------

Statistical analysis description:

Race group-Other. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	26.1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-77.3
upper limit	129.5

## Secondary: Changes From Baseline in CD4+ Cell Counts at Week 144 by Subgroups

End point title	Changes From Baseline in CD4+ Cell Counts at Week 144 by Subgroups
-----------------	--

End point description:

CD4+ cells are type of white blood cells that fight infection & as HIV infection progresses, the number of these cells declines. Blood samples were collected at specified time points to assess CD4+. It was evaluated by flow cytometry. Baseline value is latest pre-dose assessment (Day 1). Change from Baseline was post-dose visit value minus Baseline value. Adjusted mean & standard error is presented for subgroups (Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, Age group, Gender and race). For each subgroup, adjusted mean is estimated mean change from Baseline in each arm calculated from ANCOVA model adjusting for following covariates/factors: treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, subgroup, treatment & relevant subgroup interaction. For CD4+ cell count subgroup, Baseline CD4+ cell count group is included as factor only. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles)

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and Week 144

End point values	DTG + 3TC - Double-blind Phase + Open- label Phase	DTG + TDF/FTC - Double-blind Phase + Open- label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	356 <sup>[120]</sup>	358 <sup>[121]</sup>		
Units: Cells per cubic millimeter				
arithmetic mean (standard error)				
Baseline plasma HIV-1 RNA, ≤100000, n=214,223	295.7 (± 14.14)	296.1 (± 13.84)		
Baseline plasma HIV-1 RNA, >100000, n=56, 64	334.3 (± 27.77)	329.6 (± 26.15)		
Baseline CD4+ cell count, ≤200, n=17, 24	290.2 (± 50.44)	272.9 (± 42.29)		
Baseline CD4+ cell count, >200, n=253, 263	304.7 (± 12.97)	306.2 (± 12.71)		
Age group, <35, n=155, 167	298.0 (± 16.59)	316.0 (± 15.93)		
Age group-1, 35 to <50, n=92, 87	305.6 (± 21.48)	302.1 (± 22.11)		
Age group-1, ≥50, n=23,33	337.4 (± 43.01)	242.2 (± 35.83)		
Female, n=43, 43	346.6 (± 31.40)	321.7 (± 31.67)		
Male, n=227, 244	295.9 (± 13.66)	300.0 (± 13.20)		
Race group, White, n=190,201	314.2 (± 14.93)	314.0 (± 14.53)		
Race group, African Am/African H., n=26, 26	243.8 (± 40.42)	295.1 (± 40.39)		
Race group, Asian, n=26, 34	244.0 (± 40.47)	264.1 (± 35.41)		
Race group, Other, n=28,26	346.2 (± 38.87)	279.9 (± 40.42)		

Notes:

[120] - ITT-E Population.

[121] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 2
Statistical analysis description:	
Baseline plasma HIV-1 RNA, >100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, Baseline plasma HIV-1 RNA, and treatment and Baseline plasma HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	4.7
Confidence interval	
level	95 %
sides	2-sided
lower limit	-69.4
upper limit	78.8

<b>Statistical analysis title</b>	Statistical Analysis 1
Statistical analysis description: Baseline plasma HIV-1 RNA, <=100000. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, Baseline plasma HIV-1 RNA, and treatment and relevant Baseline plasma HIV-1 RNA interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-0.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-39.2
upper limit	38.3

<b>Statistical analysis title</b>	Statistical Analysis 5
Statistical analysis description: Age Group, <35. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-18
Confidence interval	
level	95 %
sides	2-sided
lower limit	-63.2
upper limit	27.1

<b>Statistical analysis title</b>	Statistical Analysis 6
Statistical analysis description: Age Group, 35 to <50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	3.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-57
upper limit	64

<b>Statistical analysis title</b>	Statistical Analysis 13
Statistical analysis description: Race group-Other. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	66.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-43.7
upper limit	176.2

<b>Statistical analysis title</b>	Statistical Analysis 8
Statistical analysis description: Female. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	24.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	-62.5
upper limit	112.3



<b>Statistical analysis title</b>	Statistical Analysis 9
Statistical analysis description:	
Male. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, gender, and treatment and gender interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-4.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-41.5
upper limit	33.1

<b>Statistical analysis title</b>	Statistical Analysis 10
Statistical analysis description:	
Race group-white. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	0.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-40.6
upper limit	41.1

<b>Statistical analysis title</b>	Statistical Analysis 11
Statistical analysis description:	
Race group-African Am/African H. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-51.3

Confidence interval	
level	95 %
sides	2-sided
lower limit	-163.6
upper limit	60.9

<b>Statistical analysis title</b>	Statistical Analysis 12
-----------------------------------	-------------------------

Statistical analysis description:

Race group-Asian. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, race, and treatment and race interaction.

Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Median difference (net)
Point estimate	-20.1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-125.3
upper limit	85

<b>Statistical analysis title</b>	Statistical Analysis 4
-----------------------------------	------------------------

Statistical analysis description:

Baseline CD4+ cell count,>200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction.

Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	-1.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-37.2
upper limit	34.1

<b>Statistical analysis title</b>	Statistical Analysis 3
-----------------------------------	------------------------

Statistical analysis description:

Baseline CD4+ cell count,<=200. Following covariates were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, and treatment and Baseline CD4+ cell count interaction.

Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
-------------------	---

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	17.4
Confidence interval	
level	95 %
sides	2-sided
lower limit	-111.1
upper limit	145.8

<b>Statistical analysis title</b>	Statistical Analysis 7
Statistical analysis description:	
Age Group, >=50. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count, age, and treatment and age interaction.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Mean difference (net)
Point estimate	95.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-14.8
upper limit	205.2

### **Secondary: Change from Baseline in EuroQol – 5 Dimensions – 5 Levels (EQ-5D-5L) utility score at Weeks 4, 24, 48**

End point title	Change from Baseline in EuroQol – 5 Dimensions – 5 Levels (EQ-5D-5L) utility score at Weeks 4, 24, 48
End point description:	
EQ-5D-5L questionnaire provides a profile of participant function and a global health state rating. The five-item measure has 1 question assessing each of 5 dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression and 5 levels for each dimension including 1=no problems, 2=slight problems, 3=moderate problems, 4=severe problems and 5=extreme problems. The health state is defined by combining the levels of answers from each of the 5 questions. Each health state is referred to in terms of a 5 digit code. Health state 5 digit code is translated into utility score, which is valued up to 1 (perfect health) with lower values meaning worse state. EQ-5D-5L utility score ranges from -0.281 to 1. Higher scores indicate better health. Baseline was the latest pre-dose assessment (Day 1) and change from Baseline=post-dose value minus Baseline value. Only those participants available at the specified time points were analyzed.	
End point type	Secondary
End point timeframe:	
Baseline (Day 1) and Weeks 4, 24, 48	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[122]</sup>	358 <sup>[123]</sup>		
Units: Scores on a scale				
arithmetic mean (standard error)				
Week 4, n=349, 348	0.0130 (± 0.00362)	0.0078 (± 0.00353)		
Week 24, n=352, 351	0.0131 (± 0.00371)	0.0168 (± 0.00333)		
Week 48, n=352, 351	0.0134 (± 0.00384)	0.0129 (± 0.00349)		

Notes:

[122] - ITT-E Population.

[123] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description:	
Week 4. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count (factor), and Baseline EQ-5D utility, treatment*visit and Baseline EQ-5D utility*visit with visit as the repeated factor..	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.302
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	0.0052
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.0047
upper limit	0.0152

Statistical analysis title	Statistical Analysis 3
Statistical analysis description:	
Week48. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count (factor), and Baseline EQ-5D utility, treatment*visit and Baseline EQ-5D utility*visit with visit as the repeated factor..	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase

Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.934
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	0.0004
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.0098
upper limit	0.0106

<b>Statistical analysis title</b>	Statistical Analysis 2
-----------------------------------	------------------------

Statistical analysis description:

Week24. Following covariates/factors were adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count (factor), and Baseline EQ-5D utility, treatment\*visit and Baseline EQ-5D utility\*visit with visit as the repeated factor.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.45
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	-0.0038
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.0136
upper limit	0.006

## Secondary: Change From Baseline in EQ-5D-5L Utility Score at Week 96

End point title	Change From Baseline in EQ-5D-5L Utility Score at Week 96
-----------------	---

End point description:

EQ-5D-5L questionnaire provides a profile of participant function and a global health state rating. The five-item measure has 1 question assessing each of 5 dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression and 5 levels for each dimension including 1=no problems, 2=slight problems, 3=moderate problems, 4=severe problems and 5=extreme problems. The health state is defined by combining the levels of answers from each of the 5 questions. Each health state is referred to in terms of a 5 digit code. Health state 5 digit code is translated into utility score, which is valued up to 1 (perfect health) with lower values meaning worse state. EQ-5D-5L utility score ranges from -0.281 to 1. Higher scores indicate better health. Baseline was the latest pre-dose assessment (Day 1) and change from Baseline=post-dose value minus Baseline value. Only those participants available at the specified time points were analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and Week 96

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	352 <sup>[124]</sup>	351 <sup>[125]</sup>		
Units: Scores on a scale				
arithmetic mean (standard error)	0.0079 ( $\pm$ 0.00450)	0.0091 ( $\pm$ 0.00408)		

Notes:

[124] - ITT-E Population.

[125] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	
Week 96. Covariates adjusted: Treatment, Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, and Baseline EQ-5D utility, treatment*visit and Baseline EQ-5D utility*visit as factors and covariate, with visit as the repeated factor.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	703
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.842
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	-0.0012
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.0132
upper limit	0.0107

## Secondary: Change From Baseline in EQ-5D-5L Utility Score at Week 144

End point title	Change From Baseline in EQ-5D-5L Utility Score at Week 144
End point description:	
EQ-5D-5L questionnaire provides a profile of participant function and a global health state rating. The five-item measure has 1 question assessing each of 5 dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression and 5 levels for each dimension including 1=no problems, 2=slight problems, 3=moderate problems, 4=severe problems and 5=extreme problems. The health state is defined by combining the levels of answers from each of the 5 questions. Each health state is referred to in terms of a 5 digit code. Health state 5 digit code is translated into utility score, which is valued up to 1 (perfect health) with lower values meaning worse state. EQ-5D-5L utility score ranges from -0.281 to 1. Higher scores indicate better health. Baseline was the latest pre-dose assessment (Day 1) and change from Baseline=post-dose value minus Baseline value. Only those participants available at the specified time points were analyzed.	
End point type	Secondary

End point timeframe:

Baseline (Day 1) and Week 144

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	352 <sup>[126]</sup>	351 <sup>[127]</sup>		
Units: Scores on a scale				
arithmetic mean (standard error)	0.0143 (± 0.00388)	0.0135 (± 0.00365)		

Notes:

[126] - ITT-E Population.

[127] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description: Week 144. Covariates adjusted: Treatment, Baseline plasma HIV-1 RNA, Baseline CD4+ cell count, and Baseline EQ-5D utility, treatment*visit and Baseline EQ-5D utility*visit as factors and covariate, with visit as the repeated factor.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	703
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.879
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	0.0008
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.0097
upper limit	0.0113

## Secondary: Change from Baseline in EuroQol – 5 Dimensions – 5 Levels (EQ-5D-5L) Thermometer Scores at Weeks 4, 24 48

End point title	Change from Baseline in EuroQol – 5 Dimensions – 5 Levels (EQ-5D-5L) Thermometer Scores at Weeks 4, 24 48
-----------------	---

End point description:

EQ-5D-5L questionnaire provides a profile of participant function and a global health state rating. The five-item measure has one question assessing each of five dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression and 5 levels for each dimension including 1=no problems, 2=slight problems, 3=moderate problems, 4=severe problems and 5=extreme problems. EQ-5D-5L included EQ visual Analogue scale (EQ VAS) 'Thermometer' which provided Self-rated current health status. Score ranges from 0 (worst imaginable health state) to 100 (best imaginable health state). MMRM was run on the LOCF dataset, using the observed margins (OM) option. Baseline was the

latest pre-dose assessment value (Day 1) and change from Baseline=post-dose value minus Baseline value. Only those participants available at the specified time points were analyzed (represented by n=x in the category titles).)

End point type	Secondary
End point timeframe:	
Baseline (Day 1) and Weeks 4, 24, 48	

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	356 <sup>[128]</sup>	358 <sup>[129]</sup>		
Units: Scores on a scale				
arithmetic mean (standard error)				
Week 4, n=349, 348	2.3 (± 0.48)	1.2 (± 0.52)		
Week 24, n=352, 350	3.7 (± 0.54)	3.2 (± 0.51)		
Week 48, n=352, 350	4.3 (± 0.49)	2.8 (± 0.49)		

Notes:

[128] - ITT-E Population.

[129] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
----------------------------	------------------------

Statistical analysis description:

Week 4. Covariates adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count (factor), Baseline EQ-5D thermometer, treatment\*visit and Baseline EQ-5D thermometer\*visit with visit as the repeated factor

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.137
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	1.1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.3
upper limit	2.4

Statistical analysis title	Statistical Analysis 3
----------------------------	------------------------

Statistical analysis description:

Week48. Covariates adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count (factor), Baseline EQ-5D thermometer, treatment\*visit and Baseline EQ-5D thermometer\*visit with visit as the repeated factor.

Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind
-------------------	---



	phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.031
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	1.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.1
upper limit	2.8

<b>Statistical analysis title</b>	Statistical Analysis 2
Statistical analysis description:	
Week24. Covariates adjusted: Treatment, Baseline plasma HIV-1 RNA (factor), Baseline CD4+ cell count (factor), Baseline EQ-5D thermometer, treatment*visit and Baseline EQ-5D thermometer*visit with visit as the repeated factor	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	714
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.458
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	0.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.9
upper limit	2

## Secondary: Change From Baseline in EQ-5D-5L Thermometer Scores at Week 96

End point title	Change From Baseline in EQ-5D-5L Thermometer Scores at Week 96
End point description:	
EQ-5D-5L questionnaire provided a profile of participant function and a global health state rating. The five-item measure has one question assessing each of five dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression and 5 levels for each dimension including 1=no problems, 2=slight problems, 3=moderate problems, 4=severe problems and 5=extreme problems. EQ-5D-5L included EQ visual Analogue scale (EQ VAS) 'Thermometer' which provided Self-rated current health status. Score ranges from 0 (worst imaginable health state) to 100 (best imaginable health state). MMRM was run on the LOCF dataset, using the observed margins (OM) option. Baseline was the latest pre-dose assessment value (Day 1) and change from Baseline=post-dose value minus Baseline value. Adjusted mean and standard error is presented. Only those participants available at the specified time points were analyzed.	
End point type	Secondary

End point timeframe:

Baseline (Day 1) and Week 96

End point values	DTG + 3TC- Double blind phase	DTG + TDF/FTC- Double blind phase		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	352 <sup>[130]</sup>	350 <sup>[131]</sup>		
Units: Scores on a scale				
arithmetic mean (standard error)	4.1 (± 0.51)	2.4 (± 0.56)		

Notes:

[130] - ITT-E Population.

[131] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description:	
Week 96. Covariates adjusted: Treatment, Baseline plasma HIV-1 RNA, Baseline CD4+ cell count and Baseline EQ-5D utility, treatment*visit and Baseline EQ-5D utility*visit as factors and covariate, with visit as the repeated factor.	
Comparison groups	DTG + 3TC-Double blind phase v DTG + TDF/FTC-Double blind phase
Number of subjects included in analysis	702
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.027
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	1.7
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.2
upper limit	3.2

## Secondary: Change From Baseline in EQ-5D-5L Thermometer Scores at Week 144

End point title	Change From Baseline in EQ-5D-5L Thermometer Scores at Week 144
-----------------	---

End point description:

EQ-5D-5L questionnaire provided a profile of participant function and a global health state rating. The five-item measure has one question assessing each of five dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression and 5 levels for each dimension including 1=no problems, 2=slight problems, 3=moderate problems, 4=severe problems and 5=extreme problems. EQ-5D-5L included EQ visual Analogue scale (EQ VAS) 'Thermometer' which provided Self-rated current health status. Score ranges from 0 (worst imaginable health state) to 100 (best imaginable health state). MMRM was run on the LOCF dataset, using the observed margins (OM) option. Baseline was the latest pre-dose assessment value (Day 1) and change from Baseline=post-dose value minus Baseline value. Adjusted mean and standard error is presented. Only those participants available at the specified time points were analyzed.

End point type	Secondary
End point timeframe:	
Baseline (Day 1) and Week 144	

End point values	DTG + 3TC - Double-blind Phase + Open-label Phase	DTG + TDF/FTC - Double-blind Phase + Open-label Phase		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	352 <sup>[132]</sup>	350 <sup>[133]</sup>		
Units: Scores on a scale				
arithmetic mean (standard error)	5.2 (± 0.48)	3.0 (± 0.50)		

Notes:

[132] - ITT-E Population.

[133] - ITT-E Population.

## Statistical analyses

Statistical analysis title	Statistical Analysis
Statistical analysis description:	
Week 144. Covariates adjusted: Treatment, Baseline plasma HIV-1 RNA, Baseline CD4+ cell count and Baseline EQ-5D utility, treatment*visit and Baseline EQ-5D utility*visit as factors and covariate, with visit as the repeated factor.	
Comparison groups	DTG + 3TC - Double-blind Phase + Open-label Phase v DTG + TDF/FTC - Double-blind Phase + Open-label Phase
Number of subjects included in analysis	702
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.001
Method	MMRM
Parameter estimate	Mean difference (net)
Point estimate	2.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.9
upper limit	3.6

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

All-cause mortality, SAEs and non-SAEs were collected up to Week 148 in Double-blind Phase + Open-label Phase and from Week 148 to Week 280 in Continuation Phase

Adverse event reporting additional description:

All-cause mortality, SAEs and non-SAEs were reported for the Safety Population for the Double-blind Phase + Open-label Phase. Safety-Continuation Population was used for the Continuation Phase which comprises all participants who received at least 1 dose of study treatment after entering the Continuation Phase

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
Dictionary version	25.0

### Reporting groups

Reporting group title	DTG + 3TC-Double-blind Phase + Open-label Phase
-----------------------	---

Reporting group description:

Participants received a two-drug regimen of DTG + 3TC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + 3TC from Week 96 to Week 148 in the open-label phase.

Reporting group title	DTG + 3TC-Continuation Phase
-----------------------	------------------------------

Reporting group description:

Participants received a DTG + 3TC administered orally, once daily from Week 148 to Week 280 in the continuation phase.

Reporting group title	DTG + TDF/FTC-Double-blind Phase + Open-label Phase
-----------------------	---

Reporting group description:

Participants received a three-drug regimen of DTG + TDF/FTC FDC administered orally, once daily until Week 96 in double-blind phase and participants continued to receive DTG + TDF/FTC FDC from Week 96 to Week 148 in the open-label phase.

Serious adverse events	DTG + 3TC-Double-blind Phase + Open-label Phase	DTG + 3TC-Continuation Phase	DTG + TDF/FTC-Double-blind Phase + Open-label Phase
Total subjects affected by serious adverse events			
subjects affected / exposed	37 / 356 (10.39%)	5 / 233 (2.15%)	38 / 358 (10.61%)
number of deaths (all causes)	1	0	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
B-cell lymphoma			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anogenital warts			

subjects affected / exposed	2 / 356 (0.56%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Clear cell renal cell carcinoma			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hodgkin's disease			
subjects affected / exposed	2 / 356 (0.56%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Varicose vein			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
Abortion spontaneous			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	2 / 358 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Non-cardiac chest pain			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Chest pain			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Asthma			

subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pulmonary sarcoidosis			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Major depression			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicide attempt			
subjects affected / exposed	3 / 356 (0.84%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 3	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Substance-induced psychotic disorder			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychotic disorder			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Delirium			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicidal ideation			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bipolar I disorder			

subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Head injury			
subjects affected / exposed	1 / 356 (0.28%)	1 / 233 (0.43%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Joint dislocation			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Post lumbar puncture syndrome			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tendon rupture			
subjects affected / exposed	0 / 356 (0.00%)	1 / 233 (0.43%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Jaw fracture			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Multiple injuries			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders			
Coronary artery stenosis			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Coronary artery disease			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 1	0 / 0	0 / 0
Cardiac failure			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acute myocardial infarction			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Polyneuropathy			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Syncope			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acute polyneuropathy			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Lymphadenopathy			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Gastric ulcer haemorrhage			



subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Intestinal obstruction			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Colitis			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anal fistula			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Cholecystitis acute			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cholelithiasis			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatitis toxic			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Nephrolithiasis			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	3 / 358 (0.84%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 3
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue			

disorders			
Rhabdomyolysis			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Arthritis reactive			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Abscess limb			
subjects affected / exposed	2 / 356 (0.56%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			
subjects affected / exposed	2 / 356 (0.56%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 2	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatitis A			
subjects affected / exposed	3 / 356 (0.84%)	1 / 233 (0.43%)	4 / 358 (1.12%)
occurrences causally related to treatment / all	0 / 3	0 / 1	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Appendicitis			
subjects affected / exposed	1 / 356 (0.28%)	2 / 233 (0.86%)	4 / 358 (1.12%)
occurrences causally related to treatment / all	0 / 1	0 / 2	0 / 4
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Bacterial infection			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anal abscess			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Acute hepatitis C			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cellulitis			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	2 / 358 (0.56%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Chagoma			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Lower respiratory tract infection			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Helicobacter gastritis			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Herpes zoster			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Perineal abscess			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Encephalitis			

subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Appendicitis perforated			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Peritonitis			
subjects affected / exposed	1 / 356 (0.28%)	0 / 233 (0.00%)	0 / 358 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Erysipelas			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary tract infection			
subjects affected / exposed	0 / 356 (0.00%)	0 / 233 (0.00%)	1 / 358 (0.28%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Frequency threshold for reporting non-serious adverse events: 2 %

<b>Non-serious adverse events</b>	DTG + 3TC-Double-blind Phase + Open-label Phase	DTG + 3TC-Continuation Phase	DTG + TDF/FTC-Double-blind Phase + Open-label Phase
Total subjects affected by non-serious adverse events			
subjects affected / exposed	272 / 356 (76.40%)	46 / 233 (19.74%)	268 / 358 (74.86%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Anogenital warts			
subjects affected / exposed	14 / 356 (3.93%)	0 / 233 (0.00%)	17 / 358 (4.75%)
occurrences (all)	14	0	19
Vascular disorders			
Hypertension			
subjects affected / exposed	16 / 356 (4.49%)	5 / 233 (2.15%)	14 / 358 (3.91%)
occurrences (all)	18	9	14
Nervous system disorders			

Headache subjects affected / exposed occurrences (all)	51 / 356 (14.33%) 80	7 / 233 (3.00%) 8	47 / 358 (13.13%) 82
Dizziness subjects affected / exposed occurrences (all)	10 / 356 (2.81%) 13	0 / 233 (0.00%) 0	10 / 358 (2.79%) 10
Paraesthesia subjects affected / exposed occurrences (all)	4 / 356 (1.12%) 4	0 / 233 (0.00%) 0	9 / 358 (2.51%) 10
General disorders and administration site conditions			
Fatigue subjects affected / exposed occurrences (all)	16 / 356 (4.49%) 19	0 / 233 (0.00%) 0	14 / 358 (3.91%) 15
Pyrexia subjects affected / exposed occurrences (all)	18 / 356 (5.06%) 24	0 / 233 (0.00%) 0	12 / 358 (3.35%) 13
Influenza like illness subjects affected / exposed occurrences (all)	19 / 356 (5.34%) 20	0 / 233 (0.00%) 0	16 / 358 (4.47%) 25
Immune system disorders			
Seasonal allergy subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 10	0 / 233 (0.00%) 0	8 / 358 (2.23%) 12
Gastrointestinal disorders			
Diarrhoea subjects affected / exposed occurrences (all)	50 / 356 (14.04%) 69	0 / 233 (0.00%) 0	53 / 358 (14.80%) 73
Nausea subjects affected / exposed occurrences (all)	13 / 356 (3.65%) 13	0 / 233 (0.00%) 0	32 / 358 (8.94%) 37
Dyspepsia subjects affected / exposed occurrences (all)	13 / 356 (3.65%) 20	0 / 233 (0.00%) 0	12 / 358 (3.35%) 13
Haemorrhoids subjects affected / exposed occurrences (all)	21 / 356 (5.90%) 25	0 / 233 (0.00%) 0	16 / 358 (4.47%) 17

Abdominal pain subjects affected / exposed occurrences (all)	16 / 356 (4.49%) 16	0 / 233 (0.00%) 0	16 / 358 (4.47%) 19
Abdominal pain upper subjects affected / exposed occurrences (all)	6 / 356 (1.69%) 9	0 / 233 (0.00%) 0	10 / 358 (2.79%) 12
Vomiting subjects affected / exposed occurrences (all)	11 / 356 (3.09%) 12	0 / 233 (0.00%) 0	9 / 358 (2.51%) 10
Gastritis subjects affected / exposed occurrences (all)	10 / 356 (2.81%) 11	0 / 233 (0.00%) 0	6 / 358 (1.68%) 6
Toothache subjects affected / exposed occurrences (all)	11 / 356 (3.09%) 15	0 / 233 (0.00%) 0	9 / 358 (2.51%) 11
Gastrooesophageal reflux disease subjects affected / exposed occurrences (all)	10 / 356 (2.81%) 10	0 / 233 (0.00%) 0	7 / 358 (1.96%) 7
Respiratory, thoracic and mediastinal disorders			
Oropharyngeal pain subjects affected / exposed occurrences (all)	12 / 356 (3.37%) 13	0 / 233 (0.00%) 0	15 / 358 (4.19%) 15
Cough subjects affected / exposed occurrences (all)	13 / 356 (3.65%) 13	0 / 233 (0.00%) 0	19 / 358 (5.31%) 19
Rhinitis allergic subjects affected / exposed occurrences (all)	15 / 356 (4.21%) 24	0 / 233 (0.00%) 0	5 / 358 (1.40%) 5
Respiratory disorder subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 11	0 / 233 (0.00%) 0	5 / 358 (1.40%) 7
Nasal congestion subjects affected / exposed occurrences (all)	3 / 356 (0.84%) 3	0 / 233 (0.00%) 0	8 / 358 (2.23%) 8
Asthma			

subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 8	0 / 233 (0.00%) 0	3 / 358 (0.84%) 3
Skin and subcutaneous tissue disorders			
Pruritus			
subjects affected / exposed	9 / 356 (2.53%)	0 / 233 (0.00%)	7 / 358 (1.96%)
occurrences (all)	9	0	8
Rash			
subjects affected / exposed	11 / 356 (3.09%)	0 / 233 (0.00%)	8 / 358 (2.23%)
occurrences (all)	11	0	13
Psychiatric disorders			
Insomnia			
subjects affected / exposed	29 / 356 (8.15%)	0 / 233 (0.00%)	36 / 358 (10.06%)
occurrences (all)	33	0	40
Anxiety			
subjects affected / exposed	19 / 356 (5.34%)	0 / 233 (0.00%)	15 / 358 (4.19%)
occurrences (all)	24	0	18
Depression			
subjects affected / exposed	18 / 356 (5.06%)	0 / 233 (0.00%)	17 / 358 (4.75%)
occurrences (all)	20	0	18
Musculoskeletal and connective tissue disorders			
Back pain			
subjects affected / exposed	22 / 356 (6.18%)	0 / 233 (0.00%)	25 / 358 (6.98%)
occurrences (all)	31	0	28
Arthralgia			
subjects affected / exposed	7 / 356 (1.97%)	0 / 233 (0.00%)	22 / 358 (6.15%)
occurrences (all)	8	0	24
Myalgia			
subjects affected / exposed	5 / 356 (1.40%)	0 / 233 (0.00%)	9 / 358 (2.51%)
occurrences (all)	6	0	9
Pain in extremity			
subjects affected / exposed	8 / 356 (2.25%)	0 / 233 (0.00%)	6 / 358 (1.68%)
occurrences (all)	10	0	6
Musculoskeletal pain			
subjects affected / exposed	4 / 356 (1.12%)	0 / 233 (0.00%)	10 / 358 (2.79%)
occurrences (all)	4	0	11
Infections and infestations			

Upper respiratory tract infection subjects affected / exposed occurrences (all)	37 / 356 (10.39%) 50	7 / 233 (3.00%) 8	33 / 358 (9.22%) 41
Nasopharyngitis subjects affected / exposed occurrences (all)	49 / 356 (13.76%) 74	7 / 233 (3.00%) 8	58 / 358 (16.20%) 84
Influenza subjects affected / exposed occurrences (all)	24 / 356 (6.74%) 28	0 / 233 (0.00%) 0	18 / 358 (5.03%) 20
Bronchitis subjects affected / exposed occurrences (all)	30 / 356 (8.43%) 36	0 / 233 (0.00%) 0	21 / 358 (5.87%) 23
Syphilis subjects affected / exposed occurrences (all)	38 / 356 (10.67%) 45	6 / 233 (2.58%) 6	40 / 358 (11.17%) 47
Pharyngitis subjects affected / exposed occurrences (all)	38 / 356 (10.67%) 49	6 / 233 (2.58%) 8	30 / 358 (8.38%) 43
Herpes zoster subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 8	0 / 233 (0.00%) 0	13 / 358 (3.63%) 13
Tonsillitis subjects affected / exposed occurrences (all)	13 / 356 (3.65%) 15	0 / 233 (0.00%) 0	12 / 358 (3.35%) 13
Gonorrhoea subjects affected / exposed occurrences (all)	16 / 356 (4.49%) 22	0 / 233 (0.00%) 0	13 / 358 (3.63%) 18
Gastroenteritis subjects affected / exposed occurrences (all)	17 / 356 (4.78%) 19	0 / 233 (0.00%) 0	15 / 358 (4.19%) 15
Sinusitis subjects affected / exposed occurrences (all)	15 / 356 (4.21%) 17	0 / 233 (0.00%) 0	15 / 358 (4.19%) 20
Pharyngotonsillitis subjects affected / exposed occurrences (all)	15 / 356 (4.21%) 26	0 / 233 (0.00%) 0	12 / 358 (3.35%) 18



Respiratory tract infection viral subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 10	0 / 233 (0.00%) 0	9 / 358 (2.51%) 15
Respiratory tract infection subjects affected / exposed occurrences (all)	9 / 356 (2.53%) 13	0 / 233 (0.00%) 0	10 / 358 (2.79%) 19
Suspected COVID-19 subjects affected / exposed occurrences (all)	0 / 356 (0.00%) 0	8 / 233 (3.43%) 8	0 / 358 (0.00%) 0
COVID-19 subjects affected / exposed occurrences (all)	0 / 356 (0.00%) 0	16 / 233 (6.87%) 18	0 / 358 (0.00%) 0
Conjunctivitis subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 8	0 / 233 (0.00%) 0	6 / 358 (1.68%) 6
Anal chlamydia infection subjects affected / exposed occurrences (all)	9 / 356 (2.53%) 16	0 / 233 (0.00%) 0	5 / 358 (1.40%) 5
Proctitis gonococcal subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 12	0 / 233 (0.00%) 0	9 / 358 (2.51%) 12
Oral herpes subjects affected / exposed occurrences (all)	9 / 356 (2.53%) 11	0 / 233 (0.00%) 0	8 / 358 (2.23%) 9
Acute sinusitis subjects affected / exposed occurrences (all)	4 / 356 (1.12%) 4	0 / 233 (0.00%) 0	8 / 358 (2.23%) 9
Tooth infection subjects affected / exposed occurrences (all)	10 / 356 (2.81%) 10	0 / 233 (0.00%) 0	7 / 358 (1.96%) 7
Oropharyngeal gonococcal infection subjects affected / exposed occurrences (all)	8 / 356 (2.25%) 9	0 / 233 (0.00%) 0	7 / 358 (1.96%) 8
Chlamydial infection subjects affected / exposed occurrences (all)	14 / 356 (3.93%) 15	0 / 233 (0.00%) 0	10 / 358 (2.79%) 10

Urethritis			
subjects affected / exposed	9 / 356 (2.53%)	0 / 233 (0.00%)	13 / 358 (3.63%)
occurrences (all)	11	0	13
Urinary tract infection			
subjects affected / exposed	9 / 356 (2.53%)	0 / 233 (0.00%)	13 / 358 (3.63%)
occurrences (all)	11	0	13
Metabolism and nutrition disorders			
Vitamin D deficiency			
subjects affected / exposed	25 / 356 (7.02%)	0 / 233 (0.00%)	22 / 358 (6.15%)
occurrences (all)	25	0	23

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
01 December 2017	<p>Amendment No. 1: The double barrier method of contraception (male condom combined with a vaginal spermicide) was added in this study as a permitted method for preventing pregnancy in females of reproductive potential. Exclusion criterion #15 (limitations on investigational drug use) was broadened to include additional countries as needed. Inclusion of Portugal was required by the Portuguese National Ethics Committee for Clinical Research. Assessment of weight at Weeks 96 and 144 was added to monitor the incidence of significant weight gain with dolutegravir use. Assessment of inflammation biomarkers (IL-6, hs-CRP) at Day 1, and Weeks 48, 96 and 144, was added as a new exploratory endpoint. Assessment of telomere length at Day 1, and Weeks 96 and 144, was added as a new exploratory endpoint. For clarification purposes, the 'peripheral blood mononuclear cell (PBMC)' sample in Time and Events table and HIV-1 Exploratory Analyses was renamed as a 'whole blood' sample. The Day 1 'PBMC' sample (now named 'whole blood' sample) originally designated for virology use was additionally designated for telomere length measurement, where possible. Additional whole blood samples were added for measurement of telomere length at Week 96 and Week 144. A description of commercial image dolutegravir tablets was added to Investigational Product and Other Study Treatment to allow use of commercial material as well as clinical trial material during the study. The physical description for open-label lamivudine was corrected. Standard procedures for forwarding pregnancy information to the Antiretroviral Pregnancy Register were added. For clarification purposes, the AE severity gradings in were updated to be consistent. This change has no impact on the investigator's evaluation of adverse events. Minor revisions were made to the text to provide updated information, correct errors and improve accuracy and consistency.</p>
14 June 2018	<p>Changes were made to the protocol to manage and mitigate risks following identification of a potential safety issue related to neural tube defects in infants born to women with exposure to dolutegravir at the time of conception. Changes were also made to update the references to the DTG IB to reflect the most current versions.</p> <ul style="list-style-type: none"><li>• The Risk Assessment table (Section 4.6.1) was updated to include language regarding risk and mitigation of neural tube defects.</li><li>• The withdrawal criteria (Section 5.4) were updated to include a reminder that females of reproductive potential who change their minds and desire to be pregnant, or who state they no longer are willing to comply with the approved pregnancy avoidance methods, should also be withdrawn from the study.</li><li>• The Time and Events table (Section 7.1). was updated to include a reminder for investigators to check at every visit that females of reproductive potential are avoiding pregnancy.</li><li>• The modified list of highly effective methods for avoiding pregnancy in females of reproductive potential (FRP) (Section 12.9.1) was updated to exclude the double barrier method of contraception, which does not meet updated GSK/ViiV criteria for a highly effective method.</li></ul>

Notes:

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported