



## Clinical trial results:

### A Multicenter, Open-label, Single-arm, Two-step Study to Evaluate the Safety and Single-dose Pharmacokinetics of Famciclovir and Multiple-dose Safety After Administration of Famciclovir Oral Pediatric Formulation to Children 1 to 12 Years of Age With Herpes Simplex Infection

#### Summary

EudraCT number	2015-004441-17
Trial protocol	Outside EU/EEA
Global end of trial date	07 December 2007

#### Results information

Result version number	v1 (current)
This version publication date	19 December 2016
First version publication date	19 December 2016

#### Trial information

##### Trial identification

Sponsor protocol code	CFAM810B2303
-----------------------	--------------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT00098059
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	Novartis Pharma AG
Sponsor organisation address	CH 4002, Basel, Switzerland,
Public contact	Clinical Disclosure Office, Novartis Pharma AG, +41 613241111,
Scientific contact	Clinical Disclosure Office, Novartis Pharma AG, +41 613241111,

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	07 December 2007
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	07 December 2007
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

The main objective of the trial were:

Part A: To evaluate the safety and tolerability, and pharmacokinetics (PK) of a single dose of famciclovir oral pediatric formulation in children from 1 to 12 years of age with herpes simplex virus (HSV) infection, in order to define the dose in this age group which gives a similar exposure to a 500 milligrams (mg) dose in adults,

Part B: To evaluate the safety and tolerability of multiple doses of famciclovir pediatric formulation administered twice-daily for 7 days in subjects from 1 to 12 years of age who had HSV infection.

Protection of trial subjects:

The study was in compliance with the ethical principles derived from the Declaration of Helsinki and in compliance with all International Conference on Harmonization (ICH) Good Clinical Practice (GCP) Guidelines. All the local regulatory requirements pertinent to safety of trial subjects were followed.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	05 February 2005
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	United States: 63
Country: Number of subjects enrolled	Panama: 11
Worldwide total number of subjects	74
EEA total number of subjects	0

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	17
Children (2-11 years)	53
Adolescents (12-17 years)	4

Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

The study was conducted at USA (6 centers):Part A; USA (8 centers), Panama (2 centers): Part B

### Pre-assignment

Screening details:

A total of 74 subjects (Part A: 27 subjects, Part B: 47 subjects) were enrolled in the study.

### Period 1

Period 1 title	Overall period (overall period)
Is this the baseline period?	Yes
Allocation method	Non-randomised - controlled
Blinding used	Not blinded

Blinding implementation details:

The study was open label study, hence no blinding was performed

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Famciclovir: Single dose (Part A)

Arm description:

Subjects were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kilogram (kg) body weight with a dose escalation up to a maximum dose of 500 mg. Subjects who had weight  $\geq 40$  kg received a famciclovir dose of 500 mg.

Arm type	Experimental
Investigational medicinal product name	Famciclovir
Investigational medicinal product code	FAM810
Other name	Famvir
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

Subjects were orally administered with a single-dose closest to 12.5 mg/kg body weight with a maximum dose of 500 mg.

<b>Arm title</b>	Famciclovir: Multiple doses (Part B)
------------------	--------------------------------------

Arm description:

Subjects were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy twice daily (b.i.d.) with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight  $\geq 40$  kg received a famciclovir dose of 500 mg.

Arm type	Experimental
Investigational medicinal product name	Famciclovir
Investigational medicinal product code	FAM810
Other name	Famvir
Pharmaceutical forms	Capsule
Routes of administration	Oral use

Dosage and administration details:

Subjects were orally administered with a body weight stratified dose ranged from 150 mg to 500 mg b.i.d. of famciclovir sprinkle capsules with a dose separation of 12 hours for a period of 7 days.

<b>Number of subjects in period 1</b>	Famciclovir: Single dose (Part A)	Famciclovir: Multiple doses (Part B)
Started	27	47
Completed	27	46
Not completed	0	1
Protocol deviation	-	1

## Baseline characteristics

### Reporting groups

Reporting group title	Famciclovir: Single dose (Part A)
-----------------------	-----------------------------------

Reporting group description:

Subjects were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kilogram (kg) body weight with a dose escalation up to a maximum dose of 500 mg. Subjects who had weight  $\geq 40$  kg received a famciclovir dose of 500 mg.

Reporting group title	Famciclovir: Multiple doses (Part B)
-----------------------	--------------------------------------

Reporting group description:

Subjects were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy twice daily (b.i.d.) with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight  $\geq 40$  kg received a famciclovir dose of 500 mg.

Reporting group values	Famciclovir: Single dose (Part A)	Famciclovir: Multiple doses (Part B)	Total
Number of subjects	27	47	74
Age categorical Units: Subjects			
Infants and toddlers (28 days-23 months)	4	13	17
Children (2-11 years)	21	32	53
Adolescents (12-17 years)	2	2	4
Age continuous Units: years			
arithmetic mean	6	5	
standard deviation	$\pm 4.61$	$\pm 3.74$	-
Gender categorical Units: Subjects			
Female	17	24	41
Male	10	23	33

### Subject analysis sets

Subject analysis set title	Subjects aged: 1 to <2 years (Part A)
----------------------------	---------------------------------------

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

All subjects aged between 1 to <2 years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.

Subject analysis set title	Subjects aged: 2 to <6 years (Part A)
----------------------------	---------------------------------------

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

All subjects aged between 2 to <6 years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.

Subject analysis set title	Subjects aged: 6 to $\leq 12$ years (Part A)
----------------------------	--

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

All subjects aged between 6 to <12 years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.

Subject analysis set title	Subjects aged: 13 to $\leq 18$ years (Part A)
----------------------------	---

Subject analysis set type	Sub-group analysis
---------------------------	--------------------

Subject analysis set description:

All subjects aged between 13 to <18 years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.

Subject analysis set title	Subjects aged: 1 to <2 years (Part B)
Subject analysis set type	Sub-group analysis

Subject analysis set description:

All subjects aged between 1 to <2 years were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight  $\geq$  40 kg received a famciclovir dose of 500 mg.

Subject analysis set title	Subjects aged: 2 to <6 years (Part B)
Subject analysis set type	Sub-group analysis

Subject analysis set description:

All subjects aged between 2 to <6 years were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight  $\geq$  40 kg received a famciclovir dose of 500 mg.

Subject analysis set title	Subjects aged: 6 to $\leq$ 12 years (Part B)
Subject analysis set type	Sub-group analysis

Subject analysis set description:

All subjects aged between 6 to <12 years were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight  $\geq$  40 kg received a famciclovir dose of 500 mg.

Reporting group values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to $\leq$ 12 years (Part A)
Number of subjects	4	13	8
Age categorical Units: Subjects			
Infants and toddlers (28 days-23 months)			
Children (2-11 years)			
Adolescents (12-17 years)			
Age continuous Units: years			
arithmetic mean	1	3.5	10.3
standard deviation	$\pm$ 0	$\pm$ 1.2	$\pm$ 0.89
Gender categorical Units: Subjects			
Female	1	11	4
Male	3	2	4

Reporting group values	Subjects aged: 13 to $\leq$ 18 years (Part A)	Subjects aged: 1 to <2 years (Part B)	Subjects aged: 2 to <6 years (Part B)
Number of subjects	2	13	16
Age categorical Units: Subjects			
Infants and toddlers (28 days-23 months)			
Children (2-11 years)			
Adolescents (12-17 years)			

Age continuous Units: years arithmetic mean standard deviation	16 ± 1.41	1 ± 0	3.5 ± 1.03
Gender categorical Units: Subjects			
Female	1	7	10
Male	1	6	6

<b>Reporting group values</b>	Subjects aged: 6 to <=12 years (Part B)		
Number of subjects	18		
Age categorical Units: Subjects			
Infants and toddlers (28 days-23 months) Children (2-11 years) Adolescents (12-17 years)			
Age continuous Units: years arithmetic mean standard deviation	9.1 ± 2.14		
Gender categorical Units: Subjects			
Female	7		
Male	11		



## End points

### End points reporting groups

Reporting group title	Famciclovir: Single dose (Part A)
Reporting group description: Subjects were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kilogram (kg) body weight with a dose escalation up to a maximum dose of 500 mg. Subjects who had weight $\geq 40$ kg received a famciclovir dose of 500 mg.	
Reporting group title	Famciclovir: Multiple doses (Part B)
Reporting group description: Subjects were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy twice daily (b.i.d.) with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight $\geq 40$ kg received a famciclovir dose of 500 mg.	
Subject analysis set title	Subjects aged: 1 to $<2$ years (Part A)
Subject analysis set type	Sub-group analysis
Subject analysis set description: All subjects aged between 1 to $<2$ years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.	
Subject analysis set title	Subjects aged: 2 to $<6$ years (Part A)
Subject analysis set type	Sub-group analysis
Subject analysis set description: All subjects aged between 2 to $<6$ years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.	
Subject analysis set title	Subjects aged: 6 to $\leq 12$ years (Part A)
Subject analysis set type	Sub-group analysis
Subject analysis set description: All subjects aged between 6 to $<12$ years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.	
Subject analysis set title	Subjects aged: 13 to $\leq 18$ years (Part A)
Subject analysis set type	Sub-group analysis
Subject analysis set description: All subjects aged between 13 to $<18$ years were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg weight with a maximum dose of 500 mg.	
Subject analysis set title	Subjects aged: 1 to $<2$ years (Part B)
Subject analysis set type	Sub-group analysis
Subject analysis set description: All subjects aged between 1 to $<2$ years were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight $\geq 40$ kg received a famciclovir dose of 500 mg.	
Subject analysis set title	Subjects aged: 2 to $<6$ years (Part B)
Subject analysis set type	Sub-group analysis
Subject analysis set description: All subjects aged between 2 to $<6$ years were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight $\geq 40$ kg received a famciclovir dose of 500 mg.	
Subject analysis set title	Subjects aged: 6 to $\leq 12$ years (Part B)
Subject analysis set type	Sub-group analysis
Subject analysis set description: All subjects aged between 6 to $<12$ years were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight $\geq 40$ kg received a famciclovir dose of 500 mg.	

up to a maximum of 500 mg b.i.d. Subjects who had weight  $\geq$  40 kg received a famciclovir dose of 500 mg.

### Primary: Time to reach maximum plasma concentration (Tmax) of famciclovir during Part A

End point title	Time to reach maximum plasma concentration (Tmax) of famciclovir during Part A <sup>[1]</sup>
-----------------	---

End point description:

Tmax was defined as the time to reach maximum plasma concentration. Penciclovir (metabolite of famciclovir) plasma concentrations were determined by using a validated liquid chromatography/tandem mass spectrometry (LC/MS/MS) method. The limit of quantification was 0.15 microgram ( $\mu$ g)/ millilitre (mL) for both compounds. The analysis was performed in pharmacokinetic analysis set (PAS) population, defined as all subjects who received at least one scheduled dose of famciclovir and provided all primary PK parameters in at least one treatment period.

End point type	Primary
----------------	---------

End point timeframe:

At pre-dose, 1, 2, 3, 4, and 5 hours post treatment administration

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to $\leq$ 12 years (Part A)	Subjects aged: 13 to $\leq$ 18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: hours				
median (full range (min-max))	1.21 (1 to 1.5)	1.07 (1 to 4.03)	1 (1 to 2.07)	1.47 (0.97 to 1.97)

### Statistical analyses

No statistical analyses for this end point

### Primary: Maximum plasma concentration (Cmax) of famciclovir during Part A

End point title	Maximum plasma concentration (Cmax) of famciclovir during Part A <sup>[2]</sup>
-----------------	---

End point description:

Cmax was defined as the maximum plasma concentration. Penciclovir (metabolite of famciclovir) plasma concentrations were determined by using a validated LC/MS/MS method. The limit of quantification was 0.15  $\mu$ g/mL for both compounds. The analysis was performed in PAS population.

End point type	Primary
----------------	---------

End point timeframe:

At pre-dose, 1, 2, 3, 4, and 5 hours post treatment administration

Notes:

[2] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to <=12 years (Part A)	Subjects aged: 13 to <=18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: microgram( $\mu$ g)/milliliter(mL)				
arithmetic mean (full range (min-max))	2.84 (1.42 to 4.47)	2.44 (0.42 to 3.81)	2.82 (1.52 to 3.79)	1.89 (1.06 to 2.72)

## Statistical analyses

No statistical analyses for this end point

### Primary: Area under the plasma concentration time curve from time zero to the time point of the last measurable concentration (AUC 0-tlast) of famciclovir during Part A

End point title	Area under the plasma concentration time curve from time zero to the time point of the last measurable concentration (AUC 0-tlast) of famciclovir during Part A <sup>[3]</sup>
-----------------	--

End point description:

AUC 0-tlast was defined as the area under the plasma concentration time curve from time zero to the time point of the last measurable concentration. Penciclovir (metabolite of famciclovir) plasma concentrations were determined by using a validated LC/MS/MS method. The limit of quantification was 0.15  $\mu$ g/mL for both compounds. The analysis was performed in PAS population.

End point type	Primary
----------------	---------

End point timeframe:

At pre-dose, 1, 2, 3, 4, and 5 hours post treatment administration

Notes:

[3] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to <=12 years (Part A)	Subjects aged: 13 to <=18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: (microgram( $\mu$ g)/milliliter(mL))*hour(h))				
arithmetic mean (full range (min-max))	5.73 (3.02 to 8.45)	5.71 (1.63 to 8.17)	6.98 (4.72 to 8.66)	4.81 (3.57 to 6.06)

## Statistical analyses

No statistical analyses for this end point

### Primary: Area under the plasma concentration time curve from time zero to infinity (AUC 0-infinity) of famciclovir during Part A

End point title	Area under the plasma concentration time curve from time zero to infinity (AUC 0-infinity) of famciclovir during Part A <sup>[4]</sup>
-----------------	--

End point description:

AUC 0-infinity was defined as the area under the plasma concentration time curve from time zero to infinity. Penciclovir (metabolite of famciclovir) plasma concentrations were determined by using a

validated LC/MS/MS method. The limit of quantification was 0.15 µg/mL for both compounds. The analysis was performed in PAS population.

End point type	Primary
----------------	---------

End point timeframe:

At pre-dose, 1, 2, 3, 4, and 5 hours post treatment administration

Notes:

[4] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to <=12 years (Part A)	Subjects aged: 13 to <=18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: µg/mL*h				
arithmetic mean (full range (min-max))	6.17 (3.43 to 8.99)	6.85 (3.19 to 9.12)	8.15 (6.49 to 9.71)	5.93 (4.84 to 7.01)

## Statistical analyses

No statistical analyses for this end point

## Primary: Terminal elimination half-life (T<sub>1/2</sub>) of famciclovir during Part A

End point title	Terminal elimination half-life (T <sub>1/2</sub> ) of famciclovir during Part
-----------------	---

End point description:

T<sub>1/2</sub> was defined as the terminal elimination half-life. Penciclovir (metabolite of famciclovir) plasma concentrations were determined by using a validated LC/MS/MS method. The limit of quantification was 0.15 µg/mL for both compounds. The analysis was performed in PAS population.

End point type	Primary
----------------	---------

End point timeframe:

At pre-dose, 1, 2, 3, 4, and 5 hours post treatment administration

Notes:

[5] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to <=12 years (Part A)	Subjects aged: 13 to <=18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: Hours				
arithmetic mean (full range (min-max))	1.09 (1.01 to 1.18)	1.36 (1.1 to 1.7)	1.6 (1.3 to 2.11)	1.86 (1.6 to 2.12)

## Statistical analyses

No statistical analyses for this end point

**Primary: Apparent oral clearance (if AUC 0-infinity is available) (CL/F) of famciclovir during Part A**

End point title	Apparent oral clearance (if AUC 0-infinity is available) (CL/F) of famciclovir during Part A <sup>[6]</sup>
-----------------	---

End point description:

CL/F was defined as the apparent oral clearance. Penciclovir (metabolite of famciclovir) plasma concentrations were determined by using a validated LC/MS/MS method. The limit of quantification was 0.15 µg/mL for both compounds. The analysis was performed in PAS population.

End point type	Primary
----------------	---------

End point timeframe:

At pre-dose, 1, 2, 3, 4, and 5 hours post treatment administration

Notes:

[6] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to ≤12 years (Part A)	Subjects aged: 13 to ≤18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: Lites(L)/Hour(h)				
arithmetic mean (full range (min-max))	20.8 (11 to 28.8)	25.1 (18.1 to 33.3)	43.7 (32.4 to 60.8)	68.8 (56.2 to 81.5)

**Statistical analyses**

No statistical analyses for this end point

**Primary: Number of subjects with Adverse Events (AEs), Serious Adverse Events(SAEs), AE leading to discontinuation and who died**

End point title	Number of subjects with Adverse Events (AEs), Serious Adverse Events(SAEs), AE leading to discontinuation and who died <sup>[7]</sup>
-----------------	---

End point description:

AEs are defined as any unfavorable and unintended diagnosis, symptom, sign (including an abnormal laboratory finding), syndrome or disease which either occurs during study, having been absent at baseline, or, if present at baseline, appears to worsen. Serious adverse events are any untoward medical occurrences that result in death, are life threatening, require (or prolong) hospitalization, cause persistent or significant disability/incapacity, result in congenital anomalies or birth defects, or are other conditions which in judgment of investigators represent significant hazards. The analysis was performed on safety set population defined as subjects who received any dose (including partial dose) of study medication and had at least one post-baseline safety or acceptability assessment.

End point type	Primary
----------------	---------

End point timeframe:

From Day 1 up to Day 15

Notes:

[7] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to <=12 years (Part A)	Subjects aged: 13 to <=18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: Subjects				
AEs	0	2	1	2
SAEs	0	0	0	0
Deaths	0	0	0	0
AEs leading to discontinuation	0	0	0	0
AE requiring concomitant medication	0	0	0	0

End point values	Subjects aged: 1 to <2 years (Part B)	Subjects aged: 2 to <6 years (Part B)	Subjects aged: 6 to <=12 years (Part B)	
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	
Number of subjects analysed	13	16	18	
Units: Subjects				
AEs	6	8	12	
SAEs	0	0	0	
Deaths	0	0	0	
AEs leading to discontinuation	0	0	0	
AE requiring concomitant medication	4	1	3	

## Statistical analyses

No statistical analyses for this end point

## Primary: Number of subjects with clinically significant laboratory abnormalities

End point title	Number of subjects with clinically significant laboratory abnormalities <sup>[8]</sup>
-----------------	--

End point description:

Subjects with laboratory values outside the defined normal range were graded as clinically significant laboratory abnormalities. Laboratory values were assessed according to the National Cancer Institute-Common terminology criteria for Adverse Events (NCI-CTCAE). Hematology and clinical chemistry were performed. The analysis was performed on the safety set.

End point type	Primary
----------------	---------

End point timeframe:

From Day 1 to Day 15

Notes:

[8] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: only summary stats available

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to <=12 years (Part A)	Subjects aged: 13 to <=18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: Subjects				
Hematology	0	0	0	0
Clinical chemistry	0	0	0	0

End point values	Subjects aged: 1 to <2 years (Part B)	Subjects aged: 2 to <6 years (Part B)	Subjects aged: 6 to <=12 years (Part B)	
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	
Number of subjects analysed	13	16	18	
Units: Subjects				
Hematology	0	0	0	
Clinical chemistry	0	0	0	

## Statistical analyses

No statistical analyses for this end point

## Secondary: Assessment of acceptability of pediatric formulation during Part A

End point title	Assessment of acceptability of pediatric formulation during Part A
-----------------	--

End point description:

Assessment of acceptability was done using a modified, 5-point facial hedonic scale, subjects or caregivers were asked to complete an assessment of study medication. Subjects marked the scale with their response of choice. The scale represented a balance of choices from 'I did not like it' to 'I like it very much' with a mid-point of neither like nor dislike, with qualitative evaluation. The analysis was performed on the safety set.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1

End point values	Subjects aged: 1 to <2 years (Part A)	Subjects aged: 2 to <6 years (Part A)	Subjects aged: 6 to <=12 years (Part A)	Subjects aged: 13 to <=18 years (Part A)
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	Subject analysis set
Number of subjects analysed	4	13	8	2
Units: Subjects				
Bitter	1	3	4	1
Sweet	2	5	4	0
Other	1	5	1	1
Very badly / Unacceptable	1	0	0	0
Badly but accepted	0	4	0	2
Neither good nor bad	1	1	4	0

Well accepted	0	5	2	0
Very well accepted	2	3	2	0

## Statistical analyses

No statistical analyses for this end point

## Secondary: Assessment of acceptability of pediatric formulation during Part B

End point title	Assessment of acceptability of pediatric formulation during Part B
-----------------	--

End point description:

Assessment of acceptability was done using a modified, 5-point facial hedonic scale, subjects or caregivers were asked to complete an assessment of study medication. Subjects marked the scale with their response of choice. The scale represented a balance of choices from 'I did not like it' to 'I like it very much' with a mid-point of neither like nor dislike, with qualitative evaluation. The analysis was performed on the safety set. Here, 'clinic' and 'home' signifies post-first dose in clinic and home respectively and day 8 have been reported for post-last dose at home.

End point type	Secondary
----------------	-----------

End point timeframe:

From Day 1 to Day 8

End point values	Subjects aged: 1 to <2 years (Part B)	Subjects aged: 2 to <6 years (Part B)	Subjects aged: 6 to <=12 years (Part B)	
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	
Number of subjects analysed	13	16	18	
Units: Subjects				
Bitter	3	6	9	
Sweet	3	5	2	
Other	7	5	7	
Very badly/unacceptable:Day 1(clinic)	0	2	0	
Badly but accepted: Day 1(clinic)	9	4	7	
Neither good nor bad: Day 1(clinic)	2	1	2	
Well accepted: Day 1(clinic)	1	4	5	
Very well accepted: Day 1(clinic)	1	5	4	
Very badly/unacceptable:Day 1(home)	1	0	0	
Badly but accepted: Day 1(home)	7	5	4	
Neither good nor bad: Day 1(home)	2	2	4	
Well accepted: Day 1(home)	2	5	5	
Very well accepted: Day 1(home)	1	3	5	
Very badly/unacceptable: EoS/early discontinuation	0	1	0	
Badly but accepted: EoS/early discontinuation	7	4	3	
Neither good nor bad: EoS/early discontinuation	3	3	3	
Well accepted: EoS/early discontinuation	1	3	5	



Very well accepted: EoS/early discontinuation	2	4	6	
---	---	---	---	--

## Statistical analyses

---

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Serious Adverse Events are monitored from date of First Subject First Visit (FSFV) until Last Subject Last Visit (LSLV). All other adverse events are monitored from First Subject First Treatment until LSLV

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	10.1
--------------------	------

### Reporting groups

Reporting group title	Famciclovir: Single dose (Part A)
-----------------------	-----------------------------------

Reporting group description:

Subjects were orally administered with a single body weight stratified dose of famciclovir of 12.5 mg/kg body weight with a dose escalation up to a maximum dose of 500 mg. Subjects who had weight  $\geq 40$  kg received a famciclovir dose of 500 mg.

Reporting group title	Famciclovir: Multiple doses (Part B)
-----------------------	--------------------------------------

Reporting group description:

Subjects were orally administered with famciclovir sprinkle capsules alone or with concomitant oral anti-herpes therapy b.i.d. with a dose separation of 12 hours for a period of 7 days. Subjects were administered with a body weight stratified daily dose in 8-step from 150 mg b.i.d. up to a maximum of 500 mg b.i.d. Subjects who had weight  $\geq 40$  kg received a famciclovir dose of 500 mg.

Serious adverse events	Famciclovir: Single dose (Part A)	Famciclovir: Multiple doses (Part B)	
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 27 (0.00%)	0 / 47 (0.00%)	
number of deaths (all causes)	0	0	
number of deaths resulting from adverse events	0	0	

Frequency threshold for reporting non-serious adverse events: 2 %

Non-serious adverse events	Famciclovir: Single dose (Part A)	Famciclovir: Multiple doses (Part B)	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	5 / 27 (18.52%)	26 / 47 (55.32%)	
Investigations			
Blood uric acid increased			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Injury, poisoning and procedural complications			

Arthropod bite subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Contusion subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Skin laceration subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Vascular disorders Haematoma subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Flushing subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Nervous system disorders Clumsiness subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Dizziness subjects affected / exposed occurrences (all)	1 / 27 (3.70%) 1	0 / 47 (0.00%) 0	
Headache subjects affected / exposed occurrences (all)	2 / 27 (7.41%) 2	4 / 47 (8.51%) 4	
Psychomotor hyperactivity subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Somnolence subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 2	
General disorders and administration site conditions Chest pain subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	

Irritability subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Pyrexia subjects affected / exposed occurrences (all)	1 / 27 (3.70%) 1	2 / 47 (4.26%) 2	
Vessel puncture site haematoma subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Gastrointestinal disorders			
Abdominal pain upper subjects affected / exposed occurrences (all)	2 / 27 (7.41%) 2	2 / 47 (4.26%) 3	
Diarrhoea subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	4 / 47 (8.51%) 4	
Flatulence subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Gingival bleeding subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Nausea subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	3 / 47 (6.38%) 3	
Vomiting subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	5 / 47 (10.64%) 6	
Respiratory, thoracic and mediastinal disorders			
Cough subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	3 / 47 (6.38%) 3	
Dysphonia subjects affected / exposed occurrences (all)	0 / 27 (0.00%) 0	1 / 47 (2.13%) 1	
Nasal congestion			

subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Pharyngolaryngeal pain			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Rhinorrhoea			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Skin and subcutaneous tissue disorders			
Blister			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Dry skin			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Pruritus			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Rash			
subjects affected / exposed	1 / 27 (3.70%)	0 / 47 (0.00%)	
occurrences (all)	1	0	
Infections and infestations			
Furuncle			
subjects affected / exposed	1 / 27 (3.70%)	0 / 47 (0.00%)	
occurrences (all)	1	0	
Conjunctivitis viral			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Gastroenteritis			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Otitis media			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	
Metabolism and nutrition disorders			

Hypokalaemia			
subjects affected / exposed	0 / 27 (0.00%)	1 / 47 (2.13%)	
occurrences (all)	0	1	

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
07 July 2006	Adjusted the dosing scheme in Part A and Part B according to the interim results from Part A of both studies (i.e. CFAM810B2303 and CFAM810B2304) and discussions with the FDA. It also added a criterion to exclude subjects who weighed <9 kg.
17 November 2006	Requirement for PK and multiple-dose safety assessments in a cohort of subjects, 12 to 18 years of age and the upper age limit of the pediatric subjects to be studied from 12 to 18 years old was amended. The amendment created a new cohort for these adolescent subjects.
18 June 2007	Cancelled the requirement for inclusion of adolescent subjects in the study by deleting the new cohort (which had been designated as cohort 4 to include subjects 12 to 18 years of age), amending the upper age limit of the pediatric subjects to be studied from 18 back to 12 years, and deleting the requirement for single-dose PK for 3 to 4 subjects in cohort 4. The amendment also readjusted the number of subjects per cohort in Part B to compensate for the removal of cohort 4.

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported