



## Clinical trial results:

**A Phase 1/2 multicenter, open-label study to determine the recommended dose and regimen of Durvalumab (MEDI4736) in combination with Lenalidomide (LEN) with and without Dexamethasone (DEX) in subjects with newly diagnosed multiple myeloma (NDMM)**

### Summary

EudraCT number	2015-004831-11
Trial protocol	DE ES DK FI NL
Global end of trial date	06 September 2022

### Results information

Result version number	v1
This version publication date	17 September 2023
First version publication date	17 September 2023

### Trial information

#### Trial identification

Sponsor protocol code	MEDI4736-MM-002
-----------------------	-----------------

#### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	-
WHO universal trial number (UTN)	-

Notes:

### Sponsors

Sponsor organisation name	Bristol-Myers Squibb
Sponsor organisation address	Chaussée de la Hulpe 185, Brussels, Belgium, 1170
Public contact	EU Study Start-Up Unit, Bristol-Myers Squibb International Corporation, Clinical.Trials@bms.com
Scientific contact	Bristol-Myers Squibb Study Director, Bristol-Myers Squibb, Clinical.Trials@bms.com

Notes:

### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	16 March 2021
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	06 September 2022
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

Determine the recommended dose and regimen of durvalumab in combination with LEN with and without DEX in subjects with newly diagnosed multiple myeloma (NDMM).

Protection of trial subjects:

The study was in compliance with the ethical principles derived from the Declaration of Helsinki and in compliance with all International Conference on Harmonization Good Clinical Practice Guidelines. All the local regulatory requirements pertinent to safety of trial participants were followed.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	25 April 2016
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Canada: 13
Country: Number of subjects enrolled	Denmark: 7
Country: Number of subjects enrolled	Finland: 5
Country: Number of subjects enrolled	Germany: 1
Country: Number of subjects enrolled	Italy: 7
Country: Number of subjects enrolled	Spain: 15
Country: Number of subjects enrolled	Netherlands: 4
Country: Number of subjects enrolled	United States: 4
Worldwide total number of subjects	56
EEA total number of subjects	39

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0

Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	17
From 65 to 84 years	39
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details: -

### Pre-assignment

Screening details:

56 subjects treated

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Not blinded

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Cohort A: High risk, TNE

Arm description:

High risk, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$  75 years old) or 20 mg/day ( $>$  75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle.

Arm type	Experimental
Investigational medicinal product name	Durvalumab
Investigational medicinal product code	MEDI4736
Other name	
Pharmaceutical forms	Solution for infusion
Routes of administration	Intravenous use

Dosage and administration details:

1500 mg on day 1 of each 28-day cycle

Investigational medicinal product name	Dexamethasone
Investigational medicinal product code	DEX
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

40 mg/day ( $\leq$  75 years old) or 20 mg/day ( $>$  75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle.

Investigational medicinal product name	Lenalidomide
Investigational medicinal product code	LEN
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

25 mg/day on days 1 to 21 of each 28-day treatment cycle

<b>Arm title</b>	Cohort B: $\geq$ 65 years old, TNE
------------------	------------------------------------

Arm description:

$\geq$  65 years old, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were not high risk were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$  75 years old) or 20 mg/day ( $>$  75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle, up to 12 cycles.

Arm type	Experimental
Investigational medicinal product name	Durvalumab
Investigational medicinal product code	MEDI4736
Other name	
Pharmaceutical forms	Solution for infusion
Routes of administration	Intravenous use

Dosage and administration details:

1500 mg on day 1 of each 28-day cycle

Investigational medicinal product name	Dexamethasone
Investigational medicinal product code	DEX
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

40 mg/day ( $\leq$  75 years old) or 20 mg/day ( $>$  75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle.

Investigational medicinal product name	Lenalidomide
Investigational medicinal product code	LEN
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

25 mg/day on days 1 to 21 of each 28-day treatment cycle

<b>Arm title</b>	Cohort C: High risk, Post-transplant
------------------	--------------------------------------

Arm description:

High risk, post-transplant NDMM participants were administered the following as maintenance therapy: • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 10 mg/day on Days 1 to 21 of each 28-day treatment cycle.

Arm type	Experimental
Investigational medicinal product name	Lenalidomide
Investigational medicinal product code	LEN
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

10 mg/day on days 1 to 21 of each 28-day treatment cycle

Investigational medicinal product name	Durvalumab
Investigational medicinal product code	MEDI4736
Other name	
Pharmaceutical forms	Solution for infusion
Routes of administration	Intravenous use

Dosage and administration details:

1500 mg on day 1 of each 28-day cycle

<b>Number of subjects in period 1</b>	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant
Started	25	10	21
Completed	0	0	0
Not completed	25	10	21
Consent withdrawn by subject	2	-	-
Adverse event, non-fatal	2	-	1
Progressive Disease	2	1	-
Full Clinical Hold by FDA	19	9	20

## Baseline characteristics

### Reporting groups

Reporting group title	Cohort A: High risk, TNE
Reporting group description:	
High risk, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$ 75 years old) or 20 mg/day ( $>$ 75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle.	
Reporting group title	Cohort B: $\geq$ 65 years old, TNE
Reporting group description:	
$\geq$ 65 years old, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were not high risk were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$ 75 years old) or 20 mg/day ( $>$ 75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle, up to 12 cycles.	
Reporting group title	Cohort C: High risk, Post-transplant
Reporting group description:	
High risk, post-transplant NDMM participants were administered the following as maintenance therapy: • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 10 mg/day on Days 1 to 21 of each 28-day treatment cycle.	

Reporting group values	Cohort A: High risk, TNE	Cohort B: $\geq$ 65 years old, TNE	Cohort C: High risk, Post-transplant
Number of subjects	25	10	21
Age Categorical			
Units: participants			
$\leq$ 75 years	12	7	20
$>$ 75 years	13	3	1
Age Continuous			
Units: years			
arithmetic mean	75.4	72.5	59.6
standard deviation	$\pm$ 4.17	$\pm$ 5.25	$\pm$ 7.52
Sex: Female, Male			
Units: participants			
Female	15	0	4
Male	10	10	17
Race (NIH/OMB)			
Units: Subjects			
American Indian or Alaska Native	0	0	0
Asian	1	1	0
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	1	0	0
White	21	9	17
More than one race	0	0	0
Unknown or Not Reported	2	0	4

Reporting group values	Total		
Number of subjects	56		

Age Categorical			
Units: participants			
<=75 years	39		
>75 years	17		
Age Continuous			
Units: years			
arithmetic mean			
standard deviation	-		
Sex: Female, Male			
Units: participants			
Female	19		
Male	37		
Race (NIH/OMB)			
Units: Subjects			
American Indian or Alaska Native	0		
Asian	2		
Native Hawaiian or Other Pacific Islander	0		
Black or African American	1		
White	47		
More than one race	0		
Unknown or Not Reported	6		



## End points

### End points reporting groups

Reporting group title	Cohort A: High risk, TNE
Reporting group description: High risk, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$ 75 years old) or 20 mg/day ( $>$ 75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle.	
Reporting group title	Cohort B: $\geq$ 65 years old, TNE
Reporting group description: $\geq$ 65 years old, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were not high risk were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$ 75 years old) or 20 mg/day ( $>$ 75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle, up to 12 cycles.	
Reporting group title	Cohort C: High risk, Post-transplant
Reporting group description: High risk, post-transplant NDMM participants were administered the following as maintenance therapy: • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 10 mg/day on Days 1 to 21 of each 28-day treatment cycle.	

### Primary: Participants with Dose-Limiting Toxicities (DLTs) During the Dose-Determining Timeframe (Day 1 – Day 28)

End point title	Participants with Dose-Limiting Toxicities (DLTs) During the Dose-Determining Timeframe (Day 1 – Day 28) <sup>[1]</sup>
End point description: A Dose Review Team (DRT) evaluated DLTs and, if applicable, other data to determine the recommended dose (RD) of durvalumab to use in the Expansion Period. The DRT included sponsor personnel, investigators and outside consultants. A DLT was defined as: a. Grade 4 neutropenia for $\geq$ 5 days. b. Grade 3 neutropenia associated with fever ( $\geq$ 38.5°C / 101.3°F) of any duration. c. Grade 4 thrombocytopenia or Grade 3 thrombocytopenia with bleeding, or platelets transfusion. d. Grade 4 hematologic toxicity that does not resolve to baseline level $\leq$ 72 hours. e. Grade 4 anemia, unexplained by underlying disease. f. Any nonhematologic toxicity Grade $\geq$ 3 except for alopecia and nausea. g. Treatment interruption $\geq$ 2 weeks due to AE. If $\leq$ 1 of the 6 initial participants in each cohort experience a DLT during cycle 1, the RD was durvalumab 1500 mg; If $\geq$ 2 of the 6 initial participants in any cohort experience a DLT during cycle 1, the maximum tolerated dose (MTD) was exceeded and the	
End point type	Primary
End point timeframe: First treatment cycle: Day 1 to Day 28	

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only summary analyses planned for this endpoint.

End point values	Cohort A: High risk, TNE	Cohort B: $\geq$ 65 years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	6	7	6	
Units: Count of participants	1	1	0	

## Statistical analyses

No statistical analyses for this end point

## Secondary: Participants with Treatment Emergent Adverse Events (TEAE)

End point title	Participants with Treatment Emergent Adverse Events (TEAE)
-----------------	--

End point description:

An AE is any noxious, unintended, or untoward medical occurrence that may appear or worsen during the course of a study. A TEAE includes AEs between the first dose date of either study drug and 90 days after the last dose of study drug. A serious AE is any AE occurring at any dose that: • Results in death; • Is life-threatening; • Requires or prolongs existing inpatient hospitalization; • Results in persistent or significant disability/incapacity; • Is a congenital anomaly/birth defect; • Constitutes an important medical event. The Investigator assessed the relationship of each AE to study drug and graded the severity according to the National Cancer Institute (NCI) Common Terminology Criteria for Adverse Events (CTCAE, Version 4.03): - Grade 1 = Mild - Grade 2 = Moderate (some limitation in activity; no/minimal medical intervention required) - Grade 3 = Severe (limitation in activity; medical intervention required) - Grade 4 = Life-threatening - Grade 5 = Death 99999=NA

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 up to Week 84 (the longer of 90 days after discontinuing treatment with DURVA, or 28 days after the last dose of LEN or dex)

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	10	21	
Units: Count of participants				
$\geq 1$ TEAE	24	9	18	
$\geq 1$ treatment-related TEAE	22	8	17	
$\geq 1$ related to DURVA	16	4	15	
$\geq 1$ related to LEN	22	8	15	
$\geq 1$ related to dex	16	6	99999	
$\geq 1$ TEAE of severity grade 3 or 4	19	8	7	
$\geq 1$ severity 3/4, related to study drug	13	8	6	
$\geq 1$ severity 3/4, related to DURVA	7	4	4	
$\geq 1$ severity 3/4, related to LEN	11	8	4	
$\geq 1$ severity 3/4, related to dex	5	2	99999	
$\geq 1$ Grade 5 TEAE (Death)	1	0	1	
$\geq 1$ Grade 5 TEAE related to study drug	0	0	1	
$\geq 1$ Grade 5 TEAE related to DURVA	0	0	1	
$\geq 1$ Grade 5 TEAE related to LEN	0	0	0	
$\geq 1$ Grade 5 TEAE related to dex	0	0	99999	
$\geq 1$ Serious TEAE	12	6	4	
$\geq 1$ Serious TEAE related to study drug	8	4	4	
$\geq 1$ Serious TEAE related to DURVA	5	1	4	

>=1 Serious TEAE related to LEN	6	3	2	
>=1 Serious TEAE related to dex	5	1	99999	
>=1 TEAE leading to discontinuation of study drug	3	0	1	
>= TEAE leading to discontinuation of DURVA	2	0	1	
>= TEAE leading to discontinuation of LEN	3	0	0	
>= TEAE leading to discontinuation of dex	2	0	99999	
>=1 leading to delay/reduction/interruption drug	14	7	6	
>=1 leading to dose delay of DURVA	10	3	4	
>=1 leading to infusion interruption of DURVA	0	0	1	
>=1 leading to dose reduction of LEN	2	2	3	
>=1 leading to interruption of LEN	13	5	5	
>=1 leading to dose reduction of dex	1	3	99999	
>=1 leading to interruption of dex	9	3	99999	

## Statistical analyses

No statistical analyses for this end point

## Secondary: Overall Response Rate (ORR) for Cohorts A and B: Percentage of Participants Who Achieved a Partial Response or Better According to the International Myeloma Working Group (IMWG) Uniform Response Criteria

End point title	Overall Response Rate (ORR) for Cohorts A and B: Percentage of Participants Who Achieved a Partial Response or Better According to the International Myeloma Working Group (IMWG) Uniform Response Criteria
-----------------	---

End point description:

Tumor response, including progressive disease, was assessed by the investigators and captured the best assessment of response during the treatment period. ORR was defined as partial response (PR) or better which includes PR, very good partial response (VGPR), complete response (CR), or stringent complete response (sCR). A PR required  $\geq 50\%$  reduction of serum M-Protein and reduction in 24-hour urinary M-protein by  $\geq 90\%$  or to  $< 200$  mg per 24 hours. If present at baseline, a  $\geq 50\%$  reduction in the size of soft tissue plasmacytomas was also required. sCR required - a negative immunofixation of serum and urine and - disappearance of any soft tissue plasmacytomas and -  $\leq 5\%$  plasma cells in bone marrow and normal free light-chain (FLC) ratio and - absence of clonal plasma cells by immunohistochemistry or 2- to 4-color flow cytometry.

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 of each cycle starting with Cycle 2 up to Cycle 17 plus one week for the end of treatment visit (Day 29 up to Week 73)

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	24	10	0 <sup>[2]</sup>	
Units: percentage of participants				
number (confidence interval 80%)	66.7 (51.6 to 79.5)	50.0 (26.7 to 73.3)	( to )	

Notes:

[2] - 0 subjects analyzed.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Response Improvement Rate (RIR) for Cohort C: Percentage of Participants Achieving a Response Improved from Cycle 1 Day 1 as Assessed by the Investigators Using the International Myeloma Working Group (IMWG) Uniform Response Criteria

End point title	Response Improvement Rate (RIR) for Cohort C: Percentage of Participants Achieving a Response Improved from Cycle 1 Day 1 as Assessed by the Investigators Using the International Myeloma Working Group (IMWG) Uniform Response Criteria
-----------------	---

End point description:

Response Improvement Rate is defined as the percentage of participants who achieved a response from treatment as compared to the pre-autologous stem cell transplantation [ASCT] diseases measurement used as baseline for response assessment. IMWG response categories could be stable disease (SD), partial response (PR), very good partial response (VGPR), complete response (CR), or stringent complete response (sCR), as long as it represented an improvement compared to prior to transplant. 99999=NA

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Cycle 1 Day 1); Treatment: Day 1 of each cycle starting with Cycle 2 up to Cycle 15 plus one week for the end of treatment visit (Day 29 up to Week 61)

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	0 <sup>[3]</sup>	0 <sup>[4]</sup>	21	
Units: percentage of participants				
number (confidence interval 80%)	( to )	( to )	0 (-99999 to 99999)	

Notes:

[3] - 0 subjects analyzed

[4] - 0 subjects analyzed

### Statistical analyses

No statistical analyses for this end point

### Secondary: Time to Response (for Cohorts A and B)

End point title	Time to Response (for Cohorts A and B)
-----------------	--

End point description:

Time to response (for responders only, per IMWG Uniform Response Criteria) is calculated as the time from the first date of dosing of study medication to the first date of documented response (PR or better).

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 of each cycle starting with Cycle 2 up to Cycle 17 plus one week for the end of treatment visit (Day 29 up to Week 73)

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	16	5	0 <sup>[5]</sup>	
Units: weeks				
median (full range (min-max))	4.20 (3.9 to 23.1)	4.10 (4.0 to 13.0)	( to )	

Notes:

[5] - 0 subjects analyzed

## Statistical analyses

No statistical analyses for this end point

## Secondary: Kaplan-Meier Estimates for Duration of Response (for Cohort A and B)

End point title	Kaplan-Meier Estimates for Duration of Response (for Cohort A and B)
-----------------	--

End point description:

Duration of response (for responders only) was defined as the time from earliest date of documented response (PR or better) to the earliest date of disease progression (DP) as determined by the investigator per IMWG Uniform Response criteria or death during study treatment, whichever occurred first. 99999=NA

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 of each cycle starting with Cycle 2 up to Cycle 17 plus one week for the end of treatment visit (Day 29 up to Week 73)

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	16	5	0 <sup>[6]</sup>	
Units: months				
median (confidence interval 80%)	10.3 (7.2 to 99999)	99999 (99999 to 99999)	( to )	

Notes:

[6] - 0 subjects analyzed

## Statistical analyses

**Secondary: Durvalumab (DURVA) Serum Pharmacokinetic (PK) Parameters in Cycle 1: Area Under the Concentration-time Curve from Time Zero to the Last Measured Time Point (AUC0-last)**

End point title	Durvalumab (DURVA) Serum Pharmacokinetic (PK) Parameters in Cycle 1: Area Under the Concentration-time Curve from Time Zero to the Last Measured Time Point (AUC0-last)
-----------------	---

## End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \times \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

## End point timeframe:

pre-infusion (-60 to -5 minutes prior to dose), end of infusion (EOI), 4 hours, 168 hours (Day 8), 336 hours (Day 15) and 504 hours (Day 22) after administration of DURVA on Day 1

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	23	10	20	
Units: day* $\mu$ g/L				
geometric mean (geometric coefficient of variation)	3535033.014 ( $\pm 39.8$ )	3582905.960 ( $\pm 21.0$ )	4026520.655 ( $\pm 39.1$ )	

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Area Under the Concentration-time Curve from Time Zero to Infinity (AUC0-inf)**

End point title	Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Area Under the Concentration-time Curve from Time Zero to Infinity (AUC0-inf)
-----------------	--

## End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \times \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

## End point timeframe:

pre-infusion (-60 to -5 minutes prior to dose), end of infusion (EOI), 4 hours, 168 hours (Day 8), 336 hours (Day 15) and 504 hours (Day 22) after administration of DURVA on Day 1

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	23	9	19	
Units: day* $\mu\text{g/L}$				
geometric mean (geometric coefficient of variation)	4944601.671 ( $\pm 60.9$ )	5532568.144 ( $\pm 62.5$ )	6531541.670 ( $\pm 36.4$ )	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Time to Maximum Observed Concentration (Tmax)

End point title	Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Time to Maximum Observed Concentration (Tmax)
-----------------	--

End point description:

Time to maximum observed concentration of Durvalumab (DURVA) after multiple doses on day 1 obtained from the observed concentration versus time data

End point type	Secondary
----------------	-----------

End point timeframe:

pre-infusion (-60 to -5 minutes prior to dose), end of infusion (EOI), 4 hours, 168 hours (Day 8), 336 hours (Day 15) and 504 hours (Day 22) after administration of DURVA on Day 1

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	23	10	20	
Units: day				
median (full range (min-max))	0.051 (0.04 to 0.22)	0.106 (0.04 to 0.22)	0.180 (0.04 to 0.22)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Maximum Observed Concentration (Cmax)

End point title	Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Maximum Observed Concentration (Cmax)
-----------------	--

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $\text{CV\%} = 100 \times \sqrt{\text{EXP}(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

pre-infusion (-60 to -5 minutes prior to dose), end of infusion (EOI), 4 hours, 168 hours (Day 8), 336 hours (Day 15) and 504 hours (Day 22) after administration of DURVA on Day 1

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	23	10	20	
Units: $\mu\text{g/L}$				
geometric mean (geometric coefficient of variation)	449280.231 ( $\pm 36.7$ )	452827.419 ( $\pm 25.5$ )	482602.748 ( $\pm 39.9$ )	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Clearance (CL)

End point title	Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Clearance (CL)
-----------------	---

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $\text{CV\%} = 100 \times \text{SQRT}(\text{EXP}(\sigma^2) - 1)$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

pre-infusion (-60 to -5 minutes prior to dose), end of infusion (EOI), 4 hours, 168 hours (Day 8), 336 hours (Day 15) and 504 hours (Day 22) after administration of DURVA on Day 1

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	23	9	19	
Units: L/day				
geometric mean (geometric coefficient of variation)	0.303 ( $\pm 60.9$ )	0.271 ( $\pm 62.5$ )	0.230 ( $\pm 36.4$ )	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Volume of Distribution (Vz)



End point title	Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Volume of Distribution (Vz)
End point description: Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows: $CV\% = 100 \cdot \sqrt{\exp(\sigma^2) - 1}$ , where $\sigma^2$ denotes the variance of the log-transformed values.	
End point type	Secondary
End point timeframe: pre-infusion (-60 to -5 minutes prior to dose), end of infusion (EOI), 4 hours, 168 hours (Day 8), 336 hours (Day 15) and 504 hours (Day 22) after administration of DURVA on Day 1	

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	23	9	19	
Units: liters				
geometric mean (geometric coefficient of variation)	4.244 ( $\pm$ 33.5)	4.582 ( $\pm$ 33.3)	4.563 ( $\pm$ 42.9)	

## Statistical analyses

No statistical analyses for this end point

## Secondary: Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Terminal Elimination Half-life (t<sub>1/2</sub>)

End point title	Durvalumab (DURVA) Serum PK Parameters in Cycle 1: Terminal Elimination Half-life (t <sub>1/2</sub> )
End point description: Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows: $CV\% = 100 \cdot \sqrt{\exp(\sigma^2) - 1}$ , where $\sigma^2$ denotes the variance of the log-transformed values.	
End point type	Secondary
End point timeframe: pre-infusion (-60 to -5 minutes prior to dose), end of infusion (EOI), 4 hours, 168 hours (Day 8), 336 hours (Day 15) and 504 hours (Day 22) after administration of DURVA on Day 1	

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	23	9	19	
Units: day				
geometric mean (geometric coefficient of variation)	10.984 ( $\pm$ 52.1)	13.376 ( $\pm$ 90.9)	15.844 ( $\pm$ 29.8)	

## Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Area Under the Concentration-time Curve from Time Zero to the Last Measured Time Point (AUC0-last)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Area Under the Concentration-time Curve from Time Zero to the Last Measured Time Point (AUC0-last)
-----------------	--

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \times \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

Cycle 1 Day 1: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	24	10	21	
Units: hour*ng/mL				
geometric mean (geometric coefficient of variation)	1468.102 ( $\pm$ 78.6)	1442.549 ( $\pm$ 29.9)	591.085 ( $\pm$ 22.3)	

## Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Area Under the Concentration-time Curve from Time Zero to Infinity (AUC0-inf)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Area Under the Concentration-time Curve from Time Zero to Infinity (AUC0-inf)
-----------------	---

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \times \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

Cycle 1 Day 1: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	13	3	12	
Units: hour*ng/mL				
geometric mean (geometric coefficient of variation)	2534.684 ( $\pm$ 48.6)	2011.889 ( $\pm$ 26.8)	768.104 ( $\pm$ 33.5)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Maximum Observed Concentration (Cmax)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Maximum Observed Concentration (Cmax)
-----------------	---

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 * \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

Cycle 1 Day 1: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	24	10	21	
Units: ng/mL				
geometric mean (geometric coefficient of variation)	395.774 ( $\pm$ 56.2)	354.018 ( $\pm$ 34.7)	161.372 ( $\pm$ 33.9)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Time to Maximum Observed Concentration (Tmax)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Time to Maximum Observed Concentration (Tmax)
End point description: Time to maximum observed concentration of Lenalidomide (LEN) after multiple doses on day 1 obtained from the observed concentration versus time data	
End point type	Secondary
End point timeframe: Cycle 1 Day 1: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose	

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	24	10	21	
Units: hour				
median (full range (min-max))	1.050 (0.42 to 4.08)	1.925 (0.50 to 4.00)	1.150 (0.43 to 4.17)	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Terminal Elimination Half-life (t<sub>1/2</sub>)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Terminal Elimination Half-life (t <sub>1/2</sub> )
End point description: Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows: $CV\% = 100 \times \sqrt{\exp(\sigma^2) - 1}$ , where $\sigma^2$ denotes the variance of the log-transformed values.	
End point type	Secondary
End point timeframe: Cycle 1 Day 1: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose	

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	13	3	12	
Units: hour				
geometric mean (geometric coefficient of variation)	3.477 ( $\pm 61.3$ )	3.051 ( $\pm 16.8$ )	2.883 ( $\pm 29.2$ )	

## Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Apparent Clearance (CL/F)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Apparent Clearance (CL/F)
-----------------	---

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \cdot \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

Cycle 1 Day 1: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	13	3	12	
Units: L/hour				
geometric mean (geometric coefficient of variation)	8.838 ( $\pm$ 52.7)	12.426 ( $\pm$ 26.8)	13.019 ( $\pm$ 33.5)	

## Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Apparent Volume of Distribution (Vz/F)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 1: Apparent Volume of Distribution (Vz/F)
-----------------	--

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \cdot \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

Cycle 1 Day 1: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	13	3	12	
Units: liters				
geometric mean (geometric coefficient of variation)	44.329 ( $\pm 25.4$ )	54.689 ( $\pm 29.7$ )	54.157 ( $\pm 22.3$ )	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma PK parameters in Cycle 1 Day 15: Area Under the Concentration-time Curve from Time Zero to the Last Measured Time Point (AUC0-last)

End point title	Lenalidomide (LEN) Plasma PK parameters in Cycle 1 Day 15: Area Under the Concentration-time Curve from Time Zero to the Last Measured Time Point (AUC0-last)
-----------------	---

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \times \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$  denotes the variance of the log-transformed values.

End point type	Secondary
----------------	-----------

End point timeframe:

Cycle 1 Day 15: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	18	9	16	
Units: hour*ng/mL				
geometric mean (geometric coefficient of variation)	1911.739 ( $\pm 65.5$ )	1754.349 ( $\pm 34.3$ )	629.151 ( $\pm 16.8$ )	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 15: Maximum Observed Concentration (Cmax)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 15: Maximum Observed Concentration (Cmax)
-----------------	--

End point description:

Geometric mean was obtained by computing the arithmetic mean of the logarithm-transformed values of concentration/PK parameters and then using the exponentiation to return the computation to the original scales. Geometric CV% was calculated as follows:  $CV\% = 100 \times \sqrt{\exp(\sigma^2) - 1}$ , where  $\sigma^2$

denotes the variance of the log-transformed values.

End point type	Secondary
End point timeframe:	
Cycle 1 Day 15: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose	

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	18	9	16	
Units: ng/mL				
geometric mean (geometric coefficient of variation)	409.764 ( $\pm 66.0$ )	452.850 ( $\pm 31.7$ )	171.235 ( $\pm 33.6$ )	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Participants Who Had Either Disease Progression or Death

End point title	Participants Who Had Either Disease Progression or Death
-----------------	--

End point description:

This outcome was originally defined as a Kaplan-Meier estimate of progression-free survival (PFS) which estimated the time between first date of dosing of study medication and disease progression, as determined by the investigator using the IMWG Uniform Response Criteria, or death during study treatment, whichever occurred earlier. However due to the early study termination and limited follow-up time, the majority of participants were censored for PFS analysis. Data reported instead represent the number of participants who died during study treatment or had disease progression within 90 days of the last dose of durvalumab.

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 up to Week 84

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	10	21	
Units: Count of participants	4	1	3	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Participants Who Developed Anti-drug Antibody Against Durvalumab

End point title	Participants Who Developed Anti-drug Antibody Against Durvalumab
End point description: The number of participants who develop antidrug antibody against durvalumab at any of the sampling timepoints during the study.	
End point type	Secondary
End point timeframe: Pre-dose samples on Day 1 of cycles 1, 2, 4, 6, 10, and 14 (study days 1, 29, 85, 141, 253, 393)	

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	10	21	
Units: Count of participants	0	1	0	

### Statistical analyses

No statistical analyses for this end point

### Secondary: Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 15: Time to Maximum Observed Concentration (Tmax)

End point title	Lenalidomide (LEN) Plasma Pharmacokinetic (PK) Parameters in Cycle 1 Day 15: Time to Maximum Observed Concentration (Tmax)
End point description: Time to maximum observed concentration of Lenalidomide (LEN) after multiple doses on day 15 obtained from the observed concentration versus time data	
End point type	Secondary
End point timeframe: Cycle 1 Day 15: pre-dose, 0.5, 1, 2, 4, and 8 hours post LEN dose	

End point values	Cohort A: High risk, TNE	Cohort B: $\geq 65$ years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	18	9	16	
Units: hour				
median (full range (min-max))	2.000 (1.00 to 8.00)	1.000 (0.55 to 2.00)	1.042 (0.47 to 4.00)	

### Statistical analyses

No statistical analyses for this end point



---

**Secondary: Participants Who Died Up To Data Cut-off Date (15 December 2017)**

---

End point title	Participants Who Died Up To Data Cut-off Date (15 December 2017)
-----------------	--

End point description:

This outcome was originally defined as a Kaplan-Meier estimate of overall survival (OS) and was defined as the time between first date of dosing of study medication and death due to any cause. However due to the early study termination and limited follow-up time, the majority of participants were censored for OS analysis. Data reported instead represent the number of participants who died due to any cause from Day 1 up to data cut-off.

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 up to Week 87

---

End point values	Cohort A: High risk, TNE	Cohort B: >=65 years old, TNE	Cohort C: High risk, Post-transplant	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	25	10	21	
Units: Count of participants	2	0	1	

---

**Statistical analyses**

---

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

SAEs and NSAEs assessed from first dose up to week 84 (the longer of 90 days after d/c treatment with DURVA, or 28 days after the last dose of LEN or DEX). Deaths (all-causes) assessed from first dose to study completion (Up to approximately 88 months)

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	22.0
--------------------	------

### Reporting groups

Reporting group title	Cohort A: High Risk, TNE
-----------------------	--------------------------

Reporting group description:

High risk, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$  75 years old) or 20 mg/day ( $>$  75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle.

Reporting group title	Cohort B: $\geq$ 65 Years Old, TNE
-----------------------	------------------------------------

Reporting group description:

$\geq$  65 years old, transplant non-eligible [TNE], newly diagnosed multiple myeloma (NDMM) participants who were not high risk were administered • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 25 mg/day (adjust per the creatinine clearance [CrCl] value) on Days 1 to 21 of each 28-day treatment cycle • Oral dexamethasone (dex) 40 mg/day ( $\leq$  75 years old) or 20 mg/day ( $>$  75 years old) on Days 1, 8, 15, and 22 of each 28-day cycle, up to 12 cycles.

Reporting group title	Cohort C: High Risk, Post-transplant
-----------------------	--------------------------------------

Reporting group description:

High risk, post-transplant NDMM participants were administered the following as maintenance therapy: • Intravenous (IV) durvalumab at 1500 mg on Day 1 of each 28-day cycle • Oral lenalidomide (LEN) 10 mg/day on Days 1 to 21 of each 28-day treatment cycle.

Serious adverse events	Cohort A: High Risk, TNE	Cohort B: $\geq$ 65 Years Old, TNE	Cohort C: High Risk, Post-transplant
Total subjects affected by serious adverse events			
subjects affected / exposed	12 / 25 (48.00%)	6 / 10 (60.00%)	4 / 21 (19.05%)
number of deaths (all causes)	9	3	5
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Adenocarcinoma of colon			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metastases to liver			

subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metastases to lung			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Hypotension			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Asthenia			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General physical health deterioration			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyrexia			
subjects affected / exposed	2 / 25 (8.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	2 / 3	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			
Cytokine release syndrome			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	1 / 1
Respiratory, thoracic and mediastinal disorders			
Pulmonary embolism			

subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Acute pulmonary oedema			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 1	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Pubis fracture			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Cerebrovascular accident			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Transient ischaemic attack			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Febrile neutropenia			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences causally related to treatment / all	1 / 1	0 / 0	2 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Vomiting			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diarrhoea			

subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Cholecystitis acute			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Renal failure			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Endocrine disorders			
Hyperthyroidism			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Adrenal insufficiency			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Inappropriate antidiuretic hormone secretion			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Muscular weakness			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pathological fracture			

subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Clostridium difficile infection			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cystitis klebsiella			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Influenza			
subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Osteomyelitis			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Parainfluenzae virus infection			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumocystis jirovecii pneumonia			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			

subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urosepsis			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urinary tract infection			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Upper respiratory tract infection			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Diabetes mellitus			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Diabetic ketoacidosis			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tumour lysis syndrome			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	<b>Cohort A: High Risk, TNE</b>	<b>Cohort B: &gt;=65 Years Old, TNE</b>	<b>Cohort C: High Risk, Post-transplant</b>
Total subjects affected by non-serious adverse events			
subjects affected / exposed	24 / 25 (96.00%)	9 / 10 (90.00%)	18 / 21 (85.71%)
<b>Vascular disorders</b>			
Deep vein thrombosis			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Hypotension			
subjects affected / exposed	2 / 25 (8.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences (all)	2	0	0
Phlebitis			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
<b>General disorders and administration site conditions</b>			
Fatigue			
subjects affected / exposed	4 / 25 (16.00%)	3 / 10 (30.00%)	7 / 21 (33.33%)
occurrences (all)	4	3	8
Chills			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	1	0	2
Asthenia			
subjects affected / exposed	7 / 25 (28.00%)	2 / 10 (20.00%)	1 / 21 (4.76%)
occurrences (all)	7	2	1
General physical health deterioration			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Peripheral swelling			
subjects affected / exposed	2 / 25 (8.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	2	1	0
Oedema peripheral			
subjects affected / exposed	8 / 25 (32.00%)	2 / 10 (20.00%)	0 / 21 (0.00%)
occurrences (all)	8	2	0
Non-cardiac chest pain			
subjects affected / exposed	2 / 25 (8.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences (all)	2	0	0
Influenza like illness			



subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	1	0	3
Pyrexia			
subjects affected / exposed	3 / 25 (12.00%)	4 / 10 (40.00%)	3 / 21 (14.29%)
occurrences (all)	4	5	3
Immune system disorders			
Seasonal allergy			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Respiratory, thoracic and mediastinal disorders			
Productive cough			
subjects affected / exposed	2 / 25 (8.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	2	2	1
Oropharyngeal pain			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	3 / 21 (14.29%)
occurrences (all)	0	1	3
Nasal congestion			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
Epistaxis			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Dyspnoea			
subjects affected / exposed	2 / 25 (8.00%)	3 / 10 (30.00%)	0 / 21 (0.00%)
occurrences (all)	2	3	0
Dysphonia			
subjects affected / exposed	1 / 25 (4.00%)	2 / 10 (20.00%)	0 / 21 (0.00%)
occurrences (all)	1	2	0
Cough			
subjects affected / exposed	3 / 25 (12.00%)	2 / 10 (20.00%)	5 / 21 (23.81%)
occurrences (all)	3	2	5
Respiratory tract congestion			
subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	1	1	0
Rhinorrhoea			

subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 1	1 / 10 (10.00%) 1	1 / 21 (4.76%) 1
Upper-airway cough syndrome subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 1	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Psychiatric disorders			
Insomnia subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	4 / 10 (40.00%) 4	3 / 21 (14.29%) 3
Depression subjects affected / exposed occurrences (all)	4 / 25 (16.00%) 4	0 / 10 (0.00%) 0	2 / 21 (9.52%) 2
Investigations			
Alanine aminotransferase increased subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	0 / 10 (0.00%) 0	1 / 21 (4.76%) 2
Amylase increased subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 1	1 / 10 (10.00%) 2	0 / 21 (0.00%) 0
Aspartate aminotransferase increased subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Blood alkaline phosphatase increased subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 3	0 / 10 (0.00%) 0	1 / 21 (4.76%) 1
Blood creatinine increased subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	0 / 10 (0.00%) 0	1 / 21 (4.76%) 1
Glomerular filtration rate decreased subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Lipase increased subjects affected / exposed occurrences (all)	3 / 25 (12.00%) 5	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Prostatic specific antigen increased			

subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Weight decreased subjects affected / exposed occurrences (all)	5 / 25 (20.00%) 5	1 / 10 (10.00%) 1	3 / 21 (14.29%) 3
Injury, poisoning and procedural complications			
Infusion related reaction subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 1	0 / 10 (0.00%) 0	2 / 21 (9.52%) 2
Fall subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 3	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Contusion subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Tooth fracture subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	0 / 10 (0.00%) 0	3 / 21 (14.29%) 3
Cardiac disorders			
Atrial flutter subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Atrial fibrillation subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Cardiac failure congestive subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Nervous system disorders			
Disturbance in attention subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Dizziness subjects affected / exposed occurrences (all)	4 / 25 (16.00%) 4	1 / 10 (10.00%) 1	3 / 21 (14.29%) 4
Peripheral sensory neuropathy			

subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	2 / 21 (9.52%)
occurrences (all)	1	1	2
Paraesthesia			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	1	0	3
Headache			
subjects affected / exposed	1 / 25 (4.00%)	2 / 10 (20.00%)	1 / 21 (4.76%)
occurrences (all)	1	2	1
Dysgeusia			
subjects affected / exposed	3 / 25 (12.00%)	1 / 10 (10.00%)	2 / 21 (9.52%)
occurrences (all)	3	1	2
Tremor			
subjects affected / exposed	3 / 25 (12.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	3	1	0
Blood and lymphatic system disorders			
Leukopenia			
subjects affected / exposed	3 / 25 (12.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	7	1	1
Neutropenia			
subjects affected / exposed	11 / 25 (44.00%)	5 / 10 (50.00%)	3 / 21 (14.29%)
occurrences (all)	21	11	6
Anaemia			
subjects affected / exposed	13 / 25 (52.00%)	2 / 10 (20.00%)	3 / 21 (14.29%)
occurrences (all)	20	2	3
Thrombocytopenia			
subjects affected / exposed	8 / 25 (32.00%)	2 / 10 (20.00%)	4 / 21 (19.05%)
occurrences (all)	16	3	6
Ear and labyrinth disorders			
Ear pruritus			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
Abdominal pain upper			

subjects affected / exposed	3 / 25 (12.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	3	1	0
Frequent bowel movements			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Dry mouth			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	1	0	2
Diarrhoea			
subjects affected / exposed	13 / 25 (52.00%)	2 / 10 (20.00%)	5 / 21 (23.81%)
occurrences (all)	19	3	5
Constipation			
subjects affected / exposed	9 / 25 (36.00%)	4 / 10 (40.00%)	4 / 21 (19.05%)
occurrences (all)	12	5	4
Anal fissure			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Gastrooesophageal reflux disease			
subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	1	1	0
Vomiting			
subjects affected / exposed	5 / 25 (20.00%)	2 / 10 (20.00%)	2 / 21 (9.52%)
occurrences (all)	6	2	2
Nausea			
subjects affected / exposed	7 / 25 (28.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	11	0	3
Haemorrhoids			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	2	0
Hepatobiliary disorders			
Hepatobiliary disease			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	3	0
Skin and subcutaneous tissue disorders			
Night sweats			

subjects affected / exposed occurrences (all)	3 / 25 (12.00%) 3	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Pruritus			
subjects affected / exposed occurrences (all)	4 / 25 (16.00%) 4	1 / 10 (10.00%) 1	2 / 21 (9.52%) 3
Rash			
subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	1 / 10 (10.00%) 1	2 / 21 (9.52%) 2
Dry skin			
subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 1	0 / 10 (0.00%) 0	5 / 21 (23.81%) 7
Erythema			
subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 2	1 / 10 (10.00%) 3	0 / 21 (0.00%) 0
Rash maculo-papular			
subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	0 / 10 (0.00%) 0	2 / 21 (9.52%) 2
Rash generalised			
subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Rash erythematous			
subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Solar dermatitis			
subjects affected / exposed occurrences (all)	0 / 25 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Renal and urinary disorders			
Haematuria			
subjects affected / exposed occurrences (all)	2 / 25 (8.00%) 2	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Dysuria			
subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 1	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Nocturia			
subjects affected / exposed occurrences (all)	1 / 25 (4.00%) 1	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0

Polyuria			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Pollakiuria			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Renal failure			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Urine odour abnormal			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Urethral stenosis			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Endocrine disorders			
Hypothyroidism			
subjects affected / exposed	2 / 25 (8.00%)	0 / 10 (0.00%)	4 / 21 (19.05%)
occurrences (all)	2	0	4
Hyperthyroidism			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	6 / 21 (28.57%)
occurrences (all)	2	0	6
Adrenal insufficiency			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	3 / 25 (12.00%)	0 / 10 (0.00%)	4 / 21 (19.05%)
occurrences (all)	3	0	4
Back pain			
subjects affected / exposed	6 / 25 (24.00%)	2 / 10 (20.00%)	2 / 21 (9.52%)
occurrences (all)	6	2	2
Bone pain			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	0	0	2
Intervertebral disc degeneration			

subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Flank pain			
subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	1	1	0
Pathological fracture			
subjects affected / exposed	2 / 25 (8.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	2	1	1
Pain in extremity			
subjects affected / exposed	3 / 25 (12.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	4	1	1
Neck pain			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	0	0	2
Musculoskeletal pain			
subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	1	1	1
Musculoskeletal chest pain			
subjects affected / exposed	4 / 25 (16.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	4	1	1
Muscle spasms			
subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	2 / 21 (9.52%)
occurrences (all)	1	1	2
Muscle contracture			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Intervertebral disc protrusion			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Infections and infestations			
Respiratory tract infection			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
Pneumonia			
subjects affected / exposed	2 / 25 (8.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences (all)	2	0	0



Oral candidiasis			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	2
Nasopharyngitis			
subjects affected / exposed	0 / 25 (0.00%)	0 / 10 (0.00%)	4 / 21 (19.05%)
occurrences (all)	0	0	7
Influenza			
subjects affected / exposed	1 / 25 (4.00%)	1 / 10 (10.00%)	2 / 21 (9.52%)
occurrences (all)	1	1	2
Herpes simplex			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Gastroenteritis			
subjects affected / exposed	2 / 25 (8.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences (all)	2	0	0
Candida infection			
subjects affected / exposed	0 / 25 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Upper respiratory tract infection			
subjects affected / exposed	1 / 25 (4.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	2	0	4
Urinary tract infection			
subjects affected / exposed	5 / 25 (20.00%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences (all)	9	0	0
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	4 / 25 (16.00%)	3 / 10 (30.00%)	2 / 21 (9.52%)
occurrences (all)	4	3	2
Dehydration			
subjects affected / exposed	2 / 25 (8.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences (all)	4	0	1
Hypocalcaemia			
subjects affected / exposed	2 / 25 (8.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	2	1	0
Hypokalaemia			

subjects affected / exposed	4 / 25 (16.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	7	1	0

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
27 June 2016	Table of Events Update
06 April 2017	Inclusion Criteria Update
05 December 2019	Clinical Hold on the Study Update

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

Study was placed on full clinical hold by the US FDA on 05 Sep 2017. Study was closed for further enrollment and subjects were discontinued from all treatments. Subjects were followed for SPMs for 5 years after last subject was enrolled per protocol.

Notes: