



Clinical trial results:

A Phase Ib/II Study Evaluating the Safety and Efficacy of Obinutuzumab in Combination with Atezolizumab Plus Polatuzumab Vedotin in Patients with Relapsed or Refractory Follicular Lymphoma and Rituximab in Combination with Atezolizumab Plus Polatuzumab Vedotin in Patients with Relapsed or Refractory Diffuse Large B-Cell Lymphoma

Summary

EudraCT number	2015-004845-25
Trial protocol	DE PL
Global end of trial date	07 October 2019

Results information

Result version number	v2
This version publication date	26 September 2020
First version publication date	21 September 2019
Version creation reason	

Trial information

Trial identification

Sponsor protocol code	BO29561
-----------------------	---------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT02729896
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	F. Hoffmann-La Roche AG
Sponsor organisation address	Grenzacherstrasse 124, Basel, Switzerland, CH-4070
Public contact	F. Hoffmann-La Roche AG, F. Hoffmann-La Roche AG, 41 616878333, global.trial_information@roche.com
Scientific contact	F. Hoffmann-La Roche AG, F. Hoffmann-La Roche AG, 41 616878333, globa.trial_information@roche.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	07 October 2019
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	07 October 2019
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

To Evaluate the Safety and Efficacy of Obinutuzumab in Combination with Atezolizumab plus Polatuzumab Vedotin in Patients with Relapsed or Refractory Follicular Lymphoma and Rituximab in Combination with Atezolizumab plus Polatuzumab Vedotin in Patients with Relapsed or Refractory Diffuse Large B Cell Lymphoma.

Protection of trial subjects:

All study subjects were required to read and sign an Informed Consent Form.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	09 November 2016
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Germany: 11
Country: Number of subjects enrolled	Poland: 14
Country: Number of subjects enrolled	United States: 11
Worldwide total number of subjects	36
EEA total number of subjects	25

Notes:

Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	20
From 65 to 84 years	15

85 years and over	1
-------------------	---

Subject disposition

Recruitment

Recruitment details: -

Pre-assignment

Screening details:

56 patients were screened and 36 patients enrolled and dosed in the study.

Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Non-randomised - controlled
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
------------------------------	-----

Arm title	FL Cohort 1.4 mg
------------------	------------------

Arm description:

During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.4 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants. Due to unexpected toxicity, recruitment was stopped, and atezolizumab discontinued in all treated patients. This combination will not be further developed.

Arm type	Experimental
Investigational medicinal product name	Atezolizumab
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1200 mg

Investigational medicinal product name	Obinutuzumab
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1000 mg

Investigational medicinal product name	Polatuzumab Vedotin
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1.4 mg

Arm title	FL Cohort 1.8 mg
------------------	------------------

Arm description:

During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.8 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants. Due to unexpected toxicity, recruitment was stopped, and atezolizumab discontinued in all

treated patients. This combination will not be further developed.

Arm type	Experimental
Investigational medicinal product name	Atezolizumab
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1200 mg

Investigational medicinal product name	Obinutuzumab
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1000 mg

Investigational medicinal product name	Polatuzumab Vedotin
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1.8 mg

Arm title	Safety Run-in Phase
------------------	---------------------

Arm description:

For DLBCL, during the induction treatment Cycles 1-6 (21-day cycles): participants received rituximab on Day 1 and Pola on Day 1.

Arm type	Experimental
Investigational medicinal product name	Atezolizumab
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1200 mg

Investigational medicinal product name	Polatuzumab Vedotin
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

1.4 mg or 1.8 mg

Investigational medicinal product name	Rituximab
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Infusion
Routes of administration	Intravenous use

Dosage and administration details:

375 mg/m²

Number of subjects in period 1	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase
Started	3	10	23
Completed	1	1	1
Not completed	2	9	22
Physician decision	1	2	3
Withdrawal By Subject	1	-	3
Death	-	2	13
Progressive Disease	-	1	-
Not Specified	-	2	-
Lost to follow-up	-	2	3

Baseline characteristics

Reporting groups

Reporting group title	FL Cohort 1.4 mg
Reporting group description:	
During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.4 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants. Due to unexpected toxicity, recruitment was stopped, and atezolizumab discontinued in all treated patients. This combination will not be further developed.	
Reporting group title	FL Cohort 1.8 mg
Reporting group description:	
During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.8 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants. Due to unexpected toxicity, recruitment was stopped, and atezolizumab discontinued in all treated patients. This combination will not be further developed.	
Reporting group title	Safety Run-in Phase
Reporting group description:	
For DLBCL, during the induction treatment Cycles 1-6 (21-day cycles): participants received rituximab on Day 1 and Pola on Day 1.	

Reporting group values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase
Number of subjects	3	10	23
Age Categorical			
Units: Subjects			
<=18 years	0	0	0
Between 18 and 65 years	2	6	12
>=65 years	1	4	11
Age Continuous			
Units: Years			
arithmetic mean	52.3	57.7	64.3
standard deviation	± 13.6	± 11.7	± 15.3
Sex: Female, Male			
Units: Subjects			
Female	0	4	9
Male	3	6	14
Race/Ethnicity, Customized			
Units: Subjects			
Hispanic or Latino	0	1	3
Not Hispanic or Latino	3	9	18
Unknown	0	0	2
Race/Ethnicity, Customized			
Units: Subjects			
Other	0	1	2
White	3	9	21

Reporting group values	Total		
Number of subjects	36		

Age Categorical Units: Subjects			
<=18 years	0		
Between 18 and 65 years	20		
>=65 years	16		
Age Continuous Units: Years			
arithmetic mean			
standard deviation	-		
Sex: Female, Male Units: Subjects			
Female	13		
Male	23		
Race/Ethnicity, Customized Units: Subjects			
Hispanic or Latino	4		
Not Hispanic or Latino	30		
Unknown	2		
Race/Ethnicity, Customized Units: Subjects			
Other	3		
White	33		

End points

End points reporting groups

Reporting group title	FL Cohort 1.4 mg
Reporting group description: During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.4 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants. Due to unexpected toxicity, recruitment was stopped, and atezolizumab discontinued in all treated patients. This combination will not be further developed.	
Reporting group title	FL Cohort 1.8 mg
Reporting group description: During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.8 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants. Due to unexpected toxicity, recruitment was stopped, and atezolizumab discontinued in all treated patients. This combination will not be further developed.	
Reporting group title	Safety Run-in Phase
Reporting group description: For DLBCL, during the induction treatment Cycles 1-6 (21-day cycles): participants received rituximab on Day 1 and Pola on Day 1.	

Primary: Percentage of Participants with CR at EOI, as Determined by the investigator on the Basis of Positron Emission Tomography and Computed Tomography (PET-CT) Scan

End point title	Percentage of Participants with CR at EOI, as Determined by the investigator on the Basis of Positron Emission Tomography and Computed Tomography (PET-CT) Scan ^[1]
End point description: Tumor response assessment was performed by the investigator according to modified Lugano classification using PET/CT scan. CR was defined as a score of 1 (no uptake above background), 2 (uptake \leq mediastinum), or 3 (uptake $<$ mediastinum but \leq liver) with or without a residual mass on PET 5-PS, for lymph nodes and extralymphatic sites; no new lesions; no evidence of FDG-avid disease in bone marrow; and normal/IHC-negative bone marrow morphology. 90% confidence interval (CI) for percentage of responders was calculated using Clopper-Pearson method. All PET evaluable 1L FL and 1L DLBCL patients with at least one dose of atezolizumab were included in efficacy population.	
End point type	Primary
End point timeframe: Within 6 to 8 weeks after Day 1 of Cycle 6 (up to approximately 6 months)	

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Statistical analyses not available for this endpoint

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	7	16	
Units: Percentage of participants				
number (not applicable)	33.33	14.30	12.5	

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of Participants with CR at EOI, as Determined by Investigator on the Basis of CT Scans Alone

End point title	Percentage of Participants with CR at EOI, as Determined by Investigator on the Basis of CT Scans Alone
-----------------	---

End point description:

Tumor response assessment was performed by investigator according to modified Lugano classification using computed tomography (CT) scan. OR: a response of CR or PR. CR: Target nodes/nodal masses regressed to ≤ 1.5 cm in LD; no extralymphatic sites of disease; organ enlargement regressed to normal; no new lesions; normal/IHC-negative bone marrow morphology. All CT evaluable 1L FL and 1L DLBCL patients with at least one dose of atezolizumab were included in efficacy population.

End point type	Secondary
----------------	-----------

End point timeframe:

Within 6 to 8 weeks after Day 1 of Cycle 6 (up to approximately 6 months)

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	7	16	
Units: Percentage of participants				
number (not applicable)	0.00	57.14	12.50	

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of Participants with Objective Response (CR + PR) at EOI, as Determined by the Investigator on the Basis of PET-CT Scans

End point title	Percentage of Participants with Objective Response (CR + PR) at EOI, as Determined by the Investigator on the Basis of PET-CT Scans
-----------------	---

End point description:

Tumor response assessment was performed by investigator according to modified Lugano classification using PET/CT scan. OR: a response of CR or PR. CR: a score of 1 (no uptake above background), 2 (uptake \leq mediastinum), or 3 (uptake $<$ mediastinum but \leq liver) with or without a residual mass on PET 5-PS, for lymph nodes & extralymphatic sites; no new lesions; no evidence of FDG-avid disease in bone marrow; and normal/IHC-negative bone marrow morphology. PR with a score 4 (uptake moderately greater than $>$ liver) or 5 (uptake markedly $>$ liver and/or new lesions) with reduced uptake compared with baseline and residual mass(es) of any size on PET 5-PS for lymph nodes and extralymphatic sites; no new lesions; and reduced residual uptake in bone marrow compared with baseline. All PET evaluable 1L FL and 1L DLBCL patients with at least one dose of atezolizumab were included in efficacy population

End point type	Secondary
End point timeframe:	
Within 6 to 8 weeks after Day 1 of Cycle 6 (up to approximately 6 months)	

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	7	16	
Units: Percentage of participants				
number (not applicable)	33.33	57.14	25.00	

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of Participants with Objective Response (CR + PR) at EOI, as Determined by the Investigator on the Basis of CT Scans Alone

End point title	Percentage of Participants with Objective Response (CR + PR) at EOI, as Determined by the Investigator on the Basis of CT Scans Alone
-----------------	---

End point description:

Tumor response assessment was performed by investigator according to modified Lugano classification using CT scan. OR: a response of CR or PR. CR: Target nodes/nodal masses regressed to ≤ 1.5 cm in LDi; no extralymphatic sites of disease; organ enlargement regressed to normal; no new lesions; normal/IHC-negative bone marrow morphology. PR with ≥ 50 percent decrease in SPD of up to six target measurable nodes and extranodal sites, a 5 mm x 5 mm default value when lesions were too small, 0 x0 mm value when lesions were no longer visible, actual measurements were used for nodes greater than 5 mm x 5 mm for lymph nodes & extralymphatic sites; absent/normal, regression for non-measured lesion; spleen enlargement regression by > 50 percent; no new lesions; reduced residual uptake in bone marrow compared to baseline. All CT evaluable 1L FL and 1L DLBCL patients with at least one dose of atezolizumab were included in efficacy population.

End point type	Secondary
End point timeframe:	
Within 6 to 8 weeks after Day 1 of Cycle 6 (up to approximately 6 months)	

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	7	16	
Units: Percentage of participants				
number (not applicable)	33.33	57.14	25.00	

Statistical analyses

Secondary: Percentage of Participants with Best Response of CR or PR during the Study, as Determined by the Investigator on the Basis of CT Scans Alone

End point title	Percentage of Participants with Best Response of CR or PR during the Study, as Determined by the Investigator on the Basis of CT Scans Alone
-----------------	--

End point description:

Tumor response assessment was performed by investigator according to modified Lugano classification using CT scan. OR: a response of CR or PR. CR: Target nodes/nodal masses regressed to ≤ 1.5 cm in LD; no extralymphatic sites of disease; organ enlargement regressed to normal; no new lesions; normal/IHC-negative bone marrow morphology. PR with ≥ 50 percent decrease in SPD of up to six target measurable nodes and extranodal sites, a 5 mm x 5 mm default value when lesions were too small, 0 x 0 mm value when lesions were no longer visible, actual measurements were used for nodes greater than 5 mm x 5 mm for lymph nodes & extralymphatic sites; absent/normal, regression for non-measured lesion; spleen enlargement regression by > 50 percent; no new lesions; reduced residual uptake in bone marrow compared to baseline. All CT evaluable 1L FL and 1L DLBCL patients with at least one dose of atezolizumab were included in efficacy population.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to approximately 4 years

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	7	16	
Units: Percentage of participants				
number (not applicable)				
Complete response	33.3	14.3	12.5	
Partial response	0	42.9	12.5	
Stable disease	33.3	14.3	6.3	
Progressive disease	33.3	14.3	31.3	
Not available	0	14.3	37.5	

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of Participants with Adverse Events and Serious Adverse Events

End point title	Percentage of Participants with Adverse Events and Serious Adverse Events
-----------------	---

End point description:

An AE was any untoward medical occurrence in a participant administered a pharmaceutical product and which did not necessarily have to have a causal relationship with the treatment. An adverse event was therefore any unfavourable and unintended sign (including an abnormal laboratory finding, for example), symptom, or disease temporally associated with the use of a pharmaceutical product, whether or not considered related to the pharmaceutical product. Grading was completed according to the CTCAE, version 4.0 for severity and tumor flare reactions were graded according to NCI CTCAE v3.0.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to approximately 4 years

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	10	21	
Units: Participants				
number (not applicable)	100.00	100.00	81.00	

Statistical analyses

No statistical analyses for this end point

Secondary: Serum Obinutuzumab Concentration

End point title	Serum Obinutuzumab Concentration ^[2]
End point description: pre-dose (0 hr), 30 min after EOI on Day 1 Cycle 1; pre-dose (within 5 hr), 30 min after EOI on Day 1 of Cycles 2, 4, 6; maintenance phase: pre-dose (within 5 hr) on Day 1 of Months 1, 7, 13, 19; anytime during treatment discontinuation visit, 120 days after the last dose, and 1 year after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 50 mg/hr and decreases every 30 min to maximum of 400 mg/hr). 99999 represents the upper limit of confidence interval was not estimable due to the low number of subjects within events.	
End point type	Secondary
End point timeframe: Pre-dose (0 hr) up to approximately 4 years	

Notes:

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: Only arms treated with Obinutuzumab were included in this endpoint

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	3	10		
Units: mcg/mL				
arithmetic mean (standard deviation)				
Induction cycle 1 Day 1 Pre-dose	99999 (± 99999)	99999 (± 99999)		
Induction cycle 1 Day 1 Post-dose	308 (± 41.6)	295 (± 171)		
Induction cycle 2 Day 1 Pre-dose	351 (± 81.3)	403 (± 143)		
Induction cycle 2 Day 1 Post-dose	616 (± 115)	672 (± 130)		
Induction cycle 4 Day 1 Pre-dose	433 (± 322)	311 (± 177)		
Induction cycle 4 Day 1 Post-dose	583 (± 189)	619 (± 160)		
Induction cycle 6 Day 1 Pre-dose	288 (± 123)	308 (± 194)		
Induction cycle 6 Day 1 Post-dose	514 (± 121)	605 (± 161)		
Maintenance Month 1 Day 1 Pre-dose	221 (± 99999)	171 (± 131)		
Maintenance Month 7 Pre-dose	90.4 (± 58.8)	83.8 (± 59.6)		
Maintenance Month 13 Pre-dose	78.8 (± 42.8)	158 (± 1.41)		

Study drug completion or early discontinuation	0 (\pm 0)	37.2 (\pm 99999)		
PK and Immunogenicity follow up	110 (\pm 120)	15.4 (\pm 3.20)		

Statistical analyses

No statistical analyses for this end point

Secondary: Serum Rituximab Concentration

End point title	Serum Rituximab Concentration ^[3]
-----------------	--

End point description:

pre-dose (0 hr), 30 min after EOI on Day 1 Cycle 1; pre-dose (within 5 hr) on Day 1 of Cycles 2, 4; pre-dose (within 5 hr), 30 min after EOI on Day 1 of Cycle 6; anytime during treatment discontinuation visit, 120 days after the last dose, and 1 year after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 50 mg/hr and increases every 30 min to maximum of 400 mg/hr)

End point type	Secondary
----------------	-----------

End point timeframe:

Pre-dose (0 hr) up to approximately 4 years

Notes:

[3] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Only arms treated with Obinutuzumab were included in this endpoint

End point values	Safety Run-in Phase			
Subject group type	Reporting group			
Number of subjects analysed	21			
Units: mcg/mL				
arithmetic mean (standard deviation)				
Induction Cycle 1 Day 1 Pre-dose	30.4 (\pm 28.8)			
Induction Cycle 1 Day 1 Post-dose	197 (\pm 56.9)			
Induction Cycle 2 Day 1 Pre-dose	43.5 (\pm 28.7)			
Induction Cycle 4 Day 1 Pre-dose	84.8 (\pm 35.7)			
Induction Cycle 6 Day 1 Pre-dose	101 (\pm 45.5)			
Induction Cycle 6 Day 1 Post-dose	239 (\pm 42.8)			

Statistical analyses

No statistical analyses for this end point

Secondary: Serum Atezo Concentration

End point title	Serum Atezo Concentration
-----------------	---------------------------

End point description:

pre-dose (within 5 hr), 30 min after EOI on Day 1 of Cycles 2, 3, 4; pre-dose (within 5 hr) on Day 1 of Cycle 6; maintenance phase: pre-dose (within 5 hr) on Day 1 of Month 1; 30 min after EOI on Day 2 of Month 1; pre-dose (within 5 hr) on Day 1 of Month 4, 7, 13, 19; anytime during treatment discontinuation visit, 120 days after the last dose, and 1-2 years after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 60 min and decreases to 30 min). 99999 represents the upper limit of confidence interval was not estimable due to the low number of subjects within events.

End point type	Secondary
End point timeframe:	
Pre-dose (0 hr) up to approximately 4 years	

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	10	21	
Units: mcg/mL				
arithmetic mean (standard deviation)				
Induction Cycle 2 Day 1 Pre-dose	99999 (± 99999)	99999 (± 99999)	99999 (± 99999)	
Induction Cycle 2 Day 1 Post-dose	332 (± 56.7)	365 (± 80.7)	355 (± 78.8)	
Induction Cycle 3 Day 1 Pre-dose	86.9 (± 28.8)	78.7 (± 18.4)	79.7 (± 30.6)	
Induction Cycle 4 Day 1 Pre-dose	139 (± 33.1)	125 (± 47.3)	139 (± 56.4)	
Induction Cycle 4 Day 1 Post-dose	462 (± 82.0)	450 (± 136)	489 (± 131)	
Induction Cycle 6 Day 1 Pre-dose	194 (± 67.7)	181 (± 112)	180 (± 90.6)	
Maintenance Month 1 Day 1 Pre-dose	100 (± 66.0)	70.7 (± 55.5)	0 (± 0)	
Maintenance Month 1 Day 2 Post-dose	577 (± 144)	473 (± 177)	0 (± 0)	
Maintenance Month 4 Pre-dose	240 (± 101)	140 (± 141)	0 (± 0)	
Maintenance Month 7 Pre-dose	226 (± 134)	221 (± 90.5)	0 (± 0)	
Maintenance Month 13 Pre-dose	106 (± 83.0)	0 (± 0)	0 (± 0)	
Study drug completion or early discontinuation	0 (± 0)	127 (± 99999)	93.0 (± 14.2)	
Consolidation Month 1 Day 1 Pre-dose	0 (± 0)	0 (± 0)	132 (± 43.3)	
Consolidation Month 1 Day 2 Post-dose	0 (± 0)	0 (± 0)	503 (± 214)	
PK and Immunogenicity followup 120D	61.4 (± 60.3)	16.8 (± 26.8)	19.9 (± 0.778)	
PK and Immunogenicity followup 1 Year	0 (± 0)	0 (± 0)	0 (± 0)	

Statistical analyses

No statistical analyses for this end point

Secondary: Serum Pola Concentration

End point title	Serum Pola Concentration
End point description:	
pre-dose (0 hr) on Day 1 Cycle 1; pre-dose (within 5 hr) on Day 1 of Cycles 2, 4; maintenance phase: pre-dose (within 5 hr) on Day 1 of Months 1; anytime during treatment discontinuation visit, 120 days after the last dose, and 1 year after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 90 min and decreases to 30 min). 99999 represents the upper limit of confidence interval was not estimable due to the low number of subjects within events.	
End point type	Secondary
End point timeframe:	
Pre-dose (0 hr) up to approximately 4 years	

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	10	21	
Units: mcg/mL				
arithmetic mean (standard deviation)				
Pre Day 120 Follow-up	0 (± 0)	0.136 (± 99999)	2.91 (± 99999)	
Induction Cycle 1 Day 1 Pre-dose	0 (± 0)	0.847 (± 99999)	35.9 (± 99999)	
Induction Cycle 2 Day 1 Pre-dose	2.54 (± 0.918)	2.62 (± 1.22)	3.37 (± 3.60)	
Induction Cycle 4 Day 1 Pre-dose	4.73 (± 2.10)	4.27 (± 1.63)	4.92 (± 2.65)	
Maintenance Month 1 Day 1 Pre-dose	1.30 (± 0.696)	1.63 (± 1.75)	0 (± 0)	
Study drug completion or early discontinuation	0 (± 0)	0 (± 0)	0.716 (± 0.200)	

Statistical analyses

No statistical analyses for this end point

Secondary: Incidence of Human Anti-Human Antibodies (HAHAs) to Obinutuzumab

End point title	Incidence of Human Anti-Human Antibodies (HAHAs) to Obinutuzumab ^[4]
-----------------	---

End point description:

Pre-dose (0 hr) on Day 1 of Cycle 1, 6, anytime during treatment discontinuation visit, 120 days after the last dose, and 1 year after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 50 mg/hr and increased every 30 min to maximum of 400 mg/hr)

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to approximately 4 years

Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: Only arms treated with Obinutuzumab were included in this endpoint

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	3	10		
Units: Percentage of participants	0	0		

Statistical analyses

No statistical analyses for this end point

Secondary: Incidence of Human Anti-Chimeric Antibodies (HACAs) to Rituximab

End point title	Incidence of Human Anti-Chimeric Antibodies (HACAs) to Rituximab ^[5]
-----------------	---

End point description:

Pre-dose (0 hr) on Day 1 of Cycle 1, 2, 4, 6, anytime during treatment discontinuation visit, 120 days

after the last dose, and 1 year after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 50 mg/hr and increased every 30 min to maximum of 400 mg/hr)

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline to approximately 4 years

Notes:

[5] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Only arms treated with Obinutuzumab were included in this endpoint

End point values	Safety Run-in Phase			
Subject group type	Reporting group			
Number of subjects analysed	21			
Units: Percentage of participants				
number (not applicable)				
Baseline, positive	5.6			
Baseline, negative	94.4			
Induction Cycle 2 Day 1, positive	5.6			
Induction Cycle 2 Day 1, negative	94.4			
Induction Cycle 4 Day 1	100			
Induction Cycle 6 Day 1	100			
Study drug completion	100			

Statistical analyses

No statistical analyses for this end point

Secondary: Incidence of Anti-Therapeutic Antibodies (ATAs) to Atezo

End point title	Incidence of Anti-Therapeutic Antibodies (ATAs) to Atezo
-----------------	--

End point description:

Pre-dose (0 hr) on Day 1 of Cycle 2, 3, 4, 6, Month 1, 4, 7, 13 and 19, anytime during treatment discontinuation visit, 120 days after the last dose, and 1 year after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 60 min and decreases to 30 min)

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline to approximately 4 years

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	10	21	
Units: Percentage of participants				
number (not applicable)				
Induction, Cycle 2 Day 1, positive	0	0	5.3	
Induction, Cycle 2 Day 1, negative	0	0	94.7	
Induction, Cycle 3 Day 1, positive	0	0	7.1	
Induction, Cycle 3 Day 1, negative	0	0	92.9	

Induction, Cycle 4 Day 1, negative	0	0	100.0	
Induction, Cycle 6 Day 1, negative	0	0	100.0	
Consolidation Month 1 Day 1, negative	0	0	100.0	
Study drug completion, negative	0	0	100.0	
Atezo PK Immunogenicity Follow up (120D), negative	0	0	100.0	

Statistical analyses

No statistical analyses for this end point

Secondary: Incidence of ATAs to Pola

End point title	Incidence of ATAs to Pola
End point description: Pre-dose (0 hr) on Day 1 of Cycle 1, 2, 4, anytime during treatment discontinuation visit, 120 days after the last dose, and 1 year after the last dose up to approximately 4 years (1 cycle=21 days; infusion rate: starts with 90 min and decreases to 30 min)	
End point type	Secondary
End point timeframe: Baseline to approximately 4 years	

End point values	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	3	10	21	
Units: Percentage of participants	0	0	0	

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

From randomization to end of study (approximately 4 years)

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	22.1
--------------------	------

Reporting groups

Reporting group title	FL Cohort 1.4 mg
-----------------------	------------------

Reporting group description:

During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.4 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants.

Reporting group title	FL Cohort 1.8 mg
-----------------------	------------------

Reporting group description:

During the induction treatment Cycle 1 (21-day cycles): participants received obinutuzumab on Days 1, 8, and 15 and Pola on Day 1; Cycles 2-6: participants received 1000 mg of obinutuzumab on Day 1, 1200 mg of Atezo on Day 1, and 1.8 mg of Pola on Day 1. This was followed by obinutuzumab on Day 1 of every other month starting with Month 1 for 24 months, during maintenance treatment for FL participants

Reporting group title	Safety Run-in Phase
-----------------------	---------------------

Reporting group description:

For DLBCL, during the induction treatment Cycles 1-6 (21-day cycles): participants received rituximab on Day 1 and Pola on Day 1.

Serious adverse events	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 3 (0.00%)	4 / 10 (40.00%)	2 / 21 (9.52%)
number of deaths (all causes)	0	2	12
number of deaths resulting from adverse events			
Investigations			
C-REACTIVE PROTEIN INCREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
GUILLAIN-BARRE SYNDROME			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

LETHARGY			
subjects affected / exposed	0 / 3 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
ANAEMIA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
FEBRILE NEUTROPENIA			
subjects affected / exposed	0 / 3 (0.00%)	2 / 10 (20.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
DEATH			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	1 / 1	0 / 0
PYREXIA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
ABDOMINAL PAIN			
subjects affected / exposed	0 / 3 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
STOMATITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
PLEURAL EFFUSION			

subjects affected / exposed	0 / 3 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 1
PNEUMONITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
DERMATITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ERYTHEMA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
BRONCHOPULMONARY ASPERGILLOSIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 0
ERYSIPELAS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
SEPSIS			
subjects affected / exposed	0 / 3 (0.00%)	0 / 10 (0.00%)	1 / 21 (4.76%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Frequency threshold for reporting non-serious adverse events: 5 %

Non-serious adverse events	FL Cohort 1.4 mg	FL Cohort 1.8 mg	Safety Run-in Phase
Total subjects affected by non-serious adverse events subjects affected / exposed	3 / 3 (100.00%)	10 / 10 (100.00%)	15 / 21 (71.43%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps) BASAL CELL CARCINOMA subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
Vascular disorders HYPERTENSION subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
THROMBOSIS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
General disorders and administration site conditions CHEST PAIN subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
FACE OEDEMA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
FATIGUE subjects affected / exposed occurrences (all)	2 / 3 (66.67%) 2	3 / 10 (30.00%) 3	1 / 21 (4.76%) 1
INFLUENZA LIKE ILLNESS subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
INFUSION SITE EXTRAVASATION subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
OEDEMA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
OEDEMA PERIPHERAL subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	3 / 10 (30.00%) 3	5 / 21 (23.81%) 5

PERIPHERAL SWELLING subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 3	1 / 21 (4.76%) 1
PYREXIA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	4 / 10 (40.00%) 6	0 / 21 (0.00%) 0
Respiratory, thoracic and mediastinal disorders			
COUGH subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	2 / 10 (20.00%) 3	2 / 21 (9.52%) 3
DYSPNOEA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 10 (0.00%) 0	2 / 21 (9.52%) 2
DYSPNOEA EXERTIONAL subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 10 (0.00%) 0	2 / 21 (9.52%) 2
HICCUPS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
LARYNGEAL OEDEMA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
PULMONARY EMBOLISM subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 2	0 / 21 (0.00%) 0
PULMONARY THROMBOSIS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Psychiatric disorders			
DEPRESSION subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 10 (0.00%) 0	1 / 21 (4.76%) 1
INSOMNIA subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	0 / 10 (0.00%) 0	0 / 21 (0.00%) 0
ANXIETY			

subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
Investigations			
ALANINE AMINOTRANSFERASE INCREASED			
subjects affected / exposed	1 / 3 (33.33%)	2 / 10 (20.00%)	0 / 21 (0.00%)
occurrences (all)	1	2	0
BLOOD CREATININE INCREASED			
subjects affected / exposed	0 / 3 (0.00%)	2 / 10 (20.00%)	0 / 21 (0.00%)
occurrences (all)	0	2	0
BLOOD LACTATE DEHYDROGENASE INCREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
C-REACTIVE PROTEIN INCREASED			
subjects affected / exposed	0 / 3 (0.00%)	2 / 10 (20.00%)	0 / 21 (0.00%)
occurrences (all)	0	5	0
GAMMA-GLUTAMYLTRANSFERASE INCREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
LIPASE INCREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
NEUTROPHIL COUNT DECREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	2	0
PROCALCITONIN INCREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
THYROXINE DECREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
TRI-IODOTHYRONINE DECREASED			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
WEIGHT DECREASED			

subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	4 / 10 (40.00%) 4	0 / 21 (0.00%) 0
WHITE BLOOD CELL COUNT DECREASED subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Injury, poisoning and procedural complications INFUSION RELATED REACTION subjects affected / exposed occurrences (all)	1 / 3 (33.33%) 1	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Cardiac disorders SINUS TACHYCARDIA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Nervous system disorders DIZZINESS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	2 / 21 (9.52%) 2
FACIAL PARALYSIS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
HEADACHE subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 2	1 / 21 (4.76%) 1
HYPOAESTHESIA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 2	1 / 21 (4.76%) 1
NEUROPATHY PERIPHERAL subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 2	0 / 21 (0.00%) 0
PARAESTHESIA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 2	0 / 21 (0.00%) 0
POLYNEUROPATHY subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Blood and lymphatic system disorders			

ANAEMIA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	2 / 21 (9.52%)
occurrences (all)	0	1	2
ANAEMIA FOLATE DEFICIENCY			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
LEUKOPENIA			
subjects affected / exposed	0 / 3 (0.00%)	2 / 10 (20.00%)	1 / 21 (4.76%)
occurrences (all)	0	5	1
NEUTROPENIA			
subjects affected / exposed	1 / 3 (33.33%)	4 / 10 (40.00%)	3 / 21 (14.29%)
occurrences (all)	1	7	4
THROMBOCYTOPENIA			
subjects affected / exposed	0 / 3 (0.00%)	4 / 10 (40.00%)	0 / 21 (0.00%)
occurrences (all)	0	5	0
Eye disorders			
DRY EYE			
subjects affected / exposed	0 / 3 (0.00%)	2 / 10 (20.00%)	1 / 21 (4.76%)
occurrences (all)	0	2	1
GLAUCOMA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
SCLERITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
CATARACT NUCLEAR			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
VITREOUS DEGENERATION			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Gastrointestinal disorders			
ABDOMINAL PAIN			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	2	1
CONSTIPATION			

subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
DIARRHOEA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	3 / 10 (30.00%) 3	0 / 21 (0.00%) 0
DRY MOUTH subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	1 / 21 (4.76%) 1
DYSPEPSIA subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
GASTROESOPHAGEAL REFLUX DISEASE subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
STOMATITIS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
VOMITING subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Skin and subcutaneous tissue disorders NIGHT SWEATS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
RASH subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 4	0 / 21 (0.00%) 0
Renal and urinary disorders ACUTE KIDNEY INJURY subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	0 / 10 (0.00%) 0	2 / 21 (9.52%) 3
Endocrine disorders AUTOIMMUNE THYROIDITIS subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
HYPOTHYROIDISM			

subjects affected / exposed occurrences (all)	0 / 3 (0.00%) 0	1 / 10 (10.00%) 1	0 / 21 (0.00%) 0
Musculoskeletal and connective tissue disorders			
ARTHRALGIA			
subjects affected / exposed	0 / 3 (0.00%)	0 / 10 (0.00%)	2 / 21 (9.52%)
occurrences (all)	0	0	2
BACK PAIN			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
BONE PAIN			
subjects affected / exposed	1 / 3 (33.33%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	1	1	0
MUSCULOSKELETAL PAIN			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
SACROILIITIS			
subjects affected / exposed	1 / 3 (33.33%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences (all)	1	0	0
TENDONITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Infections and infestations			
BACTERAEMIA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	2	0
BACTERIAL INFECTION			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
BRONCHOPULMONARY ASPERGILLOSIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
CONJUNCTIVITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
CYTOMEGALOVIRUS INFECTION			

subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
LARYNGITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
NASOPHARYNGITIS			
subjects affected / exposed	1 / 3 (33.33%)	2 / 10 (20.00%)	0 / 21 (0.00%)
occurrences (all)	2	2	0
ORAL FUNGAL INFECTION			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	2	0
PELVIC ABSCESS			
subjects affected / exposed	1 / 3 (33.33%)	0 / 10 (0.00%)	0 / 21 (0.00%)
occurrences (all)	1	0	0
PNEUMONIA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
RHINITIS			
subjects affected / exposed	1 / 3 (33.33%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	4	1	0
SINUSITIS			
subjects affected / exposed	2 / 3 (66.67%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	2	1	0
UPPER RESPIRATORY TRACT INFECTION			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	3 / 21 (14.29%)
occurrences (all)	0	2	3
URINARY TRACT INFECTION			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	2	0
VIRAL INFECTION			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
VIRAL UPPER RESPIRATORY TRACT INFECTION			

subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
ACUTE SINUSITIS			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
CONJUNCTIVITIS VIRAL			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
Metabolism and nutrition disorders			
DECREASED APPETITE			
subjects affected / exposed	0 / 3 (0.00%)	3 / 10 (30.00%)	1 / 21 (4.76%)
occurrences (all)	0	3	1
FLUID RETENTION			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
HYPERGLYCAEMIA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	0 / 21 (0.00%)
occurrences (all)	0	1	0
HYPOALBUMINAEMIA			
subjects affected / exposed	0 / 3 (0.00%)	1 / 10 (10.00%)	1 / 21 (4.76%)
occurrences (all)	0	1	1
HYPOKALAEMIA			
subjects affected / exposed	0 / 3 (0.00%)	3 / 10 (30.00%)	1 / 21 (4.76%)
occurrences (all)	0	3	1

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
14 January 2016	Version 2: The protocol has been revised to update the DLT window and to clarify inclusion/exclusion and stopping criteria. Additional minor changes have been made to improve clarity and consistency.
28 June 2016	Version 3: The inclusion criterion on contraception requirements for women of childbearing potential has been revised to include the duration of contraception for atezolizumab and polatuzumab vedotin. Enrollment rules into the dose escalation phase have been updated, for patients' safety considerations. A sequential enrollment instead of a parallel enrollment will be used for each of the two dosing groups,
17 November 2016	Version 4: "Immune-mediated" was revised to "immune-related" when referring to adverse events. Cut-off date for Atezolizumab was changed to 15 December 2016. The title has been revised. Study treatment has been modified to include Rituximab (R00452294) in combination with atezolizumab plus polatuzumab vedotin in patients with relapsed or refractory diffuse large B-cell lymphoma.
04 May 2017	Version 5: The protocol has been modified to prohibit use of the term "sudden death" on the Adverse Event eCRF, unless it is combined with the presumed cause of death. Language has been added to clarify that the Sponsor will review all protocol deviations, and clarification has been added that prospective requests to deviate from the protocol are not allowed.
21 December 2017	Version 6: Clinical Data) has been updated with the most recent efficacy and safety results from Study G029383. Aligned the protocol with the current atezolizumab Investigator's Brochure, Version 10. A few exclusion criteria updated.
01 May 2018	Version 7: The study design and treatment schedule has been revised to reflect discontinuation of atezolizumab in patients still receiving study treatment. In addition, the post-induction (rituximab + atezolizumab consolidation) treatment phase has been removed from the treatment schedule (rituximab + atezolizumab + polatuzumab vedotin [R+Atezo+Pola] treatment group) of patients with RR DLBCL.
07 November 2018	Version 8: Lists of risks for atezolizumab-associated adverse events were revised to include nephritis. Regular Internal Monitoring Committee assessments stopped taking place as no new safety signals identified. Language was changed allowing patients still under treatment to enter the extension study. Medical Monitor information was updated. Survival follow-up period for assessment of new anti-lymphoma treatment re-added. PK sampling one year after last dose for polatuzumab vedotin was removed.

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

Limitations of the trial such as small numbers of subjects analysed or technical problems leading to unreliable data.

A safety signal was observed during the study in two patients in the R/R FL cohort. As a result, enrollment was permanently discontinued and atezolizumab was discontinued for all patients still receiving study treatment.

Notes: