



Clinical trial results:

A Phase 3, Open-Label, Multicenter Study of ALXN1210 in Children and Adolescents With Atypical Hemolytic Uremic Syndrome (aHUS)

Summary

EudraCT number	2016-002499-29
Trial protocol	GB DE AT SE ES BE IT FR
Global end of trial date	20 December 2022

Results information

Result version number	v2 (current)
This version publication date	02 July 2023
First version publication date	16 July 2022
Version creation reason	

Trial information

Trial identification

Sponsor protocol code	ALXN1210-aHUS-312
-----------------------	-------------------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT03131219
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	Alexion Pharmaceuticals Inc.
Sponsor organisation address	100 College Street, New Haven, CT, United States, 06510
Public contact	Alexion Europe SAS European Clinical Trial Information, Alexion Pharmaceuticals Inc., +33 787148158, clinicaltrials.eu@alexion.com
Scientific contact	Alexion Europe SAS European Clinical Trial Information, Alexion Pharmaceuticals Inc., +33 787148158, clinicaltrials.eu@alexion.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	Yes
EMA paediatric investigation plan number(s)	EMA-001943-PIP01-16
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	Yes

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	06 March 2020
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	20 December 2022
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

The purpose of the study is to assess the efficacy of ravulizumab to control disease activity in children and adolescents with aHUS who have not previously used a complement inhibitor (complement inhibitor treatment-naïve), as well as in complement inhibitor-experienced (eculizumab-experienced) adolescent participants.

Protection of trial subjects:

This trial was conducted in compliance with Good Clinical Practice.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	01 September 2017
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	United States: 14
Country: Number of subjects enrolled	Germany: 1
Country: Number of subjects enrolled	Korea, Republic of: 6
Country: Number of subjects enrolled	Spain: 5
Country: Number of subjects enrolled	Belgium: 2
Country: Number of subjects enrolled	Japan: 5
Country: Number of subjects enrolled	United Kingdom: 1
Worldwide total number of subjects	34
EEA total number of subjects	8

Notes:

Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	7
Children (2-11 years)	17

Adolescents (12-17 years)	10
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

Subject disposition

Recruitment

Recruitment details: -

Pre-assignment

Screening details:

This is an ongoing study and the data presented are the study results from the 26-week Initial Evaluation Period and data from the Extension Period through the data cutoff date at 03 Dec 2019.

Period 1

Period 1 title	Initial Evaluation Period
Is this the baseline period?	Yes
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
Arm title	Complement Inhibitor Treatment Naïve

Arm description:

Complement inhibitor treatment-naïve participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.

Arm type	Experimental
Investigational medicinal product name	Ravulizumab
Investigational medicinal product code	ALXN1210
Other name	
Pharmaceutical forms	Concentrate for solution for infusion
Routes of administration	Intravenous use

Dosage and administration details:

Participants received ravulizumab at prespecified dose and timepoints.

Arm title	Eculizumab Experienced
------------------	------------------------

Arm description:

Eculizumab-experienced participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.

Arm type	Experimental
Investigational medicinal product name	Ravulizumab
Investigational medicinal product code	ALXN1210
Other name	
Pharmaceutical forms	Concentrate for solution for infusion
Routes of administration	Intravenous use

Dosage and administration details:

Participants received ravulizumab at prespecified dose and timepoints.

Number of subjects in period 1	Complement Inhibitor Treatment Naïve	Eculizumab Experienced
Started	24	10
Received At Least 1 Dose of Study Drug	24	10
Completed	17	10
Not completed	7	0
Physician decision	1	-
Adverse event	2	-
Deemed ineligible post treatment	3	-
Protocol deviation	1	-

Period 2

Period 2 title	Extension Period
Is this the baseline period?	No
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
Arm title	Complement Inhibitor Treatment Naïve

Arm description:

Complement inhibitor treatment-naïve participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.

Arm type	Experimental
Investigational medicinal product name	Ravulizumab
Investigational medicinal product code	ALXN1210
Other name	
Pharmaceutical forms	Concentrate for solution for infusion
Routes of administration	Intravenous use

Dosage and administration details:

Participants received ALXN1210 at prespecified dose and timepoints.

Arm title	Eculizumab Experienced
------------------	------------------------

Arm description:

Eculizumab-experienced participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.

Arm type	Experimental
Investigational medicinal product name	Ravulizumab
Investigational medicinal product code	ALXN1210
Other name	
Pharmaceutical forms	Concentrate for solution for infusion
Routes of administration	Intravenous use

Dosage and administration details:

Participants received ALXN1210 at prespecified dose and timepoints.

Number of subjects in period 2	Complement Inhibitor Treatment Naïve	Eculizumab Experienced
Started	17	10
Received At Least 1 Dose of Study Drug	17	10
Completed	16	10
Not completed	1	0
Physician decision	1	-

Baseline characteristics

Reporting groups

Reporting group title	Complement Inhibitor Treatment Naïve
Reporting group description:	
Complement inhibitor treatment-naïve participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.	
Reporting group title	Ecuzumab Experienced
Reporting group description:	
Ecuzumab-experienced participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.	

Reporting group values	Complement Inhibitor Treatment Naïve	Ecuzumab Experienced	Total
Number of subjects	24	10	34
Age categorical Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	0	0	0
Infants and toddlers (28 days-23 months)	6	1	7
Children (2-11 years)	15	2	17
Adolescents (12-17 years)	3	7	10
Adults (18-64 years)	0	0	0
From 65-84 years	0	0	0
85 years and over	0	0	0
Age Continuous Units: years			
arithmetic mean	6.0	11.0	-
standard deviation	± 4.79	± 4.97	-
Sex: Female, Male Units: participants			
Female	14	1	15
Male	10	9	19
Baseline Estimated Glomerular Filtration Rate (eGFR) Units: mL/min/1.73 m ²			
arithmetic mean	26.4	104.90	-
standard deviation	± 21.17	± 29.545	-

End points

End points reporting groups

Reporting group title	Complement Inhibitor Treatment Naïve
Reporting group description: Complement inhibitor treatment-naïve participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.	
Reporting group title	Eculizumab Experienced
Reporting group description: Eculizumab-experienced participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.	
Reporting group title	Complement Inhibitor Treatment Naïve
Reporting group description: Complement inhibitor treatment-naïve participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.	
Reporting group title	Eculizumab Experienced
Reporting group description: Eculizumab-experienced participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.	

Primary: Percentage Of Complement Inhibitor Treatment-naïve Participants With Complete Thrombotic Microangiopathy (TMA) Response at Week 26

End point title	Percentage Of Complement Inhibitor Treatment-naïve Participants With Complete Thrombotic Microangiopathy (TMA) Response at Week 26 ^{[1][2]}
End point description: Complete TMA response during the 26-week Initial Evaluation Period is a composite endpoint that required normalization of hematological parameters (platelet count and lactate dehydrogenase) and improvement in kidney function ($\geq 25\%$ reduction in serum creatinine from baseline); for participants on dialysis, baseline was established at least 6 days after the end of dialysis. Participants had to meet these criteria for 2 separate assessments obtained at least 4 weeks apart, and any measurement in between. To be considered a responder, the latest time point a participant could first meet the response criteria was 28 days before the Week 26 (Day 183) assessment. Percentage based on the responders among treated participants. Full Analysis Set: all participants who received at least 1 dose of ravulizumab, had at least 1 post-baseline efficacy assessment, and met pre-specified eligibility criteria. Here, Overall 'Number of Participants Analyzed' = those who were evaluable for this endpoint.	
End point type	Primary
End point timeframe: Week 26	

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analysis was planned to be reported for this endpoint.

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Endpoint was planned to be reported for "Complement Inhibitor Treatment Naïve" arm only.

End point values	Complement Inhibitor Treatment Naïve			
Subject group type	Reporting group			
Number of subjects analysed	18			
Units: percentage of participants				
number (confidence interval 95%)				
Complete TMA response	77.8 (52.4 to 93.6)			
Platelet count normalization	94.4 (72.7 to 99.9)			
LDH normalization	88.9 (65.3 to 98.6)			
≥25% improvement in serum creatinine from baseline	83.3 (58.6 to 96.4)			

Statistical analyses

No statistical analyses for this end point

Secondary: Time To Complete TMA Response In Complement Inhibitor Treatment-naïve Participants

End point title	Time To Complete TMA Response In Complement Inhibitor Treatment-naïve Participants ^[3]
-----------------	---

End point description:

Participants that did not have a response were censored at the date of last visit or study discontinuation at the time when the analysis was performed. The time to complete TMA Response is reported in days. The time of the event of a confirmed complete TMA response was considered the first time point at which all the criteria for complete TMA response were met. Participants had to meet all complete TMA response criteria at 2 separate assessments obtained at least 4 weeks (28 days) apart, and any measurement in between. Full Analysis Set (FAS): all participants who received at least 1 dose of ravulizumab, had at least 1 post-baseline efficacy assessment, and met pre-specified eligibility criteria. Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this endpoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline through at least Week 52 and up to Week 111

Notes:

[3] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: Endpoint was planned to be reported for "Complement Inhibitor Treatment Naïve" arm only.

End point values	Complement Inhibitor Treatment Naïve			
Subject group type	Reporting group			
Number of subjects analysed	18			
Units: days				
median (inter-quartile range (Q1-Q3))	30.0 (22.0 to 88.0)			

Statistical analyses

No statistical analyses for this end point

Secondary: Proportion Of Complement Inhibitor Treatment-naïve Participants With Complete TMA Response at Week 52

End point title	Proportion Of Complement Inhibitor Treatment-naïve Participants With Complete TMA Response at Week 52 ^[4]
-----------------	--

End point description:

The proportion of participants considered responders, along with a 2-sided 95% CI based on exact confidence limits using the Clopper Pearson method is reported. Full Analysis Set (FAS): all participants who received at least 1 dose of ravulizumab, had at least 1 post-baseline efficacy assessment, met pre-specified eligibility criteria, and had evaluable data at specified timepoint. Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this endpoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Week 52

Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Endpoint was planned to be reported for "Complement Inhibitor Treatment Naïve" arm only.

End point values	Complement Inhibitor Treatment Naïve			
Subject group type	Reporting group			
Number of subjects analysed	18			
Units: proportion of participants				
number (confidence interval 95%)				
Complete TMA responder	0.882 (0.636 to 0.985)			
Platelet count normalization	0.882 (0.636 to 0.985)			
LDH normalization	1.000 (0.805 to 1.000)			
≥25% improvement in serum creatinine from baseline	1.000 (0.805 to 1.000)			

Statistical analyses

No statistical analyses for this end point

Secondary: Participants Who Do Not Require Dialysis at Weeks 26 and 52

End point title	Participants Who Do Not Require Dialysis at Weeks 26 and 52
-----------------	---

End point description:

For participants requiring dialysis within 5 days prior to ALXN1210 treatment initiation, the number of participants no longer requiring dialysis is reported. Full Analysis Set (FAS): all participants who received at least 1 dose of ravulizumab, had at least 1 post-baseline efficacy assessment, met pre-specified eligibility criteria, and had evaluable data at specified timepoint. Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this outcome measure and 'Number Analyzed' signifies participants evaluable for specified categories.

End point type	Secondary
----------------	-----------

End point timeframe:
Week 26 and Week 52

End point values	Complement Inhibitor Treatment Naïve	Eculizumab Experienced		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	6	0 ^[5]		
Units: participants				
Week 26	5			
Week 52	6			

Notes:

[5] - No participants were analyzed for this reporting group.

Statistical analyses

No statistical analyses for this end point

Secondary: Change From Baseline In eGFR At Weeks 26 and 52

End point title	Change From Baseline In eGFR At Weeks 26 and 52
-----------------	---

End point description:

Kidney function evaluated by eGFR was summarized at baseline and the Week 26 and Week 52 time points using descriptive statistics for continuous variables for the observed value, as well as the change from baseline. The baseline value was defined as the average of the values from the assessments performed prior to the first study drug infusion (these could include results from Screening and the Day 1 visit). A value of 10 mL/min/1.73 m² for eGFR was imputed for participants requiring dialysis for acute kidney injury. The observed value and change from baseline are reported in mL/min/1.73 m². An increase indicated improvement in kidney function. Analysis population was Full Analysis Set. Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this outcome measure and 'Number Analyzed' signifies participants evaluable for specified categories.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Week 26 and Week 52

End point values	Complement Inhibitor Treatment Naïve	Eculizumab Experienced		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	18	10		
Units: mL/min/1.73 m ²				
median (full range (min-max))				
Baseline (n=17, 10)	22.0 (10 to 84)	99.75 (54 to 136.5)		
Change From Baseline at Week 26 (n=17, 10)	80.0 (0 to 222)	-2.00 (-94 to 18)		
Change From Baseline at Week 52 (n=16, 10)	94.0 (10 to 230)	-3.00 (-20 to 9)		

Statistical analyses

No statistical analyses for this end point

Secondary: Participants With Change From Baseline In CKD Stage At Weeks 26 and 52

End point title	Participants With Change From Baseline In CKD Stage At Weeks 26 and 52
End point description: The CKD stage (classified based on the National Kidney Foundation CKD Stage) is presented as the change from baseline in the participants that Improved (excluding those with Stage 1 [normal renal function] at baseline as they cannot improve), Worsened (excluding those with Stage 5 at baseline as they cannot worsen), and Stayed the Same, compared to the CKD stage at baseline. Baseline was derived based on the last available eGFR before starting treatment. Stage 5 was considered the worst category, while Stage 1 was considered the best category. A 2-sided 95% CI for the proportion, based on exact confidence limits using the Clopper-Pearson method, was provided for each category. Analysis population was Full Analysis Set. Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this endpoint and 'Number Analyzed' signifies participants evaluable for specified categories.	
End point type	Secondary
End point timeframe: Baseline, Week 26, and Week 52	

End point values	Complement Inhibitor Treatment Naïve	Eculizumab Experienced		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	18	10		
Units: participants				
Week 26, Improved (n=17, 10)	15	0		
Week 26, Worsened (n=17, 10)	0	3		
Week 26, Stayed the Same (n=17, 10)	2	7		
Week 52, Improved (n=16, 10)	16	0		
Week 52, Worsened (n=11, 10)	0	0		
Week 52, Stayed the Same (n=16, 10)	0	10		

Statistical analyses

No statistical analyses for this end point

Secondary: Change From Baseline In Platelet Count At Weeks 26 and 52

End point title	Change From Baseline In Platelet Count At Weeks 26 and 52
-----------------	---

End point description:

The hematologic TMA parameter of platelet count was summarized at baseline and at Week 26 and Week 52 using descriptive statistics for continuous variables for the change from baseline. Results are reported in platelets*10⁹/liter (L) blood. Full Analysis Set (FAS): all participants who received at least 1 dose of ravulizumab, had at least 1 post-baseline efficacy assessment, met pre-specified eligibility criteria, and had evaluable data at the specified time point (Week 26 or Week 52). Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this endpoint and 'Number Analyzed' signifies participants evaluable for specified categories.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Week 26 and Week 52

End point values	Complement Inhibitor Treatment Naïve	Eculizumab Experienced		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	18	10		
Units: platelets*10 ⁹ /L				
median (full range (min-max))				
Baseline (n=18, 10)	51.25 (14 to 125)	281.75 (207 to 415.5)		
Change from Baseline at Week 26 (n=18, 10)	247.00 (57.5 to 368.5)	-2.25 (-74.5 to 123.5)		
Change from Baseline at Week 52 (n=17, 10)	213.00 (19.5 to 471.5)	-34.75 (-109 to 109)		

Statistical analyses

No statistical analyses for this end point

Secondary: Change From Baseline In LDH At Weeks 26 and 52

End point title	Change From Baseline In LDH At Weeks 26 and 52
-----------------	--

End point description:

The hematologic TMA parameter of serum LDH was summarized at baseline and at Week 26 and Week 52 using descriptive statistics for continuous variables for the change from baseline. Results are reported in units (U)/L serum. Full Analysis Set (FAS): all participants who received at least 1 dose of ravulizumab, had at least 1 post-baseline efficacy assessment, met pre-specified eligibility criteria, and had evaluable data at the specified timepoint (Week 26 or Week 52). Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this endpoint and 'Number Analyzed' signifies participants evaluable for specified categories.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Week 26 and Week 52

End point values	Complement Inhibitor Treatment Naïve	Eculizumab Experienced		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	18	10		
Units: U/L				
median (full range (min-max))				
Baseline (n=17, 10)	1963.00 (772 to 4985)	206.50 (138.5 to 356)		
Change From Baseline at Week 26 (n=16, 10)	-1851.50 (-4713 to -513)	-8.50 (-50.5 to 50.5)		
Change From Baseline at Week 52 (n=16, 10)	-1825.50 (-4724 to -579)	-17.50 (-34.5 to 29.5)		

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage Of Complement Inhibitor Treatment-naïve Participants With An Increase From Baseline In Hemoglobin ≥ 20 g/L Through Week 26 and Week 52

End point title	Percentage Of Complement Inhibitor Treatment-naïve Participants With An Increase From Baseline In Hemoglobin ≥ 20 g/L Through Week 26 and Week 52 ^[6]
-----------------	---

End point description:

The percentage of participants with an increase from baseline in hemoglobin ≥ 20 g/L, observed at 2 separate assessments obtained at least 4 weeks (28 days) apart, and any measurement in between, was assessed through Week 26 and Week 52 and is presented as the percentage of responders, along with a 2-sided 95% CI. The 95% CIs are based on exact confidence limits using the Clopper-Pearson method. To be considered a responder during the 26-week and 52-week Extension Periods, the latest time point a participant could first meet the response criteria was 28 days before the respective Week 26 and Week 52 assessments (components of the response maintained for at least 28 days). Analysis population was Full Analysis Set. Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this endpoint.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline through Week 26 and through Week 52

Notes:

[6] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: Endpoint was planned to be reported for "Complement Inhibitor Treatment Naïve" arm only.

End point values	Complement Inhibitor Treatment Naïve			
Subject group type	Reporting group			
Number of subjects analysed	18			
Units: percentage of participants				
number (confidence interval 95%)				
Week 26	100 (80.5 to 100)			
Week 52	94.1 (71.3 to 99.9)			

Statistical analyses

No statistical analyses for this end point

Secondary: Change From Baseline In Hemoglobin At Weeks 26 and 52

End point title	Change From Baseline In Hemoglobin At Weeks 26 and 52
-----------------	---

End point description:

The hematologic TMA parameter of hemoglobin level was summarized at baseline and at Week 26 and Week 52 using descriptive statistics for continuous variables for the change from baseline. Results are reported in grams (g)/L blood. Full Analysis Set (FAS): all participants who received at least 1 dose of ravulizumab, had at least 1 post-baseline efficacy assessment, met pre-specified eligibility criteria, and had evaluable data at the specified timepoint (Week 26 or Week 52). Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this endpoint and 'Number Analyzed' signifies participants evaluable for specified categories.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Week 26 and Week 52

End point values	Complement Inhibitor Treatment Naïve	Eculizumab Experienced		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	18	10		
Units: g/L				
median (full range (min-max))				
Baseline (n=18, 10)	74.25 (32 to 106)	132.00 (114.5 to 148)		
Change From Baseline at Week 26 (n=15, 10)	46.50 (26.5 to 86)	-3.50 (-19.5 to 8)		
Change From Baseline at Week 52 (n=17, 10)	51.50 (-19 to 80)	5.50 (-7.5 to 13.5)		

Statistical analyses

No statistical analyses for this end point

Secondary: Change From Baseline In Quality Of Life As Measured By The Functional Assessment Of Chronic Illness Therapy (FACIT)-Fatigue Version 4 Questionnaire (Participants ≥5 Years Of Age) At Weeks 26 and 52

End point title	Change From Baseline In Quality Of Life As Measured By The Functional Assessment Of Chronic Illness Therapy (FACIT)-Fatigue Version 4 Questionnaire (Participants ≥5 Years Of Age) At Weeks 26 and 52
-----------------	---

End point description:

Quality of life was assessed in participants >5 years of age by the Pediatric FACIT-Fatigue Questionnaire (reported by participants who were ≥8 years of age at the time of enrollment; caregiver reported or caregiver assistance for participants who were 5 to <8 years of age at the time of enrollment). The FACIT Fatigue data were summarized at baseline and each post baseline time point using descriptive statistics for continuous variables for the observed value as well as the change from baseline. The FACIT Fatigue Version 4 questionnaire at baseline and each post-infusion time point was scored using standard scoring algorithms. The score ranges from 0 to 52, with a higher score indicating less fatigue. An increase in score indicated an improvement in quality of life. Analysis population was Full Analysis Set. Here, Overall 'Number of Participants Analyzed' signifies those who were evaluable for this outcome measure.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Week 26 and Week 52

End point values	Complement Inhibitor Treatment Naïve	Eculizumab Experienced		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	9	8		
Units: units on a scale				
median (full range (min-max))				
Baseline	35.00 (4 to 44)	50.00 (42 to 52)		
Change From Baseline at Week 26	10.00 (4 to 48)	0.00 (-5 to 3)		
Change From Baseline at Week 52	9.00 (3 to 47)	-1.00 (-7 to 2)		

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

From the beginning of the initial evaluation period (Day 1) through data cutoff (at least 52 weeks and up to a maximum of 111 weeks of treatment, representing 36.2 patient-years of exposure).

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	25.1
--------------------	------

Reporting groups

Reporting group title	Complement Inhibitor Treatment Naïve
-----------------------	--------------------------------------

Reporting group description:

Complement inhibitor treatment-naïve participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.

Reporting group title	Eculizumab Experienced
-----------------------	------------------------

Reporting group description:

Eculizumab-experienced participants received weight-based doses of ravulizumab during the 26-week Initial Evaluation Period. After the Initial Evaluation Period, participants rolled over into an Extension Period in which all participants continued their weight-based maintenance dose of ravulizumab.

Serious adverse events	Complement Inhibitor Treatment Naïve	Eculizumab Experienced	
Total subjects affected by serious adverse events			
subjects affected / exposed	16 / 24 (66.67%)	1 / 10 (10.00%)	
number of deaths (all causes)	0	0	
number of deaths resulting from adverse events			
Injury, poisoning and procedural complications			
Subdural haematoma			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vascular disorders			
Hypertension			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hypertensive urgency			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Congenital, familial and genetic disorders			
Atrial septal defect			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
Headache			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Posterior reversible encephalopathy syndrome			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	4 / 24 (16.67%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	2 / 6	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			
Diarrhoea			
subjects affected / exposed	3 / 24 (12.50%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Abdominal pain			

subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Vomiting			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pancreatitis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Respiratory, thoracic and mediastinal disorders			
Pulmonary haemorrhage			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal and urinary disorders			
Glomerulonephritis membranoproliferative			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Proteinuria			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Musculoskeletal and connective tissue disorders			
Myalgia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Juvenile idiopathic arthritis			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
Bronchitis			
subjects affected / exposed	1 / 24 (4.17%)	1 / 10 (10.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis viral			
subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cytomegalovirus enteritis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Escherichia bacteraemia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Escherichia pyelonephritis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Human bocavirus infection			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pharyngitis			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pneumonia			
subjects affected / exposed	1 / 24 (4.17%)	1 / 10 (10.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Viral infection			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Viral pharyngitis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastroenteritis rotavirus			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Upper respiratory tract infection			
subjects affected / exposed	1 / 24 (4.17%)	1 / 10 (10.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin infection			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 2	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Pyelonephritis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urinary tract infection			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	1 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Metabolism and nutrition disorders			
Decreased appetite			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 0 %

Non-serious adverse events	Complement Inhibitor Treatment Naïve	Eculizumab Experienced	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	22 / 24 (91.67%)	10 / 10 (100.00%)	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Melanocytic naevus			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Vascular disorders			
Hypertension			
subjects affected / exposed	6 / 24 (25.00%)	1 / 10 (10.00%)	
occurrences (all)	7	1	
Hypotension			
subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Hot flush			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	12 / 24 (50.00%)	1 / 10 (10.00%)	
occurrences (all)	25	1	
Fatigue			

subjects affected / exposed	2 / 24 (8.33%)	1 / 10 (10.00%)	
occurrences (all)	3	1	
Oedema			
subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Device related thrombosis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Vascular device occlusion			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Dysplasia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Face oedema			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Peripheral swelling			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Vaccination site reaction			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Pain			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Chills			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Immune system disorders			
Drug hypersensitivity			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	4	0	
Reproductive system and breast disorders			

Penile erythema			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Respiratory, thoracic and mediastinal disorders			
Sinus disorder			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Rhinitis allergic			
subjects affected / exposed	1 / 24 (4.17%)	1 / 10 (10.00%)	
occurrences (all)	1	1	
Epistaxis			
subjects affected / exposed	2 / 24 (8.33%)	2 / 10 (20.00%)	
occurrences (all)	5	2	
Oropharyngeal pain			
subjects affected / exposed	2 / 24 (8.33%)	3 / 10 (30.00%)	
occurrences (all)	3	3	
Productive cough			
subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Dyspnoea			
subjects affected / exposed	3 / 24 (12.50%)	0 / 10 (0.00%)	
occurrences (all)	3	0	
Nasal congestion			
subjects affected / exposed	4 / 24 (16.67%)	0 / 10 (0.00%)	
occurrences (all)	4	0	
Rhinorrhoea			
subjects affected / exposed	4 / 24 (16.67%)	0 / 10 (0.00%)	
occurrences (all)	7	0	
Cough			
subjects affected / exposed	5 / 24 (20.83%)	2 / 10 (20.00%)	
occurrences (all)	10	2	
Wheezing			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Bronchospasm			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Pharyngeal erythema			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Pulmonary oedema			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Pleural effusion			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Tachypnoea			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Upper-airway cough syndrome			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Psychiatric disorders			
Confusional state			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Depressed mood			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Insomnia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Attention deficit hyperactivity disorder			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Product issues			
Device occlusion			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Device malfunction			

subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Investigations			
Influenza A virus test positive subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Vitamin D decreased subjects affected / exposed occurrences (all)	3 / 24 (12.50%) 5	0 / 10 (0.00%) 0	
Clostridium test positive subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Lymphocyte count decreased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 3	0 / 10 (0.00%) 0	
Amylase increased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Salmonella test positive subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
White blood cell count increased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Blood iron decreased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Blood potassium increased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Crystal urine present subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Neutrophil count increased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	

Transaminases increased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Weight decreased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
White blood cell count decreased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Neutrophil count decreased subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 6	0 / 10 (0.00%) 0	
Injury, poisoning and procedural complications			
Contusion subjects affected / exposed occurrences (all)	5 / 24 (20.83%) 10	1 / 10 (10.00%) 1	
Head injury subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	0 / 10 (0.00%) 0	
Skin abrasion subjects affected / exposed occurrences (all)	3 / 24 (12.50%) 3	0 / 10 (0.00%) 0	
Ligament sprain subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	0 / 10 (0.00%) 0	
Arthropod sting subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 4	0 / 10 (0.00%) 0	
Limb injury subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 2	0 / 10 (0.00%) 0	
Eye injury subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Radial head dislocation			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Thermal burn			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Torus fracture			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Wound			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Lip injury			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Scratch			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Traumatic haematoma			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Cardiac disorders			
Cardiomegaly			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Left ventricular hypertrophy			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Angina pectoris			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Tachycardia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Ventricular extrasystoles			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	

Nervous system disorders			
Headache			
subjects affected / exposed	8 / 24 (33.33%)	1 / 10 (10.00%)	
occurrences (all)	26	1	
Dizziness			
subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences (all)	3	0	
Central nervous system lesion			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Seizure			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Cerebral disorder			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Lethargy			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Paraesthesia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Blood and lymphatic system disorders			
Lymphadenopathy			
subjects affected / exposed	2 / 24 (8.33%)	1 / 10 (10.00%)	
occurrences (all)	2	1	
Anaemia			
subjects affected / exposed	3 / 24 (12.50%)	0 / 10 (0.00%)	
occurrences (all)	6	0	
Neutropenia			
subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Macrocytosis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Increased tendency to bruise			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Leukopenia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Ear and labyrinth disorders			
Ear pain			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
External ear inflammation			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Eye disorders			
Photophobia			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Lacrimation increased			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Eye discharge			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Amblyopia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Hypermetropia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Periorbital oedema			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Strabismus			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Conjunctival oedema			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Eye pain			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Ocular hyperaemia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Vision blurred			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Gastrointestinal disorders			
Vomiting			
subjects affected / exposed	8 / 24 (33.33%)	2 / 10 (20.00%)	
occurrences (all)	30	3	
Dyspepsia			
subjects affected / exposed	1 / 24 (4.17%)	1 / 10 (10.00%)	
occurrences (all)	3	1	
Diarrhoea			
subjects affected / exposed	7 / 24 (29.17%)	2 / 10 (20.00%)	
occurrences (all)	14	2	
Abdominal pain			
subjects affected / exposed	5 / 24 (20.83%)	1 / 10 (10.00%)	
occurrences (all)	10	1	
Abdominal distension			
subjects affected / exposed	2 / 24 (8.33%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Nausea			
subjects affected / exposed	4 / 24 (16.67%)	0 / 10 (0.00%)	
occurrences (all)	12	0	
Constipation			
subjects affected / exposed	5 / 24 (20.83%)	0 / 10 (0.00%)	
occurrences (all)	9	0	
Pancreatitis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	

Abdominal pain upper			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Gastrointestinal disorder			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Acetonaemic vomiting			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Breath odour			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Gingival bleeding			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Mouth haemorrhage			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Rectal prolapse			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Duodenal ulcer			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Faecaloma			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Gastric ulcer			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Mouth ulceration			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Odynophagia			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	2	

Lip dry subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Tooth disorder subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Skin and subcutaneous tissue disorders			
Dermatitis diaper subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	0 / 10 (0.00%) 0	
Rash subjects affected / exposed occurrences (all)	4 / 24 (16.67%) 5	0 / 10 (0.00%) 0	
Erythema subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Asteatosis subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Dry skin subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 5	0 / 10 (0.00%) 0	
Skin exfoliation subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 2	0 / 10 (0.00%) 0	
Alopecia subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 2	0 / 10 (0.00%) 0	
Dermatitis contact subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Hyperhidrosis subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Keratosis pilaris			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Scab			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Skin hypopigmentation			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Erythema multiforme			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Nail discolouration			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Rash papular			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Miliaria			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Pruritus			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Eczema			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Renal and urinary disorders			
Acute kidney injury			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Chronic kidney disease			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Hydronephrosis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	

Proteinuria			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Haematuria			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Endocrine disorders			
Hypothyroidism			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Musculoskeletal and connective tissue disorders			
Tenosynovitis			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Pain in extremity			
subjects affected / exposed	2 / 24 (8.33%)	1 / 10 (10.00%)	
occurrences (all)	2	2	
Osteochondrosis			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Back pain			
subjects affected / exposed	3 / 24 (12.50%)	1 / 10 (10.00%)	
occurrences (all)	4	1	
Myalgia			
subjects affected / exposed	3 / 24 (12.50%)	0 / 10 (0.00%)	
occurrences (all)	7	0	
Arthralgia			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	2	0	
Mobility decreased			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Juvenile idiopathic arthritis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Osteopenia			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Tendonitis			
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	
Infections and infestations			
Nasopharyngitis			
subjects affected / exposed	7 / 24 (29.17%)	2 / 10 (20.00%)	
occurrences (all)	14	2	
Otitis media			
subjects affected / exposed	1 / 24 (4.17%)	2 / 10 (20.00%)	
occurrences (all)	1	3	
Viral infection			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	2	
Pneumonia			
subjects affected / exposed	2 / 24 (8.33%)	1 / 10 (10.00%)	
occurrences (all)	2	1	
Influenza			
subjects affected / exposed	1 / 24 (4.17%)	1 / 10 (10.00%)	
occurrences (all)	1	1	
Impetigo			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	1	
Hand-foot-and-mouth disease			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	2	
Bronchitis			
subjects affected / exposed	0 / 24 (0.00%)	1 / 10 (10.00%)	
occurrences (all)	0	4	
Viral upper respiratory tract infection			
subjects affected / exposed	2 / 24 (8.33%)	2 / 10 (20.00%)	
occurrences (all)	5	3	
Pharyngitis			
subjects affected / exposed	2 / 24 (8.33%)	3 / 10 (30.00%)	
occurrences (all)	2	3	

Upper respiratory tract infection subjects affected / exposed occurrences (all)	4 / 24 (16.67%) 6	4 / 10 (40.00%) 23
Sinusitis subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	0 / 10 (0.00%) 0
Conjunctivitis subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	0 / 10 (0.00%) 0
Tonsillitis subjects affected / exposed occurrences (all)	3 / 24 (12.50%) 4	0 / 10 (0.00%) 0
Rhinovirus infection subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 5	0 / 10 (0.00%) 0
Otitis externa subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 3	0 / 10 (0.00%) 0
Gastroenteritis subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	2 / 10 (20.00%) 2
COVID-19 subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	0 / 10 (0.00%) 0
Pharyngotonsillitis subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 4	1 / 10 (10.00%) 1
Oral herpes subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 3	0 / 10 (0.00%) 0
Ear infection subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 2	0 / 10 (0.00%) 0
Body tinea subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 2	0 / 10 (0.00%) 0

Skin infection		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	2	0
Vaginal infection		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	2	0
Enterobiasis		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Gastroenteritis viral		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Laryngitis		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Molluscum contagiosum		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Myringitis		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Pharyngitis streptococcal		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Urinary tract infection		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Cellulitis		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Cytomegalovirus enteritis		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0
Fungal skin infection		
subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)
occurrences (all)	1	0

Soft tissue infection subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Croup infectious subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 2	1 / 10 (10.00%) 2	
Onychomycosis subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Tinea pedis subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Metabolism and nutrition disorders			
Dehydration subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 4	2 / 10 (20.00%) 2	
Iron deficiency subjects affected / exposed occurrences (all)	3 / 24 (12.50%) 3	0 / 10 (0.00%) 0	
Decreased appetite subjects affected / exposed occurrences (all)	2 / 24 (8.33%) 2	0 / 10 (0.00%) 0	
Hypoglycaemia subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Hypercholesterolaemia subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Vitamin D deficiency subjects affected / exposed occurrences (all)	1 / 24 (4.17%) 1	0 / 10 (0.00%) 0	
Underweight subjects affected / exposed occurrences (all)	0 / 24 (0.00%) 0	1 / 10 (10.00%) 1	
Acetonaemia			

subjects affected / exposed	1 / 24 (4.17%)	0 / 10 (0.00%)	
occurrences (all)	1	0	

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
23 August 2018	<ul style="list-style-type: none">•The loading dose for patients 5 to < 10 kg was increased from 300 milligrams (mg) to 600 mg.•The entry criteria were revised to allow enrollment into Cohort 2 of adolescent participants previously treated with eculizumab for at least the past 90 days.•Revised the entry criterion to allow LDH and hemoglobin results obtained during the Screening Period or within 28 days prior to the start of the Screening Period.•New objectives and endpoints were added to evaluate the safety and efficacy of ALXN1210 in Cohort 2.•Added 2 interim analyses: 1) when 12 to 14 complement inhibitor treatment-naïve participants (i.e., Cohort 1) have completed or withdrawn from the end of the 26-week Initial Evaluation Period; and 2) when all study participants have completed or withdrawn from the 26-week Initial Evaluation Period.•The Screening Period was clarified as 28 days for Cohort 2.•Added a provision to allow a supplemental dose of ALXN1210 to be administered to a participant if the Investigator and Sponsor mutually agree that a participant will potentially benefit.•Added a provision to allow for a dose to be administered as 2 separate infusions no more than approximately 24 hours apart if the Investigator and Sponsor mutually agree that the infusion volume (120 mL) of the loading dose for patients ≥ 5 to < 10 kg (600 mg) was too high for an individual participant.•The study sample size was increased to align with the planned sample size for each age category.• Statistical language was clarified to indicate that the analyses for Cohort 1 and Cohort 2 would be conducted and reported separately.•To reduce the participant data collection burden, removed the exploratory endpoints of Additional Signs or Symptoms of aHUS and Healthcare Resource Utilization.
16 July 2019	<ul style="list-style-type: none">•Increased duration of the Extension Period from 2 years to 4.5 years or until the product is registered or approved (in accordance with country-specific regulations), whichever occurs first, to allow additional safety, pharmacokinetic/pharmacodynamic (PK/PD), and immunogenicity evaluations.•Revised schedule of assessments to align with the increased duration of the Extension Period.•To improve clarity, to differentiate early termination of participants from study versus discontinuation of ravulizumab treatment with continuation in the study (for monitoring visits) in the Extension Period.•Added criteria on TMA recurrence and guidance on retreatment with ravulizumab for participants who discontinue ravulizumab and remain in the study in the Extension Period.•Expanded secondary efficacy and PK/PD endpoints to include evaluation of participants who discontinue study drug as well as those who resume treatment.•The language in the Data Monitoring Committee was aligned with other protocols at a program level.

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported