



## Clinical trial results:

### Clinical assessment of fluticasone propionate/ salmeterol xinafoate HFA MDI in 6-month to 4-year-old Japanese patients with bronchial asthma

#### Summary

EudraCT number	2016-003479-22
Trial protocol	Outside EU/EEA
Global end of trial date	03 October 2016

#### Results information

Result version number	v2 (current)
This version publication date	13 April 2017
First version publication date	21 December 2016
Version creation reason	

#### Trial information

##### Trial identification

Sponsor protocol code	200860
-----------------------	--------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	-
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	GlaxoSmithKline
Sponsor organisation address	980 Great West Road, Brentford, Middlesex, United Kingdom,
Public contact	GSK Response Center, GlaxoSmithKline, 1 866-435-7343,
Scientific contact	GSK Response Center, GlaxoSmithKline, 1 866-435-7343,

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

---

**Results analysis stage**

Analysis stage	Final
Date of interim/final analysis	16 December 2016
Is this the analysis of the primary completion data?	Yes
Primary completion date	14 June 2016
Global end of trial reached?	Yes
Global end of trial date	03 October 2016
Was the trial ended prematurely?	No

Notes:

---

**General information about the trial**

Main objective of the trial:

To evaluate the efficacy and safety of fluticasone propionate (FP)/ salmeterol xinafoate (SLM) hydrofluoroalkane (HFA) MDI 50/25 µg 1 or 2 inhalation bid for 8 weeks in comparison with FP HFA MDI 50 µg 1 or 2 inhalation bid in 6-month to 4-year-old Japanese patients with bronchial asthma.

Protection of trial subjects:

Not applicable

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	26 May 2014
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

---

**Population of trial subjects**

---

**Subjects enrolled per country**

Country: Number of subjects enrolled	Japan: 370
Worldwide total number of subjects	370
EEA total number of subjects	0

Notes:

---

**Subjects enrolled per age group**

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	67
Children (2-11 years)	303
Adolescents (12-17 years)	0
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

Study evaluated the efficacy and safety of fluticasone propionate (FP)/salmeterol xinafoate hydrofluoroalkane (SLM HFA) twice-daily (BID) via metered-dose inhaler (MDI) for 8 weeks in comparison with FP HFA in 6-months to 4-years-old Japanese participants (par.) with infantile bronchial asthma.

### Pre-assignment

Screening details:

Eligible par. at screening entered a 2-week run-in period to receive FP HFA MDI 50 µg, followed by 8-week double-blind treatment period (TP) 1 to receive FP/SLM HFA MDI 50/25 µg or FP HFA MDI 50 µg. In TP2, par. received FP/SLM HFA MDI 50/25 µg for 16 weeks (open-label phase). The total duration of the study was 27 weeks with follow-up.

### Period 1

Period 1 title	Period 1: 8 weeks
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Investigator, Monitor, Subject, Data analyst

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	FP HFA 50 µg

Arm description:

In TP1, participants were randomized to receive one or two inhalations of FP HFA 50 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.

Arm type	Active comparator
Investigational medicinal product name	Fluticasone propionate (FP) HFA
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Inhalation solution
Routes of administration	Inhalation use

Dosage and administration details:

FP HFA was administered via pressurized Metered-Dose Inhaler at 50 µg BID (one or two inhalations given using AeroChamber Plus with face mask) for 2 weeks in run-in period and 8 weeks in TP1.

<b>Arm title</b>	FP/SLM HFA 50/25 µg
------------------	---------------------

Arm description:

In TP1, participants were randomized to receive one or two inhalations of FP/SLM HFA 50/25 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.

Arm type	Experimental
Investigational medicinal product name	Fluticasone propionate (FP)/salmeterol xinafoate (SLM) HFA
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Inhalation solution
Routes of administration	Inhalation use

Dosage and administration details:

FP/SLM HFA was administered via pressurized Metered-Dose Inhaler at 50/25µg BID (one or two inhalations given using AeroChamber Plus with face mask) for 16 weeks in TP2.

Number of subjects in period 1 <sup>[1]</sup>	FP HFA 50 µg	FP/SLM HFA 50/25 µg
Started	150	150
Completed	142	148
Not completed	8	2
Consent withdrawn by subject	1	-
Adverse event, non-fatal	1	-
Par. Reached Stopping Criteria	5	2
Protocol deviation	1	-

Notes:

[1] - The number of subjects reported to be in the baseline period are not the same as the worldwide number enrolled in the trial. It is expected that these numbers will be the same.

Justification: A total of 370 participants were enrolled of which 300 participants were randomized.

## Period 2

Period 2 title	Period 2: 16 weeks
Is this the baseline period?	No
Allocation method	Randomised - controlled
Blinding used	Not blinded

## Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	FP 50 µg - FP/SLM 50/25 µg

Arm description:

1 or 2 inhalations of FP/SLM HFA MDI 50/25 µg were administered twice daily in TP2 to those participants who received FP HFA MDI 50 µg in TP1.

Arm type	Experimental
Investigational medicinal product name	Fluticasone propionate (FP)/salmeterol xinafoate (SLM) HFA
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Inhalation solution
Routes of administration	Inhalation use

Dosage and administration details:

FP/SLM HFA was administered via pressurized Metered-Dose Inhaler at 50/25µg BID (one or two inhalations given using AeroChamber Plus with face mask) for 16 weeks in TP2.

<b>Arm title</b>	FP/SLM 50/25 µg - FP/SLM 50/25 µg
------------------	-----------------------------------

Arm description:

1 or 2 inhalations of FP/SLM HFA MDI 50/25 µg were administered twice daily in TP2 to those participants who received FP/SLM HFA MDI 50/25 µg in TP1.

Arm type	Experimental
Investigational medicinal product name	Fluticasone propionate (FP)/salmeterol xinafoate (SLM) HFA
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Inhalation solution
Routes of administration	Inhalation use

Dosage and administration details:

FP/SLM HFA was administered via pressurized Metered-Dose Inhaler at 50/25µg BID (one or two inhalations given using AeroChamber Plus with face mask) for 16 weeks in TP2.

<b>Number of subjects in period 2<sup>[2]</sup></b>	<b>FP 50 µg - FP/SLM 50/25 µg</b>	<b>FP/SLM 50/25 µg - FP/SLM 50/25 µg</b>
Started	141	147
Completed	132	136
Not completed	9	11
Physician decision	2	-
Consent withdrawn by subject	1	1
Adverse event, non-fatal	4	5
Par. Reached Stopping Criteria	2	5

---

Notes:

[2] - The number of subjects starting the period is not consistent with the number completing the preceding period. It is expected the number of subjects starting the subsequent period will be the same as the number completing the preceding period.

Justification: The number of participants that started Treatment Period 2 are not equal to the number that completed Treatment Period 1 due to one participant in each treatment arm didn't transfer to Treatment Period 2.

## Baseline characteristics

### Reporting groups

Reporting group title	FP HFA 50 µg
-----------------------	--------------

Reporting group description:

In TP1, participants were randomized to receive one or two inhalations of FP HFA 50 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.

Reporting group title	FP/SLM HFA 50/25 µg
-----------------------	---------------------

Reporting group description:

In TP1, participants were randomized to receive one or two inhalations of FP/SLM HFA 50/25 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.

Reporting group values	FP HFA 50 µg	FP/SLM HFA 50/25 µg	Total
Number of subjects	150	150	300
Age categorical Units: Subjects			

Age continuous			
Age continuous description			
Units: months			
arithmetic mean	38.4	40.5	
standard deviation	± 14.1	± 14.07	-
Gender categorical			
Gender categorical description			
Units: Subjects			
Female	60	55	115
Male	90	95	185
Race/Ethnicity, Customized			
Units: Subjects			
Asian - Japanese Heritage	150	150	300

## End points

### End points reporting groups

Reporting group title	FP HFA 50 µg
Reporting group description: In TP1, participants were randomized to receive one or two inhalations of FP HFA 50 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.	
Reporting group title	FP/SLM HFA 50/25 µg
Reporting group description: In TP1, participants were randomized to receive one or two inhalations of FP/SLM HFA 50/25 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.	
Reporting group title	FP 50 µg - FP/SLM 50/25 µg
Reporting group description: 1 or 2 inhalations of FP/SLM HFA MDI 50/25 µg were administered twice daily in TP2 to those participants who received FP HFA MDI 50 µg in TP1.	
Reporting group title	FP/SLM 50/25 µg - FP/SLM 50/25 µg
Reporting group description: 1 or 2 inhalations of FP/SLM HFA MDI 50/25 µg were administered twice daily in TP2 to those participants who received FP/SLM HFA MDI 50/25 µg in TP1.	

### Primary: Mean change from Baseline in total asthma symptom score (daytime plus night time) at the end of the Treatment Period 1 (TP1)

End point title	Mean change from Baseline in total asthma symptom score (daytime plus night time) at the end of the Treatment Period 1 (TP1)
End point description: The participant's parent or legally acceptable representative made entries asthma symptom experienced by the participant in a patient diary twice daily (day time and night time) in the form of scores on a 4-point rating scale from Baseline (Week -1) until end of TP1 (Week 8). Scores ranged from 0 to 3(0: one, 1: mild, 2: moderate, 3: severe) and maximum score is 6 per day. The Baseline value is a mean value of the last 7 consecutive days during the run-in period (excluding the day of Visit 2 [Randomization]). The end of the TP1 value is a mean value of the last 7 consecutive days during the TP1 (excluding the last day of the TP1). Change from Baseline is the difference between the value of the endpoint at the time point of interest and the Baseline value. Participants who completed TP1 and completed their diary were analyzed.	
End point type	Primary
End point timeframe: Baseline and Week 8	

End point values	FP HFA 50 µg	FP/SLM HFA 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	142 <sup>[1]</sup>	148 <sup>[2]</sup>		
Units: Scores on a scale				
least squares mean (standard error)	-3.01 (± 0.545)	-3.97 (± 0.534)		

Notes:

[1] - ITT Population: all randomized par. who received at least one dose of study medication.

[2] - ITT Population: all randomized par. who received at least one dose of study medication.

## Statistical analyses

<b>Statistical analysis title</b>	Statistical analysis 1
Comparison groups	FP HFA 50 µg v FP/SLM HFA 50/25 µg
Number of subjects included in analysis	290
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.206
Method	ANCOVA
Parameter estimate	Difference in Least square means
Point estimate	-0.97
Confidence interval	
level	95 %
sides	2-sided
lower limit	-2.47
upper limit	0.54

## Secondary: Mean change from Baseline in night-time asthma symptoms score at the end of Treatment Period 1 (TP1)

End point title	Mean change from Baseline in night-time asthma symptoms score at the end of Treatment Period 1 (TP1)
End point description: The participant's parent or legally acceptable representative recorded asthma symptoms experienced by the participant during the night in a patient diary in the form of scores on a 4-point rating scale from Baseline (Week -1) until end of TP1 (Week 8). Scores ranged from 0 to 3(0: one, 1: mild, 2: moderate, 3: severe) and maximum score is 3 per day. Change from Baseline in the asthma symptom scores at night time at the end of TP1 was analyzed. The Baseline value is a mean value of the last 7 consecutive days during the run-in period (excluding the day of Visit 2 [Randomization]). The end of the TP1 value is a mean value of the last 7 consecutive days during the TP1 (excluding the last day of the TP1). Change from Baseline is the difference between the value of the endpoint at the time point of interest and the Baseline value. Participants who completed TP1 and completed their diary were analyzed.	
End point type	Secondary
End point timeframe: Baseline and Week 8	

<b>End point values</b>	FP HFA 50 µg	FP/SLM HFA 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	142 <sup>[3]</sup>	148 <sup>[4]</sup>		
Units: Scores on a scale				
least squares mean (standard error)	-1.61 (± 0.292)	-2.1 (± 0.286)		

Notes:

[3] - ITT Population

[4] - ITT Population

## Statistical analyses

<b>Statistical analysis title</b>	Statistical analysis 1
Comparison groups	FP/SLM HFA 50/25 µg v FP HFA 50 µg



Number of subjects included in analysis	290
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.235
Method	ANCOVA
Parameter estimate	Difference in Least square means
Point estimate	-0.49
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.29
upper limit	0.32

### Secondary: Mean change from Baseline in daytime asthma symptoms score at the end of Treatment Period 1 (TP1)

End point title	Mean change from Baseline in daytime asthma symptoms score at the end of Treatment Period 1 (TP1)
End point description:	The participant's parent or legally acceptable representative recorded asthma symptoms experienced by the participant during the day in a patient diary in the form of scores on a 4-point rating scale from Baseline (Week -1) until end of TP1 (Week 8). Scores ranged from 0 to 3(0: one, 1: mild, 2: moderate, 3: severe) and maximum score is 3 per day. Change from Baseline in the asthma symptom scores at day time at the end of TP1 was analyzed. The Baseline value is a mean value of the last 7 consecutive days during the run-in period (excluding the day of Visit 2 [Randomization]). The end of the TP1 value is a mean value of the last 7 consecutive days during the TP1 (excluding the last day of the TP1). Change from Baseline is the difference between the value of the endpoint at the time point of interest and the Baseline value. Participants who completed TP1 and completed their diary were analyzed.
End point type	Secondary
End point timeframe:	
Baseline and Week 8	

End point values	FP HFA 50 µg	FP/SLM HFA 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	142 <sup>[5]</sup>	148 <sup>[6]</sup>		
Units: Scores on a scale				
least squares mean (standard error)	-1.39 (± 0.287)	-1.87 (± 0.281)		

Notes:

[5] - ITT Population

[6] - ITT Population

### Statistical analyses

Statistical analysis title	Statistical analysis 1
Comparison groups	FP HFA 50 µg v FP/SLM HFA 50/25 µg

Number of subjects included in analysis	290
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.236
Method	ANCOVA
Parameter estimate	Difference in Least-Square means
Point estimate	-0.48
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.27
upper limit	0.31

### Secondary: Number of participants with at least one asthma exacerbation in Treatment Period 1 (TP1)

End point title	Number of participants with at least one asthma exacerbation in Treatment Period 1 (TP1)
-----------------	--

End point description:

The definition of exacerbations was amended during the study. <Original> An exacerbation is defined as deterioration of asthma requiring the use of systemic corticosteroids (oral, parenteral, or depot) for at least 3 days or an in-patient hospitalization or emergency department visit due to asthma that required systemic corticosteroids. <Amendment> An asthma exacerbation is defined as deterioration of asthma requiring the use of prednisone or hydrocortisone equivalent systemic corticosteroids for at least 3 days, or requiring the use of dexamethasone or betametasone equivalent systemic corticosteroids (oral, intravenous or intramuscular), or requiring the use of systemic depot corticosteroids once, or an in-patient hospitalization that required treatment for respiratory symptom with wheezing, or emergency department visit due to asthma that required intravenous systemic corticosteroids.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to 8 weeks

End point values	FP HFA 50 µg	FP/SLM HFA 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	147 <sup>[7]</sup>	150 <sup>[8]</sup>		
Units: Participants	8	4		

Notes:

[7] - ITT Population

[8] - ITT Population

### Statistical analyses

Statistical analysis title	Statistical analysis 1
Comparison groups	FP/SLM HFA 50/25 µg v FP HFA 50 µg

Number of subjects included in analysis	297
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Odds ratio (OR)
Point estimate	0.47
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.14
upper limit	1.6

### Secondary: Mean change from Baseline in Japanese Pediatric Asthma Control Program (JPAC) score at the end of Treatment Period 1 (TP1)

End point title	Mean change from Baseline in Japanese Pediatric Asthma Control Program (JPAC) score at the end of Treatment Period 1 (TP1)
-----------------	--

#### End point description:

Severity and control statuses based on Japanese pediatric guideline for the treatment and management of asthma (JPGl) can be assessed according to JPAC. Theoretically range of JPAC score was 0 (poor control) to 18 (complete control) point. JPAC questionnaire was recorded at Baseline (Week -2) and Week 8 by the participant's parent or legally acceptable representative who knew the participant's asthma for the last month. Change from Baseline is the difference between the value of the endpoint at the time point of interest and the Baseline value. Participants who completed TP1 were analyzed.

End point type	Secondary
End point timeframe:	
Baseline and Week 8	

End point values	FP HFA 50 µg	FP/SLM HFA 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	142 <sup>[9]</sup>	148 <sup>[10]</sup>		
Units: Scores on a scale				
least squares mean (standard error)	-0.3 (± 0.25)	0.4 (± 0.24)		

Notes:

[9] - ITT Population

[10] - ITT Population

### Statistical analyses

Statistical analysis title	Statistical analysis 1
Comparison groups	FP HFA 50 µg v FP/SLM HFA 50/25 µg
Number of subjects included in analysis	290
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.041
Method	ANCOVA
Parameter estimate	Difference in Least-Square Means
Point estimate	0.7

Confidence interval	
level	95 %
sides	2-sided
lower limit	0
upper limit	1.4

### Secondary: Mean change from Baseline in use of rescue medication (number of occasions used during a 24-hour period) in Treatment Period 1 (TP1)

End point title	Mean change from Baseline in use of rescue medication (number of occasions used during a 24-hour period) in Treatment Period 1 (TP1)
-----------------	--

#### End point description:

The number of inhalations of rescue salbutamol inhalation aerosol (medication used to relieve symptoms immediately) used during the day and night was recorded by the participant's parent or legally acceptable representative twice daily in a patient diary from Baseline (Week -1) until Week 8. A 24-hour period in which a participant's responses to both the morning and evening assessments indicated no use of rescue medication was considered as rescue free. Participants who were rescue free for 24-hour periods during the 8 weeks in TP1 were assessed. The Baseline value was derived from the last 7 days of the patient diary prior to the randomization of the participant. Change from Baseline is the difference between the value of the endpoint at the time point of interest and the Baseline value. Participants who completed TP1 and completed their diary were analyzed.

End point type	Secondary
----------------	-----------

#### End point timeframe:

Baseline and Week 8

End point values	FP HFA 50 µg	FP/SLM HFA 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	142 <sup>[11]</sup>	148 <sup>[12]</sup>		
Units: Occasions per 24 hours				
least squares mean (standard error)	0.07 (± 0.048)	0.01 (± 0.047)		

#### Notes:

[11] - ITT Population

[12] - ITT Population

### Statistical analyses

Statistical analysis title	Statistical analysis 1
Comparison groups	FP/SLM HFA 50/25 µg v FP HFA 50 µg
Number of subjects included in analysis	290
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.335
Method	ANCOVA
Parameter estimate	Difference in Least Square Means
Point estimate	-0.06

Confidence interval	
level	95 %
sides	2-sided
lower limit	-0.2
upper limit	0.07

### Secondary: Mean change from Baseline in use of rescue medication (percentage of days with rescue-free 24-hour period) at the end of Treatment Period 1 (TP1)

End point title	Mean change from Baseline in use of rescue medication (percentage of days with rescue-free 24-hour period) at the end of Treatment Period 1 (TP1)
-----------------	---

#### End point description:

The number of inhalations of rescue salbutamol inhalation aerosol (medication used to relieve symptoms immediately) used during the day and night was recorded by the participant's parent or legally acceptable representative twice daily in a patient diary. A 24-hour period in which a participant's responses to both the morning and evening assessments indicated no use of rescue medication was considered as rescue free. Participants who were rescue free for 24-hour periods during the 8-week Treatment Period were assessed. The Baseline value was derived from the last 7 days of the patient diary prior to the randomization of the participant. Change from Baseline is the difference between the value of the endpoint at the time point of interest and the Baseline value. Participants who completed TP1 and completed their diary were analyzed.

End point type	Secondary
----------------	-----------

#### End point timeframe:

Baseline and Week 8

End point values	FP HFA 50 µg	FP/SLM HFA 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	142 <sup>[13]</sup>	148 <sup>[14]</sup>		
Units: Percentage of days				
least squares mean (standard error)	-2.9 (± 2.16)	-0.3 (± 2.11)		

#### Notes:

[13] - ITT Population

[14] - ITT Population

### Statistical analyses

Statistical analysis title	Statistical analysis 1
Comparison groups	FP HFA 50 µg v FP/SLM HFA 50/25 µg
Number of subjects included in analysis	290
Analysis specification	Pre-specified
Analysis type	other
P-value	= 0.389
Method	ANCOVA
Parameter estimate	Difference in Least Square Means
Point estimate	2.6

Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.3
upper limit	8.6

---

**Other pre-specified: Mean change from Baseline in total asthma symptom score (daytime plus night time) at the end of the Treatment Period 2 (TP2)**

---

End point title	Mean change from Baseline in total asthma symptom score (daytime plus night time) at the end of the Treatment Period 2 (TP2)
-----------------	--

End point description:

The participant's parent or legally acceptable representative recorded asthma symptoms experienced by the participant in a patient diary twice daily (daytime and night time) in the form of scores on a 4-point rating scale from Baseline (Week -1) until end of TP2 (Week 24). Scores ranged from 0 (none) to 3 (severe) and maximum score is 6 per day. Change from Baseline in the asthma symptom scores at daytime plus night time at the end of TP2 was analyzed. The Baseline value is a mean value of the last 7 consecutive days during the run-in period (excluding the day of Visit 2 [Randomization]). The end of the TP2 value is a mean value of the last 7 consecutive days during the TP2 (excluding the last day of the TP2). Change from Baseline is the difference between the value of the endpoint at the time point of interest and the Baseline value. Participants who received at least one dose of open-label medication in the TP2 were analyzed.

End point type	Other pre-specified
----------------	---------------------

End point timeframe:

Baseline and Week 24

---

End point values	FP 50 µg - FP/SLM 50/25 µg	FP/SLM 50/25 µg - FP/SLM 50/25 µg		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	141 <sup>[15]</sup>	147 <sup>[16]</sup>		
Units: Scores on a scale				
arithmetic mean (standard deviation)	-5.29 (± 6.422)	-6.1 (± 7.665)		

Notes:

[15] - ITT Population

[16] - ITT Population

**Statistical analyses**

---

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

All on-treatment serious adverse events (SAEs) and non-serious AEs were collected for 25 weeks in Treatment Period 1 and Treatment Period 2.

Adverse event reporting additional description:

On-treatment AEs and SAEs are reported for the ITT Population.

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	19.1
--------------------	------

### Reporting groups

Reporting group title	Period 1 - FP HFA 50 µg
-----------------------	-------------------------

Reporting group description:

In TP1, participants were randomized to receive one or two inhalations of FP HFA 50 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.

Reporting group title	Period 1 - FP/SLM HFA 50/25 µg
-----------------------	--------------------------------

Reporting group description:

In TP1, participants were randomized to receive one or two inhalations of FP/SLM HFA 50/25 µg BID for 8 weeks via a MDI using AeroChamber Plus with face mask. Salbutamol was provided as a rescue medication.

Reporting group title	Period 2 - FP/SLM HFA 50/25 µg
-----------------------	--------------------------------

Reporting group description:

1 or 2 inhalations of FP/SLM HFA MDI 50/25 µg were administered twice daily in TP2 to those participants who received FP HFA MDI 50 µg or FP/SLM HFA MDI 50/25 µg in TP1.

Serious adverse events	Period 1 - FP HFA 50 µg	Period 1 - FP/SLM HFA 50/25 µg	Period 2 - FP/SLM HFA 50/25 µg
Total subjects affected by serious adverse events			
subjects affected / exposed	5 / 150 (3.33%)	1 / 150 (0.67%)	20 / 288 (6.94%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			
Injury, poisoning and procedural complications			
Skull fracture			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skull fractured base			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Subdural haemorrhage			

subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Febrile convulsion			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Asthma			
subjects affected / exposed	4 / 150 (2.67%)	0 / 150 (0.00%)	8 / 288 (2.78%)
occurrences causally related to treatment / all	0 / 4	0 / 0	0 / 8
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
Henoch-Schonlein purpura			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Bronchitis			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Upper respiratory tract infection			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia			



subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	7 / 288 (2.43%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 7
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Herpangina			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Otitis media acute			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pneumonia bacterial			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pseudocroup			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tracheobronchitis mycoplasmal			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Viral pharyngitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Frequency threshold for reporting non-serious adverse events: 0 %

<b>Non-serious adverse events</b>	Period 1 - FP HFA 50 µg	Period 1 - FP/SLM HFA 50/25 µg	Period 2 - FP/SLM HFA 50/25 µg
Total subjects affected by non-serious adverse events			
subjects affected / exposed	109 / 150 (72.67%)	111 / 150 (74.00%)	262 / 288 (90.97%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Skin papilloma			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	0	0	2
Penile neoplasm			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	7 / 150 (4.67%)	6 / 150 (4.00%)	10 / 288 (3.47%)
occurrences (all)	8	7	11
Non-cardiac chest pain			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Pain			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Immune system disorders			
Allergy to animal			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	3 / 288 (1.04%)
occurrences (all)	0	0	3
Food allergy			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Reproductive system and breast disorders			
Balanoposthitis			
subjects affected / exposed	1 / 150 (0.67%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	1	1	0
Respiratory, thoracic and mediastinal disorders			
Upper respiratory tract inflammation			
subjects affected / exposed	18 / 150 (12.00%)	10 / 150 (6.67%)	34 / 288 (11.81%)
occurrences (all)	25	13	63
Asthma			

subjects affected / exposed	10 / 150 (6.67%)	4 / 150 (2.67%)	27 / 288 (9.38%)
occurrences (all)	10	4	32
Rhinorrhoea			
subjects affected / exposed	0 / 150 (0.00%)	4 / 150 (2.67%)	7 / 288 (2.43%)
occurrences (all)	0	5	7
Cough			
subjects affected / exposed	2 / 150 (1.33%)	1 / 150 (0.67%)	2 / 288 (0.69%)
occurrences (all)	2	1	3
Epistaxis			
subjects affected / exposed	1 / 150 (0.67%)	2 / 150 (1.33%)	1 / 288 (0.35%)
occurrences (all)	2	2	1
Infantile asthma			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0
Nasal obstruction			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences (all)	0	1	1
Productive cough			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences (all)	0	1	1
Rhinitis allergic			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	5 / 288 (1.74%)
occurrences (all)	0	1	5
Dysphonia			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Nasal congestion			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Wheezing			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Investigations			
Blood cortisol decreased			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	3 / 288 (1.04%)
occurrences (all)	1	0	3

Neutrophil count increased subjects affected / exposed occurrences (all)	1 / 150 (0.67%) 1	0 / 150 (0.00%) 0	0 / 288 (0.00%) 0
Electrocardiogram QT prolonged subjects affected / exposed occurrences (all)	0 / 150 (0.00%) 0	0 / 150 (0.00%) 0	2 / 288 (0.69%) 2
Alanine aminotransferase increased subjects affected / exposed occurrences (all)	0 / 150 (0.00%) 0	0 / 150 (0.00%) 0	1 / 288 (0.35%) 1
Aspartate aminotransferase increased subjects affected / exposed occurrences (all)	0 / 150 (0.00%) 0	0 / 150 (0.00%) 0	1 / 288 (0.35%) 1
Cardiac murmur subjects affected / exposed occurrences (all)	0 / 150 (0.00%) 0	0 / 150 (0.00%) 0	1 / 288 (0.35%) 1
Injury, poisoning and procedural complications			
Arthropod bite subjects affected / exposed occurrences (all)	2 / 150 (1.33%) 2	2 / 150 (1.33%) 2	5 / 288 (1.74%) 5
Arthropod sting subjects affected / exposed occurrences (all)	3 / 150 (2.00%) 3	0 / 150 (0.00%) 0	7 / 288 (2.43%) 8
Contusion subjects affected / exposed occurrences (all)	1 / 150 (0.67%) 1	1 / 150 (0.67%) 1	2 / 288 (0.69%) 3
Fall subjects affected / exposed occurrences (all)	1 / 150 (0.67%) 1	1 / 150 (0.67%) 1	1 / 288 (0.35%) 1
Laceration subjects affected / exposed occurrences (all)	0 / 150 (0.00%) 0	2 / 150 (1.33%) 2	2 / 288 (0.69%) 2
Bite subjects affected / exposed occurrences (all)	1 / 150 (0.67%) 1	0 / 150 (0.00%) 0	0 / 288 (0.00%) 0
Joint dislocation			

subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Skin abrasion			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences (all)	0	1	1
Thermal burn			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Tongue injury			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0
Wound			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	0	0	2
Chillblains			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Closed globe injury			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Ear abrasion			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Nail injury			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Scar			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Scratch			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Conjunctival hyperaemia			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	2
Nervous system disorders			

Febrile convulsion subjects affected / exposed occurrences (all)	0 / 150 (0.00%) 0	1 / 150 (0.67%) 1	1 / 288 (0.35%) 2
Blood and lymphatic system disorders Anaemia subjects affected / exposed occurrences (all)	0 / 150 (0.00%) 0	0 / 150 (0.00%) 0	2 / 288 (0.69%) 2
Ear and labyrinth disorders External ear disorder subjects affected / exposed occurrences (all)  Ear pain subjects affected / exposed occurrences (all)	1 / 150 (0.67%) 1  0 / 150 (0.00%) 0	0 / 150 (0.00%) 0  0 / 150 (0.00%) 0	0 / 288 (0.00%) 0  1 / 288 (0.35%) 1
Eye disorders Conjunctivitis allergic subjects affected / exposed occurrences (all)  Eye discharge subjects affected / exposed occurrences (all)  Keratitis subjects affected / exposed occurrences (all)  Seasonal allergy subjects affected / exposed occurrences (all)  Blepharitis subjects affected / exposed occurrences (all)  Eye pruritus subjects affected / exposed occurrences (all)  Chalazion subjects affected / exposed occurrences (all)  Eczema eyelids	1 / 150 (0.67%) 1  1 / 150 (0.67%) 1  0 / 150 (0.00%) 0  1 / 150 (0.67%) 1  0 / 150 (0.00%) 0  0 / 150 (0.00%) 0  0 / 150 (0.00%) 0  0 / 150 (0.00%) 0	3 / 150 (2.00%) 3  0 / 150 (0.00%) 0  1 / 150 (0.67%) 1  2 / 150 (1.33%) 2  0 / 150 (0.00%) 0  0 / 150 (0.00%) 0  0 / 150 (0.00%) 0  0 / 150 (0.00%) 0	9 / 288 (3.13%) 9  3 / 288 (1.04%) 3  0 / 288 (0.00%) 0  3 / 288 (1.04%) 3  2 / 288 (0.69%) 2  1 / 288 (0.35%) 1

subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Eyelids pruritus			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Gastrointestinal disorders			
Diarrhoea			
subjects affected / exposed	1 / 150 (0.67%)	5 / 150 (3.33%)	14 / 288 (4.86%)
occurrences (all)	1	5	16
Vomiting			
subjects affected / exposed	4 / 150 (2.67%)	2 / 150 (1.33%)	11 / 288 (3.82%)
occurrences (all)	4	2	12
Stomatitis			
subjects affected / exposed	1 / 150 (0.67%)	2 / 150 (1.33%)	1 / 288 (0.35%)
occurrences (all)	1	2	1
Cheilitis			
subjects affected / exposed	0 / 150 (0.00%)	2 / 150 (1.33%)	1 / 288 (0.35%)
occurrences (all)	0	2	1
Abdominal pain			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	1	0	2
Anal fissure			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	1	0	1
Anal inflammation			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0
Aphthous ulcer			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	1	0	2
Constipation			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	11 / 288 (3.82%)
occurrences (all)	0	1	11
Enteritis			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences (all)	0	1	1

Enterocolitis			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Faeces hard			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0
Mouth haemorrhage			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Dental caries			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	0	0	2
Anal pruritus			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Dyspepsia			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	2
Gastritis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Haemorrhoids			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Mouth ulceration			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Proctitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Skin and subcutaneous tissue disorders			
Urticaria			
subjects affected / exposed	2 / 150 (1.33%)	5 / 150 (3.33%)	10 / 288 (3.47%)
occurrences (all)	2	5	11
Miliaria			



subjects affected / exposed	4 / 150 (2.67%)	2 / 150 (1.33%)	6 / 288 (2.08%)
occurrences (all)	4	2	6
Eczema			
subjects affected / exposed	2 / 150 (1.33%)	2 / 150 (1.33%)	12 / 288 (4.17%)
occurrences (all)	3	2	12
Dermatitis diaper			
subjects affected / exposed	1 / 150 (0.67%)	2 / 150 (1.33%)	6 / 288 (2.08%)
occurrences (all)	1	2	6
Dry skin			
subjects affected / exposed	1 / 150 (0.67%)	1 / 150 (0.67%)	4 / 288 (1.39%)
occurrences (all)	1	1	4
Rash			
subjects affected / exposed	1 / 150 (0.67%)	1 / 150 (0.67%)	6 / 288 (2.08%)
occurrences (all)	1	1	7
Dermatitis atopic			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	1	0	1
Dermatitis contact			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	4 / 288 (1.39%)
occurrences (all)	1	0	4
Idiopathic urticaria			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Onychoclasia			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Dermatitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	0	0	2
Pruritus			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	0	0	2
Blister			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Cold urticaria			

subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Eczema asteatotic			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Hyperkeratosis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Keratosis pilaris			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Rash vesicular			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Musculoskeletal and connective tissue disorders			
Periarthritis			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Myalgia			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Infections and infestations			
Upper respiratory tract infection			
subjects affected / exposed	17 / 150 (11.33%)	28 / 150 (18.67%)	56 / 288 (19.44%)
occurrences (all)	19	35	91
Nasopharyngitis			
subjects affected / exposed	24 / 150 (16.00%)	18 / 150 (12.00%)	68 / 288 (23.61%)
occurrences (all)	33	22	107
Bronchitis			
subjects affected / exposed	13 / 150 (8.67%)	14 / 150 (9.33%)	42 / 288 (14.58%)
occurrences (all)	15	18	58
Gastroenteritis			
subjects affected / exposed	13 / 150 (8.67%)	11 / 150 (7.33%)	43 / 288 (14.93%)
occurrences (all)	13	14	48
Pharyngitis			

subjects affected / exposed	9 / 150 (6.00%)	11 / 150 (7.33%)	39 / 288 (13.54%)
occurrences (all)	12	14	54
Hand-foot-and-mouth disease			
subjects affected / exposed	4 / 150 (2.67%)	9 / 150 (6.00%)	9 / 288 (3.13%)
occurrences (all)	4	9	9
Impetigo			
subjects affected / exposed	3 / 150 (2.00%)	5 / 150 (3.33%)	7 / 288 (2.43%)
occurrences (all)	3	5	9
Herpangina			
subjects affected / exposed	4 / 150 (2.67%)	3 / 150 (2.00%)	1 / 288 (0.35%)
occurrences (all)	5	3	1
Influenza			
subjects affected / exposed	4 / 150 (2.67%)	3 / 150 (2.00%)	38 / 288 (13.19%)
occurrences (all)	4	3	40
Otitis media			
subjects affected / exposed	4 / 150 (2.67%)	3 / 150 (2.00%)	17 / 288 (5.90%)
occurrences (all)	4	3	18
Sinusitis			
subjects affected / exposed	4 / 150 (2.67%)	3 / 150 (2.00%)	20 / 288 (6.94%)
occurrences (all)	4	3	20
Molluscum contagiosum			
subjects affected / exposed	3 / 150 (2.00%)	1 / 150 (0.67%)	8 / 288 (2.78%)
occurrences (all)	3	1	8
Bacterial rhinitis			
subjects affected / exposed	2 / 150 (1.33%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences (all)	2	1	2
Conjunctivitis			
subjects affected / exposed	0 / 150 (0.00%)	3 / 150 (2.00%)	10 / 288 (3.47%)
occurrences (all)	0	3	11
Exanthema subitum			
subjects affected / exposed	2 / 150 (1.33%)	1 / 150 (0.67%)	3 / 288 (1.04%)
occurrences (all)	2	1	3
Gastroenteritis viral			
subjects affected / exposed	1 / 150 (0.67%)	2 / 150 (1.33%)	2 / 288 (0.69%)
occurrences (all)	1	2	2
Mumps			

subjects affected / exposed	2 / 150 (1.33%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences (all)	2	1	1
Pneumonia			
subjects affected / exposed	1 / 150 (0.67%)	2 / 150 (1.33%)	5 / 288 (1.74%)
occurrences (all)	1	2	5
Streptococcal infection			
subjects affected / exposed	2 / 150 (1.33%)	1 / 150 (0.67%)	7 / 288 (2.43%)
occurrences (all)	2	1	7
Varicella			
subjects affected / exposed	2 / 150 (1.33%)	1 / 150 (0.67%)	6 / 288 (2.08%)
occurrences (all)	2	1	6
Beta haemolytic streptococcal infection			
subjects affected / exposed	1 / 150 (0.67%)	1 / 150 (0.67%)	3 / 288 (1.04%)
occurrences (all)	1	1	3
Conjunctivitis bacterial			
subjects affected / exposed	1 / 150 (0.67%)	1 / 150 (0.67%)	1 / 288 (0.35%)
occurrences (all)	1	1	1
Otitis media acute			
subjects affected / exposed	0 / 150 (0.00%)	2 / 150 (1.33%)	5 / 288 (1.74%)
occurrences (all)	0	2	5
Croup infectious			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	3 / 288 (1.04%)
occurrences (all)	0	1	3
Campylobacter infection			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Erythema infectiosum			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	1	0	1
Gastroenteritis norovirus			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0
Gastroenteritis rotavirus			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0

Hordeolum			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	2 / 288 (0.69%)
occurrences (all)	0	1	2
Pharyngitis streptococcal			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	1	0	1
Respiratory syncytial virus bronchitis			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0
Respiratory syncytial virus infection			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	5 / 288 (1.74%)
occurrences (all)	1	0	5
Rotavirus infection			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Tonsillitis			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	11 / 288 (3.82%)
occurrences (all)	1	0	11
Tonsillitis bacterial			
subjects affected / exposed	1 / 150 (0.67%)	0 / 150 (0.00%)	0 / 288 (0.00%)
occurrences (all)	1	0	0
Viral upper respiratory tract infection			
subjects affected / exposed	0 / 150 (0.00%)	1 / 150 (0.67%)	0 / 288 (0.00%)
occurrences (all)	0	1	0
Adenovirus infection			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	6 / 288 (2.08%)
occurrences (all)	0	0	7
Lower respiratory tract infection			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	3 / 288 (1.04%)
occurrences (all)	0	0	3
Rhinitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	3 / 288 (1.04%)
occurrences (all)	0	0	3
Chronic sinusitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	0	0	2

Oral herpes			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	2 / 288 (0.69%)
occurrences (all)	0	0	2
Abscess limb			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Cystitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Dermatitis infected			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Genital infection fungal			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Gingivitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Herpes virus infection			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Herpes zoster			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Laryngitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Localised infection			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Metapneumovirus infection			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Otitis externa			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	2

Parotitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Parvovirus infection			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Pharyngoconjunctival fever of children			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Rash pustular			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Tinea pedis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Tonsillitis streptococcal			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Viral pharyngitis			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1
Metabolism and nutrition disorders			
Dehydration			
subjects affected / exposed	0 / 150 (0.00%)	0 / 150 (0.00%)	1 / 288 (0.35%)
occurrences (all)	0	0	1

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
30 January 2015	The primary purpose of this amendment is to change the wording related to the definition of inclusion criteria, re-screening criteria, permitted medications and non- drug therapies, prohibited medications and non-drug therapies, asthma exacerbation, withdrawal criteria and rescue medication and to clarify ambiguous description based on the comments from Safety Review Team.
26 January 2016	The primary objectives of this amendment is to set up interim analyses with the view to posting and disclosing study result summary on the clinical trial registries within 6 months after primary compression achievement of the last subject, change administrative aspects of the trail, and make adjustments to ambiguous descriptions.

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported