



## Clinical trial results:

### A Phase 3, Double-blind, Parallel-group Study to Evaluate the Efficacy and Safety of Tezacaftor in Combination With Ivacaftor in Subjects Aged 6 Through 11 Years With Cystic Fibrosis, Homozygous or Heterozygous for the F508del-CFTR Mutation

#### Summary

EudraCT number	2016-004479-35
Trial protocol	IE DK BE DE GB PL FR
Global end of trial date	21 December 2018

#### Results information

Result version number	v1
This version publication date	09 July 2019
First version publication date	09 July 2019

#### Trial information

##### Trial identification

Sponsor protocol code	VX16-661-115
-----------------------	--------------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT03559062
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	Vertex Pharmaceuticals Incorporated
Sponsor organisation address	50 Northern Avenue, Boston, Massachusetts, United States,
Public contact	Medical Monitor, Vertex Pharmaceuticals Incorporated, +1 617 341 6777, medicalinfo@vrtx.com
Scientific contact	Medical Monitor, Vertex Pharmaceuticals Incorporated, +1 617 341 6777, medicalinfo@vrtx.com

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	Yes
EMA paediatric investigation plan number(s)	EMA-001640-PIP01-14
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	Yes

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	01 February 2019
Is this the analysis of the primary completion data?	Yes
Primary completion date	21 December 2018
Global end of trial reached?	Yes
Global end of trial date	21 December 2018
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

To evaluate the efficacy of tezacaftor (TEZ) in combination with ivacaftor (IVA) in subjects with cystic fibrosis (CF) aged 6 through 11 years, homozygous or heterozygous for F508del.

Protection of trial subjects:

The study was conducted in accordance with the ethical principles stated in the Declaration of Helsinki and the International Council on Harmonization (ICH) Guideline for Good Clinical Practice (GCP).

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	17 May 2018
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Australia: 14
Country: Number of subjects enrolled	Germany: 15
Country: Number of subjects enrolled	France: 8
Country: Number of subjects enrolled	Switzerland: 8
Country: Number of subjects enrolled	United Kingdom: 8
Country: Number of subjects enrolled	Denmark: 6
Country: Number of subjects enrolled	Belgium: 5
Country: Number of subjects enrolled	Ireland: 4
Country: Number of subjects enrolled	Poland: 1
Worldwide total number of subjects	69
EEA total number of subjects	47

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23	0

months)	
Children (2-11 years)	69
Adolescents (12-17 years)	0
Adults (18-64 years)	0
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details: -

### Pre-assignment

Screening details:

A total of 69 subjects were randomized, out of which 67 subjects received study drug and were included in subject disposition and baseline characteristics section.

### Period 1

Period 1 title	Overall Period (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator, Carer, Assessor

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Placebo

Arm description:

Subjects with genotype F/F received placebo matched to TEZ/IVA fixed dose combination (FDC) in the morning and placebo matched to IVA in the evening for 8 weeks.

Arm type	Blinding arm
Investigational medicinal product name	Placebo (matched to TEZ/IVA)
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Subjects received placebo matched to TEZ/IVA once daily in the morning.

Investigational medicinal product name	Placebo (matched to IVA)
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Subjects received placebo matched to IVA once daily in the evening.

<b>Arm title</b>	TEZ/IVA
------------------	---------

Arm description:

Subjects with genotype F/F received TEZ/IVA FDC in the morning and IVA in the evening for 8 weeks. Subjects with genotype F/RF received TEZ/IVA FDC and placebo matched to IVA in the morning and IVA in the evening for 8 weeks.

Arm type	Experimental
Investigational medicinal product name	TEZ/IVA
Investigational medicinal product code	VX-661/VX-770
Other name	Tezacaftor/Ivacaftor fixed dose combination
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Subjects received TEZ/IVA once daily in the morning.

Investigational medicinal product name	IVA
Investigational medicinal product code	VX-770
Other name	Ivacaftor
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Subjects received IVA once daily in the evening.

Investigational medicinal product name	Placebo (matched to IVA)
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Subjects received placebo matched to IVA once daily in the morning.

<b>Arm title</b>	Ivacaftor
------------------	-----------

Arm description:

Subjects with genotype F/RF received placebo matched to TEZ/IVA FDC in the morning and IVA in morning and evening for 8 weeks.

Arm type	Blinding arm
Investigational medicinal product name	IVA
Investigational medicinal product code	VX-770
Other name	Ivacaftor
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Subjects received IVA every 12 hours.

Investigational medicinal product name	Placebo (matched to TEZ/IVA)
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Subjects received placebo matched to TEZ/IVA once daily in the morning.

<b>Number of subjects in period 1<sup>[1]</sup></b>	Placebo	TEZ/IVA	Ivacaftor
Started	10	54	3
Completed	10	53	3
Not completed	0	1	0
Other	-	1	-

Notes:

[1] - The number of subjects reported to be in the baseline period are not the same as the worldwide number enrolled in the trial. It is expected that these numbers will be the same.

Justification: A total of 69 subjects were randomized, out of which 67 subjects received study drug and were included in subject disposition and baseline characteristics section.

## Baseline characteristics

### Reporting groups

Reporting group title	Placebo
Reporting group description: Subjects with genotype F/F received placebo matched to TEZ/IVA fixed dose combination (FDC) in the morning and placebo matched to IVA in the evening for 8 weeks.	
Reporting group title	TEZ/IVA
Reporting group description: Subjects with genotype F/F received TEZ/IVA FDC in the morning and IVA in the evening for 8 weeks. Subjects with genotype F/RF received TEZ/IVA FDC and placebo matched to IVA in the morning and IVA in the evening for 8 weeks.	
Reporting group title	Ivacaftor
Reporting group description: Subjects with genotype F/RF received placebo matched to TEZ/IVA FDC in the morning and IVA in morning and evening for 8 weeks.	

Reporting group values	Placebo	TEZ/IVA	Ivacaftor
Number of subjects	10	54	3
Age categorical Units: Subjects			

Age continuous Units: years arithmetic mean standard deviation	9.0 ± 1.7	8.5 ± 1.7	9.0 ± 1.7
Gender categorical Units: Subjects			
Female	6	29	2
Male	4	25	1

Reporting group values	Total		
Number of subjects	67		
Age categorical Units: Subjects			

Age continuous Units: years arithmetic mean standard deviation	-		
Gender categorical Units: Subjects			
Female	37		
Male	30		

## End points

### End points reporting groups

Reporting group title	Placebo
Reporting group description: Subjects with genotype F/F received placebo matched to TEZ/IVA fixed dose combination (FDC) in the morning and placebo matched to IVA in the evening for 8 weeks.	
Reporting group title	TEZ/IVA
Reporting group description: Subjects with genotype F/F received TEZ/IVA FDC in the morning and IVA in the evening for 8 weeks. Subjects with genotype F/RF received TEZ/IVA FDC and placebo matched to IVA in the morning and IVA in the evening for 8 weeks.	
Reporting group title	Ivacaftor
Reporting group description: Subjects with genotype F/RF received placebo matched to TEZ/IVA FDC in the morning and IVA in morning and evening for 8 weeks.	

### Primary: Absolute Change in Lung Clearance Index 2.5

End point title	Absolute Change in Lung Clearance Index 2.5 <sup>[1]</sup> [2]
End point description:	

End point type	Primary
End point timeframe: From baseline through Week 8	

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The study was designed to perform within treatment group comparison. Because single group within treatment comparisons cannot be reported in the EudraCT database, no statistical analyses are reported.

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is applicable for only TEZ/IVA.

End point values	TEZ/IVA			
Subject group type	Reporting group			
Number of subjects analysed	54			
Units: Lung clearance index				
least squares mean (standard error)	-0.51 ( $\pm$ 0.11)			

### Statistical analyses

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

From first dose of study drug up to safety follow-up visit

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	21.1
--------------------	------

### Reporting groups

Reporting group title	Placebo
-----------------------	---------

Reporting group description:

Subjects with genotype F/F received placebo matched to TEZ/IVA FDC in the morning and placebo matched to IVA in the evening for 8 weeks.

Reporting group title	TEZ/IVA
-----------------------	---------

Reporting group description:

Subjects with genotype F/F received TEZ/IVA FDC in the morning and IVA in the evening for 8 weeks.  
Subjects with genotype F/RF received TEZ/IVA FDC and placebo matched to IVA in the morning and IVA in the evening for 8 weeks.

Reporting group title	Ivacaftor
-----------------------	-----------

Reporting group description:

Subjects with genotype F/RF received placebo matched to TEZ/IVA FDC in the morning and IVA in morning and evening for 8 weeks.

Serious adverse events	Placebo	TEZ/IVA	Ivacaftor
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 10 (0.00%)	0 / 54 (0.00%)	0 / 3 (0.00%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			

Frequency threshold for reporting non-serious adverse events: 5 %

Non-serious adverse events	Placebo	TEZ/IVA	Ivacaftor
Total subjects affected by non-serious adverse events			
subjects affected / exposed	8 / 10 (80.00%)	31 / 54 (57.41%)	2 / 3 (66.67%)
Investigations			
Alanine aminotransferase increased			
subjects affected / exposed	1 / 10 (10.00%)	0 / 54 (0.00%)	0 / 3 (0.00%)
occurrences (all)	2	0	0
Aspartate aminotransferase increased			



subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 1	0 / 54 (0.00%) 0	0 / 3 (0.00%) 0
Injury, poisoning and procedural complications			
Thermal burn			
subjects affected / exposed	1 / 10 (10.00%)	0 / 54 (0.00%)	0 / 3 (0.00%)
occurrences (all)	1	0	0
Arthropod bite			
subjects affected / exposed	0 / 10 (0.00%)	0 / 54 (0.00%)	1 / 3 (33.33%)
occurrences (all)	0	0	1
Nervous system disorders			
Headache			
subjects affected / exposed	1 / 10 (10.00%)	8 / 54 (14.81%)	0 / 3 (0.00%)
occurrences (all)	1	11	0
Gastrointestinal disorders			
Vomiting			
subjects affected / exposed	0 / 10 (0.00%)	4 / 54 (7.41%)	0 / 3 (0.00%)
occurrences (all)	0	4	0
Abdominal pain			
subjects affected / exposed	0 / 10 (0.00%)	3 / 54 (5.56%)	0 / 3 (0.00%)
occurrences (all)	0	3	0
Respiratory, thoracic and mediastinal disorders			
Cough			
subjects affected / exposed	1 / 10 (10.00%)	9 / 54 (16.67%)	0 / 3 (0.00%)
occurrences (all)	1	10	0
Sputum increased			
subjects affected / exposed	1 / 10 (10.00%)	0 / 54 (0.00%)	0 / 3 (0.00%)
occurrences (all)	1	0	0
Rhinorrhoea			
subjects affected / exposed	0 / 10 (0.00%)	3 / 54 (5.56%)	1 / 3 (33.33%)
occurrences (all)	0	4	1
Productive cough			
subjects affected / exposed	1 / 10 (10.00%)	7 / 54 (12.96%)	0 / 3 (0.00%)
occurrences (all)	1	9	0
Nasal congestion			
subjects affected / exposed	0 / 10 (0.00%)	3 / 54 (5.56%)	0 / 3 (0.00%)
occurrences (all)	0	3	0

Throat irritation subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 1	0 / 54 (0.00%) 0	0 / 3 (0.00%) 0
Skin and subcutaneous tissue disorders			
Rash subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 1	0 / 54 (0.00%) 0	0 / 3 (0.00%) 0
Blister subjects affected / exposed occurrences (all)	0 / 10 (0.00%) 0	0 / 54 (0.00%) 0	1 / 3 (33.33%) 1
Musculoskeletal and connective tissue disorders			
Arthritis subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 1	0 / 54 (0.00%) 0	0 / 3 (0.00%) 0
Infections and infestations			
Infective pulmonary exacerbation of cystic fibrosis subjects affected / exposed occurrences (all)	2 / 10 (20.00%) 4	3 / 54 (5.56%) 3	0 / 3 (0.00%) 0
Bacterial disease carrier subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 1	0 / 54 (0.00%) 0	0 / 3 (0.00%) 0
Gastroenteritis subjects affected / exposed occurrences (all)	0 / 10 (0.00%) 0	4 / 54 (7.41%) 5	0 / 3 (0.00%) 0
Nasopharyngitis subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 1	5 / 54 (9.26%) 5	0 / 3 (0.00%) 0
Otitis media subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 2	0 / 54 (0.00%) 0	0 / 3 (0.00%) 0
Rhinitis subjects affected / exposed occurrences (all)	1 / 10 (10.00%) 1	0 / 54 (0.00%) 0	0 / 3 (0.00%) 0
Impetigo subjects affected / exposed occurrences (all)	0 / 10 (0.00%) 0	0 / 54 (0.00%) 0	1 / 3 (33.33%) 1

Upper respiratory tract infection subjects affected / exposed	0 / 10 (0.00%)	0 / 54 (0.00%)	1 / 3 (33.33%)
occurrences (all)	0	0	2
Viral upper respiratory tract infection subjects affected / exposed	0 / 10 (0.00%)	0 / 54 (0.00%)	1 / 3 (33.33%)
occurrences (all)	0	0	1

## **More information**

### **Substantial protocol amendments (globally)**

Were there any global substantial amendments to the protocol? No

---

### **Interruptions (globally)**

Were there any global interruptions to the trial? No

### **Limitations and caveats**

None reported