

**Table 1****Morphological characteristics**

|  |                              | <b>Week 0</b> | <b>Week 3</b> | <b>Week 6</b> | <b><i>P, between<br/>Week 0 and<br/>Week 6</i></b> |
|--|------------------------------|---------------|---------------|---------------|--|
| <b>Median infundibulum diameter, <math>\mu\text{m}</math></b>                            |                              | 63            | 57            | 54            | 0.771  |
| <b>IQR</b>   |                              | 37-77         | 39-67         | 39-71         |  |
| <b>Follicle border, %</b>  | <i>Dark grey</i>             | 60.1          | 91.4          | 88.5          | <0.0001*   |
|  | <i>White hyperkeratotic</i>  | 39.9          | 8.6           | 11.5          |  |
| <b>Follicle content, %</b>   | <i>Empty</i>                 | 59.8          | 72.9          | 80.3          | <0.0001**  |
|  | <i>Amorphous grey</i>        | 25.1          | 17.4          | 11.8          |  |
|  | <i>White hyperreflective</i> | 15.1          | 9.7           | 7.9           |  |
| <b>Other characteristics, %</b>  | <i>Thickened border</i>      | 46.7          | 28.6          | 23.4          | <0.0001  |
| <b>Median epidermal thickness, <math>\mu\text{m}</math></b>                              |                              | 85.9          |               | 102.3         | 0.0003   |
| <b>IQR</b>   |                              | 74-99         |               | 88-117        |  |
| <b>Blood flow</b>  |                              | 0.15          |               | 0.18          | 0.454  |
| <b>*Differences between the distribution of follicle content at Week 0 versus Week 6</b> |                              |               |               |               |  |
| <b>** Differences in the distribution of follicle border at Week 0 versus Week 6</b>     |                              |               |               |               |  |