



Clinical trial results:

A PHASE 3, MULTICENTER, RANDOMIZED, DOUBLE-BLIND, PARALLEL-ARM, PLACEBO-CONTROLLED MAINTENANCE STUDY OF MIRIKIZUMAB IN PATIENTS WITH MODERATELY TO SEVERELY ACTIVE ULCERATIVE COLITIS (LUCENT 2)

Summary

EudraCT number	2017-003238-96
Trial protocol	DE GB NL LV CZ LT ES BE HU AT SK DK HR IT RO
Global end of trial date	17 February 2025

Results information

Result version number	v2 (current)
This version publication date	04 March 2026
First version publication date	15 December 2022
Version creation reason	<ul style="list-style-type: none">New data added to full data set Final results to be added with ME2 Data

Trial information

Trial identification

Sponsor protocol code	I6T-MC-AMBG
-----------------------	-------------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT03524092
WHO universal trial number (UTN)	-
Other trial identifiers	Trial Number: 16823

Notes:

Sponsors

Sponsor organisation name	Eli Lilly and Company
Sponsor organisation address	Lilly Corporate Center, Indianapolis, IN, United States, 46285
Public contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 877CTLilly,
Scientific contact	Available Mon - Fri 9 AM - 5 PM EST, Eli Lilly and Company, 1 8772854559,

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	17 February 2025
Is this the analysis of the primary completion data?	Yes
Primary completion date	03 November 2021
Global end of trial reached?	Yes
Global end of trial date	17 February 2025
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

The purpose of this study is to evaluate the efficacy and safety of mirikizumab as maintenance therapy in participants who completed as clinical responders in the prior 12-week induction study LUCENT-1 (NCT03518086).

Protection of trial subjects:

This study was conducted in accordance with International Conference on Harmonization (ICH) Good Clinical Practice, and the principles of the Declaration of Helsinki, in addition to following the laws and regulations of the country or countries in which a study is conducted.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	19 October 2018
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Argentina: 11
Country: Number of subjects enrolled	Australia: 12
Country: Number of subjects enrolled	Austria: 8
Country: Number of subjects enrolled	Belgium: 9
Country: Number of subjects enrolled	Canada: 31
Country: Number of subjects enrolled	China: 166
Country: Number of subjects enrolled	Czechia: 52
Country: Number of subjects enrolled	Denmark: 7
Country: Number of subjects enrolled	France: 57
Country: Number of subjects enrolled	Germany: 37
Country: Number of subjects enrolled	Hungary: 23
Country: Number of subjects enrolled	India: 72
Country: Number of subjects enrolled	Ireland: 1
Country: Number of subjects enrolled	Israel: 15
Country: Number of subjects enrolled	Italy: 36
Country: Number of subjects enrolled	Japan: 123
Country: Number of subjects enrolled	Latvia: 29
Country: Number of subjects enrolled	Lithuania: 22
Country: Number of subjects enrolled	Malaysia: 6

Country: Number of subjects enrolled	Mexico: 10
Country: Number of subjects enrolled	Netherlands: 10
Country: Number of subjects enrolled	Poland: 122
Country: Number of subjects enrolled	Romania: 14
Country: Number of subjects enrolled	Russian Federation: 100
Country: Number of subjects enrolled	Serbia: 21
Country: Number of subjects enrolled	Slovakia: 21
Country: Number of subjects enrolled	Korea, Republic of: 25
Country: Number of subjects enrolled	Spain: 20
Country: Number of subjects enrolled	Switzerland: 10
Country: Number of subjects enrolled	Taiwan: 3
Country: Number of subjects enrolled	Turkey: 8
Country: Number of subjects enrolled	Ukraine: 93
Country: Number of subjects enrolled	United Kingdom: 14
Country: Number of subjects enrolled	United States: 140
Worldwide total number of subjects	1328
EEA total number of subjects	468

Notes:

Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	1216
From 65 to 84 years	112
85 years and over	0

Subject disposition

Recruitment

Recruitment details:

A total of 1177 Participants (pts) enrolled in study of which 716 participants entered Blinded Maintenance Period as Induction Responder (IR) and 461 pts entered into open label extended Induction period as Induction Nonresponders. China Maximized Extended Enrollment (ME2): This is an extension phase of the main study, with an additional 151 pts.

Pre-assignment

Screening details:

This study designed to evaluate safety and efficacy of mirikizumab (miri) in achieving remission at Week (Wk) 40 in pts who completed 12 wk induction study I6T-MC-AMAN(NCT03518086) as clinical responders to mirikizumab.

As pre-specified in analysis plan, outcome measures were not reported for ME2 arms/groups, as no formal efficacy was pre-specified

Period 1

Period 1 title	Blinded Maintenance Period (Wk 0-40)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

Arms

Are arms mutually exclusive?	Yes
Arm title	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)

Arm description:

Participants who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.

Arm type	Placebo
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

PBO SC every 4 weeks (Q4W).

Arm title	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
------------------	--

Arm description:

Participants who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Arm type	Experimental
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

200 mg mirikizumab SC Q4W.

Arm title	Maintenance Period: PBO IR - PBO SC
------------------	-------------------------------------

Arm description:

Participants who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Arm type	PBO IR
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

PBO SC every 4 weeks (Q4W).

Arm title	Maintenance Period: Miri IR - PBO SC - ME2 Cohort
------------------	---

Arm description:

Participants in the ME2 Cohort who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.

Arm type	Experimental
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

PBO SC Q4W (ME2 Cohort).

Arm title	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort
------------------	---

Arm description:

Participants in the ME2 Cohort who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Arm type	Experimental
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

200 mg mirikizumab SC Q4W (ME2 Cohort).

Arm title	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
------------------	--

Arm description:

Participants in the ME2 Cohort who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Arm type	Experimental
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

PBO SC every Q4W (ME2 Cohort).

Number of subjects in period 1^[1]	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC	Maintenance Period: PBO IR - PBO SC
Started	192	389	135
Received at Least One Dose of Study Drug	192	389	135
Completed	119	347	90
Not completed	73	42	45
Consent withdrawn by subject	7	8	6
Physician decision	-	1	-
Adverse event, non-fatal	16	6	1
Pregnancy	-	-	1
Protocol Violation	-	1	1
Loss of Response Rescue Period	42	19	29
Lost to follow-up	-	1	-
Lack of efficacy	8	6	7

Number of subjects in period 1^[1]	Maintenance Period: Miri IR - PBO SC - ME2 Cohort	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
Started	21	52	12
Received at Least One Dose of Study Drug	21	52	12
Completed	12	46	8
Not completed	9	6	4
Consent withdrawn by subject	1	-	-
Physician decision	-	-	-
Adverse event, non-fatal	2	4	-
Pregnancy	-	-	-
Protocol Violation	-	-	-
Loss of Response Rescue Period	6	2	4
Lost to follow-up	-	-	-
Lack of efficacy	-	-	-

Notes:

[1] - The number of subjects reported to be in the baseline period are not the same as the worldwide number enrolled in the trial. It is expected that these numbers will be the same.

Justification: Data is reported separately for Blinded Maintenance Period as Induction Responder (IR) and open label extended Induction period as Induction Nonresponders.

Period 2

Period 2 title	Open Label Extended Induction (Wk 0-12)
Is this the baseline period?	No
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
------------------------------	-----

Arm title	Extended Induction: Induction Nonresponders - 300mg Miri IV
------------------	---

Arm description:

Participants who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.

Arm type	Induction Nonresponders
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Intravenous use

Dosage and administration details:

300 mg mirikizumab IV Q4W for 3 doses.

Arm title	Extended Induction: Induction Nonresponders-300mg Miri IV-ME2
------------------	---

Arm description:

Participants in the ME2 Cohort who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.

Arm type	Experimental
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Intravenous use

Dosage and administration details:

300 mg mirikizumab IV Q4W for 3 doses (ME2 Cohort).

Number of subjects in period 2 ^[2]	Extended Induction: Induction Nonresponders - 300mg Miri IV	Extended Induction: Induction Nonresponders- 300mg Miri IV-ME2
Started	461	66
Completed	271	33
Not completed	190	33
Physician decision	1	-
Consent withdrawn by subject	6	5
Lack of capacity due to lack of staff	1	-
Adverse event, non-fatal	11	1
Covid-19 related study disruption	4	-
Protocol Violation	5	1
Lack of efficacy	162	26

Notes:

[2] - The number of subjects starting the period is not consistent with the number completing the preceding period. It is expected the number of subjects starting the subsequent period will be the same as the number completing the preceding period.

Justification: Included only participants who entered open label extended induction period.

Period 3

Period 3 title	Open Label Maintenance Period (Wk 12-40)
Is this the baseline period?	No
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
------------------------------	-----

Arm title	Open Label Maintenance Period (Wk 12-40)
------------------	--

Arm description:

Participants who initially did not respond to induction study (LUCENT-1), but responded to extended induction therapy at Week 12 of LUCENT-2 (delayed responders), received 200 mg mirikizumab SC Q4W during open label maintenance period from Week 12 until Week 40 or until early termination (rescue was not available for these participants).

Arm type	Open Label Maintenance Period
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

200 mg mirikizumab SC Q4W.

Arm title	Open Label Maintenance:Delayed Responders- 200mg Miri SC-ME2
------------------	--

Arm description:

Participants in the ME2 Cohort who initially did not respond to induction study (LUCENT-1), but responded to extended induction therapy at Week 12 of LUCENT-2 (delayed responders), received 200 mg mirikizumab SC Q4W during open label maintenance period from Week 12 until Week 40 or until early termination (rescue was not available for these participants).

Arm type	Experimental
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

200 mg mirikizumab SC Q4W (ME2 Cohort).

Number of subjects in period 3	Open Label Maintenance Period (Wk 12-40)	Open Label Maintenance:Delayed Responders-200mg Miri SC- ME2
Started	271	33
Completed	256	30
Not completed	15	3
Consent withdrawn by subject	-	2

Adverse event, non-fatal	4	1
Covid-19 related study disruption	1	-
Protocol Violation	1	-
Lack of efficacy	9	-

Period 4

Period 4 title	Loss of Response Rescue Period
Is this the baseline period?	No
Allocation method	Not applicable
Blinding used	Not blinded

Arms

Are arms mutually exclusive?	Yes
Arm title	Loss of Response (LOR) Rescue Period:LOR Cohort-300 mg Miri IV

Arm description:

Participants who received blinded PBO SC or blinded 200 mg mirikizumab SC Q4W during maintenance period and experienced a loss of response at or after Week 12, received rescue therapy with open label 300 mg mirikizumab intravenous (IV) Q4W for 3 doses.

Arm type	LOR Rescue Period
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Intravenous use

Dosage and administration details:

300 mg mirikizumab IV Q4W.

Arm title	LOR Rescue Period:LOR Cohort-300 mg Miri IV - ME2 Cohort
------------------	--

Arm description:

Participants in the ME2 Cohort who received blinded PBO SC or blinded 200 mg mirikizumab SC Q4W during maintenance period and experienced a loss of response at or after Week 12, received rescue therapy with open label 300 mg mirikizumab intravenous (IV) Q4W for 3 doses.

Arm type	Experimental
Investigational medicinal product name	Mirikizumab
Investigational medicinal product code	
Other name	LY3074828
Pharmaceutical forms	Solution for injection
Routes of administration	Intravenous use

Dosage and administration details:

300 mg mirikizumab IV Q4W (ME2 Cohort).

Number of subjects in period 4^[3]	Loss of Response (LOR) Rescue Period:LOR Cohort-300 mg Miri IV	LOR Rescue Period:LOR Cohort-300 mg Miri IV -
Started	90	12
Completed	81	9
Not completed	9	3
Consent withdrawn by subject	1	-
Adverse event, non-fatal	2	-
Pregnancy	1	-
Rolled Over to Study I6TMCAMAP (NCT03519945)	-	2
Lack of efficacy	5	1

Notes:

[3] - The number of subjects starting the period is not consistent with the number completing the preceding period. It is expected the number of subjects starting the subsequent period will be the same as the number completing the preceding period.

Justification: Included only participants who experienced loss of response and were eligible for loss of response rescue induction.

Baseline characteristics

Reporting groups

Reporting group title	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)
Reporting group description: Participants who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Reporting group description: Participants who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: PBO IR - PBO SC
Reporting group description: Participants who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: Miri IR - PBO SC - ME2 Cohort
Reporting group description: Participants in the ME2 Cohort who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort
Reporting group description: Participants in the ME2 Cohort who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
Reporting group description: Participants in the ME2 Cohort who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	

Reporting group values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC	Maintenance Period: PBO IR - PBO SC
Number of subjects	192	389	135
Age categorical Units: Subjects			
Age continuous			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: years			
arithmetic mean	41.20	43.30	40.80
standard deviation	± 12.88	± 14.13	± 13.40
Gender categorical			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
Female	78	160	61
Male	114	229	74

Ethnicity			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1). Ethnicity data collected only for United States (US) participants.			
Units: Subjects			
Hispanic or Latino	2	12	2
Not Hispanic or Latino	18	33	8
Unknown or Not Reported	172	344	125
Race			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
American Indian or Alaska Native	1	3	2
Asian	51	93	28
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	0	6	0
White	138	285	103
More than one race	0	0	2
Unknown or Not Reported	2	2	0
Region of Enrollment			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
Argentina	1	5	0
Australia	0	3	0
Austria	2	2	0
Belgium	1	2	2
Canada	3	5	2
China	3	4	1
Czechia	7	19	11
Denmark	3	3	0
France	10	18	6
Germany	5	13	3
Hungary	1	8	2
India	18	25	16
Ireland	0	1	0
Israel	3	6	1
Italy	3	12	4
Japan	25	47	8
Latvia	8	10	3
Lithuania	6	4	5
Malaysia	0	3	0
Mexico	1	3	2
Netherlands	1	6	0
Poland	15	32	13
Romania	4	2	0
Russia	19	36	16
Serbia	1	9	4
Slovakia	6	7	1
South Korea	3	8	1
Spain	1	6	4
Switzerland	0	4	0
Taiwan	1	1	0
Turkey	0	5	2

Ukraine	18	31	16
United Kingdom	2	4	2
United States	21	45	10

Reporting group values	Maintenance Period: Miri IR - PBO SC - ME2 Cohort	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
Number of subjects	21	52	12
Age categorical			
Units: Subjects			

Age continuous			
----------------	--	--	--

All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).

Units: years			
arithmetic mean	49.10	42.50	56.20
standard deviation	± 10.71	± 13.90	± 14.01

Gender categorical			
--------------------	--	--	--

All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).

Units: Subjects			
Female	6	23	5
Male	15	29	7

Ethnicity			
-----------	--	--	--

All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1). Ethnicity data collected only for United States (US) participants.

Units: Subjects			
Hispanic or Latino			
Not Hispanic or Latino			
Unknown or Not Reported	21	52	12

Race			
------	--	--	--

All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).

Units: Subjects			
American Indian or Alaska Native	0	0	0
Asian	21	52	12
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	0	0	0
White	0	0	0
More than one race	0	0	0
Unknown or Not Reported	0	0	0

Region of Enrollment			
----------------------	--	--	--

All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).

Units: Subjects			
Argentina	0	0	0
Australia	0	0	0
Austria	0	0	0
Belgium	0	0	0
Canada	0	0	0
China	21	52	12
Czechia	0	0	0
Denmark	0	0	0
France	0	0	0
Germany	0	0	0
Hungary	0	0	0

India	0	0	0
Ireland	0	0	0
Israel	0	0	0
Italy	0	0	0
Japan	0	0	0
Latvia	0	0	0
Lithuania	0	0	0
Malaysia	0	0	0
Mexico	0	0	0
Netherlands	0	0	0
Poland	0	0	0
Romania	0	0	0
Russia	0	0	0
Serbia	0	0	0
Slovakia	0	0	0
South Korea	0	0	0
Spain	0	0	0
Switzerland	0	0	0
Taiwan	0	0	0
Turkey	0	0	0
Ukraine	0	0	0
United Kingdom	0	0	0
United States	0	0	0

Reporting group values	Total		
Number of subjects	801		
Age categorical			
Units: Subjects			

Age continuous			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: years			
arithmetic mean			
standard deviation	-		
Gender categorical			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
Female	333		
Male	468		
Ethnicity			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1). Ethnicity data collected only for United States (US) participants.			
Units: Subjects			
Hispanic or Latino	16		
Not Hispanic or Latino	59		
Unknown or Not Reported	726		
Race			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
American Indian or Alaska Native	6		
Asian	257		

Native Hawaiian or Other Pacific Islander	0		
Black or African American	6		
White	526		
More than one race	2		
Unknown or Not Reported	4		
Region of Enrollment			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
Argentina	6		
Australia	3		
Austria	4		
Belgium	5		
Canada	10		
China	93		
Czechia	37		
Denmark	6		
France	34		
Germany	21		
Hungary	11		
India	59		
Ireland	1		
Israel	10		
Italy	19		
Japan	80		
Latvia	21		
Lithuania	15		
Malaysia	3		
Mexico	6		
Netherlands	7		
Poland	60		
Romania	6		
Russia	71		
Serbia	14		
Slovakia	14		
South Korea	12		
Spain	11		
Switzerland	4		
Taiwan	2		
Turkey	7		
Ukraine	65		
United Kingdom	8		
United States	76		

Subject analysis sets

Subject analysis set title	200 Milligram (mg) Miri SC
Subject analysis set type	Per protocol
Subject analysis set description: Participants who received 200 mg mirikizumab SC every Q4W.	
Subject analysis set title	Extended Induction: Induction Nonresponders - 300mg Miri IV
Subject analysis set type	Intention-to-treat

Subject analysis set description:

Participants who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.

Subject analysis set title	Extended Induction Induction Nonresponders-300mg Miri IV - ME2
Subject analysis set type	Intention-to-treat

Subject analysis set description:

Participants in the ME2 Cohort who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.

Reporting group values	200 Milligram (mg) Miri SC	Extended Induction: Induction Nonresponders - 300mg Miri IV	Extended Induction Induction Nonresponders- 300mg Miri IV - ME2
Number of subjects	652	461	66
Age categorical			
Units: Subjects			

Age continuous			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: years			
arithmetic mean		43.40	44.40
standard deviation	±	± 13.86	± 14.82
Gender categorical			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
Female		166	18
Male		295	48
Ethnicity			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1). Ethnicity data collected only for United States (US) participants.			
Units: Subjects			
Hispanic or Latino		11	0
Not Hispanic or Latino		53	0
Unknown or Not Reported		397	66
Race			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
American Indian or Alaska Native		4	0
Asian		87	66
Native Hawaiian or Other Pacific Islander		1	0
Black or African American		6	0
White		358	0
More than one race		1	0
Unknown or Not Reported		4	0
Region of Enrollment			
All randomized participants who are responders to Miri and PBO in induction study (LUCENT-1).			
Units: Subjects			
Argentina		5	0
Australia		9	0
Austria		4	0
Belgium		4	0

Canada		21	0
China		7	66
Czechia		15	0
Denmark		1	0
France		23	0
Germany		16	0
Hungary		12	0
India		13	0
Ireland		0	0
Israel		5	0
Italy		17	0
Japan		43	0
Latvia		8	0
Lithuania		7	0
Malaysia		3	0
Mexico		4	0
Netherlands		3	0
Poland		62	0
Romania		8	0
Russia		29	0
Serbia		7	0
Slovakia		7	0
South Korea		13	0
Spain		9	0
Switzerland		6	0
Taiwan		1	0
Turkey		1	0
Ukraine		28	0
United Kingdom		6	0
United States		64	0

End points

End points reporting groups

Reporting group title	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)
Reporting group description: Participants who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Reporting group description: Participants who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: PBO IR - PBO SC
Reporting group description: Participants who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: Miri IR - PBO SC - ME2 Cohort
Reporting group description: Participants in the ME2 Cohort who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort
Reporting group description: Participants in the ME2 Cohort who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	
Reporting group title	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
Reporting group description: Participants in the ME2 Cohort who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.	
Reporting group title	Extended Induction: Induction Nonresponders - 300mg Miri IV
Reporting group description: Participants who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.	
Reporting group title	Extended Induction: Induction Nonresponders-300mg Miri IV-ME2
Reporting group description: Participants in the ME2 Cohort who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.	
Reporting group title	Open Label Maintenance Period (Wk 12-40)
Reporting group description: Participants who initially did not respond to induction study (LUCENT-1), but responded to extended induction therapy at Week 12 of LUCENT-2 (delayed responders), received 200 mg mirikizumab SC Q4W during open label maintenance period from Week 12 until Week 40 or until early termination (rescue was not available for these participants).	
Reporting group title	Open Label Maintenance:Delayed Responders- 200mg Miri SC-ME2
Reporting group description: Participants in the ME2 Cohort who initially did not respond to induction study (LUCENT-1), but responded to extended induction therapy at Week 12 of LUCENT-2 (delayed responders), received 200 mg mirikizumab SC Q4W during open label maintenance period from Week 12 until Week 40 or until	

early termination (rescue was not available for these participants).

Reporting group title	Loss of Response (LOR) Rescue Period:LOR Cohort-300 mg Miri IV
-----------------------	--

Reporting group description:

Participants who received blinded PBO SC or blinded 200 mg mirikizumab SC Q4W during maintenance period and experienced a loss of response at or after Week 12, received rescue therapy with open label 300 mg mirikizumab intravenous (IV) Q4W for 3 doses.

Reporting group title	LOR Rescue Period:LOR Cohort-300 mg Miri IV - ME2 Cohort
-----------------------	--

Reporting group description:

Participants in the ME2 Cohort who received blinded PBO SC or blinded 200 mg mirikizumab SC Q4W during maintenance period and experienced a loss of response at or after Week 12, received rescue therapy with open label 300 mg mirikizumab intravenous (IV) Q4W for 3 doses.

Subject analysis set title	200 Milligram (mg) Miri SC
----------------------------	----------------------------

Subject analysis set type	Per protocol
---------------------------	--------------

Subject analysis set description:

Participants who received 200 mg mirikizumab SC every Q4W.

Subject analysis set title	Extended Induction: Induction Nonresponders - 300mg Miri IV
----------------------------	---

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

Participants who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.

Subject analysis set title	Extended Induction Induction Nonresponders-300mg Miri IV - ME2
----------------------------	--

Subject analysis set type	Intention-to-treat
---------------------------	--------------------

Subject analysis set description:

Participants in the ME2 Cohort who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.

Primary: Percentage of Participants in Clinical Remission at Week 40 (Mirikizumab Induction Responders)

End point title	Percentage of Participants in Clinical Remission at Week 40 (Mirikizumab Induction Responders) ^[1]
-----------------	---

End point description:

Clinical remission at week 40 is defined as achieving a 9-point modified Mayo score for rectal bleeding=0, stool frequency=0 or 1 with ≥ 1 point decrease from baseline, and endoscopy=0 or 1 (excluding friability). Stool Frequency Subscore, based on the participant's diary and scored from 0 (normal number of stools) to 3 (5 or more stools than normal); Rectal Bleeding Subscore, based on the participant's diary and scored from 0 (no blood) to 3 (blood only passed); Endoscopy Subscore, based on central reading of colonoscopy or sigmoidoscopy and scored from 0 (normal or inactive disease) to 3 (severe disease, spontaneous bleeding, ulceration). Analysis population description (APD): Modified Intention-to-treat population (mITT): All randomized participants who received at least one dose of study drug and who had the modified Mayo score measured correctly at baseline. Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Primary
----------------	---------

End point timeframe:

Week 40

Notes:

[1] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: percentage of participants				
number (confidence interval 95%)	25.1 (18.8 to 31.5)	49.9 (44.7 to 55.0)		

Statistical analyses

Statistical analysis title	Percentage of Participants in Clinical Remission
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	Cochran-Mantel-Haenszel
Parameter estimate	Risk difference (RD)
Point estimate	23.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	15.2
upper limit	31.2

Secondary: Percentage of Participants in Endoscopic Remission at Week 40 (Mirikizumab Induction Responders)

End point title	Percentage of Participants in Endoscopic Remission at Week 40 (Mirikizumab Induction Responders) ^[2]
-----------------	---

End point description:

Endoscopic remission at week 40 is defined as achieving a Mayo endoscopic subscore of 0 or 1 (excluding friability) at Week 40. Endoscopy subscore is based on colonoscopy or sigmoidoscopy and scored from 0 (normal or inactive disease) to 3 (severe disease, spontaneous bleeding, ulceration). APD: Modified Intention-to-treat population (mITT): All randomized participants who received at least one dose of study drug and who had the modified Mayo score measured correctly at baseline. Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
End point timeframe:	
Week 40	

Notes:

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: percentage of participants				
number (confidence interval 95%)	29.1 (22.4 to 35.7)	58.6 (53.6 to 63.7)		

Statistical analyses

Statistical analysis title	Percentage of Participants in Endoscopic Remission
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	Cochran-Mantel-Haenszel
Parameter estimate	Risk difference (RD)
Point estimate	28.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	20.2
upper limit	36.8

Secondary: Percentage of Participants with Histologic Remission at Week 40 (Mirikizumab Induction Responders)

End point title	Percentage of Participants with Histologic Remission at Week 40 (Mirikizumab Induction Responders) ^[3]
-----------------	---

End point description:

Histologic remission was assessed using the Geboes histologic scoring system developed for assessment of histologic disease activity in ulcerative colitis. Remission was defined as Geboes histological subscore of 0 for grades: 2b (lamina propria neutrophils), and 3 (neutrophils in epithelium), and 4 (crypt destruction), and 5 (erosion or ulceration). APD: Modified Intention-to-treat population (mITT): All randomized participants (pts) who received at least one dose of study drug and who had the modified Mayo score measured correctly at baseline. Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
----------------	-----------

End point timeframe:

Week 40

Notes:

[3] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: percentage of participants				
number (confidence interval 95%)	24.6 (18.3 to 30.9)	48.5 (43.4 to 53.6)		

Statistical analyses

Statistical analysis title	Percentage of Pts with Histologic Remission
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	Cochran-Mantel-Haenszel
Parameter estimate	Risk difference (RD)
Point estimate	22.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	14.5
upper limit	30.5

Secondary: Percentage of Participants in Symptomatic Remission at Week 40 (Mirikizumab Induction Responders)

End point title	Percentage of Participants in Symptomatic Remission at Week 40 (Mirikizumab Induction Responders) ^[4]
-----------------	--

End point description:

Symptomatic remission at week 40 is defined as a Mayo score for rectal bleeding=0, stool frequency=0 or 1 with ≥ 1 point decrease from baseline. Stool frequency subscore, based on the participant's diary and scored from 0 (normal number of stools) to 3 (5 or more stools than normal). Rectal bleeding subscore, based on the participant's diary and scored from 0 (no blood seen) to 3 (blood alone passed). APD: Modified Intention-to-treat population (mITT): All randomized participants who received at least one dose of study drug and who had the modified Mayo score measured correctly at baseline. Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
----------------	-----------

End point timeframe:

Week 40

Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: percentage of participants				
number (confidence interval 95%)	39.7 (32.5 to 46.8)	71.0 (66.3 to 75.6)		

Statistical analyses

Statistical analysis title	Percentage of Pts in Symptomatic Remission
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	Cochran-Mantel-Haenszel
Parameter estimate	Risk difference (RD)
Point estimate	30.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	21.9
upper limit	38.6

Secondary: Percentage of Participants in Endoscopic Response at Week 40 (Mirikizumab Induction Responders)

End point title	Percentage of Participants in Endoscopic Response at Week 40 (Mirikizumab Induction Responders) ^[5]
-----------------	--

End point description:

Endoscopic response at week 40 is defined as achieving at least a 1 point decrease from baseline in the Mayo endoscopic subscore. APD: Modified Intention-to-treat population (mITT): All randomized participants who received at least one dose of study drug and who had the modified Mayo score measured correctly at baseline. Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
----------------	-----------

End point timeframe:

Week 40

Notes:

[5] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: percentage of participants				
number (confidence interval 95%)	40.8 (33.6 to 48.0)	72.6 (68.0 to 77.2)		

Statistical analyses

Statistical analysis title	Percentage of Participants in Endoscopic Response
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	Cochran-Mantel-Haenszel
Parameter estimate	Risk difference (RD)
Point estimate	31
Confidence interval	
level	95 %
sides	2-sided
lower limit	22.4
upper limit	39.6

Secondary: Percentage of Participants in Clinical Response at Week 40 (Mirikizumab Induction Responders)

End point title	Percentage of Participants in Clinical Response at Week 40 (Mirikizumab Induction Responders) ^[6]
-----------------	--

End point description:

Clinical response at week 40 is defined as a decrease in the 9-point modified Mayo score (MMS) [rectal bleeding, stool frequency and the endoscopic findings] inclusive of ≥ 2 points and $\geq 30\%$ from baseline with either a decrease of rectal bleeding subscore of ≥ 1 or rectal bleeding subscore of 0 or 1. The MMS is a composite score of ulcerative colitis disease activity calculated as the sum of three subscores: Stool frequency subscore, based on the participant's diary and scored from 0 (normal number of stools) to 3 (5 or more stools than normal); Rectal bleeding subscore, based on the participant's diary and scored from 0 (no blood seen) to 3 (blood alone passed); Endoscopy subscore, based on colonoscopy or sigmoidoscopy and scored from 0 (normal or inactive disease) to 3 (severe disease, spontaneous bleeding, ulceration). APD: mITT: All randomized participants who received at least one dose of study drug and who had the modified Mayo score measured correctly at baseline.

End point type	Secondary
----------------	-----------

End point timeframe:

Week 40

Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

Notes:

[6] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: percentage of participants				
number (confidence interval 95%)	49.2 (41.8 to 56.5)	80.3 (76.2 to 84.4)		

Statistical analyses

Statistical analysis title	Percentage of Participants in Clinical Response
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	Cochran-Mantel-Haenszel
Parameter estimate	Risk difference (RD)
Point estimate	30.6
Confidence interval	
level	95 %
sides	2-sided
lower limit	22.3
upper limit	38.9

Secondary: Change from Baseline to Week 40 in Health Related Quality of Life: Inflammatory Bowel Disease Questionnaire (IBDQ) Total Score (Mirikizumab Induction Responders)

End point title	Change from Baseline to Week 40 in Health Related Quality of Life: Inflammatory Bowel Disease Questionnaire (IBDQ) Total Score (Mirikizumab Induction Responders) ^[7]
-----------------	--

End point description:

The IBDQ is a 32-item participant-completed questionnaire that measures 4 aspects of subjects' lives: symptoms directly related to the primary bowel disturbance, systemic symptoms, emotional function, and social function. Responses are graded on a 7-point. Likert scale in which 7 denotes "not a problem at all" and 1 denotes "a very severe problem." Scores range from 32 to 224; a higher score indicates a better quality of life. Least square (LS) Mean was calculated using analysis of covariance (ANCOVA) model for post-baseline measures: The ANCOVA model includes: treatment, baseline value, prior biologic or tofacitinib failure (yes/no), baseline corticosteroid use (yes/no), clinical remission status (yes/no) at AMAN Week 12, and region (North America/Europe/Other). Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
----------------	-----------

End point timeframe:

Induction Baseline, Week 40

APD: mITT: All randomized participants who received at least 1 dose of study drug and had a baseline and at least one post-baseline IBDQ measurement.

Notes:

[7] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: score on a scale				
least squares mean (standard error)	24.51 (\pm 2.767)	49.75 (\pm 2.102)		

Statistical analyses

Statistical analysis title	Change from Baseline to Week 40 in IBDQ
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001 [8]
Method	ANCOVA
Parameter estimate	LS Mean difference (Final Values)
Point estimate	25.24
Confidence interval	
level	95 %
sides	2-sided
lower limit	19.16
upper limit	31.32
Variability estimate	Standard error of the mean
Dispersion value	3.094

Notes:

[8] - ANCOVA with modified baseline observation carried forward (mBOCF).

Secondary: Change from Baseline to Week 40 in Fecal Calprotectin (Mirikizumab Induction Responders)

End point title	Change from Baseline to Week 40 in Fecal Calprotectin (Mirikizumab Induction Responders) ^[9]
-----------------	---

End point description:

Fecal calprotectin is an indicator of inflammation in the colon with higher levels indicative of higher levels of inflammation. Least square (LS) Mean was calculated using ANCOVA model for post-baseline measures: The ANCOVA model includes treatment, baseline value, prior biologic or tofacitinib failure (yes/no), corticosteroid use (yes/no) at AMAN baseline, region (North America/Europe/Other), Clinical Remission status (yes/no) at AMAN Week 12. APD: Modified Intention-to-treat population (mITT): All randomized participants who received at least 1 dose of study drug and had a baseline and at least one

post-baseline fecal calprotectin measurement. Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
----------------	-----------

End point timeframe:

Induction Baseline, Week 40

Notes:

[9] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: milligram per kilogram (mg/kg)				
least squares mean (standard error)	-1155.82 (\pm 221.394)	-1995.47 (\pm 172.443)		

Statistical analyses

Statistical analysis title	Change from Baseline in Fecal Calprotectin
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	544
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001 ^[10]
Method	ANCOVA
Parameter estimate	LS Mean difference (Final Values)
Point estimate	-839.64
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1323.08
upper limit	-356.21
Variability estimate	Standard error of the mean
Dispersion value	245.99

Notes:

[10] - ANCOVA with modified baseline observation carried forward (mBOCF).

Secondary: Change From Baseline to Week 40 in Bowel Urgency Based on the Urgency Numeric Rating Scale (NRS) (Mirikizumab Induction Responders)

End point title	Change From Baseline to Week 40 in Bowel Urgency Based on the Urgency Numeric Rating Scale (NRS) (Mirikizumab Induction Responders) ^[11]
-----------------	---

End point description:

The Urgency NRS is a single participant reported item that measures the severity for the urgency (sudden or immediate need) to have a bowel movement in the past 24 hours using an 11-point NRS ranging from 0 (no urgency) to 10 (worst possible urgency). Higher scores indicate more severe

urgency. Least square (LS) Mean was calculated using mixed model repeated measures (MMRM) model for post-baseline measures: The MMRM model includes treatment, baseline value, visit, interaction of baseline value-by-visit, interaction of treatment-by-visit, prior biologic or tofacitinib failure (yes/no), baseline corticosteroid use (yes/no), clinical remission status (yes/no) at AMAN Week 12, and region (North America/Europe/Other). Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
----------------	-----------

End point timeframe:

Induction Baseline, Week 40

APD: Modified Intention-to-treat population (mITT): All randomized participants who received at least 1 dose of study drug and had a baseline and at least one post-baseline urgency NRS measurement.

Notes:

[11] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	104	316		
Units: score on a scale				
least squares mean (standard error)	-2.74 (± 0.202)	-3.80 (± 0.139)		

Statistical analyses

Statistical analysis title	Change From Baseline to Week 40 in NRS
Comparison groups	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC) v Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC
Number of subjects included in analysis	420
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.001
Method	Mixed models analysis
Parameter estimate	LS Mean difference (Final Values)
Point estimate	-1.06
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.51
upper limit	-0.61
Variability estimate	Standard error of the mean
Dispersion value	0.228

Secondary: Percentage of Participants Hospitalized for Ulcerative Colitis (UC) (Mirikizumab Induction Responders)

End point title	Percentage of Participants Hospitalized for Ulcerative Colitis
-----------------	--

End point description:

Percentage of participants hospitalized for UC. Only hospitalizations associated with an adverse event with ≥ 24 hours stay were recorded. APD: Modified Intention-to-treat population (mITT): All randomized participants who received at least one dose of study drug. Participants were analysed per their assigned treatment arm regardless of the treatment they actually received.

End point type	Secondary
----------------	-----------

End point timeframe:

Week 40

Notes:

[12] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is reported only for randomized arms as described in the protocol.

End point values	Maintenance Period: Miri IR - Placebo (PBO) Subcutaneous (SC)	Maintenance Period: Miri IR - 200 Milligram (mg) Miri SC		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	365		
Units: percentage of participants				
number (not applicable)	1.1	0		

Statistical analyses

No statistical analyses for this end point

Secondary: Pharmacokinetics (PK): Clearance of Mirikizumab

End point title	Pharmacokinetics (PK): Clearance of Mirikizumab
-----------------	---

End point description:

Clearance of mirikizumab was evaluated. APD: All randomized participants who received at least one dose of study drug subcutaneously (both induction responders and nonresponders) and had evaluable PK data.

End point type	Secondary
----------------	-----------

End point timeframe:

Predose: Weeks 0, 4, 12, 24 and 40

End point values	200 Milligram (mg) Miri SC			
Subject group type	Subject analysis set			
Number of subjects analysed	652			
Units: Liters per Hour (L/h)				
geometric mean (geometric coefficient of variation)	0.0487 (\pm 54)			

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

Baseline Up to week 56

Adverse event reporting additional description:

All randomized participants who received at least one dose of study drug. Gender-specific events occurring exclusively in female participants have had the number of participants at risk adjusted accordingly.

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	28.0
--------------------	------

Reporting groups

Reporting group title	Maintenance Period: PBO IR - PBO SC
-----------------------	-------------------------------------

Reporting group description:

Participants who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Reporting group title	Maintenance Period: Miri IR - 200 mg Miri SC
-----------------------	--

Reporting group description:

Participants who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Reporting group title	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
-----------------------	--

Reporting group description:

Participants in the ME2 Cohort who were responders to blinded placebo at Week 12 in induction study (LUCENT-1) continue to receive blinded placebo SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Reporting group title	Maintenance Period: Miri IR - PBO SC - ME2 Cohort
-----------------------	---

Reporting group description:

Participants in the ME2 Cohort who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.

Reporting group title	Open Label Maintenance: Delayed Responders-200 mg Miri SC-ME2
-----------------------	---

Reporting group description:

Participants in the ME2 Cohort who initially did not respond to induction study (LUCENT-1), but responded to extended induction therapy at Week 12 of LUCENT-2 (delayed responders), received 200 mg mirikizumab SC Q4W during open label maintenance period from Week 12 until Week 40 or until early termination (rescue was not available for these participants).

Reporting group title	LOR Rescue Period: LOR Cohort-300 mg Miri IV
-----------------------	--

Reporting group description:

Participants who received blinded PBO SC or blinded 200 mg mirikizumab SC Q4W during maintenance period and experienced a loss of response at or after Week 12, received rescue therapy with open label 300 mg mirikizumab IV Q4W for 3 doses.

Reporting group title	LOR Rescue Period: LOR Cohort-300 mg Miri IV - ME2 Cohort
-----------------------	---

Reporting group description:

Participants in the ME2 Cohort who received blinded PBO SC or blinded 200 mg mirikizumab SC Q4W during maintenance period and experienced a loss of response at or after Week 12, received rescue therapy with open label 300 mg mirikizumab IV Q4W for 3 doses.

Reporting group title	Extended Induction: Induction Nonresponders - 300mg Miri IV
-----------------------	---

Reporting group description:

Participants who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period

from Week 0 of LUCENT-2 until Week 12.

Reporting group title	Extended Induction: Induction Nonresponders-300mg Miri IV-ME2
-----------------------	---

Reporting group description:

Participants in the ME2 Cohort who were nonresponders to blinded mirikizumab or placebo in induction study (LUCENT-1), received additional 3 doses of open label 300 mg mirikizumab IV Q4W during extended induction period from Week 0 of LUCENT-2 until Week 12.

Reporting group title	Open Label Maintenance: Delayed Responders - 200 mg Miri SC
-----------------------	---

Reporting group description:

Participants who initially did not respond to induction study (LUCENT-1) but responded to extended induction therapy at Week 12 of LUCENT-2 (delayed responders), received 200 mg mirikizumab SC Q4W during open label maintenance period from Week 12 until Week 40 or until early termination (rescue was not available for these participants).

Reporting group title	Maintenance Period: Miri IR - PBO SC
-----------------------	--------------------------------------

Reporting group description:

Participants who were responders to blinded mirikizumab (miri) at Week 12 in induction study (LUCENT-1) randomized to withdraw from mirikizumab and start receiving PBO SC every 4 weeks (Q4W) from Week 0 of maintenance study (LUCENT-2) until Week 40 or until loss of response was confirmed.

Reporting group title	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort
-----------------------	---

Reporting group description:

Participants in the ME2 Cohort who were responders to blinded mirikizumab at Week 12 in induction study (LUCENT-1) randomized to continue to receive 200 mg mirikizumab SC Q4W from Week 0 of LUCENT-2 until Week 40 or until loss of response was confirmed.

Serious adverse events	Maintenance Period: PBO IR - PBO SC	Maintenance Period: Miri IR - 200 mg Miri SC	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
Total subjects affected by serious adverse events			
subjects affected / exposed	8 / 135 (5.93%)	13 / 389 (3.34%)	1 / 12 (8.33%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
adenocarcinoma of colon			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colorectal adenoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastric cancer			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
kaposi's sarcoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
lipoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
neuroendocrine tumour			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
plasmablastic lymphoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
rectal cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
thyroid cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Vascular disorders			
deep vein thrombosis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 135 (0.74%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
granulomatosis with polyangiitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypovolaemic shock			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Surgical and medical procedures			
colectomy total			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
proctocolectomy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinopexy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tooth extraction			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
abortion missed			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[1]	1 / 61 (1.64%)	0 / 160 (0.00%)	0 / 5 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			
anaphylactic reaction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders			
rectocele			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[2]	0 / 61 (0.00%)	1 / 160 (0.63%)	0 / 5 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
asthma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
depression suicidal			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

neurologic somatic symptom disorder			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
blood glucose increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
human chorionic gonadotropin increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
anastomotic fistula			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ankle fracture			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	1 / 12 (8.33%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
brain contusion			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 135 (0.74%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

compression fracture alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 135 (0.00%) 0 / 0 0 / 0	1 / 389 (0.26%) 0 / 1 0 / 0	0 / 12 (0.00%) 0 / 0 0 / 0
patella fracture alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 135 (0.74%) 0 / 1 0 / 0	0 / 389 (0.00%) 0 / 0 0 / 0	0 / 12 (0.00%) 0 / 0 0 / 0
road traffic accident alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	1 / 135 (0.74%) 0 / 1 0 / 0	0 / 389 (0.00%) 0 / 0 0 / 0	0 / 12 (0.00%) 0 / 0 0 / 0
tibia fracture alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 135 (0.00%) 0 / 0 0 / 0	0 / 389 (0.00%) 0 / 0 0 / 0	0 / 12 (0.00%) 0 / 0 0 / 0
Cardiac disorders acute coronary syndrome alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 135 (0.00%) 0 / 0 0 / 0	0 / 389 (0.00%) 0 / 0 0 / 0	0 / 12 (0.00%) 0 / 0 0 / 0
myocardial infarction alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	0 / 135 (0.00%) 0 / 0 0 / 0	0 / 389 (0.00%) 0 / 0 0 / 0	0 / 12 (0.00%) 0 / 0 0 / 0
Nervous system disorders headache alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ischaemic stroke			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
migraine			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
presyncope			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
immune thrombocytopenia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
maculopathy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinal detachment			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
anorectal polyp			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis ulcerative			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 135 (0.74%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
diarrhoea			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ileus			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
inguinal hernia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
intestinal obstruction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine perforation			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine polyp			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
peptic ulcer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
pneumoperitoneum			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

rectal haemorrhage			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
pruritus			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
xanthelasma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
nephrolithiasis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ureterolithiasis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Endocrine disorders			
autoimmune thyroiditis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			

arthralgia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
back pain			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
intervertebral disc protrusion			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
sacral pain			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 135 (0.74%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
abdominal abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
anal abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
appendicitis			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
bacillus infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
covid-19			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
covid-19 pneumonia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
campylobacter gastroenteritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
cellulitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
clostridium difficile infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

cytomegalovirus colitis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	1 / 135 (0.74%)	0 / 389 (0.00%)	0 / 12 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
diverticulitis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
escherichia infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
gastroenteritis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	1 / 135 (0.74%)	1 / 389 (0.26%)	0 / 12 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
influenza				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
klebsiella infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
large intestine infection				
alternative dictionary used: MedDRA 28.0				

subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
pneumonia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 135 (0.74%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
sepsis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
subcutaneous abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tonsillitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
hypoglycaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypokalaemia			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	0 / 12 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Maintenance Period: Miri IR - PBO SC - ME2 Cohort	Open Label Maintenance:Delayed Responders-200 mg Miri SC-ME2	LOR Rescue Period: LOR Cohort-300 mg Miri IV
Total subjects affected by serious adverse events			
subjects affected / exposed	4 / 21 (19.05%)	3 / 33 (9.09%)	3 / 90 (3.33%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
adenocarcinoma of colon			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colorectal adenoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastric cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
kaposi's sarcoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
lipoma			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
neuroendocrine tumour			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
plasmablastic lymphoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
rectal cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
thyroid cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
deep vein thrombosis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
granulomatosis with polyangiitis			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypovolaemic shock			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Surgical and medical procedures			
colectomy total			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
proctocolectomy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinopexy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tooth extraction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
abortion missed			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed ^[1]	0 / 6 (0.00%)	0 / 7 (0.00%)	0 / 43 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			
anaphylactic reaction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders			
rectocele			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[2]	0 / 6 (0.00%)	0 / 7 (0.00%)	0 / 43 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
asthma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
depression suicidal			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
neurologic somatic symptom disorder			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	1 / 90 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			

blood glucose increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
human chorionic gonadotropin increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed	0 / 21 (0.00%)	1 / 33 (3.03%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications anastomotic fistula alternative dictionary used: MedDRA 28.0 subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ankle fracture alternative dictionary used: MedDRA 28.0 subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
brain contusion alternative dictionary used: MedDRA 28.0 subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
compression fracture alternative dictionary used: MedDRA 28.0 subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
patella fracture alternative dictionary used:			

MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
road traffic accident			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tibia fracture			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders			
acute coronary syndrome			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
myocardial infarction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
headache			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 21 (4.76%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ischaemic stroke			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
migraine			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
presyncope			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
immune thrombocytopenia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
maculopathy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinal detachment			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
anorectal polyp			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis ulcerative			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 21 (4.76%)	2 / 33 (6.06%)	1 / 90 (1.11%)
occurrences causally related to treatment / all	1 / 2	0 / 2	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
diarrhoea			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ileus			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
inguinal hernia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 21 (4.76%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

intestinal obstruction alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine perforation alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine polyp alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
peptic ulcer alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
pneumoperitoneum alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
rectal haemorrhage alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders pruritus alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
xanthelasma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
nephrolithiasis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ureterolithiasis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	1 / 33 (3.03%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Endocrine disorders			
autoimmune thyroiditis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
arthralgia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	1 / 90 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
back pain			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
intervertebral disc protrusion alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
sacral pain alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
abdominal abscess alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
anal abscess alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	1 / 33 (3.03%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
appendicitis alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
bacillus infection alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
covid-19			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
covid-19 pneumonia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
campylobacter gastroenteritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
cellulitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
clostridium difficile infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 21 (4.76%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
cytomegalovirus colitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

diverticulitis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
escherichia infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
gastroenteritis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
influenza				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
klebsiella infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
large intestine infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
pneumonia				
alternative dictionary used: MedDRA 28.0				

subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	1 / 90 (1.11%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
sepsis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
subcutaneous abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tonsillitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
hypoglycaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypokalaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	LOR Rescue Period: LOR Cohort-300 mg Miri IV - ME2 Cohort	Extended Induction: Induction Nonresponders - 300mg Miri IV	Extended Induction: Induction Nonresponders- 300mg Miri IV-ME2
Total subjects affected by serious adverse events			

subjects affected / exposed	2 / 12 (16.67%)	24 / 461 (5.21%)	4 / 66 (6.06%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
adenocarcinoma of colon			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colorectal adenoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 12 (8.33%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastric cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
kaposi's sarcoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
lipoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
neuroendocrine tumour			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
plasmablastic lymphoma alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
rectal cancer alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
thyroid cancer alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
deep vein thrombosis alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
granulomatosis with polyangiitis alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypovolaemic shock alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Surgical and medical procedures			
colectomy total			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
proctocolectomy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinopexy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tooth extraction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
abortion missed			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[1]	0 / 7 (0.00%)	0 / 166 (0.00%)	0 / 18 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			
anaphylactic reaction			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders			
rectocele			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[2]	0 / 7 (0.00%)	0 / 166 (0.00%)	0 / 18 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
asthma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
depression suicidal			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
neurologic somatic symptom disorder			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
blood glucose increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
human chorionic gonadotropin			

increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
anastomotic fistula			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	1 / 66 (1.52%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ankle fracture			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
brain contusion			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
compression fracture			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
patella fracture			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
road traffic accident			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tibia fracture			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders			
acute coronary syndrome			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
myocardial infarction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
headache			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ischaemic stroke			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
migraine			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
presyncope			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
immune thrombocytopenia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
maculopathy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinal detachment			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
anorectal polyp			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis ulcerative			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 12 (8.33%)	9 / 461 (1.95%)	2 / 66 (3.03%)
occurrences causally related to treatment / all	1 / 1	0 / 9	1 / 2
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
diarrhoea			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ileus			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
inguinal hernia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
intestinal obstruction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	1 / 66 (1.52%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

large intestine perforation alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine polyp alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	1 / 66 (1.52%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
peptic ulcer alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
pneumoperitoneum alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
rectal haemorrhage alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
pruritus alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
xanthelasma alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
nephrolithiasis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ureterolithiasis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Endocrine disorders			
autoimmune thyroiditis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
arthralgia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
back pain			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
intervertebral disc protrusion			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
sacral pain			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
abdominal abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
anal abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
appendicitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
bacillus infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
covid-19			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
covid-19 pneumonia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
campylobacter gastroenteritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
cellulitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
clostridium difficile infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
cytomegalovirus colitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
diverticulitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

escherichia infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
gastroenteritis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
influenza				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
klebsiella infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
large intestine infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
pneumonia				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 12 (0.00%)	3 / 461 (0.65%)	0 / 66 (0.00%)	
occurrences causally related to treatment / all	0 / 0	1 / 3	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
sepsis				
alternative dictionary used: MedDRA 28.0				

subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
subcutaneous abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tonsillitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
hypoglycaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypokalaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Open Label Maintenance: Delayed Responders - 200 mg Miri SC	Maintenance Period: Miri IR - PBO SC	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort
Total subjects affected by serious adverse events			
subjects affected / exposed	9 / 271 (3.32%)	16 / 192 (8.33%)	7 / 52 (13.46%)
number of deaths (all causes)	0	1	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
adenocarcinoma of colon			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colorectal adenoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastric cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
kaposi's sarcoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
lipoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
neuroendocrine tumour			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
plasmablastic lymphoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	1 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

rectal cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
thyroid cancer			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
deep vein thrombosis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
granulomatosis with polyangiitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypovolaemic shock			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Surgical and medical procedures			
colectomy total			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
proctocolectomy			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinopexy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tooth extraction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
abortion missed			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[1]	0 / 111 (0.00%)	0 / 78 (0.00%)	0 / 23 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			
anaphylactic reaction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders			
rectocele			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[2]	0 / 111 (0.00%)	0 / 78 (0.00%)	0 / 23 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			

asthma alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	 0 / 271 (0.00%) 0 / 0 0 / 0	 1 / 192 (0.52%) 0 / 1 0 / 0	 0 / 52 (0.00%) 0 / 0 0 / 0
Psychiatric disorders depression suicidal alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	 0 / 271 (0.00%) 0 / 0 0 / 0	 0 / 192 (0.00%) 0 / 0 0 / 0	 0 / 52 (0.00%) 0 / 0 0 / 0
neurologic somatic symptom disorder alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	 0 / 271 (0.00%) 0 / 0 0 / 0	 0 / 192 (0.00%) 0 / 0 0 / 0	 0 / 52 (0.00%) 0 / 0 0 / 0
Investigations blood glucose increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	 0 / 271 (0.00%) 0 / 0 0 / 0	 0 / 192 (0.00%) 0 / 0 0 / 0	 0 / 52 (0.00%) 0 / 0 0 / 0
human chorionic gonadotropin increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	 0 / 271 (0.00%) 0 / 0 0 / 0	 0 / 192 (0.00%) 0 / 0 0 / 0	 0 / 52 (0.00%) 0 / 0 0 / 0
Injury, poisoning and procedural complications anastomotic fistula alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences causally related to treatment / all deaths causally related to treatment / all	 0 / 271 (0.00%) 0 / 0 0 / 0	 0 / 192 (0.00%) 0 / 0 0 / 0	 0 / 52 (0.00%) 0 / 0 0 / 0

ankle fracture alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
brain contusion alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
compression fracture alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
patella fracture alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
road traffic accident alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tibia fracture alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cardiac disorders acute coronary syndrome alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	1 / 2	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
myocardial infarction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
headache			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ischaemic stroke			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
migraine			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
presyncope			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
immune thrombocytopenia			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
maculopathy			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
retinal detachment			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
anorectal polyp			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
colitis ulcerative			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	3 / 271 (1.11%)	7 / 192 (3.65%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	2 / 3	0 / 7	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
diarrhoea			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
gastritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ileus			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
inguinal hernia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
intestinal obstruction			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine perforation			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine polyp			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

peptic ulcer alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
pneumoperitoneum alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
rectal haemorrhage alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
pruritus alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
xanthelasma alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
nephrolithiasis alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
ureterolithiasis alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Endocrine disorders			
autoimmune thyroiditis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
arthralgia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
back pain			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
intervertebral disc protrusion			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
sacral pain			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
abdominal abscess			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
anal abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
appendicitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
bacillus infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
covid-19			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 0
covid-19 pneumonia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
campylobacter gastroenteritis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

cellulitis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	0 / 52 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
clostridium difficile infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
cytomegalovirus colitis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
diverticulitis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
escherichia infection				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
gastroenteritis				
alternative dictionary used: MedDRA 28.0				
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)	
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0	
influenza				
alternative dictionary used: MedDRA 28.0				

subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	1 / 52 (1.92%)
occurrences causally related to treatment / all	0 / 1	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
klebsiella infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
large intestine infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
pneumonia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
sepsis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 271 (0.37%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
subcutaneous abscess			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
tonsillitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Metabolism and nutrition disorders			
hypoglycaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
hypokalaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	0 / 192 (0.00%)	0 / 52 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Notes:

[1] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed to this adverse event. These numbers are expected to be equal.

Justification: Gender-specific events occurring exclusively in female participants have had the number of participants at risk adjusted accordingly.

[2] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed to this adverse event. These numbers are expected to be equal.

Justification: Gender-specific events occurring exclusively in female participants have had the number of participants at risk adjusted accordingly.

Frequency threshold for reporting non-serious adverse events: 5 %

Non-serious adverse events	Maintenance Period: PBO IR - PBO SC	Maintenance Period: Miri IR - 200 mg Miri SC	Maintenance Period: PBO IR - PBO SC - ME2 Cohort
Total subjects affected by non-serious adverse events			
subjects affected / exposed	58 / 135 (42.96%)	143 / 389 (36.76%)	10 / 12 (83.33%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
uterine leiomyoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[3]	0 / 61 (0.00%)	0 / 160 (0.00%)	0 / 5 (0.00%)
occurrences (all)	0	0	0
Vascular disorders			
hypotension			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	1 / 12 (8.33%)
occurrences (all)	0	0	1
General disorders and administration site conditions			
oedema peripheral			
alternative dictionary used: MedDRA 28.0			

<p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>pyrexia</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 135 (0.00%)</p> <p>0</p> <p>6 / 135 (4.44%)</p> <p>8</p>	<p>2 / 389 (0.51%)</p> <p>2</p> <p>13 / 389 (3.34%)</p> <p>14</p>	<p>0 / 12 (0.00%)</p> <p>0</p> <p>1 / 12 (8.33%)</p> <p>1</p>
<p>Respiratory, thoracic and mediastinal disorders</p> <p>pulmonary mass</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 135 (0.00%)</p> <p>0</p>	<p>0 / 389 (0.00%)</p> <p>0</p>	<p>0 / 12 (0.00%)</p> <p>0</p>
<p>Investigations</p> <p>alanine aminotransferase increased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>aspartate aminotransferase increased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>blood alkaline phosphatase increased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>blood creatine phosphokinase increased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>weight increased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>weight decreased</p> <p>alternative dictionary used: MedDRA 28.0</p>	<p>1 / 135 (0.74%)</p> <p>1</p> <p>0 / 135 (0.00%)</p> <p>0</p> <p>0 / 135 (0.00%)</p> <p>0</p> <p>5 / 135 (3.70%)</p> <p>5</p> <p>1 / 135 (0.74%)</p> <p>1</p>	<p>4 / 389 (1.03%)</p> <p>4</p> <p>3 / 389 (0.77%)</p> <p>3</p> <p>1 / 389 (0.26%)</p> <p>1</p> <p>10 / 389 (2.57%)</p> <p>11</p> <p>1 / 389 (0.26%)</p> <p>1</p>	<p>0 / 12 (0.00%)</p> <p>0</p> <p>0 / 12 (0.00%)</p> <p>0</p> <p>0 / 12 (0.00%)</p> <p>0</p> <p>1 / 12 (8.33%)</p> <p>1</p> <p>0 / 12 (0.00%)</p> <p>0</p>

<p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>white blood cell count decreased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>2 / 135 (1.48%)</p> <p>2</p> <p>1 / 135 (0.74%)</p> <p>1</p>	<p>0 / 389 (0.00%)</p> <p>0</p> <p>3 / 389 (0.77%)</p> <p>4</p>	<p>1 / 12 (8.33%)</p> <p>1</p> <p>0 / 12 (0.00%)</p> <p>0</p>
<p>Injury, poisoning and procedural complications</p> <p>ankle fracture</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 135 (0.00%)</p> <p>0</p>	<p>0 / 389 (0.00%)</p> <p>0</p>	<p>1 / 12 (8.33%)</p> <p>1</p>
<p>Cardiac disorders</p> <p>sinus bradycardia</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>ventricular extrasystoles</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 135 (0.00%)</p> <p>0</p> <p>0 / 135 (0.00%)</p> <p>0</p>	<p>0 / 389 (0.00%)</p> <p>0</p> <p>0 / 389 (0.00%)</p> <p>0</p>	<p>1 / 12 (8.33%)</p> <p>1</p> <p>1 / 12 (8.33%)</p> <p>1</p>
<p>Nervous system disorders</p> <p>dizziness</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>headache</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>sleep deficit</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 135 (0.00%)</p> <p>0</p> <p>5 / 135 (3.70%)</p> <p>18</p> <p>0 / 135 (0.00%)</p> <p>0</p>	<p>1 / 389 (0.26%)</p> <p>1</p> <p>16 / 389 (4.11%)</p> <p>21</p> <p>0 / 389 (0.00%)</p> <p>0</p>	<p>1 / 12 (8.33%)</p> <p>1</p> <p>0 / 12 (0.00%)</p> <p>0</p> <p>1 / 12 (8.33%)</p> <p>1</p>
<p>Blood and lymphatic system disorders</p>			

<p>anaemia</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>8 / 135 (5.93%)</p> <p>11</p>	<p>8 / 389 (2.06%)</p> <p>8</p>	<p>2 / 12 (16.67%)</p> <p>3</p>
<p>Gastrointestinal disorders</p> <p>abdominal pain</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>anorectal polyp</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>colitis ulcerative</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>diarrhoea</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>haemorrhoids</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>large intestine polyp</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>3 / 135 (2.22%)</p> <p>3</p> <p>0 / 135 (0.00%)</p> <p>0</p> <p>17 / 135 (12.59%)</p> <p>18</p> <p>3 / 135 (2.22%)</p> <p>3</p> <p>2 / 135 (1.48%)</p> <p>2</p> <p>1 / 135 (0.74%)</p> <p>2</p>	<p>11 / 389 (2.83%)</p> <p>13</p> <p>0 / 389 (0.00%)</p> <p>0</p> <p>26 / 389 (6.68%)</p> <p>30</p> <p>10 / 389 (2.57%)</p> <p>10</p> <p>5 / 389 (1.29%)</p> <p>5</p> <p>3 / 389 (0.77%)</p> <p>3</p>	<p>0 / 12 (0.00%)</p> <p>0</p> <p>3 / 12 (25.00%)</p> <p>3</p> <p>0 / 12 (0.00%)</p> <p>0</p> <p>0 / 12 (0.00%)</p> <p>0</p>
<p>Skin and subcutaneous tissue disorders</p> <p>papule</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 135 (0.00%)</p> <p>0</p>	<p>0 / 389 (0.00%)</p> <p>0</p>	<p>1 / 12 (8.33%)</p> <p>1</p>
<p>Endocrine disorders</p>			

thyroiditis subacute alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 135 (0.00%) 0	0 / 389 (0.00%) 0	1 / 12 (8.33%) 1
Musculoskeletal and connective tissue disorders arthralgia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) back pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) osteoarthritis alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) neck pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	4 / 135 (2.96%) 4 4 / 135 (2.96%) 4 0 / 135 (0.00%) 0 1 / 135 (0.74%) 1	26 / 389 (6.68%) 27 6 / 389 (1.54%) 6 2 / 389 (0.51%) 2 3 / 389 (0.77%) 3	1 / 12 (8.33%) 2 1 / 12 (8.33%) 1 1 / 12 (8.33%) 1 1 / 12 (8.33%) 1
Infections and infestations covid-19 alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) bronchitis alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) clostridium difficile infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) gastrointestinal infection	5 / 135 (3.70%) 5 0 / 135 (0.00%) 0 1 / 135 (0.74%) 1	8 / 389 (2.06%) 8 4 / 389 (1.03%) 4 1 / 389 (0.26%) 1	1 / 12 (8.33%) 1 1 / 12 (8.33%) 1 0 / 12 (0.00%) 0

alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	1 / 389 (0.26%)	1 / 12 (8.33%)
occurrences (all)	0	1	1
influenza			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 135 (0.74%)	3 / 389 (0.77%)	0 / 12 (0.00%)
occurrences (all)	1	3	0
localised infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	1 / 12 (8.33%)
occurrences (all)	0	0	1
nasopharyngitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	9 / 135 (6.67%)	28 / 389 (7.20%)	0 / 12 (0.00%)
occurrences (all)	12	38	0
upper respiratory tract infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	2 / 135 (1.48%)	7 / 389 (1.80%)	1 / 12 (8.33%)
occurrences (all)	2	9	3
Metabolism and nutrition disorders			
electrolyte imbalance			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences (all)	0	0	0
hyperkalaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	1 / 12 (8.33%)
occurrences (all)	0	0	1
hypoalbuminaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 135 (0.00%)	0 / 389 (0.00%)	0 / 12 (0.00%)
occurrences (all)	0	0	0
hypokalaemia			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 135 (0.00%)	2 / 389 (0.51%)	1 / 12 (8.33%)
occurrences (all)	0	2	1

Non-serious adverse events	Maintenance Period: Miri IR - PBO SC - ME2 Cohort	Open Label Maintenance:Delayed Responders-200 mg Miri SC-ME2	LOR Rescue Period: LOR Cohort-300 mg Miri IV
Total subjects affected by non-serious adverse events			
subjects affected / exposed	16 / 21 (76.19%)	13 / 33 (39.39%)	18 / 90 (20.00%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
uterine leiomyoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[3]	0 / 6 (0.00%)	1 / 7 (14.29%)	0 / 43 (0.00%)
occurrences (all)	0	1	0
Vascular disorders			
hypotension			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences (all)	0	0	0
General disorders and administration site conditions			
oedema peripheral			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 21 (4.76%)	0 / 33 (0.00%)	1 / 90 (1.11%)
occurrences (all)	1	0	1
pyrexia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 21 (4.76%)	1 / 33 (3.03%)	1 / 90 (1.11%)
occurrences (all)	1	1	1
Respiratory, thoracic and mediastinal disorders			
pulmonary mass			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences (all)	0	0	0
Investigations			
alanine aminotransferase increased			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	2 / 21 (9.52%)	0 / 33 (0.00%)	1 / 90 (1.11%)
occurrences (all)	4	0	1
aspartate aminotransferase increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	2 / 21 (9.52%)	0 / 33 (0.00%)	1 / 90 (1.11%)
occurrences (all)	4	0	1
blood alkaline phosphatase increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	2 / 33 (6.06%)	0 / 90 (0.00%)
occurrences (all)	0	2	0
blood creatine phosphokinase increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences (all)	0	0	0
weight increased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	2 / 33 (6.06%)	0 / 90 (0.00%)
occurrences (all)	0	2	0
weight decreased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences (all)	0	0	0
white blood cell count decreased			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	1 / 33 (3.03%)	0 / 90 (0.00%)
occurrences (all)	0	2	0
Injury, poisoning and procedural complications			
ankle fracture			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 21 (0.00%)	0 / 33 (0.00%)	0 / 90 (0.00%)
occurrences (all)	0	0	0
Cardiac disorders			

sinus bradycardia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	1 / 33 (3.03%) 1	0 / 90 (0.00%) 0
ventricular extrasystoles alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	0 / 90 (0.00%) 0
Nervous system disorders dizziness alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) headache alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) sleep deficit alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0 1 / 21 (4.76%) 1 0 / 21 (0.00%) 0	0 / 33 (0.00%) 0 0 / 33 (0.00%) 0 0 / 33 (0.00%) 0	2 / 90 (2.22%) 2 5 / 90 (5.56%) 5 0 / 90 (0.00%) 0
Blood and lymphatic system disorders anaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	3 / 21 (14.29%) 3	1 / 33 (3.03%) 1	5 / 90 (5.56%) 5
Gastrointestinal disorders abdominal pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) anorectal polyp alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) colitis ulcerative	0 / 21 (0.00%) 0 0 / 21 (0.00%) 0	0 / 33 (0.00%) 0 2 / 33 (6.06%) 2	0 / 90 (0.00%) 0 0 / 90 (0.00%) 0

<p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>6 / 21 (28.57%)</p> <p>6</p>	<p>1 / 33 (3.03%)</p> <p>1</p>	<p>0 / 90 (0.00%)</p> <p>0</p>
<p>diarrhoea</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 21 (0.00%)</p> <p>0</p>	<p>2 / 33 (6.06%)</p> <p>3</p>	<p>0 / 90 (0.00%)</p> <p>0</p>
<p>haemorrhoids</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>1 / 21 (4.76%)</p> <p>1</p>	<p>0 / 33 (0.00%)</p> <p>0</p>	<p>0 / 90 (0.00%)</p> <p>0</p>
<p>large intestine polyp</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 21 (0.00%)</p> <p>0</p>	<p>1 / 33 (3.03%)</p> <p>1</p>	<p>0 / 90 (0.00%)</p> <p>0</p>
<p>Skin and subcutaneous tissue disorders</p> <p>papule</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 21 (0.00%)</p> <p>0</p>	<p>0 / 33 (0.00%)</p> <p>0</p>	<p>0 / 90 (0.00%)</p> <p>0</p>
<p>Endocrine disorders</p> <p>thyroiditis subacute</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 21 (0.00%)</p> <p>0</p>	<p>0 / 33 (0.00%)</p> <p>0</p>	<p>0 / 90 (0.00%)</p> <p>0</p>
<p>Musculoskeletal and connective tissue disorders</p> <p>arthralgia</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>back pain</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>osteoarthritis</p>	<p>1 / 21 (4.76%)</p> <p>1</p> <p>0 / 21 (0.00%)</p> <p>0</p>	<p>0 / 33 (0.00%)</p> <p>0</p> <p>0 / 33 (0.00%)</p> <p>0</p>	<p>3 / 90 (3.33%)</p> <p>5</p> <p>0 / 90 (0.00%)</p> <p>0</p>

alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	1 / 90 (1.11%) 1
neck pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	1 / 90 (1.11%) 1
Infections and infestations covid-19 alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	3 / 21 (14.29%) 3	1 / 33 (3.03%) 1	1 / 90 (1.11%) 1
bronchitis alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	1 / 33 (3.03%) 1	0 / 90 (0.00%) 0
clostridium difficile infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	2 / 21 (9.52%) 3	1 / 33 (3.03%) 1	2 / 90 (2.22%) 2
gastrointestinal infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	1 / 33 (3.03%) 1	0 / 90 (0.00%) 0
influenza alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	0 / 90 (0.00%) 0
localised infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	0 / 90 (0.00%) 0
nasopharyngitis alternative dictionary used: MedDRA 28.0			

subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	1 / 90 (1.11%) 1
upper respiratory tract infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	4 / 21 (19.05%) 5	4 / 33 (12.12%) 5	1 / 90 (1.11%) 1
Metabolism and nutrition disorders electrolyte imbalance alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	0 / 90 (0.00%) 0
hyperkalaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 21 (0.00%) 0	0 / 33 (0.00%) 0	0 / 90 (0.00%) 0
hypoalbuminaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	2 / 21 (9.52%) 2	0 / 33 (0.00%) 0	0 / 90 (0.00%) 0
hypokalaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	2 / 21 (9.52%) 4	1 / 33 (3.03%) 4	1 / 90 (1.11%) 1

Non-serious adverse events	LOR Rescue Period: LOR Cohort-300 mg Miri IV - ME2 Cohort	Extended Induction: Induction Nonresponders - 300mg Miri IV	Extended Induction: Induction Nonresponders- 300mg Miri IV-ME2
Total subjects affected by non-serious adverse events subjects affected / exposed	4 / 12 (33.33%)	74 / 461 (16.05%)	22 / 66 (33.33%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps) uterine leiomyoma alternative dictionary used: MedDRA 28.0 subjects affected / exposed ^[3] occurrences (all)	0 / 7 (0.00%) 0	0 / 166 (0.00%) 0	0 / 18 (0.00%) 0
Vascular disorders			

hypotension alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 12 (0.00%) 0	0 / 461 (0.00%) 0	0 / 66 (0.00%) 0
General disorders and administration site conditions oedema peripheral alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) pyrexia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 12 (8.33%) 1 0 / 12 (0.00%) 0	1 / 461 (0.22%) 1 3 / 461 (0.65%) 3	0 / 66 (0.00%) 0 0 / 66 (0.00%) 0
Respiratory, thoracic and mediastinal disorders pulmonary mass alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 12 (8.33%) 1	0 / 461 (0.00%) 0	0 / 66 (0.00%) 0
Investigations alanine aminotransferase increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) aspartate aminotransferase increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) blood alkaline phosphatase increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) blood creatine phosphokinase increased alternative dictionary used: MedDRA 28.0	0 / 12 (0.00%) 0 0 / 12 (0.00%) 0 0 / 12 (0.00%) 0 0 / 12 (0.00%) 0	0 / 461 (0.00%) 0 1 / 461 (0.22%) 1 1 / 461 (0.22%) 1	2 / 66 (3.03%) 2 1 / 66 (1.52%) 1 0 / 66 (0.00%) 0

<p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>weight increased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>weight decreased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>white blood cell count decreased</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 12 (0.00%)</p> <p>0</p> <p>0 / 12 (0.00%)</p> <p>0</p> <p>0 / 12 (0.00%)</p> <p>0</p> <p>0 / 12 (0.00%)</p> <p>0</p>	<p>3 / 461 (0.65%)</p> <p>3</p> <p>0 / 461 (0.00%)</p> <p>0</p> <p>0 / 461 (0.00%)</p> <p>0</p> <p>0 / 461 (0.00%)</p> <p>0</p>	<p>1 / 66 (1.52%)</p> <p>1</p> <p>0 / 66 (0.00%)</p> <p>0</p> <p>0 / 66 (0.00%)</p> <p>0</p> <p>0 / 66 (0.00%)</p> <p>0</p>
<p>Injury, poisoning and procedural complications</p> <p>ankle fracture</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 12 (0.00%)</p> <p>0</p>	<p>0 / 461 (0.00%)</p> <p>0</p>	<p>0 / 66 (0.00%)</p> <p>0</p>
<p>Cardiac disorders</p> <p>sinus bradycardia</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>ventricular extrasystoles</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p>	<p>0 / 12 (0.00%)</p> <p>0</p> <p>0 / 12 (0.00%)</p> <p>0</p>	<p>0 / 461 (0.00%)</p> <p>0</p> <p>0 / 461 (0.00%)</p> <p>0</p>	<p>0 / 66 (0.00%)</p> <p>0</p> <p>0 / 66 (0.00%)</p> <p>0</p>
<p>Nervous system disorders</p> <p>dizziness</p> <p>alternative dictionary used: MedDRA 28.0</p> <p>subjects affected / exposed</p> <p>occurrences (all)</p> <p>headache</p> <p>alternative dictionary used: MedDRA 28.0</p>	<p>0 / 12 (0.00%)</p> <p>0</p>	<p>3 / 461 (0.65%)</p> <p>3</p>	<p>1 / 66 (1.52%)</p> <p>3</p>

subjects affected / exposed occurrences (all) sleep deficit alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 12 (0.00%) 0 0 / 12 (0.00%) 0	8 / 461 (1.74%) 10 0 / 461 (0.00%) 0	1 / 66 (1.52%) 3 0 / 66 (0.00%) 0
Blood and lymphatic system disorders anaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 12 (0.00%) 0	7 / 461 (1.52%) 7	4 / 66 (6.06%) 4
Gastrointestinal disorders abdominal pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) anorectal polyp alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) colitis ulcerative alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) diarrhoea alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) haemorrhoids alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) large intestine polyp alternative dictionary used: MedDRA 28.0	0 / 12 (0.00%) 0 0 / 12 (0.00%) 0 0 / 12 (0.00%) 0 0 / 12 (0.00%) 0 1 / 12 (8.33%) 1	3 / 461 (0.65%) 3 1 / 461 (0.22%) 1 4 / 461 (0.87%) 4 2 / 461 (0.43%) 2 1 / 461 (0.22%) 1	0 / 66 (0.00%) 0 0 / 66 (0.00%) 0 1 / 66 (1.52%) 1 1 / 66 (1.52%) 1 1 / 66 (1.52%) 1

subjects affected / exposed occurrences (all)	1 / 12 (8.33%) 1	1 / 461 (0.22%) 1	1 / 66 (1.52%) 1
Skin and subcutaneous tissue disorders papule alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 12 (0.00%) 0	0 / 461 (0.00%) 0	0 / 66 (0.00%) 0
Endocrine disorders thyroiditis subacute alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 12 (0.00%) 0	0 / 461 (0.00%) 0	0 / 66 (0.00%) 0
Musculoskeletal and connective tissue disorders arthralgia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) back pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) osteoarthritis alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) neck pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 12 (0.00%) 0 0 / 12 (0.00%) 0 0 / 12 (0.00%) 0 0 / 12 (0.00%) 0	15 / 461 (3.25%) 15 0 / 461 (0.00%) 0 0 / 461 (0.00%) 0 0 / 461 (0.00%) 0	1 / 66 (1.52%) 1 1 / 66 (1.52%) 1 0 / 66 (0.00%) 0 0 / 66 (0.00%) 0
Infections and infestations covid-19 alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) bronchitis	1 / 12 (8.33%) 1	4 / 461 (0.87%) 4	4 / 66 (6.06%) 4

alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	0 / 66 (0.00%)
occurrences (all)	0	1	0
clostridium difficile infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	4 / 461 (0.87%)	3 / 66 (4.55%)
occurrences (all)	0	5	3
gastrointestinal infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences (all)	0	0	0
influenza			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	2 / 461 (0.43%)	0 / 66 (0.00%)
occurrences (all)	0	2	0
localised infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences (all)	0	0	0
nasopharyngitis			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	15 / 461 (3.25%)	0 / 66 (0.00%)
occurrences (all)	0	15	0
upper respiratory tract infection			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	3 / 461 (0.65%)	6 / 66 (9.09%)
occurrences (all)	0	3	6
Metabolism and nutrition disorders			
electrolyte imbalance			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 12 (8.33%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences (all)	1	0	0
hyperkalaemia			
alternative dictionary used: MedDRA 28.0			

subjects affected / exposed	0 / 12 (0.00%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences (all)	0	0	0
hypoalbuminaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	1 / 12 (8.33%)	0 / 461 (0.00%)	0 / 66 (0.00%)
occurrences (all)	2	0	0
hypokalaemia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 12 (0.00%)	1 / 461 (0.22%)	4 / 66 (6.06%)
occurrences (all)	0	1	4

Non-serious adverse events	Open Label Maintenance: Delayed Responders - 200 mg Miri SC	Maintenance Period: Miri IR - PBO SC	Maintenance Period: Miri IR - 200 mg Miri SC - ME2 Cohort
Total subjects affected by non-serious adverse events			
subjects affected / exposed	91 / 271 (33.58%)	82 / 192 (42.71%)	32 / 52 (61.54%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
uterine leiomyoma			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed ^[3]	0 / 111 (0.00%)	0 / 78 (0.00%)	0 / 23 (0.00%)
occurrences (all)	0	0	0
Vascular disorders			
hypotension			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	0 / 271 (0.00%)	2 / 192 (1.04%)	0 / 52 (0.00%)
occurrences (all)	0	2	0
General disorders and administration site conditions			
oedema peripheral			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	2 / 271 (0.74%)	1 / 192 (0.52%)	0 / 52 (0.00%)
occurrences (all)	3	1	0
pyrexia			
alternative dictionary used: MedDRA 28.0			
subjects affected / exposed	6 / 271 (2.21%)	5 / 192 (2.60%)	3 / 52 (5.77%)
occurrences (all)	7	7	3
Respiratory, thoracic and mediastinal disorders			

pulmonary mass alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	1 / 52 (1.92%) 1
Investigations			
alanine aminotransferase increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	1 / 192 (0.52%) 1	1 / 52 (1.92%) 1
aspartate aminotransferase increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1	0 / 192 (0.00%) 0	1 / 52 (1.92%) 1
blood alkaline phosphatase increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	2 / 271 (0.74%) 2	3 / 192 (1.56%) 3	1 / 52 (1.92%) 1
blood creatine phosphokinase increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	3 / 271 (1.11%) 3	5 / 192 (2.60%) 6	1 / 52 (1.92%) 1
weight increased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
weight decreased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
white blood cell count decreased alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1	1 / 192 (0.52%) 2	3 / 52 (5.77%) 3
Injury, poisoning and procedural complications			

ankle fracture alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
Cardiac disorders sinus bradycardia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) ventricular extrasystoles alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0 0 / 271 (0.00%) 0	0 / 192 (0.00%) 0 0 / 192 (0.00%) 0	1 / 52 (1.92%) 1 0 / 52 (0.00%) 0
Nervous system disorders dizziness alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) headache alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all) sleep deficit alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1 13 / 271 (4.80%) 21 0 / 271 (0.00%) 0	1 / 192 (0.52%) 1 2 / 192 (1.04%) 5 0 / 192 (0.00%) 0	1 / 52 (1.92%) 1 2 / 52 (3.85%) 4 0 / 52 (0.00%) 0
Blood and lymphatic system disorders anaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	9 / 271 (3.32%) 9	9 / 192 (4.69%) 12	2 / 52 (3.85%) 2
Gastrointestinal disorders abdominal pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	3 / 271 (1.11%) 5	4 / 192 (2.08%) 4	4 / 52 (7.69%) 4

anorectal polyp alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	2 / 192 (1.04%) 2	0 / 52 (0.00%) 0
colitis ulcerative alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	10 / 271 (3.69%) 11	35 / 192 (18.23%) 35	4 / 52 (7.69%) 5
diarrhoea alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	8 / 271 (2.95%) 9	1 / 192 (0.52%) 1	1 / 52 (1.92%) 1
haemorrhoids alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	6 / 271 (2.21%) 6	3 / 192 (1.56%) 3	1 / 52 (1.92%) 1
large intestine polyp alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	2 / 192 (1.04%) 2	0 / 52 (0.00%) 0
Skin and subcutaneous tissue disorders papule alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
Endocrine disorders thyroiditis subacute alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
Musculoskeletal and connective tissue disorders arthralgia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	19 / 271 (7.01%) 24	8 / 192 (4.17%) 8	1 / 52 (1.92%) 1

back pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	3 / 271 (1.11%) 3	4 / 192 (2.08%) 4	0 / 52 (0.00%) 0
osteoarthritis alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
neck pain alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1	1 / 192 (0.52%) 1	0 / 52 (0.00%) 0
Infections and infestations			
covid-19 alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	6 / 271 (2.21%) 6	4 / 192 (2.08%) 4	6 / 52 (11.54%) 6
bronchitis alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1	0 / 192 (0.00%) 0	1 / 52 (1.92%) 1
clostridium difficile infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	3 / 271 (1.11%) 4	3 / 192 (1.56%) 4	0 / 52 (0.00%) 0
gastrointestinal infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
influenza alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	4 / 52 (7.69%) 4
localised infection alternative dictionary used: MedDRA 28.0			

subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
nasopharyngitis alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	14 / 271 (5.17%) 15	11 / 192 (5.73%) 11	3 / 52 (5.77%) 3
upper respiratory tract infection alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	9 / 271 (3.32%) 9	5 / 192 (2.60%) 6	8 / 52 (15.38%) 12
Metabolism and nutrition disorders electrolyte imbalance alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0
hyperkalaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	1 / 271 (0.37%) 1	2 / 192 (1.04%) 2	0 / 52 (0.00%) 0
hypoalbuminaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	1 / 52 (1.92%) 1
hypokalaemia alternative dictionary used: MedDRA 28.0 subjects affected / exposed occurrences (all)	0 / 271 (0.00%) 0	0 / 192 (0.00%) 0	0 / 52 (0.00%) 0

Notes:

[3] - The number of subjects exposed to this adverse event is less than the total number of subjects exposed for the reporting group. These numbers are expected to be equal.

Justification: Gender-specific events occurring exclusively in female participants have had the number of participants at risk adjusted accordingly.

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
12 September 2019	<ul style="list-style-type: none">• Replaced Primary objective, major secondary objectives, added Other secondary objectives• Added: 50% biological 50% conventional failure split• Follicle stimulating hormone (FSH) testing optional in women to confirm nonchild-bearing potential• Clarified infusion-reaction related samples• Clarified collection of Endoscopy• Clarified conditions when patients perform health outcome assessments• Clarified when informed consent form (ICF) be obtained• Clarified schedule of activities• Minimized study procedures for participants• Visit window changed between AMAN and AMBG• Clarification of Inclusion Criteria• Added assessments if drug hypersensitivity event observed• Changed the term Study treatment to study drug as study drug is blinded• Clarified term "mucosal healing"• Primary endpoint terminology changed from "durable clinical remission" to "clinical remission"; "remission changed to response" with mirikizumab induction treatment for clarification of treatment• Text deleted for medical monitor consult for endoscopy findings• Added when early termination visit (ETV) endoscopy should be discussed with the medical monitor.• Added Centers for Disease Control and Prevention guidance (CDC) was for the United States (US) and the World Health Organization (WHO) was for countries outside of the US• Deleted clinical study report (CSR) coordinating investigator "with the most enrolled participant's" will be selected by the study team• Clarified use of permitted and prohibited Medications

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported