

**Clinical trial results:****A Multicenter, Randomized, Double-Blind, Active (Oseltamivir)-Controlled Study to Assess the Safety, Pharmacokinetics, and Efficacy of Baloxavir Marboxil in Otherwise Healthy Pediatric Patients 1 to <12 Years of Age With Influenza-Like Symptoms****Summary**

EudraCT number	2018-002169-21
Trial protocol	ES PL
Global end of trial date	03 April 2019

**Results information**

Result version number	v2 (current)
This version publication date	12 March 2020
First version publication date	12 October 2019
Version creation reason	

**Trial information****Trial identification**

Sponsor protocol code	CP40563
-----------------------	---------

**Additional study identifiers**

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT03629184
WHO universal trial number (UTN)	-

Notes:

**Sponsors**

Sponsor organisation name	F. Hoffmann-La Roche AG
Sponsor organisation address	Grenzacherstrasse 124, Basel, Switzerland, CH-4070
Public contact	F. Hoffmann-La Roche AG, F. Hoffmann-La Roche AG, 41 616878333, global.trial_information@roche.com
Scientific contact	F. Hoffmann-La Roche AG, F. Hoffmann-La Roche AG, 41 616878333, global.trial_information@roche.com

Notes:

**Paediatric regulatory details**

Is trial part of an agreed paediatric investigation plan (PIP)	Yes
EMA paediatric investigation plan number(s)	EMA-002440-PIP01-18
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	Yes

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	09 July 2019
Is this the analysis of the primary completion data?	Yes
Primary completion date	03 April 2019
Global end of trial reached?	Yes
Global end of trial date	03 April 2019
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

To compare the safety of a single dose of baloxavir marboxil with 5 days of oseltamivir administered twice daily.

Protection of trial subjects:

All study subjects were required to read and sign an Informed Consent Form.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	20 November 2018
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Costa Rica: 2
Country: Number of subjects enrolled	Spain: 3
Country: Number of subjects enrolled	Mexico: 1
Country: Number of subjects enrolled	Poland: 5
Country: Number of subjects enrolled	Russian Federation: 1
Country: Number of subjects enrolled	United States: 161
Worldwide total number of subjects	173
EEA total number of subjects	8

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	16
Children (2-11 years)	157
Adolescents (12-17 years)	0
Adults (18-64 years)	0

From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details: -

### Pre-assignment

Screening details:

173 participants were enrolled and dosed in this study.

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

### Arms

Are arms mutually exclusive?	Yes
------------------------------	-----

<b>Arm title</b>	Baloxavir Marboxil
------------------	--------------------

Arm description:

Participants will receive a single oral dose of baloxavir marboxil on Day 1 (based on body weight). Oseltamivir matching placebo will also be administered orally twice daily (BID) for 5 days.

Arm type	Experimental
Investigational medicinal product name	Baloxavir Marboxil
Investigational medicinal product code	
Other name	Xofluza
Pharmaceutical forms	Granules for oral suspension
Routes of administration	Oral use

Dosage and administration details:

Participants received a single oral dose of baloxavir marboxil on Day 1 (based on body weight)

Investigational medicinal product name	Oseltamivir matching placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Powder for oral suspension
Routes of administration	Oral use

Dosage and administration details:

Oseltamivir matching placebo was administered orally twice daily (BID) for 5 days

<b>Arm title</b>	Oseltamivir
------------------	-------------

Arm description:

Participants will receive oseltamivir orally BID for 5 days (based on body weight). Baloxavir marboxil matching placebo will also be administered orally on Day 1

Arm type	Active comparator
Investigational medicinal product name	Oseltamivir
Investigational medicinal product code	
Other name	Tamiflu
Pharmaceutical forms	Powder for oral suspension
Routes of administration	Oral use

Dosage and administration details:

Participants received oseltamivir orally BID for 5 days (based on body weight)

Investigational medicinal product name	Baloxavir marboxil matching placebo
Investigational medicinal product code	
Other name	

Pharmaceutical forms	Granules for oral suspension
Routes of administration	Oral use

Dosage and administration details:

Baloxavir marboxil matching placebo was administered orally on Day 1

<b>Number of subjects in period 1</b>	Baloxavir Marboxil	Osetamivir
Started	115	58
Completed	112	57
Not completed	3	1
Consent withdrawn by subject	2	1
Physician decision	1	-

## Baseline characteristics

### Reporting groups

Reporting group title	Baloxavir Marboxil
Reporting group description:	
Participants will receive a single oral dose of baloxavir marboxil on Day 1 (based on body weight). Oseltamivir matching placebo will also be administered orally twice daily (BID) for 5 days.	
Reporting group title	Oseltamivir
Reporting group description:	
Participants will receive oseltamivir orally BID for 5 days (based on body weight). Baloxavir marboxil matching placebo will also be administered orally on Day 1	

Reporting group values	Baloxavir Marboxil	Oseltamivir	Total
Number of subjects	115	58	173
Age categorical			
Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	0	0	0
Infants and toddlers (28 days-23 months)	9	7	16
Children (2-11 years)	106	51	157
Adolescents (12-17 years)	0	0	0
Adults (18-64 years)	0	0	0
From 65-84 years	0	0	0
85 years and over	0	0	0
Age Continuous			
Units: years			
arithmetic mean	6.10	6.02	
standard deviation	± 2.90	± 3.20	-
Sex: Female, Male			
Units: Subjects			
Female	60	32	92
Male	55	26	81
Race/Ethnicity, Customized			
Units: Subjects			
American Indian or Alaska Native	1	0	1
Asian	1	0	1
Black or African American	6	5	11
Native Hawaiian or other Pacific Islander	0	1	1
White	98	51	149
Multiple	4	0	4
Unknown	5	1	6
Race/Ethnicity, Customized			
Units: Subjects			
Hispanic or Latino	52	27	79
Not Hispanic or Latino	63	31	94



## End points

### End points reporting groups

Reporting group title	Baloxavir Marboxil
Reporting group description: Participants will receive a single oral dose of baloxavir marboxil on Day 1 (based on body weight). Oseltamivir matching placebo will also be administered orally twice daily (BID) for 5 days.	
Reporting group title	Oseltamivir
Reporting group description: Participants will receive oseltamivir orally BID for 5 days (based on body weight). Baloxavir marboxil matching placebo will also be administered orally on Day 1	

### Primary: Percentage of Participants with Adverse Events (AEs) and Serious Adverse Events (SAEs)

End point title	Percentage of Participants with Adverse Events (AEs) and Serious Adverse Events (SAEs) <sup>[1]</sup>
End point description: An adverse event (AE) is any untoward medical occurrence in a participant or clinical investigation participant administered a pharmaceutical product and that does not necessarily have a causal relationship with this treatment. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease temporally associated with the use of a medicinal (investigational) product, whether or not considered related to the medicinal (investigational) product. A serious adverse event (SAE) is any significant hazard, contraindication, side effect that is fatal or life-threatening, requires hospitalization or prolongation of an existing hospitalization, results in persistent or significant disability/ incapacity, is a congenital anomaly/ birth defect, is medically significant or requires intervention to prevent one or other of the outcomes listed above.	
End point type	Primary
End point timeframe: Up to Day 29	

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Descriptive stats only. No statistical analyses were pre-defined.

End point values	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	115	58		
Units: percentage of participants				
number (not applicable)				
Adverse Events (AEs)	46.1	53.4		
Serious Adverse Events (SAEs)	0	0		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Plasma Concentrations of Baloxavir Marboxil - Sparse PK Population

End point title	Plasma Concentrations of Baloxavir Marboxil - Sparse PK Population <sup>[2]</sup>
-----------------	---

End point description:

Results provided by body-weight groups for participants in the Baloxavir Marboxil arm. Values below lower limit of quantification (0.5 ng/mL) are set to zero. Here 99999 represents results data which was not estimable due to low number of events.

End point type Secondary

End point timeframe:

Days 1 (Post-Dose), 2, 4, 6 and 10

Notes:

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	97			
Units: ng/mL				
arithmetic mean (standard deviation)				
5 - <10 kg (Day 1) (n=1)	0.000 (± 99999)			
5 - <10 kg (Day 2) (n=1)	0.000 (± 99999)			
5 - <10 kg (Day 4) (n=0)	99999 (± 99999)			
5 - <10 kg (Day 6) (n=1)	0.000 (± 99999)			
5 - <10 kg (Day 10) (n=0)	99999 (± 99999)			
10 - <15 kg (Day 1) (n=16)	0.073 (± 0.2001)			
10 - <15 kg (Day 2) (n=7)	0.000 (± 0.0000)			
10 - <15 kg (Day 4) (n=9)	0.000 (± 0.0000)			
10 - <15 kg (Day 6) (n=13)	0.000 (± 0.0000)			
10 - <15 kg (Day 10)(n=3)	0.000 (± 0.0000)			
15 - <20 kg (Day 1) (n=22)	0.090 (± 0.2386)			
15 - <20 kg (Day 2) (n=13)	0.000 (± 0.0000)			
15 - <20 kg (Day 4) (n=7)	0.000 (± 0.0000)			
15 - <20 kg (Day 6) (n=17)	0.000 (± 0.0000)			
15 - <20 kg (Day 10) (n=3)	0.000 (± 0.0000)			
>=20 kg (Day 1) (n=58)	0.048 (± 0.1936)			
>=20 kg (Day 2) (n=34)	0.000 (± 0.0000)			
>=20 kg (Day 4) (n=25)	0.000 (± 0.0000)			
>=20 kg (Day 6) (n=56)	0.000 (± 0.0000)			
>=20 kg (Day 10) (n=6)	0.000 (± 0.0000)			

## Statistical analyses

No statistical analyses for this end point

### Secondary: Plasma Concentrations of S-033447 - Sparse PK Population

End point title	Plasma Concentrations of S-033447 - Sparse PK Population <sup>[3]</sup>
-----------------	---

End point description:

Results provided by body-weight groups for participants in the Baloxavir Marboxil arm. Here 99999 represents results data which was not estimable due to low number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Days 1 (Post-Dose), 2, 4, 6 and 10

Notes:

[3] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	97			
Units: ng/mL				
arithmetic mean (standard deviation)				
5 - <10 kg (Day 1) (n=1)	45.700 (± 99999)			
5 - <10 kg (Day 2) (n=1)	45.800 (± 99999)			
5 - <10 kg (Day 4) (n=0)	99999 (± 99999)			
5 - <10 kg (Day 6) (n=1)	3.110 (± 99999)			
5 - <10 kg (Day 10) (n=0)	99999 (± 99999)			
10 - <15 kg (Day 1) (n=16)	49.084 (± 53.6689)			
10 - <15 kg (Day 2) (n=7)	42.900 (± 16.5227)			
10 - <15 kg (Day 4) (n=9)	9.233 (± 5.3879)			
10 - <15 kg (Day 6) (n=13)	2.965 (± 1.6480)			
10 - <15 kg (Day 10) (n=3)	0.367 (± 0.6351)			
15 - <20 kg (Day 1) (n=22)	64.160 (± 73.6320)			
15 - <20 kg (Day 2) (n=13)	67.729 (± 46.7346)			
15 - <20 kg (Day 4) (n=7)	15.840 (± 10.8285)			
15 - <20 kg (Day 6) (n=17)	4.829 (± 3.6562)			

15 - <20 kg (Day 10) (n=3)	1.110 (± 1.9226)			
>=20 kg (Day 1) (n=58)	29.899 (± 26.1558)			
>=20 kg (Day 2) (n=34)	56.287 (± 40.4073)			
>=20 kg (Day 4) (n=25)	18.674 (± 11.2179)			
>=20 kg (Day 6) (n=56)	7.397 (± 5.0530)			
>=20 kg (Day 10) (n=6)	3.953 (± 2.2536)			

## Statistical analyses

No statistical analyses for this end point

### Secondary: Plasma Concentrations of Baloxavir Marboxil - Extensive PK Population

End point title	Plasma Concentrations of Baloxavir Marboxil - Extensive PK Population <sup>[4]</sup>
-----------------	--

End point description:

Results provided by body-weight groups for participants in the Baloxavir Marboxil arm. Values below lower limit of quantification (0.5 ng/mL) are set to zero. Here 99999 represents results data which was not estimable due to low number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Days 1 (Post-Dose), 2, 4, 6 and 10

Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	19			
Units: ng/mL				
arithmetic mean (standard deviation)				
Day 1, 0.5 - 2 hrs (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 1, 4 hrs (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 1, 6 hrs (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 2 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 4 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 6 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 10 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 1, 0.5 - 2 hrs (10 - <15 kg) (n=4)	0.000 (± 0.0000)			
Day 1, 4 hrs (10 - <15 kg) (n=2)	0.000 (± 0.0000)			

Day 1, 6 hrs (10 - <15 kg) (n=1)	0.000 (± 99999)			
Day 2 (10 - <15 kg) (n=3)	0.000 (± 0.0000)			
Day 4 (10 - <15 kg) (n=1)	0.000 (± 99999)			
Day 6 (10 - <15 kg) (n=3)	0.000 (± 0.0000)			
Day 10 (10 - <15 kg) (n=1)	0.000 (± 99999)			
Day 1, 0.5 - 2 hrs (15 - <20 kg) (n=4)	0.000 (± 0.0000)			
Day 1, 4 hrs (15 - <20 kg) (n=1)	0.000 (± 99999)			
Day 1, 6 hrs (15 - <20 kg) (n=1)	0.000 (± 99999)			
Day 2 (15 - <20 kg) (n=2)	0.000 (± 0.0000)			
Day 4 (15 - <20 kg) (n=1)	0.000 (± 99999)			
Day 6 (15 - <20 kg) (n=3)	0.000 (± 0.0000)			
Day 10 (15 - <20 kg) (n=0)	99999 (± 99999)			
Day 1, 0.5 - 2 hrs (>=20 kg) (n=10)	0.051 (± 0.1600)			
Day 1, 4 hrs (>=20 kg) (n=9)	0.062 (± 0.1863)			
Day 1, 6 hrs (>=20 kg) (n=6)	0.000 (± 0.0000)			
Day 2 (>=20 kg) (n=5)	0.000 (± 0.0000)			
Day 4 (>=20 kg) (n=4)	0.000 (± 0.0000)			
Day 6 (>=20 kg) (n=9)	0.000 (± 0.0000)			
Day 10 (>=20 kg) (n=0)	99999 (± 99999)			

## Statistical analyses

No statistical analyses for this end point

## Secondary: Plasma Concentrations of S-033447 - Extensive PK Population

End point title	Plasma Concentrations of S-033447 - Extensive PK
-----------------	--

End point description:

Results provided by body-weight groups for participants in the Baloxavir Marboxil arm. Values below lower limit of quantification (0.5 ng/mL) are set to zero. Here 99999 represents results data which was not estimable due to low number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Days 1 (Post-Dose), 2, 4, 6 and 10

Notes:

[5] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

<b>End point values</b>	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	19			
Units: ng/mL				
arithmetic mean (standard deviation)				
Day 1, 0.5 - 2 hrs (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 1, 4 hrs (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 1, 6 hrs (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 2 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 4 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 6 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 10 (5 - <10 kg) (n=0)	99999 (± 99999)			
Day 1, 0.5 - 2 hrs (10 - <15 kg) (n=4)	10.768 (± 14.7987)			
Day 1, 4 hrs (10 - <15 kg) (n=2)	49.500 (± 13.0108)			
Day 1, 6 hrs (10 - <15 kg) (n=1)	41.000 (± 99999)			
Day 2 (10 - <15 kg) (n=3)	28.933 (± 14.7514)			
Day 4 (10 - <15 kg) (n=1)	2.230 (± 99999)			
Day 6 (10 - <15 kg) (n=3)	3.131 (± 2.2828)			
Day 10 (10 - <15 kg) (n=1)	0.000 (± 99999)			
Day 1, 0.5 - 2 hrs (15 - <20 kg) (n=4)	93.883 (± 152.5431)			
Day 1, 4 hrs (15 - <20 kg) (n=1)	72.900 (± 99999)			
Day 1, 6 hrs (15 - <20 kg) (n=1)	80.300 (± 99999)			
Day 2 (15 - <20 kg) (n=2)	42.640 (± 47.6024)			
Day 4 (15 - <20 kg) (n=1)	12.200 (± 99999)			
Day 6 (15 - <20 kg) (n=3)	2.663 (± 1.5387)			
Day 10 (15 - <20 kg) (n=0)	99999 (± 99999)			
Day 1, 0.5 - 2 hrs (>=20 kg) (n=10)	19.923 (± 27.0980)			
Day 1, 4 hrs (>=20 kg) (n=9)	69.198 (± 55.7220)			
Day 1, 6 hrs (>=20 kg) (n=6)	65.527 (± 43.0799)			
Day 2 (>=20 kg) (n=5)	57.980 (± 37.8922)			
Day 4 (>=20 kg) (n=4)	21.775 (± 3.7968)			
Day 6 (>=20 kg) (n=9)	6.240 (± 3.3702)			

Day 10 ( $\geq 20$ kg) (n=0)	99999 ( $\pm$ 99999)			
------------------------------	----------------------	--	--	--

## Statistical analyses

No statistical analyses for this end point

### Secondary: Time to Alleviation of Influenza Signs and Symptoms

End point title	Time to Alleviation of Influenza Signs and Symptoms
-----------------	---

End point description:

Time to alleviation of influenza signs and symptoms is defined as the length of time taken from the start of treatment to the point at which all of the following criteria are met and remain so for at least 21.5 hours: - A score of 0 (no problem) or 1 (minor problem) for cough and nasal symptoms (items 14 and 15 of the Canadian Acute Respiratory Illness and Flu Scale [CARIFS]) - A "yes" response to the following question on the CARIFS: "Since the last assessment has the subject been able to return to day care/school, or resume his or her normal daily activity in the same way as performed prior to developing the flu?" - First return to afebrile state (tympanic temperature  $\leq 37.2$  degree Celsius [ $^{\circ}\text{C}$ ])

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 15

End point values	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	80	43		
Units: hours				
median (confidence interval 95%)	138.1 (116.6 to 163.2)	150.0 (115.0 to 165.7)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Duration of Fever

End point title	Duration of Fever
-----------------	-------------------

End point description:

Length of time taken by participants to return to afebrile state [tympanic temperature  $\leq 37.2^{\circ}\text{C}$ ] and remaining so for at least 21.5 hours.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 15

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	80	43		
Units: hours				
median (confidence interval 95%)	41.2 (24.5 to 45.7)	46.8 (30.0 to 53.5)		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Duration of Symptoms

End point title	Duration of Symptoms
-----------------	----------------------

End point description:

The clinical efficacy of baloxavir marboxil is evaluated by duration of symptoms i.e., alleviation of all symptoms as defined by a score of 0 [no problem] or 1 [minor problem] and remaining so for at least 21.5 hours, for all 18 symptoms specified in the CARIFS questionnaire).

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 15

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	80	43		
Units: hours				
median (confidence interval 95%)	66.4 (43.7 to 76.4)	67.9 (45.8 to 88.7)		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Time to Return to Normal Health and Activity

End point title	Time to Return to Normal Health and Activity
-----------------	--

End point description:

Time to Return to Normal health and activity' is identified by a 'Yes' response to the following question on the CARIFS: "Since the last assessment has the patient been able to return to day care/school, or resume his or her normal daily activity in the same way as performed prior to developing the flu?"

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 15

<b>End point values</b>	Baloxavir Marboxil	Osetamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	80	43		
Units: hours				
median (confidence interval 95%)	116.5 (94.9 to 138.0)	111.6 (80.8 to 138.3)		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Frequency of Influenza-Related Complications

End point title	Frequency of Influenza-Related Complications
End point description:	Influenza related complications include death, hospitalization, radiologically confirmed pneumonia, bronchitis, sinusitis, otitis media, encephalitis/encephalopathy, febrile seizures, myositis.
End point type	Secondary
End point timeframe:	Up to Day 29

<b>End point values</b>	Baloxavir Marboxil	Osetamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	81	43		
Units: count of events				
Total	6	4		
Death	0	0		
Hospitalization	0	0		
Sinusitis	1	0		
Otitis Media	3	3		
Pneumonia	1	0		
Bronchitis	1	0		
Encephalitis/Encephalopathy	0	0		
Febrile Seizures	0	1		
Myositis	0	0		

### Statistical analyses

No statistical analyses for this end point

**Secondary: Percentage of Participants with Influenza-Related Complications**

End point title	Percentage of Participants with Influenza-Related Complications
-----------------	---

End point description:

Influenza related complications include death, hospitalization, radiologically confirmed pneumonia, bronchitis, sinusitis, otitis media, encephalitis/encephalopathy, febrile seizures, myositis.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 29

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	81	43		
Units: percentage of participants				
number (not applicable)				
Total	7.4	7.0		
Death	0	0		
Hospitalization	0	0		
Sinusitis	1.2	0		
Otitis Media	3.7	4.7		
Pneumonia	1.2	0		
Bronchitis	1.2	0		
Encephalitis/Encephalopathy	0	0		
Febrile Seizures	0	2.3		
Myositis	0	0		

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Percentage of Participants Requiring Antibiotics**

End point title	Percentage of Participants Requiring Antibiotics
-----------------	--

End point description:

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 29

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	81	43		
Units: percentage of participants				
number (not applicable)				
Total	4.9	4.7		
Bronchitis	0	0		
Otitis Media	2.5	4.7		
Pneumonia	1.2	0		
Sinusitis	1.2	0		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Time to Cessation of Viral Shedding by Virus Titer

End point title	Time to Cessation of Viral Shedding by Virus Titer
End point description:	Time to cessation of viral shedding by virus titer is defined as the time, in hours, between the initiation of any study treatment and first time when the influenza virus titer is below the limit of detection.
End point type	Secondary
End point timeframe:	Day 1 - Day 29

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	67	37		
Units: hours				
median (confidence interval 95%)				
Virus Titer	24.2 (23.5 to 24.6)	75.8 (68.9 to 97.8)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Time to Cessation of Viral Shedding by RT-PCR

End point title	Time to Cessation of Viral Shedding by RT-PCR
End point description:	Time to cessation of viral shedding by RT-PCR, in hours, is defined as the time between the initiation of any study treatment and first time when the virus RNA by RT-PCR is below the limit of detection.
End point type	Secondary

End point timeframe:

Day 1 - Day 29

<b>End point values</b>	Baloxavir Marboxil	Osetamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	76	39		
Units: hours				
median (confidence interval 95%)	242.5 (235.8 to 262.8)	238.9 (214.0 to 286.7)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in Influenza Virus Titer at Day 2, 4, 6, 10, 15, 29

End point title	Change from Baseline in Influenza Virus Titer at Day 2, 4, 6, 10, 15, 29
-----------------	--

End point description:

Influenza virus titer (log<sub>10</sub>TCID<sub>50</sub>/ML) is the quantity of influenza virus in a given volume within the samples obtained from nasal swabs. If influenza virus titer was less than the lower limit of quantification, the virus titer was imputed as 0.749 (log<sub>10</sub>TCID<sub>50</sub>/mL). A lower value indicates lower viral titer.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, Day 2, 3 (optional), 4, 6, 10, 15 (optional), 29

<b>End point values</b>	Baloxavir Marboxil	Osetamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	67	38		
Units: log <sub>10</sub> TCID <sub>50</sub> /ML				
arithmetic mean (standard deviation)				
Baseline (Baloxvir Marboxil n=67/Osetamivir n=38)	4.43 (± 1.36)	4.27 (± 1.48)		
Day 2 (Baloxvir Marboxil n=64/Osetamivir n=37)	-3.59 (± 1.34)	-1.79 (± 1.54)		
Day 3 (Baloxvir Marboxil n=3 / Osetamivir n=2)	-2.83 (± 0.58)	-2.63 (± 0.88)		
Day 4 (Baloxvir Marboxil n=61 / Osetamivir n=31)	-3.53 (± 1.38)	-3.27 (± 1.54)		
Day 6 (Baloxvir Marboxil n=63 / Osetamivir n=35)	-3.55 (± 1.32)	-3.52 (± 1.50)		
Day 10 (Baloxvir Marboxil n=4 / Osetamivir n=4)	-3.66 (± 1.40)	-3.50 (± 1.42)		
Day 15 (Baloxvir Marboxil n=4 / Osetamivir n=4)	-3.75 (± 0.54)	-3.63 (± 1.45)		

Day 29 (Baloxvir Marboxil n=4 / Osetamivir n=4)	-3.50 (± 1.43)	-3.75 (± 1.19)		
---	----------------	----------------	--	--

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in the Amount of Virus RNA (RT-PCR) at Day 2, 4, 6, 10, 15, 29

End point title	Change from Baseline in the Amount of Virus RNA (RT-PCR) at Day 2, 4, 6, 10, 15, 29
End point description: If the amount of virus RNA was less than the lower limit of quantification, the amount of virus RNA was imputed as 2.18 for flu A and 2.93 for flu B (log10 virus particles/mL). Here 99999 represents results data which was not estimable due to low number of events.	
End point type	Secondary
End point timeframe: Baseline, Day 2, 3 (optional), 4, 6, 10, 15 (optional), 29	

End point values	Baloxavir Marboxil	Osetamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	76	40		
Units: log10 virus particles/mL				
arithmetic mean (standard deviation)				
Baseline (Baloxvir Marboxil n=76/Osetamivir n=40)	6.46 (± 1.50)	6.86 (± 1.02)		
Day 2 (Baloxvir Marboxil n=70 / Osetamivir n=39)	-1.74 (± 1.13)	-1.12 (± 1.12)		
Day 3 (Baloxvir Marboxil n=4 / Osetamivir n=2)	-1.78 (± 1.50)	-2.21 (± 0.94)		
Day 4 (Baloxvir Marboxil n=61 / Osetamivir n=30)	-2.40 (± 1.50)	-2.47 (± 1.35)		
Day 6 (Baloxvir Marboxil n=60 / Osetamivir n=30)	-2.73 (± 1.78)	-3.32 (± 1.27)		
Day 10 (Baloxvir Marboxil n=35 / Osetamivir n=15)	-3.55 (± 1.62)	-3.81 (± 1.19)		
Day 15 (Baloxvir Marboxil n=2 / Osetamivir n=1)	-1.24 (± 3.06)	-4.44 (± 99999)		
Day 29 (Baloxvir Marboxil n=1 / Osetamivir n=0)	2.18 (± 99999)	99999 (± 99999)		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of Participants with Positive Influenza Virus Titer at Day 2,

**4, 6, 10**

End point title	Percentage of Participants with Positive Influenza Virus Titer at Day 2, 4, 6, 10
End point description: The percentage is based on the number of patients with virus titer data at each time point	
End point type	Secondary
End point timeframe: Baseline, Day 2, 3 (optional), 4, 6, 10	

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	67	38		
Units: percentage of participants				
number (not applicable)				
Baseline	100	100		
Day 2	15.6	75.7		
Day 3 (optional)	33.3	50.0		
Day 4	26.2	29.0		
Day 6	12.7	5.7		
Day 10	1.6	0		

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Percentage of Participants Positive by RT-PCR at Day 2, 4, 6, 10, 15, 29**

End point title	Percentage of Participants Positive by RT-PCR at Day 2, 4, 6, 10, 15, 29
End point description: The percentage is based on the number of patients with RT-PCR data at each time point	
End point type	Secondary
End point timeframe: Day 2, 3 (optional), 4, 6, 10, 15 (optional), 29	

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	76	40		
Units: percentage of participants				
number (not applicable)				
Baseline	100	100		
Day 2	95.9	100		
Day 3 (optional)	100	100		

Day 4	89.9	97.0		
Day 6	83.1	75.7		
Day 10	46.5	44.1		
Day 15 (optional)	40.0	25.0		
Day 29	25.0	0		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Area Under the Curve in Virus Titer

End point title	Area Under the Curve in Virus Titer
-----------------	-------------------------------------

End point description:

Area under the curve (AUC) in virus titer was calculated using the trapezoidal method.

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 - Day 29

End point values	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	67	37		
Units: log[TCID/mL]*hours				
arithmetic mean (standard deviation)	-863.81 ( $\pm$ 543.37)	-849.29 ( $\pm$ 684.43)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Area Under the Curve in the Amount of Virus RNA (RT-PCR)

End point title	Area Under the Curve in the Amount of Virus RNA (RT-PCR)
-----------------	--

End point description:

AUC in virus RNA (RT-PCR) is defined as AUC of change from baseline in the amount of virus RNA (RT-PCR) from Day 1 to Day 10. AUC is calculated using the trapezoidal method similar to AUC in virus titer.

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 - Day 10

<b>End point values</b>	Baloxavir Marboxil	Oseltamivir		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	75	39		
Units: log VPs/mL*hours				
arithmetic mean (standard deviation)	-381.53 ( $\pm$ 338.53)	-353.31 ( $\pm$ 304.01)		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Area Under the Concentration to Time Curve from Time 0 to Infinity (AUC0-inf) of baloxavir marboxil

End point title	Area Under the Concentration to Time Curve from Time 0 to Infinity (AUC0-inf) of baloxavir marboxil <sup>[6]</sup>
-----------------	--

End point description:

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 10

Notes:

[6] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

<b>End point values</b>	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	0 <sup>[7]</sup>			
Units: ng.hr/ml				
arithmetic mean (standard deviation)	()			

Notes:

[7] - Not estimable as a large number of concentrations were below the lower limit of quantification

### Statistical analyses

No statistical analyses for this end point

### Secondary: Area Under the Concentration to Time Curve from Time 0 to Infinity (AUC0-inf) of S-033447

End point title	Area Under the Concentration to Time Curve from Time 0 to Infinity (AUC0-inf) of S-033447 <sup>[8]</sup>
-----------------	--

End point description:

Dose groups correspond to body-weight groups. 2mg/kg dose was used for subjects <20 kgs and 40 mg dose was used for subjects >20 kgs. Here 99999 represents results data which was not estimable due to low number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 10

Notes:

[8] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.  
Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

<b>End point values</b>	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	95			
Units: ng.hr/mL				
arithmetic mean (standard deviation)				
Non-Asian - 2 mg/kg (n=36)	4050 (± 2080)			
Asian - 40 mg (n=1)	6600 (± 99999)			
Non-Asian - 40 mg (n=58)	4390 (± 2080)			

### Statistical analyses

No statistical analyses for this end point

### Secondary: Maximum Plasma Concentration (Cmax) of baloxavir marboxil

End point title | Maximum Plasma Concentration (Cmax) of baloxavir

End point description:

End point type | Secondary

End point timeframe:

Up to Day 10

Notes:

[9] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.  
Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

<b>End point values</b>	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	0 <sup>[10]</sup>			
Units: ng/mL				
arithmetic mean (standard deviation)	( )			

Notes:

[10] - Not estimable as a large number of concentrations were below the lower limit of quantification

### Statistical analyses

No statistical analyses for this end point

### Secondary: Maximum Plasma Concentration (Cmax) of S-033447

End point title | Maximum Plasma Concentration (Cmax) of S-033447<sup>[11]</sup>

End point description:

Dose groups correspond to body-weight groups. 2mg/kg dose was used for subjects <20 kgs and 40 mg dose was used for subjects >20 kgs. Here 99999 represents results data which was not estimable due to

low number of events.

End point type	Secondary
End point timeframe:	
Up to Day 10	

Notes:

[11] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	95			
Units: ng/mL				
arithmetic mean (standard deviation)				
Non-Asian - 2 mg/kg (n=36)	109 (± 55.3)			
Asian - 40 mg (n=1)	110 (± 99999)			
Non-Asian - 40 mg (n=58)	83.2 (± 36.5)			

## Statistical analyses

No statistical analyses for this end point

## Secondary: Time to Maximum Plasma Concentration (Tmax) of baloxavir marboxil

End point title	Time to Maximum Plasma Concentration (Tmax) of baloxavir marboxil <sup>[12]</sup>
-----------------	---

End point description:

End point type	Secondary
End point timeframe:	
Up to Day 10	

Notes:

[12] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	0 <sup>[13]</sup>			
Units: hr				
arithmetic mean (standard deviation)	( )			

Notes:

[13] - Not estimable as a large number of concentrations were below the lower limit of quantification

## Statistical analyses

No statistical analyses for this end point

**Secondary: Time to Maximum Plasma Concentration (Tmax) of S-033447**

End point title	Time to Maximum Plasma Concentration (Tmax) of S-
-----------------	---

End point description:

Dose groups correspond to body-weight groups. 2mg/kg dose was used for subjects <20 kgs and 40 mg dose was used for subjects >20 kgs. Here 99999 represents results data which was not estimable due to low number of events.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Day 10

Notes:

[14] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	95			
Units: hr				
arithmetic mean (standard deviation)				
Non-Asian - 2 mg/kg (n=36)	4.12 (± 2.07)			
Asian - 40 mg (n=1)	5.50 (± 99999)			
Non-Asian - 40 mg (n=58)	5.55 (± 3.79)			

**Statistical analyses**

No statistical analyses for this end point

**Secondary: Plasma Concentrations of Baloxavir Marboxil by Dosage**

End point title	Plasma Concentrations of Baloxavir Marboxil by Dosage <sup>[15]</sup>
-----------------	---

End point description:

End point type	Secondary
----------------	-----------

End point timeframe:

24, 72, 96 and 240 hours post-dose

Notes:

[15] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	0 <sup>[16]</sup>			
Units: ng/mL				
arithmetic mean (standard deviation)	( )			

Notes:

[16] - Not estimable as a large number of concentrations were below the lower limit of quantification

## Statistical analyses

No statistical analyses for this end point

### Secondary: Plasma Concentrations of S-033447 by Dosage

End point title Plasma Concentrations of S-033447 by Dosage<sup>[17]</sup>

End point description:

Dose groups correspond to body-weight groups. 2mg/kg dose was used for subjects <20 kgs and 40 mg dose was used for subjects >20 kgs. Here 99999 represents results data which was not estimable due to low number of events.

End point type Secondary

End point timeframe:

24, 72, 96 and 240 hours post-dose

Notes:

[17] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: This endpoint is specific to the arm - Baloxavir Marboxil only.

End point values	Baloxavir Marboxil			
Subject group type	Reporting group			
Number of subjects analysed	95			
Units: ng/mL				
arithmetic mean (standard deviation)				
24 - Non-Asian - 2 mg/kg (n=36)	55.7 (± 28.1)			
24 - Asian - 40 mg (n=1)	75.2 (± 99999)			
24 - Non-Asian - 40 mg (n=58)	53.2 (± 22.4)			
72 - Non-Asian - 2 mg/kg (n=36)	13.2 (± 7.15)			
72 - Asian - 40 mg (n=1)	28.9 (± 99999)			
72 - Non-Asian - 40 mg (n=58)	17.90 (± 8.910)			
96 - Non-Asian - 2 mg/kg (n=36)	7.61 (± 4.400)			
96 - Asian - 40 mg (n=1)	19.0 (± 99999)			
96 - Non-Asian - 40 mg (n=58)	11.3 (± 6.160)			
240 - Non-Asian - 2 mg/kg (n=36)	0.989 (± 0.887)			
240 - Asian - 40 mg (n=1)	3.31 (± 99999)			
240 - Non-Asian - 40 mg (n=58)	11.3 (± 6.160)			

## Statistical analyses

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

From baseline (Day 1) until 28 days after the last dose of study drug (29 days)

Assessment type	Non-systematic
-----------------	----------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	22.0
--------------------	------

### Reporting groups

Reporting group title	Baloxavir Marboxil
-----------------------	--------------------

Reporting group description:

Participants will receive a single oral dose of baloxavir marboxil on Day 1 (based on body weight). Oseltamivir matching placebo will also be administered orally twice daily (BID) for 5 days.

Reporting group title	Oseltamivir
-----------------------	-------------

Reporting group description:

Participants will receive oseltamivir orally BID for 5 days (based on body weight). Baloxavir marboxil matching placebo will also be administered orally on Day 1

<b>Serious adverse events</b>	Baloxavir Marboxil	Oseltamivir	
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 115 (0.00%)	0 / 58 (0.00%)	
number of deaths (all causes)	0	0	
number of deaths resulting from adverse events			

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	Baloxavir Marboxil	Oseltamivir	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	15 / 115 (13.04%)	13 / 58 (22.41%)	
Gastrointestinal disorders			
Diarrhoea			
subjects affected / exposed	6 / 115 (5.22%)	1 / 58 (1.72%)	
occurrences (all)	6	1	
Vomiting			
subjects affected / exposed	7 / 115 (6.09%)	9 / 58 (15.52%)	
occurrences (all)	7	10	
Infections and infestations			

Otitis Media subjects affected / exposed occurrences (all)	3 / 115 (2.61%) 3	4 / 58 (6.90%) 5	
--	----------------------	---------------------	--

## **More information**

### **Substantial protocol amendments (globally)**

Were there any global substantial amendments to the protocol? No

---

### **Interruptions (globally)**

Were there any global interruptions to the trial? No

### **Limitations and caveats**

None reported